

Protecting Amazonia Should Focus on Protecting Indigenous, Traditional Peoples and Their Territories

Gabriel de Oliveira, Guilherme Mataveli, Carlos dos Santos, Liming He, Skye Hellenkamp, Beatriz M. Funatsu, Scott Stark, Yosio Shimabukuro

▶ To cite this version:

Gabriel de Oliveira, Guilherme Mataveli, Carlos dos Santos, Liming He, Skye Hellenkamp, et al.. Protecting Amazonia Should Focus on Protecting Indigenous, Traditional Peoples and Their Territories. Forests, 2021, 13 (1), pp.16. 10.3390/f13010016 . hal-03693412

HAL Id: hal-03693412 https://hal.science/hal-03693412v1

Submitted on 10 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Editorial

Protecting Amazonia Should Focus on Protecting Indigenous, Traditional Peoples and Their Territories

Gabriel de Oliveira ^{1,*}, Guilherme A. V. Mataveli ², Carlos A. C. dos Santos ³, Liming He ⁴, Skye E. Hellenkamp ¹, Beatriz M. Funatsu ⁵, Scott C. Stark ⁶ and Yosio E. Shimabukuro ²

- Department of Earth Sciences, University of South Alabama, Mobile, AL 36688, USA; seh1828@jagmail.southalabama.edu
- ² Earth Observation and Geoinformatics Division, National Institute for Space Research, Sao Jose dos Campos 12227-010, SP, Brazil; mataveli@alumni.usp.br (G.A.V.M.); yosio.shimabukuro@inpe.br (Y.E.S.)
- ³ Academic Unit of Atmospheric Sciences, Federal University of Campina Grande, Campina Grande 58109-970, PB, Brazil; carlos.santos@ufcg.edu.br
- ⁴ Canada Centre for Mapping and Earth Observation, Natural Resources Canada, Ottawa, ON K1S 5K2, Canada; liming.he@canada.ca
- French National Center for Scientific Research (CNRS), Université de Nantes, UMR LETG, Campus du Tertre BP 81227, CEDEX 3, 44312 Nantes, France; beatriz.funatsu@univ-nantes.fr
- 6 Department of Forestry, Michigan State University, East Lansing, MI 48824, USA; scstark@msu.edu
- Correspondence: deoliveira@southalabama.edu; Tel.: +1-(251)-591-4703

The Brazilian proverb "Uma desgraça nunca vem só" or "Misfortunes never come alone" has, unfortunately, never been more apt than in reference to the risks now facing Amazonia. Deforestation rates have been dramatically increasing in Brazil [1], with a total of 34,200 km² newly logged areas since 2019 [2]. In 2021, 13,200 km² of forest was cut down, the highest annual deforestation rate since 2006, when deforestation in Amazonia hit 14,300 km² [2]. Additionally, if deforestation alone was not enough, a severe drought in 2020 together with the COVID-19 pandemic has created a devastating synergism between forest loss, fires, which produce particulate air pollutants, and related health risks, endangering Amazonian Indigenous and traditional peoples [3]. Although it is no secret that deforestation has been dramatically increasing in Amazonia in the past few years [1,2,4,5], too little attention has been given to the plight of Indigenous lands, which have never before experienced such dramatic incursion and illegal deforestation. According to a report just released by the Brazilian National Institute for Space Research (INPE), deforestation within Indigenous lands has reached 331.1 km² in 2021 [2]. The average of the annual deforestation rates within Indigenous lands in the last three years (2019-2021) is ~81% above the decadal average rate [2]. In addition to deforestation, illegal mining is a growing threat within Indigenous lands [6–9]. As can be seen in Figure 1, deforestation within the Indigenous lands of the Brazilian Amazon in the past three years has been occurring with particular intensity in the eastern region, in the so-called "Arc of Deforestation".

Recent and emerging research is addressing the urgent need to understand increasing deforestation in Protected Areas and Indigenous lands and its impact on the health and livelihoods of Amazonian peoples and ecosystem services. de Oliveira et al. [10] evaluated the relationship between deforestation, land use and land cover (LULC) changes and fire emissions in the Apyterewa Indigenous Land, Eastern Brazilian Amazon. Combining a geographic object-based image analysis (GEOBIA) to map LULC with the Brazilian Biomass Burning Emission with Fire Radiative Power (3BEM_FRP) model, they estimated the dangerous particulate pollution (particles with a diameter less than 2.5 μ m, $PM_{2.5}$) produced by deforestation and fire. The GEOBIA approach revealed a remarkable increase in deforestation, with the conversion of primary forests to agriculture within the Apyterewa Indigenous Land reaching 200 km² between 2017 and 2019, which was clearly associated with an increase in the $PM_{2.5}$ emissions from fire. Between 2004 and 2016, the annual



Citation: de Oliveira, G.; Mataveli, G.A.V.; dos Santos, C.A.C.; He, L.; Hellenkamp, S.E.; Funatsu, B.M.; Stark, S.C.; Shimabukuro, Y.E. Protecting Amazonia Should Focus on Protecting Indigenous, Traditional Peoples and Their Territories. *Forests* 2022, *13*, 16. https://doi.org/10.3390/f13010016

Received: 9 December 2021 Accepted: 16 December 2021 Published: 23 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

Forests 2022, 13, 16 2 of 4

average emission of PM_{2.5} was estimated to be 3594 tons per year⁻¹, whereas the most recent interval of 2017–2019 had an average of 6258 tons per year⁻¹. This represented an increase of 58% in the annual average of PM_{2.5} associated with fires for the study period, contributing to respiratory health risks and the air quality crisis in Brazil in late 2019. Mataveli et al. [4] went on to combine PM_{2.5} emissions estimated from the 3BEM_FRP model and deforestation based on the MapBiomas Amazonia dataset to assess, for the first time, how deforestation drove biomass burning emissions in Amazonia over the last two decades at three scales of analysis (Amazonia-wide, countries/states and grid cell). Amazonia accounted for 48% of PM_{2.5} emitted from biomass burning in South America, whereas current deforestation rates have reached values on par with the alarming rates of the early 21st century. Emissions and deforestation were concentrated in the Eastern and Central-Southern portions of Amazonia. Amazonia-wide deforestation and emissions were linked over time. Countries/states with the greatest extent of agricultural lands were less likely to be correlated at this scale; however, this is likely because biomass burning in agricultural practices partially decouples deforestation from fire prevalence. Concentrated in regions of ongoing deforestation, in 18% of Amazonia grid cells, PM_{2.5} emissions associated with biomass burning and deforestation were significantly positively correlated. According to the authors, deforestation is an important driver of emissions in Amazonia but does not explain biomass burning alone. Future work must link climate and other non-deforestation drivers to further elucidate the factors contributing to dangerous pollution emissions from biomass burning in Amazonia.




Figure 1. Deforestation within Indigenous lands in the Brazilian Legal Amazon between 2019 and 2021. Annual deforestation rates were obtained from the National Institute for Space Research [2].

In Guerrero et al. [11], the authors pointed to another important factor related to the devastation of pristine, protected areas. They presented a temporal analysis of LULC change over a 33-year period (1985–2018) within the influence zone of the Braço Norte Hydropower Complex, Brazilian Amazon, using the Collection 4.1 level 3 LULC maps of the freely available MapBiomas dataset. Their findings showed that the most impacted LULC during this period was forest formation (from 414 km² to 287 km², a reduction of 69%), with a primary shift to pasturelands (increase of 664%, from 40 km² to 299 km²). The

Forests 2022, 13, 16 3 of 4

construction of the hydropower complex also triggered indirect impacts such as urban expansion in 2018 and an associated increase in local demand for crops. These results reveal the need for special policies that minimize the local impact of dam construction to protect important forestland, for example, through the creation of Protected Areas. Furthermore, locally adjusted parameters defining ecological-economic zoning should be adopted to balance environmental health and social benefits for Indigenous peoples, riverine population, and artisanal fishermen that depend on healthy rivers. LULC change is an essential component for monitoring environmental change and managing natural resources in areas of high natural and cultural biodiversity. Heredia-R et al. [12] conducted a revealing study in an area located in the northern Amazon of Ecuador, specifically in the Diversity and Life Zone (DLZ) of the Yasuní Biosphere Reserve (YBR). The aim of their study was to investigate the dynamics of LULC changes to support policies for environmental and sociocultural protection in the DLZ. Multitemporal satellite imagery, open-access Landsat images downloaded from the United States Geological Survey (USGS), were used to map changes in LULC divided into three time stages (1999–2009, 2009–2018, 1999–2018). The results showed that in the period 1999–2018, most of the changed LULCs were transformed into pastures in the DLZ. The authors concluded that it is imperative to improve territorial planning, to avoid conflicts between Indigenous peoples, migrant settlers, and uncontacted Indigenous peoples that live in the DLZ within the YBR.

As the studies described above show, Indigenous, traditional peoples and their territories are being significantly impacted by illegal deforestation. Even while formally protected by law, Indigenous peoples have limited resources to fight against illegal timber harvesting; therefore, deforestation is a general consequence. Indigenous lands management and monitoring has historically been focused solely on the legally bounded territory. However, this risks ignoring the pressures and land conversion dynamics building in neighboring areas, which are increasingly spilling into these Protected Areas, threatening residents and reducing the integrity and ecological functioning of the protected forest. The weakening of environmental law enforcement is coincident with the observed increase in LULC changes and deforestation in Indigenous lands [9]. Indigenous and traditional peoples are the "soul" of the Amazon, and everything should be done to protect them and their territories. Furthermore, empowering native people to control their land is one of the most effective approaches to long-term intact forest conservation [13]. To reverse the current scenario in the Amazon over the long-term, environmental education is key, as is the implementation of sustainable agricultural practices and development of conservation policies to promote forest regrowth in degraded areas. In the short term, appropriate federal government agencies in Brazil and other Amazonian nations should act swiftly to reduce illegal deforestation and combat rising anti-indigenous violence [9,14] in Indigenous lands, sending a strong message that these Protected Areas are legally controlled and monitored, in order to stop future invasions. Only such changes will enable the maintenance of environmental stability in Amazonia, contain further deforestation and degradation, and, most importantly, safeguard Indigenous and traditional peoples' long-term well-being, ways of life, and cultural heritages.

Author Contributions: Conceptualization, G.d.O. and G.A.V.M.; writing—original draft preparation, G.d.O. and G.A.V.M.; writing—review and editing, G.d.O., G.A.V.M., C.A.C.d.S., L.H., S.E.H., B.M.F., S.C.S. and Y.E.S.; visualization, G.d.O., G.A.V.M. and S.E.H. All authors have read and agreed to the published version of the manuscript.

Funding: São Paulo Research Foundation (FAPESP), grant number 2019/25701-8 (G.A.V.M.).

Conflicts of Interest: The authors declare no conflict of interest.

Forests 2022, 13, 16 4 of 4

References

1. Silva Junior, C.H.L.; Pessoa, A.C.M.; Carvalho, N.S.; Reis, J.B.C.; Anderson, L.O.; Aragao, L. The Brazilian Amazon deforestation rate in 2020 is the greatest of the decade. *Nat. Ecol. Evol.* 2021, *5*, 144–145. [CrossRef] [PubMed]

- 2. National Institute for Space Research (INPE). Monitoring of the Brazilian Amazon Deforestation by Satellite. Available online: http://terrabrasilis.dpi.inpe.br/app/dashboard/deforestation/biomes/legal_amazon/rates (accessed on 8 December 2021).
- 3. de Oliveira, G.; Chen, J.M.; Stark, S.C.; Berenguer, E.; Moutinho, P.; Artaxo, P.; Anderson, L.O.; Aragao, L. Smoke pollution's impacts in Amazonia. *Science* **2020**, *369*, *634*–*635*. [CrossRef] [PubMed]
- 4. Mataveli, G.A.V.; de Oliveira, G.; Seixas, H.T.; Pereira, G.; Stark, S.C.; Gatti, L.V.; Basso, L.S.; Tejada, G.; Cassol, H.L.G.; Anderson, L.O.; et al. Relationship between Biomass Burning Emissions and Deforestation in Amazonia over the Last Two Decades. *Forests* **2021**, *12*, 1217. [CrossRef]
- 5. Escobar, H. Brazil's deforestation is exploding—And 2020 will be worse. Science 2019. [CrossRef]
- 6. Villén-Pérez, S.; Anaya-Valenzuela, L.; Conrado da Cruz, D.; Fearnside, P.M. Mining threatens isolated indigenous peoples in the Brazilian Amazon. *Glob. Environ. Chang.* **2021**, 102398, in press. [CrossRef]
- 7. Rorato, A.C.; Camara, G.; Escada, M.I.S.; Picoli, M.C.A.; Moreira, T.; Verstegen, J.A. Brazilian amazon indigenous peoples threatened by mining bill. *Environ. Res. Lett.* **2020**, *15*, 1040a3. [CrossRef]
- 8. Rorato, A.C.; Picoli, M.C.A.; Verstegen, J.A.; Camara, G.; Silva Bezerra, F.G.; Escada, M.I.S. Environmental Threats over Amazonian Indigenous Lands. *Land* **2021**, *10*, 267. [CrossRef]
- 9. Conceição, K.V.; Chaves, M.E.D.; Picoli, M.C.A.; Sánchez, A.H.; Soares, A.R.; Mataveli, G.A.V.; Silva, D.E.; Costa, J.S.; Camara, G. Government policies endanger the indigenous peoples of the Brazilian Amazon. *Land Use Policy* **2021**, *108*, 105663. [CrossRef]
- 10. de Oliveira, G.; Chen, J.M.; Mataveli, G.A.V.; Chaves, M.E.D.; Seixas, H.T.; Cardozo, F.d.S.; Shimabukuro, Y.E.; He, L.; Stark, S.C.; dos Santos, C.A.C. Rapid Recent Deforestation Incursion in a Vulnerable Indigenous Land in the Brazilian Amazon and Fire-Driven Emissions of Fine Particulate Aerosol Pollutants. *Forests* 2020, 11, 829. [CrossRef]
- 11. Guerrero, J.V.R.; Escobar-Silva, E.V.; Chaves, M.E.D.; Mataveli, G.A.V.; Bourscheidt, V.; de Oliveira, G.; Picoli, M.C.A.; Shimabukuro, Y.E.; Moschini, L.E. Assessing Land Use and Land Cover Changes in the Direct Influence Zone of the Braço Norte Hydropower Complex, Brazilian Amazonia. *Forests* **2020**, *11*, 988. [CrossRef]
- Heredia-R, M.; Torres, B.; Cabrera-Torres, F.; Torres, E.; Díaz-Ambrona, C.G.H.; Pappalardo, S.E. Land Use and Land Cover Changes in the Diversity and Life Zone for Uncontacted Indigenous People: Deforestation Hotspots in the Yasuní Biosphere Reserve, Ecuadorian Amazon. Forests 2021, 12, 1539. [CrossRef]
- 13. Garnett, S.T.; Burgess, N.D.; Fa, J.E.; Fernández-Llamazares, Á.; Molnár, Z.; Robinson, C.J.; Watson, J.E.M.; Zander, K.K.; Austin, B.; Brondizio, E.S.; et al. A spatial overview of the global importance of Indigenous lands for conservation. *Nat. Sustain.* **2018**, *1*, 369–374. [CrossRef]
- 14. Urzedo, D.; Chatterjee, P. The Colonial Reproduction of Deforestation in the Brazilian Amazon: Violence against Indigenous Peoples for Land Development. *J. Genocide Res.* **2021**, 23, 302–324. [CrossRef]