



HAL
open science

Fast-SEA ligation: a chemoselective NCL-inspired ligation reaction effective in the nanomolar concentration range

Benjamin Grain, Benoît Snella, Jérôme Vicogne, Frédéric Capet, Birgit Wiltschi, Oleg Melnyk, Vangelis Agouridas

► To cite this version:

Benjamin Grain, Benoît Snella, Jérôme Vicogne, Frédéric Capet, Birgit Wiltschi, et al.. Fast-SEA ligation: a chemoselective NCL-inspired ligation reaction effective in the nanomolar concentration range. GFPP 22 / BPGM 5 Joint Meeting of the French and Belgian Peptides and Proteins Societies, May 2022, Port-Leucate, France. , 2022. hal-03695066

HAL Id: hal-03695066

<https://hal.science/hal-03695066v1>

Submitted on 14 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Benjamin Grain,^a Benoît Snella,^a Jérôme Vicogne,^a Frédéric Capet,^a Birgit Wiltschi,^b Oleg Melnyk,^a Vangelis Agouridas,^{a,c}

^a Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIL - Center for Infection and Immunity of Lille, F-59000 Lille, France

^b Institute of Bioprocess Science and Engineering, University of Natural Resources and Life Sciences, Vienna


^c Centrale Lille, F-59000, Lille, France

See: B. Snella, B. Grain, J. Vicogne, F. Capet, B. Wiltschi, O. Melnyk, V. Agouridas. *Angew. Chem. Int. Ed.* **2022**, *asap*

Introduction


The **Chemical Protein Synthesis** has highly benefitted from the properties of the thioester functional group, in particular with the development of the **Native Chemical Ligation** (NCL) reaction.^{1,2} However, **NCL-based modification of proteins** suffers from the moderate reactivity of classical alkyl and aryl thioesters³, especially in highly dilute media. Seeking for fast reactive thioesters surrogates, we have discovered that **oxalyl amide SEA** (^{oxo}SEA), in the form of a latent thioester precursor,⁴ could, upon activation, act as a powerful acylating agent in chemoselective ligation reactions with β -aminothiols such as the cysteine residue.

a) Principle of the Native Chemical Ligation





Rate constant of the ligation:
~ 0.3 to 4 M⁻¹ s⁻¹ for NCL
~ 30 M⁻¹ s⁻¹ for ^{oxo}SEA ligation

b) Principle of the ^{oxo}SEA Ligation




1) Synthesis of ^{oxo}SEA peptides

- Synthesis of protected lysine derivative **1** at the **gram scale** with **17% overall yield**
- Introduction of Fmoc-Lys(^{oxo}SEA)-OH **1** within peptide segments at **any position**, according to standard Fmoc-SPPS protocols



2) ^{oxo}SEA Ligation study


Investigation on the reaction of ^{oxo}SEA peptide **2** with cysteinyl peptide **3**.



^{oxo}SEA ligation with a peptide concentration of 1 mM is:


- **Efficient** (HPLC conversions to polypeptide **4** beyond 95%)
- **Fast** (Half-reaction time of 200 s)
- Optimal at a **working pH = 5,5**
- **Triggered by** reductants (TCEP)

Comparison with classical NCL rates highlighted the tremendous reactivity of the ^{oxo}SEA group, even more when decreasing peptide concentration to 50 μ M.




Determination of apparent second-order rate constant for the ^{oxo}SEA ligation in the micromolar concentration range:

- Diminution of the peptide concentration led to the increase of the rate constant
- Rate break at 150 μ M, and below this concentration value, apparent second-order rate constant thrives around 30 M⁻¹ s⁻¹




3) Nanomolar concentration range

To further explore the performance of the ^{oxo}SEA-mediated ligation in the **nanomolar concentration range**, we opted for SDS-PAGE separation followed by Western Blot as the analysis tool. This allowed us to work at the protein level by monitoring the reaction between the ^{oxo}SEA peptide derived from V5 peptidic tag **5** and the C-terminally biotinylated kringle 1 domain of the hepatocyte growth factor (K1/HGF) equipped with a N-terminal cysteine **6**.



- Detection of ligated product **7** after 200 minutes of reaction
- ^{oxo}SEA ligation effective in highly dilute solution, with peptide concentration of **270 nM**



Moreover, this ligation reaction is also efficient and chemoselective in a **complex protein medium**, as it has been successfully performed in a crude protein extract of a THP-1 whole cell lysate supplemented with 100 nM of **6** in the presence of a 15-fold excess of **5** (see article).

Conclusion

The groundbreaking reactivity of this **activatable oxalyl thioester surrogate** allows us to perform peptide and protein modification through a **chemoselective ligation reaction**. The latter proceeds extremely fast in aqueous media, under mild conditions and in **highly dilute medium** which can be as complex as a cell lysate. By being easy to introduce within peptide segments in the form of a lysine derivative, oxalyl thioesters might be of significant importance for the development of novel **crosslinking strategies** or the **polycondensation of water-soluble (bio)molecules**.

