



**HAL**  
open science

## **Adolescents and young adults with cancer: How multidisciplinary health care teams adapt their practices to better meet their specific needs**

Élise Ricadat, Karl-Leo Schwering, Sophie Fradkin, Nicolas Boissel, Isabelle Aujoulat

### ► To cite this version:

Élise Ricadat, Karl-Leo Schwering, Sophie Fradkin, Nicolas Boissel, Isabelle Aujoulat. Adolescents and young adults with cancer: How multidisciplinary health care teams adapt their practices to better meet their specific needs. *Psycho-Oncology*, 2019, 28 (7), pp.1576-1582. <10.1002/pon.5135>. <hal-03698437>

**HAL Id: hal-03698437**

**<https://hal.science/hal-03698437v1>**

Submitted on 24 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



# Psycho-Oncology

Journal of the Psychological, Social and Behavioral Dimensions of Cancer

## Adolescents and young adults with cancer: how multidisciplinary healthcare teams adapt their practices to better meet their specific needs

Journal:	<i>Psycho-Oncology</i>
Manuscript ID	PON-18-0742.R4
Wiley - Manuscript type:	Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	RICADAT, Élise; Universite Paris Diderot, IHSS/EP SCHWERING, Karl-Leo; Paris 13 University, UTRPP FRADKIN, Sophie; Universite Paris Diderot, IHSS/EP BOISSEL, Nicolas; Institut Universitaire d'Hematologie Aujoulat, Isabelle; Universite Catholique de Louvain
Keywords:	cancer, oncology, adolescents, young adults, nursing, dedicated units, qualitative methods

SCHOLARONE™  
Manuscripts

1  
2  
3 Adolescents and young adults with cancer: how multidisciplinary healthcare teams adapt their  
4 practices to better meet their specific needs  
5

6  
7 **Key words:**

8  
9 cancer, oncology, adolescent, young adult, nursing, dedicated units, qualitative methods.  
10

11  
12 **Authors**

13 **Élise RICADAT<sup>1</sup>, Karl-Leo SCHWERING<sup>2</sup>, Sophie FRADKIN<sup>1</sup>, Nicolas BOISSEL<sup>3</sup>,**

14  
15  
16 **Isabelle AUJOULAT<sup>4</sup>**

17  
18 <sup>1</sup> CRPMS EA 3522, IUH EA-3518, Paris, Paris-Diderot University, France.

19  
20 <sup>2</sup> UTRPP, Paris 13 University, Villetaneuse, France

21  
22  
23 <sup>3</sup> Haematology AYA Unit, Saint-Louis Hospital, APHP, EA-3518, Paris-Diderot University,  
24  
25 France

26  
27 <sup>4</sup> Institute of Health & Society, UCLouvain, Brussels, Belgium  
28  
29

30  
31  
32 **ABSTRACT**  
33

34 **Objective:** Dedicated adolescent and young adult (AYA) cancer units have emerged from the  
35 early 90s to address multiple challenges faced by AYA patients with cancer. Specific needs of  
36 AYA patients have been considered in an increasing number of studies. However, few describe  
37 *how* the healthcare professionals (HCPs) perceive their patients' needs and how they actually  
38 adjust their day-to-day practices to meet such needs. The purpose of this study is to identify and  
39 describe the practical methods of care and teamwork implemented by HCPs in response to what  
40 they perceive as essential to support psychosocial development of AYA patients.  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 **Methods:** Qualitative research was conducted between 2012 and 2014 with 31 HCPs from a  
51 recently created hematology AYA unit in France. The transcripts of open-ended interviews  
52 were subject to inductive analysis using *constant comparison* as recommended by the Grounded  
53 Theory methods.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Results:** Our results show how HCPs adapt their practices and care relationships to support  
4  
5 three major developmental milestones related to identity construction in AYAs: self-  
6  
7 determination and individuation from parents; gender and sexual identity; social life and  
8  
9 connectedness to peers and adults (other than parents). Our results also show how HCPs adapt  
10  
11 their practices and organizational methods to enhance the flexibility required to address their  
12  
13 young patients', thus setting consistent and high standards for the whole team. Such adaptation  
14  
15 is made possible through collaborative work and collective processes that facilitate self-  
16  
17 reflection. **Conclusions:** Our findings shed light on some meaningful young patient-friendly  
18  
19 practices of care and advocate for AYA dedicated units.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Background

Adolescence, as a transition from late childhood (12 years) to adulthood (the mid-20s up to 30), is a challenging period due to the intense physical and psychosocial changes that adolescents and young adults (AYAs) need to cope with, in order to build a valued and steady sense of self<sup>1</sup>. Cancer is a complex life-limiting condition which deeply interferes with the transition process inherent to adolescence. Having cancer is likely to hinder or delay the achievement of several important developmental milestones in adolescents: physical and psychological development, sexuality, education and vocation, family and peer relationships<sup>2-5</sup>. As a growing number of AYA patients survive into adulthood<sup>6-8</sup>, there is higher awareness of the need to comprehensively address their psychosocial and developmental needs, and to help them cope successfully with the numerous interrelated medical and psychosocial challenges due to their condition. During the last three decades, while medical care to young cancer patients has considerably improved, psychosocial care has become a major issue, with the aim to enhance quality of life in adults who have survived paediatric cancer<sup>9-12</sup>.

Several studies have addressed how far the hospitalisation experience may adversely affect the adolescence process in cancer patients, and how HCPs may try to make it the least disruptive possible<sup>13-14</sup>. Arguing that AYA patients are seen as belonging to a “no-man’s land” between paediatric and adult models of care<sup>15</sup>, the need for dedicated units to bridge this gap has emerged<sup>16-17</sup>. Although AYA units have been created since the early 90s in high-income countries, little is known about their practical organisation and what kind of adaptations are required from HCPs to best respond to their young patients’ needs<sup>18-19</sup>.

The aim of our study was to investigate how the members of a multidisciplinary team adapt their caring practices and work organisation to meet what they perceive as essential needs for the psychosocial development of AYA patients.

## METHODS

**Interviews** were conducted in an AYA haematology in-patient unit in Paris. This unit was founded in 2010 in line with the second governmental cancer action plan (2009-2013). It admits patients aged 15 to 25. A great majority of them is diagnosed with acute leukemia and high-grade lymphoma, both of which require long lasting treatments, including allogeneic stem cell transplant (HSCT). The unit is composed of 16 single bedrooms, including 4 that are equipped with high-efficiency particulate air filtration mostly used to perform HSCT. The average length of stay is about 11 days, ranging roughly from 3 days – for a short chemotherapy regimen – to 4-6 weeks for the induction phase for acute leukaemia or HSCT.

### ***Procedures***

The aim of our study was presented by the head of the AYA unit (4<sup>th</sup> author) during a regular team meeting. All staff members were invited to participate in an in-depth interview during their working hours, including night shifts. Interviews were led by two external researchers (1<sup>st</sup> and 3<sup>rd</sup> author), both clinical psychologists and PhD students at the time of the research. In line with our two research objectives, open-ended questions were used to encourage participants to reflect on their own beliefs and experiences: *“How would you define the AYA patients’ psychosocial and developmental needs? How do you see your role and that of the team as a whole in supporting these specific needs?”* Interviews were allowed to evolve as a natural discussion, following the path of every participant. The interviews lasted an average of 60 minutes [30-85], were audio-recorded with the consent of the participants and transcribed *verbatim*.

### ***Participants***

**Thirty** out of the 43 members of the staff agreed to participate (Table 1). All have been working in this unit since its creation 2 years before the study, some of them had a previous paediatric or adult oncology experience, others were at the beginning of their professional life.

### ***Data analysis***

We opted for an inductive and iterative approach for the analysis of the collected data. The initial coding of the transcripts involved the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and 5<sup>th</sup> authors, who set up a preliminary analysis grid based on an independent coding of the 4 first interviews. Referring to the Grounded Theory approach<sup>19</sup>, the rest of the transcripts were coded by the two main researchers (1<sup>st</sup> & 3<sup>rd</sup> author) in a process of *constant comparison*. Major emerging categories and sub-categories were discussed and refined until the end of the process to reach a strong consensus. This process involved 2 group discussions around preliminary results with the healthcare team.

### ***Results***

Regarding our first research objective, i.e. the AYA patients' psychosocial and developmental needs as perceived by the multidisciplinary health care team, our analysis revealed a core category – namely that of supporting identity construction in AYAs – and 3 subcategories that may be seen as 3 different facets of the process of identity construction: (i) self-determination and individuation from parents, (ii) gender and sexual identity and (iii) social life and connectedness with peers and adults (other than parents). Regarding our second research objective, i.e. to reveal how the HCPs in this unit individually and collectively adapt their practices to meet what they perceive as essential psychosocial needs of their patients, flexibility emerged as a core category, which was made possible by collaborative work and a number of collective processes that facilitate self-reflection and build a solid team cohesion. Our results are presented according to these emerging categories with examples of concrete practices to illustrate the findings.

## **SUPPORTING IDENTITY CONSTRUCTION**

The participants generally attested that AYAs go through an important process of identity construction, which involves major physical and psychological changes to be integrated. They

1  
2  
3 acknowledged that the experience of cancer largely hampers this process and advocated for the  
4 provision of in-patient care that includes ways of addressing these specific needs. They argued  
5 that an AYA patient does not only have cancer, but is a person who must achieve some  
6 important developmental milestones that need to be taken into account in the care relationship:  
7  
8 « *They remain adolescents, no matter how ill they are. An AYA patient is not an ordinary*  
9  
10 *patient* » (nurse). To mitigate cancer and treatment effects, they identified 3 major challenges  
11  
12 inherent to the identity process of any adolescent that may be particularly threatened by the  
13  
14 experience of cancer and therefore require special attention from the healthcare team.  
15  
16  
17  
18  
19  
20

21 ***Supporting individuation from parents and self-determination.*** HCP insisted on the  
22 importance of supporting the processes of self-determination and individuation as a way of  
23 compensating for the dependence created by in-patient care. Therefore, the team aims to deeply  
24 involve patients in decision-making processes during the whole duration of their  
25 hospitalisation. Parents are less systematically involved by the HCPs in their children's care, as  
26 compared to usual practices in paediatrics: « *What changes, with regards to paediatrics, is that*  
27 *the cancer diagnosis is not given first to the parents but directly disclosed to the teenager* »  
28 (doctor). HCPs assumed that the parents' presence has to be constantly discussed to support the  
29 separation process, which is even more complex in the context of cancer. According to HCPs,  
30 there is no one way of regulating the parents' presence, and individualised decisions need to be  
31 made. A concrete example is the management of extra beds for accompanying parents or  
32 relatives who ask to stay overnight. In this AYA unit, it is not a rule to have an extra bed in the  
33 room, as it is in paediatrics: « *I think that nothing should be imposed but rather proposed at*  
34 *this age. This is more important than for the younger ones.* » (doctor). Another example is how  
35 HCPs engage in trying to prevent parents from reverting to former relationships they would  
36 have had with their child at a younger age, induced by the status of being ill: « *A mother washing*  
37 *her small child doesn't shock me. However, when this happens with an AYA, I find it difficult...*  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *I try to discuss the matter with the parents and that it's not up to them to help with everyday*  
4 *things.* » (nurse aide). Regulating the distance between parents and AYAs has been  
5  
6 acknowledged by the participants as being of significant benefit to AYAs' need for  
7  
8 empowerment: « *I am thinking of situations where the parents behave in an exemplary manner*  
9  
10 *whilst being able to maintain the right distance. This has an extraordinarily beneficial effect*  
11  
12 *on both the young patient and the parent.* » (doctor).  
13  
14  
15

16  
17 ***Regarding the need to self-determine their goals and actions,*** the HCPs in our study  
18  
19 acknowledged that this is basically undermined because of the cancer treatment, which is not  
20  
21 negotiable. As a result of being aware of their young patients' need for self-determination,  
22  
23 flexibility is introduced into the manner in which the treatment is administered, to suit the young  
24  
25 patients' need to remain active decision makers, thus allowing them to develop a sense of  
26  
27 control. Non-compulsory treatments are discussed and sometimes dismissed or postponed: « *I*  
28  
29 *try to respect their preferences. If they don't want to be disturbed too early in the morning for*  
30  
31 *instance, I will accept it once but not a second time. Hence, I am not rushing them but still stand*  
32  
33 *firm.* » (nurse). The participants defined their care relationships and practices as being based on  
34  
35 an implicit rule: “allowing for negotiation *within a fixed framework*” (nurse).  
36  
37  
38  
39

40  
41 In order to allow for negotiations, there are no written internal procedures that would  
42  
43 set the rules too rigidly and hinder the necessary flexibility. In agreement with doctors, mask-  
44  
45 wearing measures are usually relaxed. External home-prepared or commercial frozen meals,  
46  
47 prohibited in other places due to infectious risks, may be permitted under certain conditions.  
48

49  
50 In their capacity as adults, HCPs saw themselves as playing a key role in supporting  
51  
52 their young patients' identity-forging process by tolerating conflicts and differing opinions:  
53  
54 « *After all, we represent the adult world for them, and perhaps it is a good thing that they know*  
55  
56 *they can't do whatever they want to do, even when they are sick* » (nurse). They acknowledged  
57  
58 negotiation and conflicts as making sense in the caring relationship: « *Teenagers are looking*  
59  
60

1  
2  
3 *for conflict. Yes, there are clashes, but these are natural things that happen because we are in*  
4 *facility that deals with adolescence. We need to remember the need for mutual respect all the*  
5 *time » (nurse).*

6  
7  
8  
9  
10 ***Supporting gender and sexual identity.*** HCPs were aware that the challenges around sexuality  
11 and body image are crucial in the process of defining one's identity during adolescence. As  
12 treatments alter the pubescent body, they perceive the importance of enhancing the patient's  
13 emerging gendered identity, much more than when working with younger or older patients:  
14 « *when I was working in paediatrics, I didn't ask myself those kinds of questions. But teenagers*  
15 *become conscious about their own bodies so I cannot take over it without asking them. They*  
16 *are a person, already an adult for some and for others still 'in progress', but at the end of the*  
17 *day, they have their say in things!* » (nurse). Consequently, HCPs are extremely respectful of  
18 their young patients' modesty and privacy. In practical terms, the AYA unit is organised with  
19 single rooms which are separated by opaque doors, thus precluding from the possibility of being  
20 observed from the outside. Moreover, for intimate aspects of care, such as hygiene-related  
21 issues or fertility preservation procedures, HCPs single out one member of their team of the  
22 same gender as the patient when they perceive that the patient feels too uncomfortable: « *Our*  
23 *male colleagues are involved in matters of care that bother boys, for instance for explaining*  
24 *the semen collection procedure. Talking to a man about this is easier than to confide in a*  
25 *woman, even more so if she is a young caregiver.* » (nurse).

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47 Finally, the staff noticed that when special care was provided to enhance well-being in  
48 addition to physical treatment, it eased the suffering stemming from a painful and injured body.  
49 Massages are offered to allow for positive experiences and to relieve anxiety and stress. HCPs  
50 were deeply convinced that attending to physical needs other than only those ones exclusively  
51 related to the treatment of cancer contributes to a better body image, self-esteem and therefore  
52 to positive identity construction.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 ***Encouraging social life and connectedness.*** HCPs acknowledged that the process of identity  
4 construction in AYAs involves developing more intense relationships with peers and  
5 maintaining their professional and social projects. They observed the disruptive effects of  
6 diagnosis and prolonged in-patient stays on AYAs' social dynamics. Despite the isolation  
7 measures mandatory in haematology, the AYA unit has been designed with the aim of opening  
8 up the boundaries of the hospital: some open living spaces encourage AYAs' social  
9 relationships and informal meetings. Various practical adaptations have been set up to reduce  
10 their feelings of isolation despite the usual hospital rules: visiting hours were extended to  
11 respect the more nocturnal hours favored by AYAs, hospital discharges are facilitated by a  
12 coordination nurse whenever the staff identifies that a patient might be suffering from social  
13 isolation, a community worker organises collective activities to encourage links between in-  
14 ward patients and coordinates the actions of several out-door organisations. One of these  
15 provides a teacher who lectures in the hospital and liaise with schools in order to facilitate the  
16 patients' pursuit of their studies. A social worker helps young adult patients already engaged in  
17 a professional activity to maintain it by liaising with their employers. This gives an indication  
18 of the numerous professions involved in the care to AYAs with cancer in this unit.

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
These concrete measures show a strong will to approach the AYA patient more  
holistically and the HCPs' conviction that caring means more than curing cancer: « *After all, I  
do not look at them as ill people in the first place, but rather as people.* » (nurse). Our  
participants conveyed that AYAs with cancer need not only efficient and evidence-based  
medical and technical treatment, but also strong links and genuine human connection, which  
they acknowledged as the core component of the care relationship: « *If they need to talk, I don't  
leave, I give them time. Interpersonal relationships are very important in this unit.* » (ward-  
aide). Such involvement was recognised as giving sense to their work. On the other hand, they  
also acknowledged that it might make them more vulnerable: « *In this unit, our caring*

1  
2  
3 *technique is based on relationships. It is not much more difficult than anything else, but it means*  
4 *that we are much more involved.* » (nurse). Reflecting on their efforts to best address their  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
young patients' needs, HCPs reported that they get personally involved beyond the boundaries  
of their traditional professional roles, thus exposing themselves to an increase risk of emotional  
burden. Reflective practices and strategies are used to process the constant adjustments that are  
required to maintain a collective cohesion and flexibility.

## **FLEXIBILITY, COLLECTIVE PROCESSES AND SELF-REFLEXION**

HCPs were aware that being individually flexible was dependent on collective strategies  
aimed at protecting their professional roles and ensuring overall coherence of staff  
decisions: « *I always try to give a consistent message, because otherwise AYA patients fall*  
*between the cracks.* » (nutritionist). In practice, they reported that formal and informal  
teamwork practices had emerged, such as feeling free to ask for help from a colleague whenever  
they face emotional difficulties with a patient: « *We must have empathy to be a good caregiver,*  
*but we also need to attend our own needs* ». *When I feel emotionally fragile with a patient, I*  
*hand him/her over to a colleague.* » (nurse). Working in pairs and mutual trust were cited as  
important team resources in which interpersonal relationships are central. Although such  
flexibility could be seen as carrying a risk of confusion, HCPs reported that sharing opinions  
ensures the team's cohesion: « *I cannot work by myself because we all have our small*  
*contribution toward building this care project.* » (social worker). Therefore, HCPs  
acknowledged the importance of the right motivation to work in such a ward: « *We are all very*  
*motivated here because facing an adolescent requires a high sense of solidarity* » (night nurse).

The weekly staff meetings aim to favour a multi-focal approach and individualisation  
of care procedures. Personalised decisions are more appropriate than decisions that would be  
taken by complying with predefined criteria according to a fixed medical protocol: « *I*  
*remember a patient, in aplasia and feverish, who wanted to leave for the weekend. From my*

1  
2  
3 *typical paediatric perspective, letting her out was out of the question... But after discussing it*  
4 *with the team, I let her go out. This enables us to do work on a case by case basis » (doctor).*

5  
6  
7  
8 Every staff member feels responsible for passing on information while optimising his/her  
9  
10 professional role: « *This multidisciplinary exchange is very constructive: it invites us to share*  
11 *different views, thus revealing aspects of a patient's experience, that I would otherwise not*  
12 *necessarily be aware of. This experience enriches me. » (nutritionist).*

13  
14  
15  
16  
17 Open-mindedness and reflexive practices were described as a strong resource that nurtures in  
18  
19 return the HCP's own professional role, and their ability to better meet the demands imposed  
20  
21 on them by AYA patients.

## 22 23 24 **DISCUSSION**

25  
26 Identity work was identified by the HCPs as the core challenge that AYAs with cancer  
27  
28 must face, along with the interconnected challenges of building one's autonomy and sexual and  
29  
30 social identities. Whereas these issues have been well studied from the patients' perspective<sup>21</sup>,  
31  
32 they have remained understudied with regard to opinions of HCPs'. Yet, HCPs are particularly  
33  
34 challenged when it comes to support such needs in the patient-provider relationship. This study  
35  
36 does not only report on individual and collective practices meant to support AYA patients'  
37  
38 psychosocial needs but also on strategies and skills HCPs should aim to develop in order to  
39  
40 meet this challenge.

41  
42  
43  
44 First, HCPs contribute concretely to individuation and self-determination targets when  
45  
46 involving patients in decision-making processes and playing a subtle role in negotiating their  
47  
48 parents' presence. Recent research suggests that AYA health communication and collaborative  
49  
50 decision-making is a sensitive issue, because the age of legal responsibility should not  
51  
52 determine who (the parents or the young patient) has to be informed<sup>22-23</sup>. Assessing as  
53  
54 comprehensively and as early as possible the patient's degree of autonomy and the quality of  
55  
56 the parent-AYA relationship should contribute to a better balance of decision-making  
57  
58  
59  
60

1  
2  
3 responsibilities as far as the patient-parent dyad is concerned. This is consistently related to an  
4  
5 important coping strategy to develop AYAs' sense of control<sup>24</sup>.  
6

7  
8 Secondly, our results highlight how HCPs pay attention to AYA patients' sexual  
9  
10 development by being sensitive and responsive to individual patients' level of sexual maturity  
11  
12 and physical development. There is some evidence that treatment-related side effects adversely  
13  
14 affect AYAs' early puberty, sexual function and development<sup>25-28</sup>. Recent research suggests  
15  
16 that impairments of body image seriously threaten the gendered identity if related to lower self-  
17  
18 esteem. Self-esteem plays a key role in the development of the sense of self as a sexual being  
19  
20 at that age<sup>29</sup>. Our study shows that HCPs support the body image with the onset of puberty by  
21  
22 being highly respectful of modesty and privacy. They promote a better body image as they offer  
23  
24 positive sensory experiences and feelings about physical impairments thanks to their careful  
25  
26 handling of physical areas affected by illness or treatments by providing massages. As shown  
27  
28 in other studies, such attitudes are likely to contribute to a better self-acceptance, thus fostering  
29  
30 enhanced relationships with peers and intimate partners<sup>30-31</sup>. We believe that it paves the way  
31  
32 for better communication regarding the emotional, romantic and sexual dimensions of their  
33  
34 quality of life, all of which are key issues in this population of patients<sup>32-33</sup>.  
35  
36  
37  
38  
39

40  
41 Another major issue for HCPs was about maintaining a bond with peers by favouring  
42  
43 outpatient hours, school attendance or the pursuit of work and projects as key elements in  
44  
45 building one's identity at that age. They adapt hospital and ward rules to reduce the disruptive  
46  
47 effects of hospitalisations. In doing so, they provide personalised adjustments which require  
48  
49 that they improve their own tolerance to conflicts. The present study shows how they develop  
50  
51 negotiation skills, patience, with mutual respect as a strategy to promote a genuine and non-  
52  
53 authoritative therapeutic bond. There is some evidence that a strong patient-oncologist alliance  
54  
55 is associated with psychosocial wellbeing and therapeutic adherence in AYA cancer patients<sup>34</sup>.  
56  
57  
58  
59  
60

1  
2  
3 Lastly, our findings highlight that a genuine personal relationship based on open-  
4 mindedness and strong motivation are to be considered subjective abilities HCPs must develop  
5 to support such an alliance. The HCPs' skills are strongly linked with the understudied concept  
6 of empathy in the patient-caregiver relationship. A recent review concludes that HCPs' empathy  
7 is associated with better psychosocial adjustment and less psychological distress in the HCPs<sup>35</sup>,  
8 although there is a lack of nursing-focused research in this field. Various sections of the  
9 transcripts obtained show deep emotional and subjective commitment as core human qualities  
10 developed by the HCPs to favour empathy: it can be seen as a main contribution to this topic.  
11 However, being empathic exposes HCPs to a "double-edged sword"<sup>36</sup>, with a clear benefit for  
12 patients potentially counterbalanced by a detrimental effect on professional stress levels,  
13 especially when patients are facing life-threatening condition<sup>37-40</sup>. It would deserve further  
14 investigation to assess how HCPs' empathy is a source of work exhaustion or distress. Our  
15 results suggest that a reflection about what it means to get personally involved in care should  
16 be introduced early in the curricula of medical and paramedical studies, and the importance of  
17 a trained and balanced workforce within the team (i.e. multidisciplinary and mixed-gendered).  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

#### 40 **Study limitations**

41  
42 The strength of our study lies in the high participation of a multidisciplinary health care  
43 team in an AYA dedicated unit, **although nurses are an over-represented professional**  
44 **category. Moreover, as this was a study conducted in a single centre, the sample of**  
45 **participants was inevitably small, and our results may not be appropriate for**  
46 **extrapolation. In addition, we opted for an iterative process of data collection and**  
47 **analysis. Yet, at this stage of our study, we are aware that we have not reached saturation**  
48 **of the data. Rather, as is typical in grounded theory, our initial research question has**  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **broadened, and there are still many issues that need to be explored to complete the**  
4  
5 **complexity of the AYA's psychosocial care.**  
6  
7  
8  
9

### 10 **Clinical implications**

11  
12 **HCPs have delivered useful recommendations to support AYA's psychosocial**  
13  
14 **development such as providing patients personalised adjustments in their caring practices**  
15  
16 **and implementing strategies to that end. As such, these can be transferable to other**  
17  
18 **similar care situations and may initiate a reflection on practical guidelines for young**  
19  
20 **patient-friendly models of care in dedicated units.**  
21  
22  
23  
24  
25

### 26 **Conflict of interest statement**

27  
28 **The author(s) declared no potential conflicts of interest with respect to the research,**  
29  
30 **authorship, and/or publication of this article.**  
31

### 32 **Acknowledgements**

33  
34 **We are grateful to all the participants who agreed to share their experience with us.**  
35  
36 **We are most grateful to INCA (Institut National du Cancer) for doctoral fellowship, and**  
37  
38 **to Novartis for their administrative support (basically interviews' transcripts).**  
39  
40 **We also thank the following colleagues for their editorial help at various stages of the**  
41  
42 **writing process: Giulietta Von Salis and Lydia Stephens.**  
43  
44  
45

### 46 **References**

---

- 47  
48 [1] Arnett JJ, Emerging adulthood: a theory of development from the late teens through the  
49 twenties, *Am Psychol*, 2000;55 (5):469–80.  
50  
51 [2] Stam H, Hartman EE, Deurloo JA, Groothoff J, Grootenhuis, Young Adult Patients with a  
52 History of Paediatric Disease: Impact on Course of Life and Transition into Adulthood, *Journal*  
53 *of Adolescent Health*, 2006, 39(1), 4-13  
54  
55 [3] Evan EE, Zeltzer LK. Psychosocial dimensions of cancer in adolescents and young adults.  
56 *Cancer*, 2006; 107:1663–71.  
57  
58  
59  
60

- 1  
2  
3 [4] Katz, A, Meeting the need for psychosocial care in young adults with cancer, 2015,  
4 Pittsburgh, PA: Oncology Nursing Society.  
5  
6 [5] Katz A, The trouble with teen and young adults with cancer, *Oncology Nursing Forum*,  
7 2015; 42 (4):327–328.  
8  
9 [6] Zebrack B, Bleyer A, Albritton K, Medearis S, Tang J, Assessing the health care needs of  
10 adolescent and young adult cancer patients and survivors, *Cancer*, 2006, 12 (107): 2915–  
11 2923.  
12  
13 [7] Zebrack B, Isaacson S, Psychosocial care of adolescent and young adult patients with  
14 cancer and survivors, *J Clin Oncol*, 2012 Apr 10;30(11):1221–6.  
15  
16 [8] Nightingale CL, Quinn GP, Shenkman EA, Curbow BA, Zebrack BJ, Krull KR, Huang  
17 IC, Health-Related Quality of Life of Young Adult Survivors of Childhood Cancer: A Review  
18 of Qualitative Studies, *Journal of adolescent and young adult oncology*, 2011; Vol 1(3):124–  
19 132.  
20  
21 [9] Morgan S, Davies S, Palmer S, Plaster M, Sex, Drugs, and Rock ‘n’ Roll: Caring for  
22 Adolescents and Young Adults with Cancer, *Journ clin oncology*, 2012, vol 28(32):4825–  
23 4830.  
24  
25 [10] Palmer S, Mitchell A, Thompson K, Sexton M, Unmet needs among adolescent cancer  
26 patients: a pilot study, *Palliat Support Care*, 2007 Jun;5(2):127–34.  
27  
28 [11] D'Agostino NM, Penney A, Zebrack B, Providing developmentally appropriate  
29 psychosocial care to adolescent and young adult cancer survivors. *Cancer*. 2011; 117(10  
30 Suppl):2329–2334.  
31  
32 [12] Clinton-McHarh, T., Carey, M., Sanson-Fisher, R., Shakeshaft, A., & Rainbird, K. (2010).  
33 Measuring the psychosocial health of adolescent & young adult (AYA) cancer survivors: A  
34 critical review. *Health & Quality of Life Outcomes*, 8(25)  
35  
36 [13] Farre A, Wood V, McDonagh JE, Parr JR, Reape D, Rapley T, Health professionals' and  
37 managers' definitions of developmentally appropriate healthcare for young people:  
38 conceptual dimensions and embedded controversies, *Arch Dis Child*, 2016 July; 101(7): 628–  
39 633.  
40  
41 [14] Palmer S, Improving care for AYA patients treated within adult hospitals: What can be  
42 done right now?, *Cancer Forum*, 2009, 33, 22–25.  
43  
44 [15] Ferrari A, Thomas D, Franklin ARK, Hayes-Lattin BM, Mascarini M, van der Graaf W,  
45 Albritton KH, Starting an Adolescent and Young Adult Program: Some Success Stories and  
46 Some Obstacles to Overcome, *Clin Oncol*, 2010, 28 (32):4850–4857.  
47  
48 [16] Dyson G, Thompson K, Palmer S, Thomas D, Schofield P, The relationship between  
49 unmet needs and distress amongst young people with cancer, *Supportive Care and Cancer*,  
50 2012 Jan;20(1):75-85  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [17] Hollis R, Morgan S, The adolescent with cancer –at the edge of no-man's land, *Lancet*  
4 *Oncol*, 2001 Jan;2(1):43–8.  
5  
6  
7 [18] Smith S, Adolescent units: an evidence-based approach to quality nursing in adolescent  
8 care, *European Journal of Oncology Nursing*, 2004;8:20–29.  
9  
10 [19] Lewis I, Morgan S, Models of care and specialized units, Bleyer WA, Barr RD, eds.  
11 *Cancer in Adolescents and Young Adults*. New York: Springer Verlag, 2007, 341–352.  
12  
13  
14 [20] Strauss A, Corbin J, *Basics of qualitative research: techniques and procedures for*  
15 *developing grounded theory*. Thousand Oaks, CA: Sage; 1998.  
16  
17 [21] Abrams AN, Muriel AC, Wiener L, *Pediatric Psychosocial Oncology: Textbook for*  
18 *Multidisciplinary Care*, New-York: Springer; 2016, chap 12: 199-219: “Adolescents and  
19 *Young Adults with Cancer: A Biopsychosocial Approach*  
20  
21  
22 [22] Hanna KM, Dashiff CJ, Stump TE, Weaver MT. Parent-Adolescent Dyads: Association  
23 of Parental Autonomy Support and Parent-Adolescent Shared Diabetes Care Responsibility,  
24 *Child Care Health Dev.*, 2013, Sept ; 39(5): 695–702  
25  
26 [23] Compas BE, Jaser SS, Dunn MJ, Rodriguez EM, Coping with Chronic Illness in  
27 *Childhood and Adolescence*, *Annu Rev Clin Psychol*. 2012 April 27; 8: 455–480  
28  
29 [24] Sadak KT, Connor C, DeLuca H, Innovative educational approaches to engage and  
30 empower the adolescent and young adult childhood cancer survivor, *Pediatr Blood Cancer*.  
31 2013 Dec;60(12):1919-21.  
32  
33 [25] Martins A, Taylor RM, Lobel B, McCann B, Soanes L, Whelan JS, Fern LA, Sex, Body  
34 Image, and relationships, A brightlight workshop on Information and Support Needs of  
35 Adolescents and Young Adults, *Journal of Adolescent and Young Adult Oncology*, 2018, vol  
36 7 (5), 1-7.  
37  
38 [26] Moules NJ, A tribe apart: Sexuality and cancer in adolescence, *Journal of Pediatric*  
39 *Oncology Nursing*, 2017; Vol 34 issue: 4: 295-308.  
40  
41 [27] Thompson AL, Long KA, Marsland AL, Impact of childhood cancer on emerging adult  
42 survivors’ romantic relationships: A qualitative account, *J Sex Med* 2013;10:65–73  
43  
44 [28] Bolte S, Zebrack B. Sexual issues in special populations: adolescents and young adults.  
45 *Semin Oncol Nurs*. 2008; 24(2):115–19.  
46  
47 [29] Evan E, Kaufman M et al., Sexual health and self-esteem in adolescents and young adults  
48 with cancer, *Cancer*, 2006, Supplement October 1, Vol. 107, 7: 1672-1679.  
49  
50 [30] Robinson L, Miedema B, Easley J, Young Adult Cancer Survivors and the Challenges of  
51 Intimacy, *Journal of Psychosocial Oncol*, 2014;32 (4): 447-462.  
52  
53 [31] Rosenberg AR, Bona K, Ketterl T, Wharton CM, Wolfe J, Baker KS, Intimacy,  
54 Substance Use, and Communication Needs During Cancer Therapy: A Report From the  
55  
56  
57  
58  
59  
60

1  
2  
3 “Resilience in Adolescents and Young Adults” Study, *J Adolesc Health*. 2017 January; 60(1):  
4 93–99  
5

6 [32] Dobinson KA, Hoyt MA, Seidler ZE, Beaumont AL, Hullmann SE, Lawsin CR, A  
7 Grounded Theory Investigation into the Psychosexual Unmet Needs of Adolescent and  
8 Young Adult Cancer Survivors, *Journal of Adolescent and Young Adult Oncology*, 2016; Vol  
9 5 (2):135-145.  
10  
11

12 [33] Thompson AL, Long KA, Marsland AL, Impact of Childhood Cancer on Emerging  
13 Adult Survivors’ Romantic Relationships: A Qualitative Account, *J Sex Med* 2013;10:65–73  
14

15 [34] Trevino KM, Fasciano K, Prigerson HG. Patient-oncologist alliance, psychosocial well-  
16 being, and treatment adherence among young adults with advanced cancer. *J Clin Oncol*.  
17 2013; 31(13):1683– 1689.  
18  
19

20 [35] Mack JW, Fasciano KM, Block SD, Communication About Prognosis With Adolescent  
21 and Young Adult Patients With Cancer: Information Needs, Prognostic Awareness, and  
22 Outcomes of Disclosure, *Journal of Clinical Oncology* 36, no. 18 (June 20 2018) 1861-1867.  
23  
24

25 [36] Lelorain S, Brédart A, Dolbeault S, Sultan S, A systematic review of the associations  
26 between empathy measures and patient outcomes in cancer care, *Psycho-Oncology*, 2012; 21:  
27 1255–1264  
28

29 [37] Abrams AN, Muriel AC, Wiener L, *Pediatric Psychosocial Oncology: Textbook for*  
30 *Multidisciplinary Care*, New-York: Springer; 2016, chap 21:367-377: “Self-Care and  
31 Sustainability for Pediatric Oncology Providers  
32  
33

34 [38] Crumpei I, Dafinoiu I, The relation of clinical empathy to secondary traumatic stress.  
35 *Procedia: Social and Behavioral Sciences*, 2011; 33:438–442  
36  
37

38 [39] Granek L, Bartels U, Scheinemann K, Labrecque M, Barrera M, Grief reactions and  
39 impact of patient death on pediatric oncologists, *Pediatr Blood Cancer*, 2015; 62(1): 134–142.  
40

41 [40] Potter P, Divanbeigi RNJ, Berger J, Norris J, Olsen RNS, Compassion fatigue and burnout:  
42 Prevalence among oncology nurses, *Clinical Journal of Oncology Nursing*, 2010, 14(5), E56.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Adolescents and young adults with cancer: how multidisciplinary healthcare teams adapt their  
4 practices to better meet their specific needs  
5

6  
7 **Key words:**

8  
9 cancer, oncology, adolescent, young adult, nursing, dedicated units, qualitative methods.  
10

11  
12 **Authors**

13  
14 **Élise RICADAT<sup>1</sup>, Karl-Leo SCHWERING<sup>2</sup>, Sophie FRADKIN<sup>1</sup>, Nicolas BOISSEL<sup>3</sup>,**

15  
16 **Isabelle AUJOULAT<sup>4</sup>**

17  
18 <sup>1</sup> CRPMS EA 3522, IUH EA-3518, Paris, Paris-Diderot University, France.

19  
20 <sup>2</sup> UTRPP, Paris 13 University, Villetaneuse, France

21  
22 <sup>3</sup> Haematology AYA Unit, Saint-Louis Hospital, APHP, EA-3518, Paris-Diderot University,  
23 France  
24  
25

26  
27 <sup>4</sup> Institute of Health & Society, UCLouvain, Brussels, Belgium  
28  
29

30  
31  
32 **ABSTRACT**  
33

34  
35 **Objective:** Dedicated adolescent and young adult (AYA) cancer units have emerged from the  
36 early 90s to address multiple challenges faced by AYA patients with cancer. Specific needs of  
37 AYA patients have been considered in an increasing number of studies. However, few describe  
38 *how* the healthcare professionals (HCPs) perceive their patients' needs and how they actually  
39 adjust their day-to-day practices to meet such needs. The purpose of this study is to identify and  
40 describe the practical methods of care and teamwork implemented by HCPs in response to what  
41 they perceive as essential to support psychosocial development of AYA patients.  
42  
43  
44  
45  
46  
47  
48  
49

50  
51 **Methods:** Qualitative research was conducted between 2012 and 2014 with 31 HCPs from a  
52 recently created hematology AYA unit in France. The transcripts of open-ended interviews  
53 were subject to inductive analysis using *constant comparison* as recommended by the Grounded  
54 Theory methods.  
55  
56  
57  
58  
59  
60

1  
2  
3 **Results:** Our results show how HCPs adapt their practices and care relationships to support  
4  
5 three major developmental milestones related to identity construction in AYAs: self-  
6  
7 determination and individuation from parents; gender and sexual identity; social life and  
8  
9 connectedness to peers and adults (other than parents). Our results also show how HCPs adapt  
10  
11 their practices and organizational methods to enhance the flexibility required to address their  
12  
13 young patients', thus setting consistent and high standards for the whole team. Such adaptation  
14  
15 is made possible through collaborative work and collective processes that facilitate self-  
16  
17 reflection. **Conclusions:** Our findings shed light on some meaningful young patient-friendly  
18  
19 practices of care and advocate for AYA dedicated units.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Background

Adolescence, as a transition from late childhood (12 years) to adulthood (the mid-20s up to 30), is a challenging period due to the intense physical and psychosocial changes that adolescents and young adults (AYAs) need to cope with, in order to build a valued and steady sense of self<sup>1</sup>. Cancer is a complex life-limiting condition which deeply interferes with the transition process inherent to adolescence. Having cancer is likely to hinder or delay the achievement of several important developmental milestones in adolescents: physical and psychological development, sexuality, education and vocation, family and peer relationships<sup>2-5</sup>. As a growing number of AYA patients survive into adulthood<sup>6-8</sup>, there is higher awareness of the need to comprehensively address their psychosocial and developmental needs, and to help them cope successfully with the numerous interrelated medical and psychosocial challenges due to their condition. During the last three decades, while medical care to young cancer patients has considerably improved, psychosocial care has become a major issue, with the aim to enhance quality of life in adults who have survived paediatric cancer<sup>9-12</sup>.

Several studies have addressed how far the hospitalisation experience may adversely affect the adolescence process in cancer patients, and how HCPs may try to make it the least disruptive possible<sup>13-14</sup>. Arguing that AYA patients are seen as belonging to a “no-man’s land” between paediatric and adult models of care<sup>15</sup>, the need for dedicated units to bridge this gap has emerged<sup>16-17</sup>. Although AYA units have been created since the early 90s in high-income countries, little is known about their practical organisation and what kind of adaptations are required from HCPs to best respond to their young patients’ needs<sup>18-19</sup>.

The aim of our study was to investigate how the members of a multidisciplinary team adapt their caring practices and work organisation to meet what they perceive as essential needs for the psychosocial development of AYA patients.

## METHODS

**Interviews** were conducted in an AYA haematology in-patient unit in Paris. This unit was founded in 2010 in line with the second governmental cancer action plan (2009-2013). It admits patients aged 15 to 25. A great majority of them is diagnosed with acute leukemia and high-grade lymphoma, both of which require long lasting treatments, including allogeneic stem cell transplant (HSCT). The unit is composed of 16 single bedrooms, including 4 that are equipped with high-efficiency particulate air filtration mostly used to perform HSCT. The average length of stay is about 11 days, ranging roughly from 3 days – for a short chemotherapy regimen – to 4-6 weeks for the induction phase for acute leukaemia or HSCT.

### ***Procedures***

The aim of our study was presented by the head of the AYA unit (4<sup>th</sup> author) during a regular team meeting. All staff members were invited to participate in an in-depth interview during their working hours, including night shifts. Interviews were led by two external researchers (1<sup>st</sup> and 3<sup>rd</sup> author), both clinical psychologists and PhD students at the time of the research. In line with our two research objectives, open-ended questions were used to encourage participants to reflect on their own beliefs and experiences: *“How would you define the AYA patients’ psychosocial and developmental needs? How do you see your role and that of the team as a whole in supporting these specific needs?”* Interviews were allowed to evolve as a natural discussion, following the path of every participant. The interviews lasted an average of 60 minutes [30-85], were audio-recorded with the consent of the participants and transcribed *verbatim*.

### ***Participants***

**Thirty** out of the 43 members of the staff agreed to participate (Table 1). All have been working in this unit since its creation 2 years before the study, some of them had a previous paediatric or adult oncology experience, others were at the beginning of their professional life.

### ***Data analysis***

We opted for an inductive and iterative approach for the analysis of the collected data. The initial coding of the transcripts involved the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and 5<sup>th</sup> authors, who set up a preliminary analysis grid based on an independent coding of the 4 first interviews. Referring to the Grounded Theory approach<sup>19</sup>, the rest of the transcripts were coded by the two main researchers (1<sup>st</sup> & 3<sup>rd</sup> author) in a process of *constant comparison*. Major emerging categories and sub-categories were discussed and refined until the end of the process to reach a strong consensus. This process involved 2 group discussions around preliminary results with the healthcare team.

### ***Results***

Regarding our first research objective, i.e. the AYA patients' psychosocial and developmental needs as perceived by the multidisciplinary health care team, our analysis revealed a core category – namely that of supporting identity construction in AYAs – and 3 subcategories that may be seen as 3 different facets of the process of identity construction: (i) self-determination and individuation from parents, (ii) gender and sexual identity and (iii) social life and connectedness with peers and adults (other than parents). Regarding our second research objective, i.e. to reveal how the HCPs in this unit individually and collectively adapt their practices to meet what they perceive as essential psychosocial needs of their patients, flexibility emerged as a core category, which was made possible by collaborative work and a number of collective processes that facilitate self-reflection and build a solid team cohesion. Our results are presented according to these emerging categories with examples of concrete practices to illustrate the findings.

## **SUPPORTING IDENTITY CONSTRUCTION**

The participants generally attested that AYAs go through an important process of identity construction, which involves major physical and psychological changes to be integrated. They

1  
2  
3 acknowledged that the experience of cancer largely hampers this process and advocated for the  
4 provision of in-patient care that includes ways of addressing these specific needs. They argued  
5 that an AYA patient does not only have cancer, but is a person who must achieve some  
6 important developmental milestones that need to be taken into account in the care relationship:  
7  
8 « *They remain adolescents, no matter how ill they are. An AYA patient is not an ordinary*  
9 *patient* » (nurse). To mitigate cancer and treatment effects, they identified 3 major challenges  
10 inherent to the identity process of any adolescent that may be particularly threatened by the  
11 experience of cancer and therefore require special attention from the healthcare team.  
12  
13

14  
15 ***Supporting individuation from parents and self-determination.*** HCP insisted on the  
16 importance of supporting the processes of self-determination and individuation as a way of  
17 compensating for the dependence created by in-patient care. Therefore, the team aims to deeply  
18 involve patients in decision-making processes during the whole duration of their  
19 hospitalisation. Parents are less systematically involved by the HCPs in their children's care, as  
20 compared to usual practices in paediatrics: « *What changes, with regards to paediatrics, is that*  
21 *the cancer diagnosis is not given first to the parents but directly disclosed to the teenager* »  
22 (doctor). HCPs assumed that the parents' presence has to be constantly discussed to support the  
23 separation process, which is even more complex in the context of cancer. According to HCPs,  
24 there is no one way of regulating the parents' presence, and individualised decisions need to be  
25 made. A concrete example is the management of extra beds for accompanying parents or  
26 relatives who ask to stay overnight. In this AYA unit, it is not a rule to have an extra bed in the  
27 room, as it is in paediatrics: « *I think that nothing should be imposed but rather proposed at*  
28 *this age. This is more important than for the younger ones.* » (doctor). Another example is how  
29 HCPs engage in trying to prevent parents from reverting to former relationships they would  
30 have had with their child at a younger age, induced by the status of being ill: « *A mother washing*  
31 *her small child doesn't shock me. However, when this happens with an AYA, I find it difficult...*  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *I try to discuss the matter with the parents and that it's not up to them to help with everyday*  
4 *things.* » (nurse aide). Regulating the distance between parents and AYAs has been  
5  
6 acknowledged by the participants as being of significant benefit to AYAs' need for  
7  
8 empowerment: « *I am thinking of situations where the parents behave in an exemplary manner*  
9  
10 *whilst being able to maintain the right distance. This has an extraordinarily beneficial effect*  
11  
12 *on both the young patient and the parent.* » (doctor).  
13  
14  
15

16  
17 ***Regarding the need to self-determine their goals and actions,*** the HCPs in our study  
18  
19 acknowledged that this is basically undermined because of the cancer treatment, which is not  
20  
21 negotiable. As a result of being aware of their young patients' need for self-determination,  
22  
23 flexibility is introduced into the manner in which the treatment is administered, to suit the young  
24  
25 patients' need to remain active decision makers, thus allowing them to develop a sense of  
26  
27 control. Non-compulsory treatments are discussed and sometimes dismissed or postponed: « *I*  
28  
29 *try to respect their preferences. If they don't want to be disturbed too early in the morning for*  
30  
31 *instance, I will accept it once but not a second time. Hence, I am not rushing them but still stand*  
32  
33 *firm.* » (nurse). The participants defined their care relationships and practices as being based on  
34  
35 an implicit rule: “allowing for negotiation *within a fixed framework*” (nurse).  
36  
37  
38  
39

40  
41 In order to allow for negotiations, there are no written internal procedures that would  
42  
43 set the rules too rigidly and hinder the necessary flexibility. In agreement with doctors, mask-  
44  
45 wearing measures are usually relaxed. External home-prepared or commercial frozen meals,  
46  
47 prohibited in other places due to infectious risks, may be permitted under certain conditions.  
48

49  
50 In their capacity as adults, HCPs saw themselves as playing a key role in supporting  
51  
52 their young patients' identity-forging process by tolerating conflicts and differing opinions:  
53  
54 « *After all, we represent the adult world for them, and perhaps it is a good thing that they know*  
55  
56 *they can't do whatever they want to do, even when they are sick* » (nurse). They acknowledged  
57  
58 negotiation and conflicts as making sense in the caring relationship: « *Teenagers are looking*  
59  
60

1  
2  
3 *for conflict. Yes, there are clashes, but these are natural things that happen because we are in*  
4 *facility that deals with adolescence. We need to remember the need for mutual respect all the*  
5 *time » (nurse).*

6  
7  
8  
9  
10 ***Supporting gender and sexual identity.*** HCPs were aware that the challenges around sexuality  
11 and body image are crucial in the process of defining one's identity during adolescence. As  
12 treatments alter the pubescent body, they perceive the importance of enhancing the patient's  
13 emerging gendered identity, much more than when working with younger or older patients:  
14 « *when I was working in paediatrics, I didn't ask myself those kinds of questions. But teenagers*  
15 *become conscious about their own bodies so I cannot take over it without asking them. They*  
16 *are a person, already an adult for some and for others still "in progress", but at the end of the*  
17 *day, they have their say in things!* » (nurse). Consequently, HCPs are extremely respectful of  
18 their young patients' modesty and privacy. In practical terms, the AYA unit is organised with  
19 single rooms which are separated by opaque doors, thus precluding from the possibility of being  
20 observed from the outside. Moreover, for intimate aspects of care, such as hygiene-related  
21 issues or fertility preservation procedures, HCPs single out one member of their team of the  
22 same gender as the patient when they perceive that the patient feels too uncomfortable: « *Our*  
23 *male colleagues are involved in matters of care that bother boys, for instance for explaining*  
24 *the semen collection procedure. Talking to a man about this is easier than to confide in a*  
25 *woman, even more so if she is a young caregiver.* » (nurse).

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47 Finally, the staff noticed that when special care was provided to enhance well-being in  
48 addition to physical treatment, it eased the suffering stemming from a painful and injured body.  
49 Massages are offered to allow for positive experiences and to relieve anxiety and stress. HCPs  
50 were deeply convinced that attending to physical needs other than only those ones exclusively  
51 related to the treatment of cancer contributes to a better body image, self-esteem and therefore  
52 to positive identity construction.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 ***Encouraging social life and connectedness.*** HCPs acknowledged that the process of identity  
4 construction in AYAs involves developing more intense relationships with peers and  
5 maintaining their professional and social projects. They observed the disruptive effects of  
6 diagnosis and prolonged in-patient stays on AYAs' social dynamics. Despite the isolation  
7 measures mandatory in haematology, the AYA unit has been designed with the aim of opening  
8 up the boundaries of the hospital: some open living spaces encourage AYAs' social  
9 relationships and informal meetings. Various practical adaptations have been set up to reduce  
10 their feelings of isolation despite the usual hospital rules: visiting hours were extended to  
11 respect the more nocturnal hours favored by AYAs, hospital discharges are facilitated by a  
12 coordination nurse whenever the staff identifies that a patient might be suffering from social  
13 isolation, a community worker organises collective activities to encourage links between in-  
14 ward patients and coordinates the actions of several out-door organisations. One of these  
15 provides a teacher who lectures in the hospital and liaise with schools in order to facilitate the  
16 patients' pursuit of their studies. A social worker helps young adult patients already engaged in  
17 a professional activity to maintain it by liaising with their employers. This gives an indication  
18 of the numerous professions involved in the care to AYAs with cancer in this unit.

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
These concrete measures show a strong will to approach the AYA patient more  
holistically and the HCPs' conviction that caring means more than curing cancer: « *After all, I  
do not look at them as ill people in the first place, but rather as people.* » (nurse). Our  
participants conveyed that AYAs with cancer need not only efficient and evidence-based  
medical and technical treatment, but also strong links and genuine human connection, which  
they acknowledged as the core component of the care relationship: « *If they need to talk, I don't  
leave, I give them time. Interpersonal relationships are very important in this unit.* » (ward-  
aide). Such involvement was recognised as giving sense to their work. On the other hand, they  
also acknowledged that it might make them more vulnerable: « *In this unit, our caring*

1  
2  
3 *technique is based on relationships. It is not much more difficult than anything else, but it means*  
4 *that we are much more involved.* » (nurse). Reflecting on their efforts to best address their  
5  
6 young patients' needs, HCPs reported that they get personally involved beyond the boundaries  
7  
8 of their traditional professional roles, thus exposing themselves to an increase risk of emotional  
9  
10 burden. Reflective practices and strategies are used to process the constant adjustments that are  
11  
12 required to maintain a collective cohesion and flexibility.  
13  
14  
15

## 16 17 18 **FLEXIBILITY, COLLECTIVE PROCESSES AND SELF-REFLEXION**

19  
20  
21 HCPs were aware that being individually flexible was dependent on collective strategies  
22  
23 aimed at protecting their professional roles and ensuring overall coherence of staff  
24  
25 decisions: « *I always try to give a consistent message, because otherwise AYA patients fall*  
26  
27 *between the cracks.* » (nutritionist). In practice, they reported that formal and informal  
28  
29 teamwork practices had emerged, such as feeling free to ask for help from a colleague whenever  
30  
31 they face emotional difficulties with a patient: « *We must have empathy to be a good caregiver,*  
32  
33 *but we also need to attend our own needs* ». *When I feel emotionally fragile with a patient, I*  
34  
35 *hand him/her over to a colleague.* » (nurse). Working in pairs and mutual trust were cited as  
36  
37 important team resources in which interpersonal relationships are central. Although such  
38  
39 flexibility could be seen as carrying a risk of confusion, HCPs reported that sharing opinions  
40  
41 ensures the team's cohesion: « *I cannot work by myself because we all have our small*  
42  
43 *contribution toward building this care project.* » (social worker). Therefore, HCPs  
44  
45 acknowledged the importance of the right motivation to work in such a ward: « *We are all very*  
46  
47 *motivated here because facing an adolescent requires a high sense of solidarity* » (night nurse).  
48  
49  
50  
51

52  
53 The weekly staff meetings aim to favour a multi-focal approach and individualisation  
54  
55 of care procedures. Personalised decisions are more appropriate than decisions that would be  
56  
57 taken by complying with predefined criteria according to a fixed medical protocol: « *I*  
58  
59 *remember a patient, in aplasia and feverish, who wanted to leave for the weekend. From my*  
60

1  
2  
3 *typical paediatric perspective, letting her out was out of the question... But after discussing it*  
4 *with the team, I let her go out. This enables us to do work on a case by case basis » (doctor).*

5  
6  
7  
8 Every staff member feels responsible for passing on information while optimising his/her  
9  
10 professional role: « *This multidisciplinary exchange is very constructive: it invites us to share*  
11 *different views, thus revealing aspects of a patient's experience, that I would otherwise not*  
12 *necessarily be aware of. This experience enriches me. » (nutritionist).*

13  
14  
15  
16  
17 Open-mindedness and reflexive practices were described as a strong resource that nurtures in  
18  
19 return the HCP's own professional role, and their ability to better meet the demands imposed  
20  
21 on them by AYA patients.  
22

## 23 **DISCUSSION**

24  
25  
26 Identity work was identified by the HCPs as the core challenge that AYAs with cancer  
27  
28 must face, along with the interconnected challenges of building one's autonomy and sexual and  
29  
30 social identities. Whereas these issues have been well studied from the patients' perspective<sup>21</sup>,  
31  
32 they have remained understudied with regard to opinions of HCPs'. Yet, HCPs are particularly  
33  
34 challenged when it comes to support such needs in the patient-provider relationship. This study  
35  
36 does not only report on individual and collective practices meant to support AYA patients'  
37  
38 psychosocial needs but also on strategies and skills HCPs should aim to develop in order to  
39  
40 meet this challenge.  
41  
42  
43

44  
45 First, HCPs contribute concretely to individuation and self-determination targets when  
46  
47 involving patients in decision-making processes and playing a subtle role in negotiating their  
48  
49 parents' presence. Recent research suggests that AYA health communication and collaborative  
50  
51 decision-making is a sensitive issue, because the age of legal responsibility should not  
52  
53 determine who (the parents or the young patient) has to be informed<sup>22-23</sup>. Assessing as  
54  
55 comprehensively and as early as possible the patient's degree of autonomy and the quality of  
56  
57 the parent-AYA relationship should contribute to a better balance of decision-making  
58  
59  
60

1  
2  
3 responsibilities as far as the patient-parent dyad is concerned. This is consistently related to an  
4  
5 important coping strategy to develop AYAs' sense of control<sup>24</sup>.  
6

7  
8 Secondly, our results highlight how HCPs pay attention to AYA patients' sexual  
9  
10 development by being sensitive and responsive to individual patients' level of sexual maturity  
11  
12 and physical development. There is some evidence that treatment-related side effects adversely  
13  
14 affect AYAs' early puberty, sexual function and development<sup>25-28</sup>. Recent research suggests  
15  
16 that impairments of body image seriously threaten the gendered identity if related to lower self-  
17  
18 esteem. Self-esteem plays a key role in the development of the sense of self as a sexual being  
19  
20 at that age<sup>29</sup>. Our study shows that HCPs support the body image with the onset of puberty by  
21  
22 being highly respectful of modesty and privacy. They promote a better body image as they offer  
23  
24 positive sensory experiences and feelings about physical impairments thanks to their careful  
25  
26 handling of physical areas affected by illness or treatments by providing massages. As shown  
27  
28 in other studies, such attitudes are likely to contribute to a better self-acceptance, thus fostering  
29  
30 enhanced relationships with peers and intimate partners<sup>30-31</sup>. We believe that it paves the way  
31  
32 for better communication regarding the emotional, romantic and sexual dimensions of their  
33  
34 quality of life, all of which are key issues in this population of patients<sup>32-33</sup>.  
35  
36  
37  
38  
39

40 Another major issue for HCPs was about maintaining a bond with peers by favouring  
41  
42 outpatient hours, school attendance or the pursuit of work and projects as key elements in  
43  
44 building one's identity at that age. They adapt hospital and ward rules to reduce the disruptive  
45  
46 effects of hospitalisations. In doing so, they provide personalised adjustments which require  
47  
48 that they improve their own tolerance to conflicts. The present study shows how they develop  
49  
50 negotiation skills, patience, with mutual respect as a strategy to promote a genuine and non-  
51  
52 authoritative therapeutic bond. There is some evidence that a strong patient-oncologist alliance  
53  
54 is associated with psychosocial wellbeing and therapeutic adherence in AYA cancer patients<sup>34</sup>.  
55  
56  
57  
58  
59  
60

1  
2  
3 Lastly, our findings highlight that a genuine personal relationship based on open-  
4 mindedness and strong motivation are to be considered subjective abilities HCPs must develop  
5 to support such an alliance. The HCPs' skills are strongly linked with the understudied concept  
6 of empathy in the patient-caregiver relationship. A recent review concludes that HCPs' empathy  
7 is associated with better psychosocial adjustment and less psychological distress in the HCPs<sup>35</sup>,  
8 although there is a lack of nursing-focused research in this field. Various sections of the  
9 transcripts obtained show deep emotional and subjective commitment as core human qualities  
10 developed by the HCPs to favour empathy: it can be seen as a main contribution to this topic.  
11 However, being empathic exposes HCPs to a "double-edged sword"<sup>36</sup>, with a clear benefit for  
12 patients potentially counterbalanced by a detrimental effect on professional stress levels,  
13 especially when patients are facing life-threatening condition<sup>37-40</sup>. It would deserve further  
14 investigation to assess how HCPs' empathy is a source of work exhaustion or distress. Our  
15 results suggest that a reflection about what it means to get personally involved in care should  
16 be introduced early in the curricula of medical and paramedical studies, and the importance of  
17 a trained and balanced workforce within the team (i.e. multidisciplinary and mixed-gendered).  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

#### 40 **Study limitations**

41  
42 The strength of our study lies in the high participation of a multidisciplinary health care  
43 team in an AYA dedicated unit, although nurses are an over-represented professional category.  
44 Moreover, as this was a study conducted in a single centre, the sample of participants was  
45 inevitably small, and our results may not be appropriate for extrapolation. In addition, we opted  
46 for an iterative process of data collection and analysis. Yet, at this stage of our study, we are  
47 aware that we have not reached saturation of the data. Rather, as is typical in grounded theory,  
48 our initial research question has broadened and there are still many issues that need to be  
49 explored to improve the AYA's psychosocial care.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Clinical implications

HCPs have delivered useful recommendations to support AYA's psychosocial development such as providing patients personalised adjustments in their caring practices and implementing strategies to that end. As such, these can be transferable to other similar care situations and may initiate a reflection on practical guidelines for young patient-friendly models of care in dedicated units.

## Conflict of interest statement

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Acknowledgements

We are grateful to all the participants who agreed to share their experience with us. We are most grateful to INCA (Institut National du Cancer) for doctoral fellowship, and to Novartis for their administrative support (basically interviews' transcripts). We also thank the following colleagues for their editorial help at various stages of the writing process: Giulietta Von Salis and Lydia Stephens.

## References

---

- [1] Arnett JJ, Emerging adulthood: a theory of development from the late teens through the twenties, *Am Psychol*, 2000;55 (5):469–80.
- [2] Stam H, Hartman EE, Deurloo JA, Groothoff J, Grootenhuis, Young Adult Patients with a History of Paediatric Disease: Impact on Course of Life and Transition into Adulthood, *Journal of Adolescent Health*, 2006, 39(1), 4-13
- [3] Evan EE, Zeltzer LK. Psychosocial dimensions of cancer in adolescents and young adults. *Cancer*, 2006; 107:1663–71.
- [4] Katz, A, Meeting the need for psychosocial care in young adults with cancer, 2015, Pittsburgh, PA: Oncology Nursing Society.
- [5] Katz A, The trouble with teen and young adults with cancer, *Oncology Nursing Forum*, 2015; 42 (4):327–328.

- 1  
2  
3 [6] Zebrack B, Bleyer A, Albritton K, Medearis S, Tang J, Assessing the health care needs of  
4 adolescent and young adult cancer patients and survivors, *Cancer*, 2006, 12 (107): 2915–  
5 2923.  
6  
7 [7] Zebrack B, Isaacson S, Psychosocial care of adolescent and young adult patients with  
8 cancer and survivors, *J Clin Oncol*, 2012 Apr 10;30(11):1221–6.  
9  
10 [8] Nightingale CL, Quinn GP, Shenkman EA, Curbow BA, Zebrack BJ, Krull KR, Huang  
11 IC, Health-Related Quality of Life of Young Adult Survivors of Childhood Cancer: A Review  
12 of Qualitative Studies, *Journal of adolescent and young adult oncology*, 2011; Vol 1(3):124–  
13 132.  
14  
15 [9] Morgan S, Davies S, Palmer S, Plaster M, Sex, Drugs, and Rock ‘n’ Roll: Caring for  
16 Adolescents and Young Adults with Cancer, *Journ clin oncology*, 2012, vol 28(32):4825–  
17 4830.  
18  
19 [10] Palmer S, Mitchell A, Thompson K, Sexton M, Unmet needs among adolescent cancer  
20 patients: a pilot study, *Palliat Support Care*, 2007 Jun;5(2):127–34.  
21  
22 [11] D'Agostino NM, Penney A, Zebrack B, Providing developmentally appropriate  
23 psychosocial care to adolescent and young adult cancer survivors. *Cancer*. 2011; 117(10  
24 Suppl):2329–2334.  
25  
26 [12] Clinton-McHarh, T., Carey, M., Sanson-Fisher, R., Shakeshaft, A., & Rainbird, K. (2010).  
27 Measuring the psychosocial health of adolescent & young adult (AYA) cancer survivors: A  
28 critical review. *Health & Quality of Life Outcomes*, 8(25)  
29  
30 [13] Farre A, Wood V, McDonagh JE, Parr JR, Reape D, Rapley T, Health professionals' and  
31 managers' definitions of developmentally appropriate healthcare for young people:  
32 conceptual dimensions and embedded controversies, *Arch Dis Child*, 2016 July; 101(7): 628–  
33 633.  
34  
35 [14] Palmer S, Improving care for AYA patients treated within adult hospitals: What can be  
36 done right now?, *Cancer Forum*, 2009, 33, 22–25.  
37  
38 [15] Ferrari A, Thomas D, Franklin ARK, Hayes-Lattin BM, Mascarin M, van der Graaf W,  
39 Albritton KH, Starting an Adolescent and Young Adult Program: Some Success Stories and  
40 Some Obstacles to Overcome, *Clin Oncol*, 2010, 28 (32):4850–4857.  
41  
42 [16] Dyson G, Thompson K, Palmer S, Thomas D, Schofield P, The relationship between  
43 unmet needs and distress amongst young people with cancer, *Supportive Care and Cancer*,  
44 2012 Jan;20(1):75-85  
45  
46 [17] Hollis R, Morgan S, The adolescent with cancer –at the edge of no-man's land, *Lancet*  
47 *Oncol*, 2001 Jan;2(1):43–8.  
48  
49 [18] Smith S, Adolescent units: an evidence-based approach to quality nursing in adolescent  
50 care, *European Journal of Oncology Nursing*, 2004;8:20–29.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 [19] Lewis I, Morgan S, Models of care and specialized units, Bleyer WA, Barr RD, eds.  
4 Cancer in Adolescents and Young Adults. New York: Springer Verlag, 2007, 341–352.  
5  
6 [20] Strauss A, Corbin J, Basics of qualitative research: techniques and procedures for  
7 developing grounded theory. Thousand Oaks, CA: Sage; 1998.  
8  
9 [21] Abrams AN, Muriel AC, Wiener L, Pediatric Psychosocial Oncology: Textbook for  
10 Multidisciplinary Care, New-York: Springer; 2016, chap 12: 199-219: “Adolescents and  
11 Young Adults with Cancer: A Biopsychosocial Approach  
12  
13 [22] Hanna KM, Dashiff CJ, Stump TE, Weaver MT. Parent-Adolescent Dyads: Association  
14 of Parental Autonomy Support and Parent-Adolescent Shared Diabetes Care Responsibility,  
15 Child Care Health Dev., 2013, Sept ; 39(5): 695–702  
16  
17 [23] Compas BE, Jaser SS, Dunn MJ, Rodriguez EM, Coping with Chronic Illness in  
18 Childhood and Adolescence, Annu Rev Clin Psychol. 2012 April 27; 8: 455–480  
19  
20 [24] Sadak KT, Connor C, DeLuca H, Innovative educational approaches to engage and  
21 empower the adolescent and young adult childhood cancer survivor, *Pediatr Blood Cancer*.  
22 2013 Dec;60(12):1919-21.  
23  
24 [25] Martins A, Taylor RM, Lobel B, McCann B, Soanes L, Whelan JS, Fern LA, Sex, Body  
25 Image, and relationships, A brightlight workshop on Information and Support Needs of  
26 Adolescents and Young Adults, *Journal of Adolescent and Young Adult Oncology*, 2018, vol  
27 7 (5), 1-7.  
28  
29 [26] Moules NJ, A tribe apart: Sexuality and cancer in adolescence, *Journal of Pediatric*  
30 *Oncology Nursing*, 2017; Vol 34 issue: 4: 295-308.  
31  
32 [27] Thompson AL, Long KA, Marsland AL, Impact of childhood cancer on emerging adult  
33 survivors’ romantic relationships: A qualitative account, *J Sex Med* 2013;10:65–73  
34  
35 [28] Bolte S, Zebrack B. Sexual issues in special populations: adolescents and young adults.  
36 *Semin Oncol Nurs*. 2008; 24(2):115–19.  
37  
38 [29] Evan E, Kaufman M et al., Sexual health and self-esteem in adolescents and young adults  
39 with cancer, *Cancer*, 2006, Supplement October 1, Vol. 107, 7: 1672-1679.  
40  
41 [30] Robinson L, Miedema B, Easley J, Young Adult Cancer Survivors and the Challenges of  
42 Intimacy, *Journal of Psychosocial Oncol*, 2014;32 (4): 447-462.  
43  
44 [31] Rosenberg AR, Bona K, Ketterl T, Wharton CM, Wolfe J, Baker KS, Intimacy,  
45 Substance Use, and Communication Needs During Cancer Therapy: A Report From the  
46 “Resilience in Adolescents and Young Adults” Study, *J Adolesc Health*. 2017 January; 60(1):  
47 93–99  
48  
49 [32] Dobinson KA, Hoyt MA, Seidler ZE, Beaumont AL, Hullmann SE, Lawsin CR, A  
50 Grounded Theory Investigation into the Psychosexual Unmet Needs of Adolescent and  
51 Young Adult Cancer Survivors, *Journal of Adolescent and Young Adult Oncology*, 2016; Vol  
52 5 (2):135-145.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 [33] Thompson AL, Long KA, Marsland AL, Impact of Childhood Cancer on Emerging  
4 Adult Survivors' Romantic Relationships: A Qualitative Account, *J Sex Med* 2013;10:65–73

5  
6 [34] Trevino KM, Fasciano K, Prigerson HG. Patient-oncologist alliance, psychosocial well-  
7 being, and treatment adherence among young adults with advanced cancer. *J Clin Oncol.*  
8 2013; 31(13):1683– 1689.  
9

10  
11 [35] Mack JW, Fasciano KM, Block SD, Communication About Prognosis With Adolescent  
12 and Young Adult Patients With Cancer: Information Needs, Prognostic Awareness, and  
13 Outcomes of Disclosure, *Journal of Clinical Oncology* 36, no. 18 (June 20 2018) 1861-1867.  
14

15  
16 [36] Lelorain S, Brédart A, Dolbeault S, Sultan S, A systematic review of the associations  
17 between empathy measures and patient outcomes in cancer care, *Psycho-Oncology*, 2012; 21:  
18 1255–1264  
19

20  
21 [37] Abrams AN, Muriel AC, Wiener L, *Pediatric Psychosocial Oncology: Textbook for*  
22 *Multidisciplinary Care*, New-York: Springer; 2016, chap 21:367-377: “Self-Care and  
23 Sustainability for Pediatric Oncology Providers  
24

25 [38] Crumpei I, Dafinoiu I, The relation of clinical empathy to secondary traumatic stress.  
26 *Procedia: Social and Behavioral Sciences*, 2011; 33:438–442  
27

28  
29 [39] Granek L, Bartels U, Scheinemann K, Labrecque M, Barrera M, Grief reactions and  
30 impact of patient death on pediatric oncologists, *Pediatr Blood Cancer*, 2015; 62(1): 134–142.  
31

32 [40] Potter P, Divanbeigi RNJ, Berger J, Norris J, Olsen RNS, Compassion fatigue and burnout:  
33 Prevalence among oncology nurses, *Clinical Journal of Oncology Nursing*, 2010, 14(5), E56.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1- Sample of the HCPs interviewed at the time of the study**

9 day nurses	10 night nurses	1 coordination nurse	2 ward-aides
1 social worker	1 kinesiologist	1 nutritionist	1 community worker
2 doctors	1 junior doctor		
1 senior health manager			

For Peer Review