



Synthesis, Identification and Antibacterial Activities of Amino Acid Schiff Base Cu(II) Complexes with Chlorinated Aromatic Moieties

Nao Otani, Antoine Fayeulle, Daisuke Nakane, Estelle Léonard, Takashiro Akitsu

► To cite this version:

Nao Otani, Antoine Fayeulle, Daisuke Nakane, Estelle Léonard, Takashiro Akitsu. Synthesis, Identification and Antibacterial Activities of Amino Acid Schiff Base Cu(II) Complexes with Chlorinated Aromatic Moieties. *Advances in Applied Microbiology*, 2022, 2 (2), pp.438-448. 10.3390/applmicrobiol2020032 . hal-03699909

HAL Id: hal-03699909

<https://hal.science/hal-03699909>

Submitted on 5 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Article

Synthesis, Identification and Antibacterial Activities of Amino Acid Schiff Base Cu(II) Complexes with Chlorinated Aromatic Moieties

Nao Otani ¹, Antoine Fayeulle ² , Daisuke Nakane ¹, Estelle Léonard ^{2,*} and Takashiro Akitsu ^{1,*}

¹ Department of Chemistry, Faculty of Science, Tokyo University of Science, 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan; akitsulab@gmail.com (N.O.); a29776@rs.tus.ac.jp (D.N.)

² ESCOM, TIMR (Integrated Transformations of Renewable Matter), Centre de Recherche Royallieu, University of Technology of Compiègne, CS 60 319, CEDEX, 60203 Compiègne, France; antoine.fayeulle@utc.fr

* Correspondence: e.leonard@escom.fr (E.L.); akitsu2@rs.tus.ac.jp (T.A.)

Abstract: Amino acid Schiff base Cu(II) complexes were synthesized under microwave irradiation using methanol as a solvent, to maximize the best conditions to obtain the attained compounds, containing aromatics possessing no, one or two chlorine atoms. The compounds' antibacterial activities were tested against Gram-positive and Gram-negative bacteria, and the most active were tested for their antioxidant activities, and as *E. coli*, in particular, was found to be sensitive to these compounds, their interaction with this bacterium was investigated. It was found that, depending on the amino acid used for the formation of the Schiff base ligand, its LogP_{o/w} mono-chlorinated or bis-chlorinated compounds are the most efficient against the tested bacteria.

Keywords: amino acid Schiff base Cu(II) complexes; chlorinated salicylaldehyde; antibacterial activity; interaction with *E. coli*



Citation: Otani, N.; Fayeulle, A.; Nakane, D.; Léonard, E.; Akitsu, T. Synthesis, Identification and Antibacterial Activities of Amino Acid Schiff Base Cu(II) Complexes with Chlorinated Aromatic Moieties. *Appl. Microbiol.* **2022**, *2*, 438–448. <https://doi.org/10.3390/applmicrobiol2020032>

Academic Editors: Seung-Chun Park and Khaled A. Shaaban

Received: 10 May 2022

Accepted: 15 June 2022

Published: 18 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Schiff bases were discovered by Hugo Schiff, and their metal complex derivatives were largely synthesized, for example, incorporating amino acids [1]. Their fields of application are varied and concern, for example, environmental sensors [2], catalysis [3,4], and anticancer [5,6] or antioxidant agents, but they also serve as antimicrobial agents, particularly against bacterial and fungal pathogens, showing that, compared to Co(II), Ni(II) or Zn(II), Cu(II) complexes show lower MIC values, revealing that these complexes present a better growth inhibitory activity [7]. Other Schiff bases' copper complexes were described from salicylaldehyde derivatives and tested as antimicrobials [8,9], but in all the previous literature on amino acid copper-complex Schiff base syntheses [10], it is noticeable that the reaction time is somehow long, lasting from several hours to days.

Since the use of microwave heating to accelerate organic chemical transformations was first reported by the group of Gedye and Giguere/Majetich in 1986 [11,12], many studies using microwaves have been published [13] about the wavelengths, frequencies and applications of various radio waves used in our daily lives. Controlled microwave heating under closed vessel conditions greatly accelerates reactions compared to conventional synthesis methods using external heating, resulting in higher yields of products, fewer unwanted side reactions and the ability to synthesize high-purity compounds with fewer raw materials [13,14]. Microwave-assisted synthesis typically utilizes “microwave dielectric heating” phenomena, such as dipolar polarization and ionic conduction mechanisms, and relies on the ability of the reacting mixture to efficiently absorb microwave energy [15]. The ability of a specific solvent to convert microwave energy into heat at a given frequency and temperature is determined by the so-called loss tangent ($\tan \delta$), expressed as the quotient, $\tan \delta = \epsilon'' / \epsilon'$, where ϵ'' is the dielectric loss, indicative of the efficiency with

which electromagnetic radiation is converted into heat, and ϵ' is the dielectric constant, describing the ability of molecules to be polarized by the electric field. A reaction medium with a high $\tan \delta$ at the standard operating frequency of a microwave synthesis reactor (2.45 GHz) is required for good absorption and, consequently, efficient heating [15]. In general, solvents used for microwave synthesis are classified as having high ($\tan \delta > 0.5$), medium ($\tan \delta 0.1\text{--}0.5$) or low ($\tan \delta < 0.1$) microwave absorption. Microwave synthesis in low-absorbing or microwave-permeable solvents are often not feasible. Among the solvents possessing a high $\tan \delta$, methanol is one of the most suitable solvents for microwave synthesis because of its high microwave absorption with a loss $\tan \delta$ of 0.659 [16]. That is why this method was previously employed for the synthesis of copper Schiff base complexes by microwave from a 2-step procedure, lowering the global reaction time to 10 min [17], leading, for example, to di-chlorinated compounds and, thanks to the synthetic optimized method, new mono-chlorinated Schiff base copper complexes were synthesized in this work, and all these compounds were tested against model bacteria to investigate how they inhibit the growth of these bacteria.

2. Materials and Methods

2.1. Compound Syntheses

Known compounds C1.1–C8.1 were prepared in the previous paper and used as described in [17]. New compounds entries C1.2–C12.2 were prepared and characterized as follows.

$\text{Cu}(\text{CH}_3\text{COO})_2 \cdot \text{H}_2\text{O}$, *L*-alanine, *L*-leucine, *L*-serine and *L*-threonine used in this study were purchased from Wako Fujifilm (Osaka, Japan), and salicylaldehyde, 3-chlorosalicylaldehyde, 4-chlorosalicylaldehyde and 5-chlorosalicylaldehyde were purchased from TCI (Tokyo, Japan). The synthesis of the compounds was conducted at 358 K using an Initiator+ microwave apparatus (Biotage, Tokyo, Japan).

2.1.1. General Procedures

To a methanol solution (20 mL) of *L*-amino acid (0.2 mmol) was added the proper chlorosalicylaldehyde (0.2 mmol), and the mixture was stirred under microwave irradiation at 358 K for 10 min. Then, a methanol solution (20 mL) of $\text{Cu}(\text{CH}_3\text{COO})_2 \cdot \text{H}_2\text{O}$ (0.0399 g, 0.2 mmol) was added and the mixture was stirred under microwave irradiation for 10 min at 358 K to produce the final product. For additional spectra, see Supplementary Materials.

2.1.2. Physical Measurements

Elemental analyses were conducted with a Perkin-Elmer 2400II CHNS/O analyzer, (Waltham, MA, USA) at the Tokyo University of Science. Infrared (IR) spectra were recorded on a JASCO FT-IR 4200 spectrophotometer (JASCO, Tokyo, Japan) in the range of $4000\text{--}400\text{ cm}^{-1}$ at 298 K. Absorption electronic (UV-Vis) spectra were measured on a JASCO V-570 spectrophotometer in the range of 800–250 nm at 298 K.

2.1.3. Preparation of C1.2

Using the general procedure with *L*-leucine (0.02623 g, 0.2 mmol) and 3-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.06985 g, 54.73%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether and ethanol, and dried in a desiccator for several days. This product was then filtered. Anal. Calcd. for $\text{C}_{13}\text{H}_{16}\text{ClCuNO}_4$: C, 44.71; H, 4.62; N, 4.01%; found: C, 45.69; H, 3.5; N, 3.84%.

2.1.4. Preparation of C2.2

Using the general procedure with *L*-leucine (0.02623 g, 0.2 mmol) and 4-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.06985 g, 32.85%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $\text{C}_{13}\text{H}_{16}\text{ClCuNO}_4$: C, 44.71; H, 4.62; N, 4.01%; found: C, 43.39; H, 2.51; N, 1.20%.

2.1.5. Preparation of C3.2

Using the general procedure with *L*-leucine (0.02623 g, 0.2 mmol) and 5-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.03918 g, 56.09%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{13}H_{16}ClCuNO_4$: C, 44.71; H, 4.62; N, 4.01%; found: C, 45.1; H, 3.71; N, 3.82%.

2.1.6. Preparation of C4.2

Using the general procedure with *L*-alanine (0.01782 g, 0.2 mmol) and 3-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.40000 g, 65.11%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{10}H_{10}ClCuNO_4$: C, 39.10; H, 3.28; N, 4.56%; found: C, 40.78; H, 2.45; N, 4.43%.

2.1.7. Preparation of C5.2

Using the general procedure with *L*-alanine (0.01782 g, 0.2 mmol) and 4-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.04146 g, 67.48%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{10}H_{10}ClCuNO_4$: C, 39.10; H, 3.28; N, 4.56%; found: C, 41.39; H, 2.32; N, 4.54%.

2.1.8. Preparation of C6.2

To a methanol solution (20 mL) of alanine (0.01782 g, 0.2 mmol) was added 4-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) and stirred at 358 K for 10 min to obtain a pale yellow color. Then, a methanol solution (20 mL) of $Cu(CH_3COO)_2 \cdot H_2O$ (0.03990 g, 0.2 mmol) was added, and the mixture was stirred for 10 min at 358 K to produce a green compound (yield 0.01060 g, 17.25%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{10}H_{10}ClCuNO_4$: C, 39.10; H, 3.28; N, 4.56%; found: C, 36.6; H, 3.13; N, 4.08%.

2.1.9. Preparation of C7.2

To a methanol solution (20 mL) of serine (0.02102 g, 0.2 mmol) was added 3-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) and stirred at 358 K for 10 min to obtain a pale yellow color. Then, a methanol solution (20 mL) of $Cu(CH_3COO)_2 \cdot H_2O$ (0.0399 g, 0.2 mmol) was added, and the mixture was stirred for 10 min at 358 K to produce a green compound (yield 0.03737 g, 58.04%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{10}H_{10}ClCuNO_5$: C, 37.16; H, 3.12; N, 4.33%; found: C, 33.17; H, 3.27; N, 7.17%.

2.1.10. Preparation of C8.2

Using the general procedure with *L*-serine (0.02102 g, 0.2 mmol) and 4-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.04404 g, 68.40%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{10}H_{10}ClCuNO_5$: C, 37.16; H, 3.12; N, 4.33%; found: C, 35.31; H, 2.88; N, 4.95%.

2.1.11. Preparation of C9.2

Using the general procedure with *L*-serine (0.02102 g, 0.2 mmol) and 5-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.04678 g, 72.65%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{10}H_{10}ClCuNO_5$: C, 37.16; H, 3.12; N, 4.33%; found: C, 34.34; H, 2.42; N, 3.17%.

2.1.12. Preparation of C10.2

Using the general procedure with *L*-threonine (0.02380 g, 0.2 mmol) and 4-chlorosalicylaldehyde (0.03131 g, 0.2 mmol) produced a green compound (yield 0.03446 g, 51.09%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{11}H_{12}ClCuNO_5$: C, 39.18; H, 3.59; N, 4.15%; found: C, 34.34; H, 3.28; N, 6.37%.

2.1.13. Preparation of C11.2

Using the general procedure with *L*-threonine (0.02382 g, 0.2 mmol) and 4-chlorosalicylaldehyde (0.03135 g, 0.2 mmol) produced a green compound (yield 0.02844 g, 42.17%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{11}H_{12}ClCuNO_5$: C, 39.18; H, 3.59; N, 4.15%; found: C, 44.48; H, 1.62; N, 6.71%.

2.1.14. Preparation of C12.2

Using the general procedure, *L*-threonine (0.02382 g, 0.2 mmol) and 4-chlorosalicylaldehyde (0.03135 g, 0.2 mmol) produced a green compound (yield 0.13912 g, 20.63%). The resulting crude compound was filtered and the precipitate was washed with diethyl ether, acetone and ethanol, then dried in a desiccator for several days. This product was filtered. Anal. Calcd. for $C_{11}H_{12}ClCuNO_5$: C, 39.18; H, 3.59; N, 4.15%; found: C, 44.62; H, 1.53; N, 14.03%.

2.2. Antimicrobial Assays

The bacterial strains used were *E. coli* ATCC 25922TM (American Type Culture Collection, Manassas, VA, USA), *M. luteus* CRBIP 107660 (Institut Pasteur, Lille, France), *S. saprophyticus* (isolated by TIMR Laboratory) and *B. subtilis* ATCC 6051 (American Type Culture Collection, Manassas, VA, USA). They were grown on trypto-casein soy solid medium (TSA, Conda, Madrid, Spain) and incubated at 30 °C in the dark (SANYO, incubator, MIR-253) for 24 h. The solid cultures were then stored in a cold room at 4 °C or used to inoculate 5 mL of liquid mineral medium 24 h before the microplate test (glucose 10 g/L, KCl 0.250 g/L, NaH_2PO_4 6.464 g/L, $Na_2HPO_4 \cdot 2H_2O$ 10.408 g/L, $MgSO_4$ 0.244g/L, NO_3NH_4 1 g/L, $MgCl_2$ 0.05 g/L; pH = 7). A suspension of bacterial inoculum with an optical density (OD) between 0.1 and 0.2 (Ultrospec10 Amersham Bioscience) was prepared from a pre-culture for 24 h. The tests were performed on 96-well microplates (Thermo Scientific, Nunc™ Edge) in triplicate. The tested products were dissolved in DMSO as 100-times-concentrated stock solutions. Blanks were composed of non-inoculated mineral medium with or without the tested compounds to withdraw their eventual variation of absorbance along with time; positive controls were conducted with inoculated mineral medium without the compound and tests were inoculated media with the tested compounds. Growth monitoring was performed with a spectrophotometer Thermo Fisher Scientific Multiskan GO (type 1510). It was set to perform OD measurements at 600 nm of each well every 15 min for 24 h to follow the growth kinetics.

The percentage of inhibition was calculated as follows:

$$\%Inhibition = \left(1 - \frac{ODvar\ growth}{ODvar\ control} \right) * 100 \quad (1)$$

ODvar growth was the difference in OD between the highest and lowest points on the growth curve in the tests. *ODvar control* was the difference in OD between the highest and lowest points on the positive control curve.

The MIC₉₅ value was considered to be the lowest tested concentration value at which at least 95% inhibition was achieved.

2.3. Antioxidant Assays

The tests were performed on 96-well microplates (Thermo Scientific, Nunc™ Edge) in triplicate. The tested products were dissolved in DMSO. Oxidation monitoring was performed with a spectrophotometer Thermo Fisher Scientific Multiskan GO (Type 1510). It was set to perform OD measurements at 450 nm of each well every 1 min for 6 h to follow the oxidation kinetics (2 µL of sodium persulfate at 5 mg/mL; 2 µL of the tested compound of 50 µL ABTS at 0.08%) in a citrate-phosphate buffer (0.1 M) for a total volume of 200 µL. The maximum oxidation curve for the positive control reached the absorbance of 0.7. The percentage of inhibition was calculated as follows:

$$\%Inhibition = (1 - (OD_{var\ test}) / (OD_{var\ control})) * 100 \quad (2)$$

$OD_{var\ test}$ was the difference in OD between the highest and the lowest points on the test curve. $OD_{var\ control}$ was the difference in OD between the highest and lowest points on the control curve.

2.4. Interaction with *E. coli*

The tested compounds in DMSO (200 µg/mL as the final concentration) were mixed in culture media with *E. coli*. Their UV-Vis spectra were recorded after filtration on a 0.2 µm PTFE syringe filter at different times of incubation at 30 °C for up to 24 h. The values were recorded depending on the complex λ_{max} values (384 nm for C4.1, 372 nm for C6.1, 369 nm for C2.2, 357 nm for C8.2, 366 nm for C9.2 and 378 nm for C11.2) and compared to the solutions without culture media.

2.5. Log $P_{o/w}$ Calculation

The theoretical Log $P_{o/w}$ were calculated thanks to SwissADME from the Swiss Institute of Bioinformatics under the CC-BY 4.0 Creative Commons 4.0 International License.


2.6. Statistical Analysis

Data are expressed as means \pm standard deviation, and the statistical significance ($p < 0.05$) was determined by one-way ANOVA with Tukey's post hoc analysis.

3. Results and Discussions

3.1. Chemistry

For the preparation of *L*-amino acid derivative Schiff base Cu(II) complexes, two-step reactions were employed, namely, (1) imine condensation of primary amine (*L*-amino acid) and aldehyde, and (2) coordination of the Cu(II) ion from an acetate source (Scheme 1). In previous work, microwave syntheses, conventional heating and mechanochemistry for the corresponding *L*-amino acid derivative Schiff base Cu (II) complexes (series 1) [17] were compared, and this compound library was enriched with various mono-chloride compounds (series 2) to better understand the structure–activity link between the amino acid functional group/number and place of the chlorine moiety, as well as the antibacterial activity.



Scheme 1. Typical reaction scheme in a solution for the *L*-amino acid derivative Schiff base Cu(II) complexes.


From the previous work (series 1), it is noteworthy that a two-step 5 + 5 (min) microwave synthesis at 85 °C is really efficient for the reaction between valine or threonine and salicylaldehyde (entries 1 and 7, 86% yield), while dichloro salicylaldehyde with threonine or alanine produced lower results (entries 6 and 8, 75% and 78% yield). The new series was then synthesized thanks to the screening antimicrobial results (see 3.2 Antibacterial Effect), and *L*-leucine (non-polar amino acid, entries 9–11) showed that the chlorine position in R³ or R⁵ produced a yield of 55% and 56%; whereas, with R⁴ = Cl, only a 33% yield was obtained. With *L*-alanine (non-polar amino acid, entries 12–14), the best result was obtained for R⁴ = Cl (67% yield); whereas, for *L*-threonine (non-charged polar amino acid, entries 18–20), the best result was obtained with R⁵ = Cl (51% yield, respectively) lowered to R³ = Cl or both amino acids (17% and 21% yields, respectively). As for the *L*-serine derivatives (non-charged polar amino acid entries 15–17), the best result was obtained with R³ = Cl (73% yield) and the lower yield was obtained for R⁵ = Cl (58%). All these results show that there is clearly an impact from the chlorine position and the amino acid functional group on the yield obtained, not really correlated to their polarity or their charge.

3.2. Antibacterial Effect

The synthesized 20 compounds were tested at a concentration of 50 µg/mL against four bacteria *Bacillus subtilis* (Gram (+), rod-shaped), *Staphylococcus saprophyticus* (Gram (+), coccus), *Micrococcus luteus* (Gram (+), coccus) and *Escherichia coli* (Gram (−), rod-shaped). These bacterial strains were chosen in order to have a good representativity of bacterial morphologies, cell grouping and cell wall structures, which are important factors influencing resistance to antibacterial molecules. Moreover, the species studied are described as human pathogens regarding *E. coli* [18] and *S. saprophyticus* [19], or in food poisoning with *B. subtilis* [20] and *M. luteus* [21]. The first group (Table 1, Cn.1, entries 1–8) was chosen for its various functions led by the amino acids (R¹ and R²), as well as the presence of 2 or no chlorides (R³ and R⁵). Then, the second group (Table 1, Cn.2, entries 9–20) was chosen according to the best antibacterial results obtained with group 1, and the mono-chlorinated positions varied (R³, R⁴ and R⁵). It was clear from the antibacterial tests of the first group that compounds C3.1, C4.1 and C6.1 (entries 3, 4 and 6) had the best antibacterial effect (Figure 1). Indeed, C3.1 was active against all the Gram (+) bacteria, while C4.1 and C6.1 were active against both Gram (+) and Gram (−) bacteria. The last compound, C8.1, was found to be active against all the Gram (+) bacteria, but totally inactive against *E. coli*. The common factor of these three compounds compared to the others (C1.1, C2.1, C5.1 and C7.1) was the presence of the bis-chlorine atoms. Moreover, focusing on the Gram (−) bacteria, C4.1 and C6.1, which were active against this strain, were exempted from oxygen moiety afforded by the amino acid part, whereas both C3.1 and C8.1 possessed a hydroxyl group. Lastly, between the two hydroxylated compounds, the methyl moiety seemed to have a deleterious effect on the activity compared to the H atom. Based on these first excellent results by C3.1, C4.1, C6.1 and C8.1, the second group (Cn.2) was synthesized, but with moderating the place of the mono-chlorine atom (R³, R⁴ and R⁵) and then tested against the four same bacteria. First, it was really interesting to observe that C6.1 mono-chlorinated isosteres (C4.2, C5.2, C6.2) were less interesting than the bis-chlorinated compound. Then, comparing C1.2, C2.2 and C3.2 led to the observation that the most active compounds against all the strains possessed chlorine in the R⁴ position. In a similar manner, comparing C7.2, C8.2 and C9.2 found that C8.2 and C9.2 were the most active compounds and, most importantly, even better than their mono-chlorinated isostere C3.1. As expected, C10.2, C11.2 and C12.2 compounds were less active as their isostere C8.1, even though the mono-chlorinated compound C11.2 was the best of them. According to this antibacterial screening result, MIC (minimum inhibition concentration) values were tested and produced the following results.

Table 1. Summary of the results for the obtained 2-step compounds.

Entry	Series	Compound	R ¹	R ²	R ³	R ⁴	R ⁵	Yield ^a
1	1	C1.1	CH ₃	CH ₃	H	H	H	86% ^b
2		C2.1	OH	H	H	H	H	65% ^b
3		C3.1	OH	H	Cl	H	Cl	49% ^b
4		C4.1	ⁱ Pr	H	Cl	H	Cl	31% ^b
5		C5.1	H	H	H	H	H	67% ^b
6		C6.1	H	H	Cl	H	Cl	75% ^b
7		C7.1	OH	CH ₃	H	H	H	86% ^b
8		C8.1	OH	CH ₃	Cl	H	Cl	78% ^b
9	2	C1.2	ⁱ Pr	H	H	H	Cl	55%
10		C2.2	ⁱ Pr	H	H	Cl	H	33%
11		C3.2	ⁱ Pr	H	Cl	H	H	56%
12		C4.2	H	H	H	H	Cl	65%
13		C5.2	H	H	H	Cl	H	67%
14		C6.2	H	H	Cl	H	H	17%
15		C7.2	OH	H	H	H	Cl	58%
16		C8.2	OH	H	H	Cl	H	68%
17		C9.2	OH	H	Cl	H	H	73%
18		C10.2	OH	CH ₃	H	H	Cl	51%
19		C11.2	OH	CH ₃	H	Cl	H	42%
20		C12.2	OH	CH ₃	Cl	H	H	21%

^a Isolated yields; ^b from previous work [17].**Figure 1.** Inhibition rate of Cn.1 and Cn.2 against *B. subtilis*, *E. coli*, *M. luteus* and *S. saprophyticus*.

3.3. MIC₉₅ and MIC₅₀ Values

MIC₉₅ values are the minimal concentration where the inhibition reaches at least 95% and MIC₅₀ values are minimal concentration where the inhibition reaches at least 50%. The results for the tested concentrations are 50, 25, 12.5, 6.25, 3.12, 1.56 and 0.78 µg/mL. From Table 2, it can be observed that, for the bacteria *S. saprophyticus*, the bis-chlorinated molecules exhibited the lowest MIC value. This was totally inverted as for one of the other Gram (+) strains, *B. subtilis*, since the mono-chlorinated compounds possessed the lowest MIC values, maybe due to the difference of cell morphology between the two bacteria. Surprisingly, for *E. coli*, which is a Gram (−) bacteria, the MIC values were the lowest, whatever the molecules, with the best values for C6.1, C2.2 and C11.2. Indeed Gram (−) bacteria are generally described as more resistant against antibacterial molecules due to their more selective cell wall and other more specific mechanisms of resistance [22]. By comparison, between all the tested molecules, on *S. saprophyticus*, *E. coli* and *M. luteus*, the compounds C8.2 and C9.2 exhibited the lowest activity. Moreover, C2.2 and C11.2 were more active against the rod-shaped molecules than the coccus.

Table 2. MIC₉₅/MIC₅₀ values (µg/mL).

Compounds	<i>S. saprophyticus</i>	<i>E. coli</i>	<i>M. luteus</i>	<i>B. subtilis</i>
C4.1	25/25	25/25	50/12.5	>50/50
C6.1	12.5/12.5	12.5/12.5	25/6.25	>50/50
C2.2	50/25	6.25/6.25	50/12.5	6.25/3.12
C8.2	50/50	50/50	>50/12.5	25/12.5
C9.2	50/50	50/50	>50/50	25/12.5
C11.2	50/50	12.5/12.5	>50/50	12.5/12.5
Streptomycin	25/6.25	-	-	-
Daptomycin	-	-	7.75/2	30/15
Polymyxin B	-	3.75/1	-	-

3.4. Antioxidant Activity

ROS (reactive oxygen species) were suspected to be the cause of the bacterial inhibition, so oxidation tests were performed with ABTS (2,2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid) and copper compounds (C4.1, C6.1, C2.2, C8.2, C9.2 and C11.2), and compared to sodium persulfate, in the dark or under a white lamp (60 watt). The result produced absolutely no oxidized ABTS species, And, as mentioned in the literature, copper Schiff-based compounds can exhibit antioxidant activity [23–25]. The antioxidant activity of the best antibacterial compounds C4.1, C6.1, C2.2, C8.2, C9.2 and C11.2 was kinetically tested against the oxidation of ABTS to the ABTS⁺ radical [26] by sodium persulfate (Figure 2).

Two groups of antiradical compounds were identified, as it was clear that the first group (C4.1 and C6.1) did not reach a 50% inhibition with a plateau of antioxidant capability from 200–250 µg/mL and over. However, the second group of compounds (C2.2, C8.2, C9.2 and C11.2) was able to reach 50% inhibition at a 250 µg/mL concentration and in the same statistical group. The conclusion may be that the mono-chlorinated compounds of these series of Schiff bases were more active against the radical oxidation of ABTS than the bis-chlorinated ones. IC₅₀ was thus calculated from this second group of compounds and compared to the standard *L*-ascorbic acid (AA) (Table 3).




Figure 2. ABTS radical inhibition by C4.1, C6.1, C8.2, C9.2 and C11.2 compounds. Bars with different letters indicate statistically significant differences among the groups.

Table 3. IC₅₀ (mM) compared to standard L-ascorbic acid.

	C2.2	C8.2	C9.2	C11.2	AA
IC ₅₀ (mM)	0.72	0.77	0.77	0.74	0.14

3.5. Interaction of Copper Complexes with *E. coli*

Copper ions and complexes are known to be biologically active due to the redox processes involved [27]. However, if *E. coli* is equipped with multiple systems to ensure safe copper handling under varying environmental conditions [28,29], in our case, this bacteria was the most sensitive to the copper Schiff base compounds. Therefore, the method of Joseph et al. [30] was adapted to evaluate the capability of the most active compounds to interact with *E. coli*. The absorption band of the complex in the culture media at their respective λ_{\max} wavelengths lowered after interacting with *E. coli* (Table 4).

Table 4. Absorbance reduction (%) of copper Schiff base compounds in solution after *E. coli* incubation.

	C4.1	C6.1	C2.2	C8.2	C9.2	C11.2
Absorbance reduction (%)	60.3	34.3	34.4	11.9	2.3	82.3

This absorbance reduction after 24 h of incubation at 30 °C can be correlated to the compound activity. Indeed, the lowest absorbance reductions (2% and 12%) were observed for C9.2 and C8.2, respectively, and for these two compounds, the MIC₉₅ values of 50 µg/mL were the highest. As for the other compounds, good absorbance reduction greater than 34% led to lower MIC₉₅ from 6.25 to 25 µg/mL. In this group of active compounds, it is important to notice that, even if C2.2 was less absorbed than C11.2 or C4.1 by *E. coli*, its better activity might be attributed to its functional groups and chlorine placement. Additionally, the theoretical LogP_{o/w} is interesting while comparing the uncomplexed ligands. Indeed, while this Log P_{o/w} for L4.1 led to C4.1 when complexed copper was high (3.31), the values for L8.2 and L9.2 were low (1.07 and 1.06, respectively). This could also explain why if C4.1 is highly absorbed by *E. coli*, its activity is not as good as the activities of C2.2, C6.1 and C11.2 (Log P_{o/w} for L2.2 = 2.69, for L6.1 = 2.37 and for L11.2 = 1.39).

4. Conclusions

Amino acid Schiff base Cu(II) complexes were synthesized under microwave irradiation using methanol as a solvent, to maximize the best conditions to obtain the obtained compounds. First, it was found that some compounds possessing bis-chlorinated moieties had better antibacterial activity compared to non-chlorinated aromatic. The place and number of chlorine atoms on the salicylaldehyde was then investigated and, depending on the amino acid used for the formation of the Schiff base ligand, its LogP_{o/w} mono-chlorinated or bis-chlorinated compounds were the most efficient against the tested bacteria. Finally, the light antioxidant effect and the sensitivity of *E. coli* towards the most active compounds led us to test their interaction capability to be absorbed by *E. coli*. Subsequently, we plan to verify the toxicity of the molecules to guide the application, depending on the results obtained for skin, kidney, liver cells.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/applmicrobiol2020032/s1>: Figures S1–S6: IR spectra, Figures S7–S12: UV-Vis spectra, Figure S13: Tentative crystal structures; Table S1: Compounds.

Author Contributions: E.L. (PA, INV, MET, WOD); A.F. (MET, VAL, WRE); N.O. (INV), D.N. (VAL), T.A. (CO, PA, SUP, WOD). (CO) Conceptualization; (DC) Data curation; (FA) Formal Analysis; (FUA) Funding acquisition; (INV) Investigation; (MET) Methodology; (PA) Project administration; (RES) Resources; (SOF) Software; (SUP) Supervision; (VAL) Validation; (VIS) Visualization; (WOD) Writing—original draft; (WRE) Writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by TIMR UTC-ESCOM, and this work was supported by a Grant-in-Aid for Scientific Research (A) KAKENHI (20H00336).

Data Availability Statement: Not applicable.

Acknowledgments: The authors thank Yuta Mitani for his help with the X-ray crystallography.

Conflicts of Interest: There are no conflicts to declare.

References

1. Arunadevi, A.; Raman, N. Biological Response of Schiff Base Metal Complexes Incorporating Amino Acids—A Short Review. *J. Coord. Chem.* **2020**, *73*, 2095–2116. [\[CrossRef\]](#)
2. Cimerman, Z.; Galic, N.; Bosner, B. The Schiff Bases of Salicylaldehyde and Aminopyridines as Highly Sensitive Analytical Reagents. *Anal. Chim. Acta* **1997**, *343*, 145–153. [\[CrossRef\]](#)
3. Al-Hussein, M.F.I.; Adam, M.S.S. Catalytic Evaluation of Copper (II) N-Salicylidene-Amino Acid Schiff Base in the Various Catalytic Processes. *Appl. Organomet. Chem.* **2020**, *34*, e5598. [\[CrossRef\]](#)
4. Chakraborty, H.; Paul, N.; Rahman, M.L. Catalytic Activities of Schiff Base Aquocomplexes of Copper(II) towards Hydrolysis of Amino Acid Esters. *Transit. Met. Chem.* **1994**, *19*, 524–526. [\[CrossRef\]](#)
5. Zehra, S.; Roisnel, T.; Arjmand, F. Enantiomeric Amino Acid Schiff Base Copper(II) Complexes as a New Class of RNA-Targeted Metallo-Intercalators: Single X-Ray Crystal Structural Details, Comparative in Vitro DNA/RNA Binding Profile, Cleavage, and Cytotoxicity. *ACS Omega* **2019**, *4*, 7691–7705. [\[CrossRef\]](#)
6. Zuo, J.; Bi, C.; Fan, Y.; Buac, D.; Nardon, C.; Daniel, K.G.; Dou, Q.P. Cellular and Computational Studies of Proteasome Inhibition and Apoptosis Induction in Human Cancer Cells by Amino Acid Schiff Base–Copper Complexes. *J. Inorg. Biochem.* **2013**, *118*, 83–93. [\[CrossRef\]](#)
7. Kumaravel, G.; Panya Utthra, P.; Raman, N. Exploiting the Biological Efficacy of Benzimidazole Based Schiff Base Complexes with L-Histidine as a Co-Ligand: Combined Molecular Docking, DNA Interaction, Antimicrobial and Cytotoxic Studies. *Bioorganic Chem.* **2018**, *77*, 269–279. [\[CrossRef\]](#)
8. Saikumari, N. Synthesis and Characterization of Amino Acid Schiff Base and Its Copper (II) Complex and Its Antimicrobial Studies. *Mater. Today Proc.* **2021**, *47*, 1777–1781. [\[CrossRef\]](#)
9. Costamagna, J.; Vargas, J.; Latorre, R.; Alvarado, A.; Mena, G. Coordination Compounds of Copper, Nickel and Iron with Schiff Bases Derived from Hydroxynaphthaldehydes and Salicylaldehydes. *Coord. Chem. Rev.* **1992**, *119*, 67–88. [\[CrossRef\]](#)
10. Nakao, Y.; Sakurai, K.; Nakahara, A. Copper (II) Chelates of Schiff Bases Derived from Salicylaldehyde and Various α -Amino Acids. *Bull. Chem. Soc. Jpn.* **1967**, *40*, 1536–1538. [\[CrossRef\]](#)
11. Gedye, R.; Smith, F.; Westaway, K.; Ali, H.; Baldisera, L.; Laberge, L.; Rousell, J. The Use of Microwave Ovens for Rapid Organic Synthesis. *Tetrahedron Lett.* **1986**, *27*, 279–282. [\[CrossRef\]](#)

12. Giguere, R.J.; Bray, T.L.; Duncan, S.M.; Majetich, G. Application of Commercial Microwave Ovens to Organic Synthesis. *Tetrahedron Lett.* **1986**, *27*, 4945–4948. [[CrossRef](#)]
13. Gedye, R. *Microwave-Enhanced Chemistry. Fundamentals, Sample Preparation and Applications* Edited by H.M. Kingston (Duquesne University) and Stephen J. Haswell (University of Hull); American Chemical Society: Washington, DC, USA, 1997; p. 772. ISBN 0-8412-3375-6.
14. Strauss, C.R. Microwave-Assisted Reactions in Organic Synthesis—Are There Any Nonthermal Microwave Effects? Response to the Highlight by N. Kuhnert. *Angew. Chem. Int. Ed.* **2002**, *41*, 3589–3591. [[CrossRef](#)]
15. Gabriel, C.; Gabriel, S.; Grant, E.H.; Grant, E.H.; Halstead, B.S.J.; Mingos, D.M.P. Dielectric Parameters Relevant to Microwave Dielectric Heating. *Chem. Soc. Rev.* **1998**, *27*, 213–224. [[CrossRef](#)]
16. Dallinger, D.; Kappe, C.O. Microwave-Assisted Synthesis in Water as Solvent. *Chem. Rev.* **2007**, *107*, 2563–2591. [[CrossRef](#)]
17. Otani, N.; Furuya, T.; Katsuumi, N.; Haraguchi, T.; Akitsu, T. Synthesis of Amino Acid Derivative Schiff Base Copper(II) Complexes by Microwave and Wet Mechanochemical Methods. *J. Indian Chem. Soc.* **2021**, *98*, 100004. [[CrossRef](#)]
18. Kaper, J.B.; Nataro, J.P.; Mobley, H.L.T. Pathogenic Escherichia Coli. *Nat. Rev. Microbiol.* **2004**, *2*, 123–140. [[CrossRef](#)]
19. Kuroda, M.; Yamashita, A.; Hirakawa, H.; Kumano, M.; Morikawa, K.; Higashide, M.; Maruyama, A.; Inose, Y.; Matoba, K.; Toh, H.; et al. Whole Genome Sequence of Staphylococcus Saprophyticus Reveals the Pathogenesis of Uncomplicated Urinary Tract Infection. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 13272–13277. [[CrossRef](#)]
20. Turnbull, P.C.B. Bacillus. In *Medical Microbiology*; Baron, S., Ed.; University of Texas Medical Branch at Galveston: Galveston, TX, USA, 1996; ISBN 978-0-9631172-1-2.
21. Zhang, R.; Zhang, Y.; Zhang, T.; Xu, M.; Wang, H.; Zhang, S.; Zhang, T.; Zhou, W.; Shi, G. Establishing a MALDI-TOF-TOF-MS Method for Rapid Identification of Three Common Gram-Positive Bacteria (*Bacillus Cereus*, *Listeria Monocytogenes*, and *Micrococcus Luteus*) Associated with Foodborne Diseases. *Food Sci. Technol* **2022**, *42*. [[CrossRef](#)]
22. Breijyeh, Z.; Jubeh, B.; Karaman, R. Resistance of Gram-Negative Bacteria to Current Antibacterial Agents and Approaches to Resolve It. *Molecules* **2020**, *25*, 1340. [[CrossRef](#)]
23. Shah, S.S.; Shah, D.; Khan, I.; Ahmad, S.; Ali, U.; ur Rahman, A. Synthesis and Antioxidant Activities of Schiff Bases and Their Complexes: An Updated Review. *Biointerf. Res. Appl. Chem.* **2020**, *10*, 6936–6963.
24. Ganji, N.; Rambabu, A.; Vamsikrishna, N.; Daravath, S. Copper (II) Complexes with Isoxazole Schiff Bases: Synthesis, Spectroscopic Investigation, DNA Binding and Nuclease Activities, Antioxidant and Antimicrobial Studies. *J. Mol. Struct.* **2018**, *1173*, 173–182. [[CrossRef](#)]
25. Soberanes, Y.; López-Gastélum, K.-A.; Moreno-Urbalejo, J.; Salazar-Medina, A.J.; del Carmen Estrada-Montoya, M.; Sugich-Miranda, R.; Hernandez-Paredes, J.; Gonzalez-Córdova, A.F.; Vallejo-Cordoba, B.; Sotelo-Mundo, R.R.; et al. Tetrameric Copper(II) Metalloacyclic Complex Bearing an Amino Acid Derived Schiff Base Ligand: Structure, Catalytic and Antioxidant Activities. *Inorg. Chem. Commun.* **2018**, *94*, 139–141. [[CrossRef](#)]
26. Ilyasov, I.R.; Beloborodov, V.L.; Selivanova, I.A. Three ABTS•+ Radical Cation-Based Approaches for the Evaluation of Antioxidant Activity: Fast-and Slow-Reacting Antioxidant Behavior. *Chem. Pap.* **2018**, *72*, 1917–1925. [[CrossRef](#)]
27. Mitra, D.; Kang, E.-T.; Neoh, K.G. Antimicrobial Copper-Based Materials and Coatings: Potential Multifaceted Biomedical Applications. *ACS Appl. Mater. Interfaces* **2020**, *12*, 21159–21182. [[CrossRef](#)]
28. Rensing, C.; Grass, G. Escherichia Coli Mechanisms of Copper Homeostasis in a Changing Environment. *FEMS Microbiol. Rev.* **2003**, *27*, 197–213. [[CrossRef](#)]
29. Vincent, M.; Duval, R.; Hartemann, P.; Engels-Deutsch, M. Contact Killing and Antimicrobial Properties of Copper. *J. Appl. Microbiol.* **2018**, *124*, 1032–1046. [[CrossRef](#)]
30. Joseph, J.; Nagashri, K.; Rani, G.A.B. Synthesis, Characterization and Antimicrobial Activities of Copper Complexes Derived from 4-Aminoantipyrine Derivatives. *J. Saudi Chem. Soc.* **2013**, *17*, 285–294. [[CrossRef](#)]