



HAL
open science

Fractional dynamics of a transmission model of Zika disease

Hamadjam Abboubakar, Raissa Kom Regonne, Kevin André Nyemb Ndjem Eone

► **To cite this version:**

Hamadjam Abboubakar, Raissa Kom Regonne, Kevin André Nyemb Ndjem Eone. Fractional dynamics of a transmission model of Zika disease. Colloque Africain sur la Recherche en Informatique et en Mathématiques Appliquées (CARI), Oct 2022, Tunis, Tunisia. hal-03702663

HAL Id: hal-03702663

<https://hal.science/hal-03702663>

Submitted on 23 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fractional dynamics of a transmission model of Zika disease

Hamadjam Abboubakar^{*1,2,3}, Raissa Kom Regonne⁴, André Kevin Nyemb Ndjem Eone²

¹Department of Computer Engineering, U.I.T of Ngaoundéré, P.O. Box 455, Ngaoundéré, Cameroon

²LASE-Laboratory, University Institute of Technology of Ngaoundéré, P.O Box 455, Cameroon

³Laboratory of BioPhysics, University of Yaoundé 1, P.O. Box 812, Yaoundé, Cameroon.

⁴National School of Agro-industrial Sciences of Ngaoundéré, P.O. Box 455, Ngaoundéré, Cameroon.

*E-mail : h.abboubakar@gmail.com

Abstract

The goal of this work is to study a ZIKA disease model using fractional derivatives in the Caputo sense. After formulating the model, we compute the basic reproduction number R_0 and give rigorous proof of the existence of equilibrium points as well as stability analysis of these equilibrium points. Then, we study the existence and the uniqueness of the solutions of the fractional model using the Banach fixed point theory. Numerical simulations are performed to validate our analytical results, as well as to see the impact of varying the fractional parameter on the disease dynamics

Keywords

ZIKA disease; Fractional model; Caputo derivative; Asymptotic stability; Numerical simulations

I INTRODUCTION

Primarily transmitted to humans through the bite of an infected *Aedes* species mosquito (*Ae. aegypti* and *Ae. albopictus*, which are the same mosquitoes that spread dengue and chikungunya viruses), Zika disease is caused by a flavivirus called Zika virus (ZIKV). It is a public health problem in many Latin American countries which are experiencing the most serious known outbreaks of Zika virus (ZIKV), and this, since 2005 [1]. In October 2015, an association between Zika virus infection and microcephaly was reported in Brazil. From then on, epidemics and evidence of transmission quickly appeared in the Americas, Africa, and other parts of the world. To date, a total of 86 countries and territories have reported evidence of mosquito-borne Zika infection. The rise of microcephaly and Guillain-Barré syndrome is correlated with the rise of the spread of the Zika virus. Zika virus was first identified in Uganda in 1947 in monkeys, and later in humans in 1952. In 2007, the Island of Yap was the land of the first large outbreak of the disease [1, 2]. Since then, outbreaks have appeared in many countries (United Republic of Tanzania, India, Indonesia, Malaysia, Pakistan, Pacific islands, Brazil...). On February 1, 2016, Zika disease became a Public Health Emergency of International Concern [1, 2]. Since there is no available vaccine to prevent or treat Zika virus infection, the principal control means are vector controls and protection means against mosquito bites [1, 2].

Mathematical modeling is a good tool for the projection and control of several infectious disease outbreaks. The current Covid-19 outbreak is an example [3, 4]. Several works to model,

project, and control recent outbreaks of Zika virus disease in the world, were published [5, 6, 7, 8]. Fractional derivative theory has been more and more used in the last decades to model natural phenomena like infectious diseases [9, 10]. The fact that classical derivatives do not permit to describe fluctuations of different classes of population involved in infectious process, and thus heterogeneity between compartment classes, do that the use of Fractional derivatives is need nowadays [11]. Note that memory effect, which represents a great advantage of fractional derivatives compared to classical derivatives, permit to predict epidemic peaks for various diseases [12, 13]. Several papers using fractional derivatives were developed for Zika transmission dynamics [14, 15, 16, 17]. Most of them use Caputo's type derivatives. In this paper, we extend the model of Yuan *et al.* in [8] replacing integer derivatives with Caputo derivative. After the computation of the basic reproduction number \mathbf{R}_0 , we prove the uniform asymptotically stability of the Zika-free equilibrium when $\mathbf{R}_0 \leq 1$, despite the fact that the latter coexists with an endemic equilibrium. For the case $\mathbf{R}_0 > 1$, we prove that the model admits two endemic equilibrium points, one of them is stable and the other is unstable. To simulate the fractional model, we construct two numerical schemes using the Adams-Bashforth Method (ABM) and the Laplace Transform Adomian Decomposition Method (LTADM), respectively.

The outline of the paper is as follows: In section II, we recall the definition of the Caputo derivative, and two precious results which is used top prove the existence and uniqueness of solutions. Section III is devoted to model formulation and mathematical results. Numerical simulation results are presented in Section IV. A conclusion round up the paper.

II USEFUL DEFINITION AND RESULTS

In this section, we present useful definitions of Caputo derivative, as well ass useful lemmas which will be used in the model analysis (see [18, 19]).

Definition 1:

Given a function $f : [a, b] \rightarrow \mathbb{R}$ of class C^n , the Caputo fractional derivative of f of order α is defined by :

$${}^C D_t^\alpha f(t) = \frac{1}{\Gamma(n - \alpha)} \int_{t_0}^t (t - \tau)^{n-\alpha-1} f^{(n)}(\tau) d\tau \quad (1)$$

where $\alpha \notin \mathbb{N}$, $n = [\alpha] + 1$.

Proposition 1:

Let $f, g : [a, b] \rightarrow \mathbb{R}$ be such that ${}^C D_t^\alpha f(t)$ and ${}^C D_t^\alpha g(t)$ exist almost everywhere and let $c_1, c_2 \in \mathbb{R}$. Then ${}^C D_t^\alpha (c_1 f(t) + c_2 g(t))$ exists almost everywhere with ${}^C D_t^\alpha (c_1 f(t) + c_2 g(t)) = c_1 \times {}^C D_t^\alpha f(t) + c_2 \times {}^C D_t^\alpha g(t)$.

Lemma 1:

If $0 < \chi < 1$ and $\mathbf{d} \in \mathbb{N}$, then there exist positive constants $W_{\chi,1}$ and $W_{\chi,2}$ only dependent on χ , such that

$$(\mathbf{d} + 1)^\chi - \mathbf{d}^\chi \leq W_{\chi,1} (\mathbf{d} + 1)^{\chi-1} \quad \text{and} \quad (\mathbf{d} + 2)^{\chi+1} - 2(\mathbf{d} + 1)^{\chi+1} + \mathbf{d}^{\chi+1} \leq W_{\chi,2} (\mathbf{d} + 1)^{\chi-1}.$$

Lemma 2:

Assume that $\mathbf{y}_{q,m} = (m - q)^{\chi-1} (q = 1, 2, \dots, m - 1)$ & $\mathbf{y}_{q,m} = 0$ for $q \geq m, \chi, \mathbf{Q}, \mathbf{h}, \mathbf{Y} >$

0, $\mathbf{k} \leq \mathbf{Y}$ & $\mathbf{k} \in \mathbb{N}$. Let $\sum_{q=\mathbf{k}}^{q=i} \mathbf{y}_{q,m} |\mathbf{e}_q| = 0$ for $\mathbf{k} > m \geq 1$. If

$$|\mathbf{e}_m| \leq \mathbf{Q} \mathbf{h}^\chi \sum_{q=1}^{m-1} \mathbf{y}_{q,m} |\mathbf{e}_q| + |\eta_0|, \quad m = 1, 2, \dots, \mathbf{k},$$

then

$$|\mathbf{e}_k| \leq \mathcal{M} |\eta_0|, \quad \mathbf{k} = 1, 2, \dots$$

where $\mathcal{M} \in \mathbb{R}_+$ independent of \mathbf{k} and \mathbf{h} .

III THE FRACTIONAL ZIKA MODEL AND ASYMPTOTIC STABILITY RESULTS

3.1 The fractional Zika model

The basic model comes from the works of Yuan *et al.* (2021) [8] which consists of a system of five Ordinary Differential Equations. The human and mosquito populations are split into three and two compartments according to their clinical status. We denote by: S_h susceptible human, I_h Infected humans, R_h recovered human, S_v susceptible mosquitoes, I_v Infected mosquitoes. Since an infected mosquito is infected throughout its life, we do not consider a removed class in this model.

$$\begin{cases} {}^C D^\eta S_h(t) &= \mu_h(H - S_h) - \left(\beta \frac{I_h}{H} + ab \frac{I_v}{H} \right) S_h, \\ {}^C D^\eta I_h(t) &= -(\gamma + \mu_h) I_h + \left(\beta \frac{I_h}{H} + ab \frac{I_v}{H} \right) S_h, \\ {}^C D^\eta R_h(t) &= -\mu_h R_h + \gamma I_h, \\ {}^C D^\eta S_v(t) &= \mu_v(M - S_v - \theta I_v) - ac \frac{I_h}{H} S_v, \\ {}^C D^\eta I_v(t) &= ac \frac{I_h}{H} S_v - \mu_v(1 - \theta) I_v, \end{cases} \quad (2)$$

with the initial conditions $S_h = S_h^0$, $I_h = I_h^0$, $R_h = R_h^0$, $S_v = S_v^0$, and $I_v = I_v^0$.

In model (2): ${}^C D^\eta$ denotes the Caputo derivative with $\eta \in (0, 1]$; $H := S_h + I_h + R_h$ represents the total human population, μ_h is the human natural mortality rate, β is the transmission coefficient from infected humans to susceptible humans, a is the mosquitoes biting rate, b is the transmission probability from an infectious mosquito to a susceptible human per bite, γ is the recovery rate of humans, $M := S_v + I_v$ is the total mosquito population, μ_v is the mortality rate of mosquitoes, θ is the proportion of congenital infections in the offspring of infected female mosquitoes, and c is the transmission probability from an infectious human to a susceptible mosquito per bite.

Note that, as in the case of mathematical modeling of infectious disease epidemics, we consider that human and mosquito populations are still constant during the Zika epidemic reported by the Guillain-Barré Syndrome (GBS) surveillance data. Using the fact that $S = H - I - R$ and $S_v = M - I_v$, the study of the model (2) is reduced to the study of the following system

$$\begin{cases} {}^C D^\eta I_h(t) &= -(\gamma + \mu_h) I_h + \left(\beta \frac{I_h}{H} + ab \frac{I_v}{H} \right) (H - I_h - R_h), \\ {}^C D^\eta R_h(t) &= -\mu_h R_h + \gamma I_h, \\ {}^C D^\eta I_v(t) &= ac \frac{I_h}{H} (M - I_v) - \mu_v(1 - \theta) I_v, \end{cases} \quad (3)$$

which define a dynamical system in the following bounded and absorbed set

$$\Sigma = \{(I_h, R_h, I_v) \in \mathbb{R}_+^3 : I_h + R_h \leq H, I_v \leq M\}.$$

3.2 The basic reproduction number and uniform asymptotic stability results

3.2.1 The basic reproduction number and stability of the Zika-free equilibrium

System (3) always has a Zika-free equilibrium $\mathcal{E}_0 = (0, 0, 0)$. Setting $\mathbf{R}_1 = \frac{\beta}{(\gamma + \mu_h)}$, $\mathbf{R}_2 = \frac{ab}{(1 - \theta)\mu_v}$, and $\mathbf{R}_3 = \frac{acM}{H(\gamma + \mu_h)}$, we can express using the next generation method [20], the basic reproduction number of system (3) as follows:

$$\mathbf{R}_0 = \frac{\sqrt{4\mathbf{R}_2\mathbf{R}_3 + \mathbf{R}_1^2} + \mathbf{R}_1}{2}. \quad (4)$$

The matrix of the system (3) evaluated at the disease-free equilibrium point \mathcal{E}_0 is given by

$$\mathcal{J}(\mathcal{E}_0) = \begin{bmatrix} -(\gamma + \mu_h) + \beta & 0 & ab \\ \gamma & -\mu_h & 0 \\ ac\frac{M}{H} & 0 & -\mu_v(1 - \theta) \end{bmatrix}. \quad (5)$$

The characteristic equation of $\mathcal{J}(\mathcal{E}_0)$ is therefore $\det(s[I - (1 - \eta)\mathcal{J}(\mathcal{E}_0)] - \eta\mathcal{J}(\mathcal{E}_0)) = 0$ [21].

Theorem 1:

The Zika-free equilibrium point \mathcal{E}_0 of the model (3) is locally asymptotically stable if $\mathcal{R}_0 < 1$ and unstable otherwise.

Proof. Setting $M/H = \alpha$, $1 - \theta = \theta_1$, $1 - \eta = \rho$, $\mathcal{M} = s[I - \rho\mathcal{J}(\mathcal{E}_0)] - \eta\mathcal{J}(\mathcal{E}_0)$. The characteristic equation $\det(\mathcal{M}) = 0$ is equivalent to $[(\mu_h\rho + 1)s + \mu_h\eta]\Phi(s) = 0$, where $\Phi(s) := e_2s^2 + e_1s + e_0$ with $e_2 = ((1 - \mathbf{R}_0)(1 - \mathbf{R}_1 + \mathbf{R}_0)k_1\theta_1\rho^2 + \theta_1\rho)\mu_v + (1 - \mathbf{R}_1)k_1\rho + 1$, $e_1 = (2(1 - \mathbf{R}_0)(1 - \mathbf{R}_1 + \mathbf{R}_0)k_1\theta_1\eta\rho + \theta_1\eta)\mu_v + (1 - \mathbf{R}_1)k_1\eta$, and $e_0 := (1 - \mathbf{R}_0)(1 - \mathbf{R}_1 + \mathbf{R}_0)k_1\theta_1\eta^2\mu_v$.

Note that $s_1 = \frac{-\eta\mu_h}{1 + \rho\mu_h} < 0$ is a solution of $\det(\mathcal{M}) = 0$. Since $\mathbf{R}_0 < 1 \implies \mathbf{R}_1 < 1$, it follows that all coefficients e_0 , e_1 , and e_2 of $\Phi(s)$ is always positive. Thus, we conclude that all solutions of the equation $\det(\mathcal{M}) = 0$ have negative real part, which means that the Zika-free equilibrium is locally asymptotically stable. This ends the proof. \square

Theorem 2:

The Zika-free equilibrium point \mathcal{E}_0 of the model (3) is uniformly asymptotically stable if $\mathcal{R}_0 < 1$ and unstable otherwise.

Proof. Let us consider the following Lyapunov function $\mathbf{L} = \frac{\mu_v(1-\theta)}{ab}I_h + I_v$. The derivative of \mathbf{L} along the trajectories of (3), using Proposition 1, gives:

$$\begin{aligned}
D^n \mathbf{L} &= \frac{\mu_v(1-\theta)}{ab} D^n I_h + D^n I_v \\
&= -\frac{\mu_v(1-\theta)}{ab} \left(\beta \frac{I_h^2}{H} + \beta \frac{I_h R_h}{H} + ab \frac{I_h I_v}{H} + ab \frac{R_h I_v}{H} \right) + ac \frac{M}{H} I_h - \xi_2 \mu_v (1-\theta) I_v \\
&\quad - ac \frac{I_h I_v}{H} + \frac{\mu_v(1-\theta)}{ab} (\beta - (\gamma + \mu_h)) I_h + \frac{\mu_v(1-\theta)}{ab} ab I_v \\
&\leq \left[\xi_1 (\beta - (\gamma + \mu_h)) + \xi_2 ac \frac{M}{H} \right] I_h + [\xi_1 ab - \xi_2 \mu_v (1-\theta)] I_v \\
&= \xi_2 \frac{(\gamma + \mu_h) \mu (1-\theta)}{ab} (\mathbf{R}_0 - 1) (\mathbf{R}_0 + 1 - \mathbf{R}_1) I_h
\end{aligned}$$

Finally, $D^n \mathbf{L} \leq 0$ if and only if $\mathbf{R}_0 \leq 1$. Thus, from [22], it follows that the Zika-free equilibrium \mathcal{E}_0 of the fractional model (2) is uniformly asymptotically stable whenever $\mathbf{R}_0 \leq 1$. \square

3.2.2 The endemic equilibrium and its uniform stability

Equating the right-hand side of system (3) by zero, and solving the obtained system in term of I_v , we have

$$I_h = \frac{\mu_v \theta_1 H I_v}{ac(H\alpha - I_v)}, R_h = \frac{\gamma}{\mu_h} I_h, \quad (6)$$

where I_v is the solutions of the following equation $I_v(\sigma_2 I_v^2 + \sigma_1 I_v + \sigma_0) = 0$, with $\sigma_2 = a^3 b^2 c (\theta_1 \mu_v k_1 + ac \mu_h)$, $\sigma_0 = R_0 (R_0 - 1) (R_1 - R_0 - 1) (R_1 - R_0) k_1^2 \theta_1^2 H^2 \mu_v^2 \mu_h$, $\sigma_1 = k_1 \theta_1 H ab \mu_v \{ \theta_1 \mu_v k_1 ((R_0 - 1) R_1 - R_0^2) + ac \mu_h [2R_0 R_1 - 2R_0^2 - R_1 + 1] \}$.

Note that σ_2 is always positive. Since $\mathbf{R}_0 - \mathbf{R}_1 = \frac{\mathbf{R}_2 \mathbf{R}_3}{\mathbf{R}_0} > 0 \implies \mathbf{R}_1 - \mathbf{R}_0 - 1 = \frac{\mathbf{R}_2 \mathbf{R}_3}{\mathbf{R}_0} - 1 = \frac{\mathbf{R}_2 \mathbf{R}_3 - \mathbf{R}_0}{\mathbf{R}_0} < 0$, it follows that the sign of σ_0 depends of the value of \mathbf{R}_0 . Indeed, $\mathbf{R}_0 > 1 \iff \sigma_0 > 0$ and $\mathbf{R}_0 < 1 \iff \sigma_0 < 0$. Also, we have $\mathbf{R}_0 > 1 \implies \sigma_1 < 0$.

Since $\mathbf{R}_0 > \mathbf{R}_1$, it follows that $\Delta > 0$. We resume the above analysis as follows:

Theorem 3:

The number of equilibrium points os system (3) depend of the basic reproduction number as follows:

1. If $\mathbf{R}_0 \leq 1$, then system (3) has two equilibrium points: the Zika-free equilibrium $\mathcal{E}_0 =$

$$(0, 0, 0) \text{ and an endemic equilibrium point } \mathcal{E}^\dagger = (I_h^\dagger, R_h^\dagger, I_v^\dagger) = \left(\frac{\mu_v \theta_1 H I_v^\dagger}{ac(H\alpha - I_v^\dagger)}, \frac{\gamma}{\mu_h} \frac{\mu_v \theta_1 H I_v^\dagger}{ac(H\alpha - I_v^\dagger)}, I_v^\dagger \right),$$

$$\text{where } I_v^\dagger = \frac{-\sigma_1 + \sqrt{\sigma_1^2 - 4\sigma_2\sigma_0}}{2},$$

2. If $\mathbf{R}_0 > 1$, then system (3) has three equilibrium points: the Zika-free equilibrium $\mathcal{E}_0 = (0, 0, 0)$ and two endemic equilibrium points $\mathcal{E}^\pm = (I_h^\pm, R_h^\pm, I_v^\pm)$, with $I_v^\pm = \frac{-\sigma_1 \pm \sqrt{\sigma_1^2 - 4\sigma_2\sigma_0}}{2}$.

The above first case ($\mathbf{R}_0 \leq 1$) means that there is a possibility that the Zika-free equilibrium co-exists with an endemic equilibrium $\mathcal{E}^\dagger = (I_h^\dagger, R_h^\dagger, I_v^\dagger)$ whenever the $\mathbf{R}_0 \leq 1$. Nevertheless, by Theorem 2, we conclude that this endemic equilibrium is always unstable whenever $\mathbf{R}_0 \leq 1$. Now, suppose that $\mathbf{R}_0 > 1$, which means that the Zika-free equilibrium $\mathcal{E}_0 = (0, 0, 0)$ becomes unstable, system (3) admits two endemic equilibrium points $\mathcal{E}^\pm = (I_h^\pm, R_h^\pm, I_v^\pm)$ and $\mathcal{E}^\ddagger = (I_h^\ddagger, R_h^\ddagger, I_v^\ddagger)$. In what follows, we will show that one of these endemic equilibrium points is stable which implies that the other is unstable. To this aim, we follow Yuan *et al.* [8] by defining the following Lyapunov function for system (3)

$$\mathcal{L} = \frac{S_h^*}{H} \left(\frac{S_h}{S_h^*} - 1 - \ln \frac{S_h}{S_h^*} \right) + \frac{I_h^*}{H} \left(\frac{I_h}{I_h^*} - 1 - \ln \frac{I_h}{I_h^*} \right) + \frac{abS_h^*I_v^*}{\mu_v\theta_1} \left(\frac{I_v}{I_v^*} - 1 - \ln \frac{I_v}{I_v^*} \right),$$

where (S_h^*, I_h^*, I_v^*) represents one of the two endemic equilibrium points \mathcal{E}^\pm .

The equilibrium equations of system (2) lead to $\mu_h(H - S_h^*) = \left(\beta \frac{I_h^*}{H} + ab \frac{I_v^*}{H} \right) S_h^*$, $(\gamma + \mu_h)I_h^* = \left(\beta \frac{I_h^*}{H} + ab \frac{I_v^*}{H} \right) S_h^*$, $ac \frac{I_h^*}{H} S_v^* = \mu_v(1 - \theta)I_v^*$. By using these equalities in the derivative of \mathcal{L} along of trajectories of (2) leads to

$$\begin{aligned} \mathcal{L}' \leq & -\frac{\mu_h}{S_h} (S_h - S_h^*)^2 - \frac{a^2bcS_h^*I_h}{\mu_v\theta_1 H^2 I_v} (I_v - I_v^*)^2 + \frac{abS_h^*I_v^*}{H} \left(3 - \frac{S_h^*}{S_h} - \frac{S_h I_h^* I_v}{S_h^* I_h I_v^*} - \frac{I_h I_v^*}{I_h^* I_v} \right) \\ & + \beta \frac{S_h^* I_h^*}{H^2} \left(2 - \frac{S_h}{S_h^*} - \frac{S_h^*}{S_h} \right), \end{aligned}$$

which, thanks to the inequality of arithmetic and geometric means, is negative. Also, $\mathcal{L}' = 0 \implies (S_h, I_h, I_v) = (S_h^*, I_h^*, I_v^*)$. Using [22], we conclude that one of the both endemic equilibrium points is globally stable. We therefore affirm the following result:

Theorem 4:

The Zika model (2) admits a unique endemic equilibrium in Σ which is globally asymptotically stable whenever $\mathbf{R}_0 > 1$.

3.3 Existence and uniqueness of solutions

The objective of this part is to show by using the theory of the fixed points the existence and the uniqueness of the solutions of the Zika fractional model (in the Caputo sense) with the initial conditions. In this part, \mathbf{B} is the Banach space for continuous functions with real values over an interval \mathbf{I} with the associated norms

$$\| (S_h, I_h, R_h, S_v, I_v) \| = \| S_h \| + \| I_h \| + \| R_h \| + \| S_v \| + \| I_v \|,$$

where $\| S_h \| = \sup \{ | S_h(t) | : t \in \mathbf{I} \}$, $\| I_h \| = \sup \{ | I_h(t) | : t \in \mathbf{I} \}$, $\| R_h \| = \sup \{ | R_h(t) | : t \in \mathbf{I} \}$, $\| S_v \| = \sup \{ | S_v(t) | : t \in \mathbf{I} \}$, $\| I_v \| = \sup \{ | I_v(t) | : t \in \mathbf{I} \}$ and $\mathbf{B} = \varepsilon(\mathbf{I}) \times \varepsilon(\mathbf{I}) \times \varepsilon(\mathbf{I}) \times \varepsilon(\mathbf{I}) \times \varepsilon(\mathbf{I})$, with $\varepsilon(\mathbf{I})$ the Banach space for continuous functions with real values on \mathbf{I} and the above associated norms. The fractional system (2) can be written in the form:

$$\begin{aligned} {}^C D_t^\eta S_h(t) &= \mathbf{G}_1(t, S_h), {}^C D_t^\eta I_h(t) = \mathbf{G}_2(t, I_h), {}^C D_t^\eta R_h(t) = \mathbf{G}_3(t, R_h), \\ {}^C D_t^\eta S_v(t) &= \mathbf{G}_4(t, S_v), {}^C D_t^\eta I_v(t) = \mathbf{G}_5(t, I_v), \end{aligned} \quad (7)$$

where $G_i(t, x)$, $i \in [0; 5] \cap \mathbb{N}$, $x \in \{S_h, I_h, R_h, S_v, I_v\}$, represent the right-hand sides of model (2).

By applying the fractional integral operator in equation (7), we obtain:

$$\begin{aligned} S_h(t) - S_h(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} G_1(\chi, S_h) d\chi, I_h(t) - I_h(0) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} G_2(\chi, I_h) d\chi, \\ R_h(t) - R_h(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} G_3(\chi, R_h) d\chi, S_v(t) - S_v(0) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} G_4(\chi, S_v) d\chi, \\ I_v(t) - I_v(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} G_5(\chi, I_v) d\chi, \end{aligned} \quad (8)$$

The kernels \mathbf{G}_i , $i = 1, 2, 3, 4, 5$, satisfy the Lipschitz and contraction conditions if the following inequality holds:

$$0 \leq V_i < 1, \quad i = 1, 2, 3, 4, 5, \quad (9)$$

where $V_1 = \left(\mu_h + \beta + ab \frac{\nu}{H} \right)$, $V_2 = (\mu_h + \gamma) + \beta$, $V_3 = \mu_h$, $V_4 = \mu_v + ac$, $V_5 = \mu_v(1 - \theta)$. We thus claim the following result

Theorem 5:

The kernel \mathbf{G}_1 satisfies the Lipschitz condition if the following inequality is respected

$$0 \leq V_1 := \left(\mu_h + \beta + ab \frac{\nu}{H} \right) \leq 1 \quad \text{with} \quad \| I_v \| = \nu. \quad (10)$$

Proof. Consider the kernel \mathbf{G}_1 , and S_h, S_{1h} any two functions. We have

$$\| \mathbf{G}_1(t, S_h) - \mathbf{G}_1(t, S_{1h}) \| = \| -\mu_h (S_h - S_{1h}) - \left(\beta \frac{I_h}{H} + ab \frac{I_v}{H} \right) (S_h - S_{1h}) \|. \quad (11)$$

Using the triangular inequality for the norms of (11) above, we get:

$$\begin{aligned} \| \mathbf{G}_1(t, S_h) - \mathbf{G}_1(t, S_{1h}) \| &\leq \| \mu_h (S_h - S_{1h}) \| + \left\| \left(\beta \frac{I_h}{H} + ab \frac{I_v}{H} \right) (S_h - S_{1h}) \right\| \\ &\leq \left(\mu_h + \beta \frac{I_h}{H} + ab \frac{I_v}{H} \right) \| (S_h(t) - S_{1h}(t)) \|, \\ &\leq \left(\mu_h + \beta + ab \frac{\| I_v \|}{H} \right) \| (S_h(t) - S_{1h}(t)) \|, \\ &= \overbrace{\left(\mu_h + \beta + ab \frac{\nu}{H} \right)}^{V_1} \| S_h(t) - S_{1h}(t) \|. \end{aligned} \quad (12)$$

Therefore, the Lipschitz condition is fulfilled for \mathbf{G}_1 , and if in addition, $0 \leq V_1 < 1$, then it is also a contraction. We get similar results for the other kernels:

$$\begin{aligned} \| \mathbf{G}_2(t, I_h) - \mathbf{G}_2(t, I_{1h}) \| &\leq V_2 \| I_h(t) - I_{1h}(t) \| \\ \| \mathbf{G}_3(t, R_h) - \mathbf{G}_3(t, R_{1h}) \| &\leq V_3 \| R_h(t) - R_{1h}(t) \| \\ \| \mathbf{G}_4(t, S_v) - \mathbf{G}_4(t, S_{1v}) \| &\leq V_4 \| S_v(t) - S_{1v}(t) \| \\ \| \mathbf{G}_5(t, I_v) - \mathbf{G}_5(t, I_{1v}) \| &\leq V_5 \| I_v(t) - I_{1v}(t) \|. \end{aligned} \quad (13)$$

□

The expressions of the equation (8) can be written recursively as:

$$\begin{aligned}
S_{hn}(t) - S_h(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} \mathbf{G}_1(\chi, S_{h(n-1)}) d\chi, \\
E_{hn}(t) - E_h(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} \mathbf{G}_2(\chi, E_{h(n-1)}) d\chi, \\
I_{hn}(t) - I_h(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} \mathbf{G}_3(\chi, I_{h(n-1)}) d\chi, \\
S_{vn}(t) - S_v(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} \mathbf{G}_4(\chi, S_{v(n-1)}) d\chi, \\
I_{vn}(t) - I_v(0) &= \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} \mathbf{G}_5(\chi, I_{v(n-1)}) d\chi,
\end{aligned} \tag{14}$$

The difference between successive terms for recursive formulas is written:

$$\begin{aligned}
\phi_{1n}(t) &= S_{hn}(t) - S_{h(n-1)}(t) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_1(\chi, S_{h(n-1)}) - \mathbf{G}_1(\chi, S_{h(n-2)})) d\chi, \\
\phi_{2n}(t) &= I_{hn}(t) - I_{h(n-1)}(t) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_2(\chi, I_{h(n-1)}) - \mathbf{G}_2(\chi, I_{h(n-2)})) d\chi, \\
\phi_{3n}(t) &= R_{hn}(t) - R_{h(n-1)}(t) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_3(\chi, R_{h(n-1)}) - \mathbf{G}_3(\chi, R_{h(n-2)})) d\chi, \\
\phi_{4n}(t) &= S_{vn}(t) - S_{v(n-1)}(t) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_4(\chi, S_{v(n-1)}) - \mathbf{G}_4(\chi, S_{v(n-2)})) d\chi, \\
\phi_{5n}(t) &= I_{vn}(t) - I_{v(n-1)}(t) = \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_5(\chi, I_{v(n-1)}) - \mathbf{G}_5(\chi, I_{v(n-2)})) d\chi,
\end{aligned} \tag{15}$$

with initial conditions:

$$S^{h0}(t) = S_h(0), \quad I^{h0}(t) = I_h(0), \quad R^{h0}(t) = R_h(0), \quad S^{v0}(t) = S_v(0), \quad I^{v0}(t) = I_v(0). \tag{16}$$

Let us take the norm of the first expression of the equation (15)

$$\begin{aligned}
\| \phi_{1n}(t) \| &= \| S_{hn}(t) - S_{h(n-1)}(t) \| \\
&= \left\| \frac{1}{\Gamma(\eta)} \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_1(\chi, S_{h(n-1)}) - \mathbf{G}_1(\chi, S_{h(n-2)})) d\chi \right\|, \\
&\leq \frac{1}{\Gamma(\eta)} \left\| \int_0^t (t - \chi)^{\eta-1} (\mathbf{G}_1(\chi, S_{h(n-1)}) - \mathbf{G}_1(\chi, S_{h(n-2)})) d\chi \right\|.
\end{aligned} \tag{17}$$

Using the Lipschitz condition we have:

$$\| S_{hn}(t) - S_{h(n-1)}(t) \| \leq \frac{1}{\Gamma(\eta)} V_1 \int_0^t (t - \chi)^{\eta-1} \| S_{h(n-1)} - S_{h(n-2)} \| d\chi. \tag{18}$$

Thus, we obtain

$$\| \phi_{1n}(t) \| \leq \frac{1}{\Gamma(\eta)} V_1 \int_0^t (t - \chi)^{\eta-1} \| \phi_{1(n-1)}(\chi) \| d\chi. \tag{19}$$

Similar reasoning makes it possible to have:

$$\begin{aligned}
\| \phi_{2n}(t) \| &\leq \frac{1}{\Gamma(\eta)} V_2 \int_0^t (t - \chi)^{\eta-1} \| \phi_{2(n-1)}(\chi) \| d\chi, \\
\| \phi_{3n}(t) \| &\leq \frac{1}{\Gamma(\eta)} V_3 \int_0^t (t - \chi)^{\eta-1} \| \phi_{3(n-1)}(\chi) \| d\chi, \\
\| \phi_{4n}(t) \| &\leq \frac{1}{\Gamma(\eta)} V_4 \int_0^t (t - \chi)^{\eta-1} \| \phi_{4(n-1)}(\chi) \| d\chi, \\
\| \phi_{5n}(t) \| &\leq \frac{1}{\Gamma(\eta)} V_5 \int_0^t (t - \chi)^{\eta-1} \| \phi_{5(n-1)}(\chi) \| d\chi.
\end{aligned} \tag{20}$$

Note that $S_{hn}(t) = \sum_{i=1}^n \phi_{1i}(t)$, $E_{hn}(t) = \sum_{i=1}^n \phi_{2i}(t)$, $I_{hn}(t) = \sum_{i=1}^n \phi_{3i}(t)$, $S_{vn}(t) = \sum_{i=1}^n \phi_{4i}(t)$, $I_{vn}(t) = \sum_{i=1}^n \phi_{5i}(t)$.

Now let's show the uniqueness of the model solution. We claim the following

Theorem 6:

The fractional model (2) admits a unique solution on $t \in [0, T]$ if the following inequality is satisfied

$$\frac{1}{\Gamma(\eta)} V_i b^\eta \leq 1, \quad i = 1, 2, \dots, 5. \tag{21}$$

Proof. The proof follows the proof of [18]. □

IV NUMERICAL SCHEME AND RESULTS

4.1 Numerical scheme

In what follows, we will use the Adams-Bashforth-Moulton method (see [15, 23]) to construct a numerical scheme for the fractional model (3). To this aim, let us consider the uniform discretization of $[0, a]$ given by $t_m = m\aleph$, $m \in [0; N] \cap \mathbb{N}$ where $\aleph = a/m$ denotes the step size. For a given approximation $\mathcal{U}_\aleph(t_i) \approx \mathcal{U}(t_i)$, the next approximation $\mathcal{U}_\aleph(t_{i+1})$ is obtain (using the predictor-corrector method) as follows:

$$\text{Predictor: } \mathcal{U}_\aleph^p(t_{m+1}) = \sum_{l=0}^{\lceil \varphi \rceil - 1} \frac{t_{n+1}^l}{l!} \mathcal{U}_0^l + \frac{1}{\Gamma(\eta)} \sum_{l=0}^m b_{l,m+1} \mathbf{G}(t_l, \mathcal{U}_\aleph(t_l));$$

$$\text{Corrector: } \mathcal{U}_\aleph(t_{m+1}) = \sum_{l=0}^{\lceil \varphi \rceil - 1} \frac{t_{n+1}^l}{l!} \mathcal{U}_0^l + \frac{\aleph^\eta}{\Gamma(2 + \eta)} \mathbf{G}(t_{l+1}, \mathcal{U}_\aleph(t_{l+1})) + \frac{\aleph^\eta}{\Gamma(2 + \eta)} \sum_{l=0}^m c_{l,m+1} \mathbf{G}(t_l, \mathcal{U}_\aleph(t_l));$$

with

$$c_{l,m+1} = \begin{cases} m^{1+\varphi} - (m - \varphi)(m + 1)^\varphi, & \text{if } l = 0, \\ (m - l + 2)^{1+\varphi} + (m - l)^{1+\varphi} - 2(m - l + 1)^{1+\varphi}, & \text{if } 1 \leq l \leq m, \\ 1 & \text{if } l = m + 1, \end{cases}$$

$$\text{and } b_{l,m+1} = \frac{\aleph^\varphi}{\varphi} [(m - l + 1)^\varphi - (m - l)^\varphi].$$

4.2 Numerical results

For numerical simulation, we consider the same parameter values which came from parameter estimation of the model (3) using the Guillain-Barré Syndrome (GBS) surveillance data (see [8]). In [8], Yuan *et al.* obtained after model calibration $a = 0.45$, $b = 0.3$, $c = 0.4$, $\theta = 0.0034$, $\gamma = 0.14$, $\mu_h = 4 \times 10^{-5}$, $\mu_v = 0.125$, $\beta = 0.05$, $\rho = 0.0024\%$, $H = 10^5$, and $M = 1.5H = 15 \times 10^4$. With these parameter values, $\mathbf{R}_0 = 1.6349$, which means that we are in endemic state. Figure 1&2 illustrate the results of Theorems 1 and 4, respectively. It is




Figure 1: Time-series of I_h and I_v with $\mathcal{R}_0 = 0.92$ (we have set $a = 0.225$) for different values of the fractional order $\eta \in \{1; 0.9; 0.8; 0.7; 0.6\}$. Initial conditions are $(I_h(0), R_h(0), I_v(0)) = (40, 60000, 100)$.




Figure 2: Time-series of I_h and I_v with $\mathcal{R}_0 = 1.63$ for different values of the fractional order $\eta \in \{1; 0.9; 0.8; 0.7; 0.6\}$. Initial conditions are $(I_h(0), R_h(0), I_v(0)) = (40, 60000, 100)$.

clear that, when $\mathbf{R}_0 < 1$, the solution approaches the Zika-free equilibrium for $\eta = 1$, and the needed time is forward delayed as the fractional order η decreases (Fig. 1), and when $\mathbf{R}_0 > 1$, the solution approaches the endemic equilibrium (Fig. 2).

Figures 1&2 also illustrates the impact of varying fractional order η on the Zika dynamics. Indeed, we see that oscillations decrease when the fractional order decreases.

V CONCLUSION

In this work, we studied a transmission model of Zika disease with fractional derivatives in the Caputo sense. After the formulation of the model which consists to replace integer derivative

with fractional derivative in the Caputo sense, we computed the basic reproduction number R_0 and gave rigorous proof of the existence of equilibrium points of both models as well as stability analysis of these equilibrium points. Then, we studied the existence and the uniqueness of the solutions of the fractional model using the Banach fixed point theory. We finally performed numerical simulations to validate our analytical results, as well as to see the impact of varying the fractional parameter η on the disease dynamics. Indeed, when the fractional order decreases, the oscillation amplitude also decreases.

REFERENCES

- [1] Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Vector-Borne Diseases (DVBD). Zika transmission. <https://www.cdc.gov/zika/prevention/transmission-methods.html>, Accessed November 12, 2021.
- [2] World Health Organization (WHO). Zika virus disease. https://www.who.int/health-topics/zika-virus-disease#tab=tab_1, Accessed November 18, 2021.
- [3] Seraphin Djaoue, Gabriel Guilsou Kolaye, Hamadjam Abboubakar, Ado Adamou Abba Ari, and Irepran Damakoa. Mathematical modeling, analysis and numerical simulation of the covid-19 transmission with mitigation of control strategies used in cameroon. *Chaos, Solitons & Fractals*, 139:110281, 2020.
- [4] Pushendra Kumar, Vedat Suat Erturk, Hamadjam Abboubakar, and Kottakkaran Sooppy Nisar. Prediction studies of the epidemic peak of coronavirus disease in brazil via new generalised caputo type fractional derivatives. *Alexandria Engineering Journal*, 60(3):3189–3204, 2021.
- [5] Yongli Cai, Zuqin Ding, Bin Yang, Zhihang Peng, and Weiming Wang. Transmission dynamics of zika virus with spatial structure—a case study in rio de janeiro, brazil. *Physica A: Statistical Mechanics and its Applications*, 514:729–740, 2019.
- [6] Parinya Suparit, Anuwat Wiratsudakul, and Charin Modchang. A mathematical model for zika virus transmission dynamics with a time-dependent mosquito biting rate. *Theoretical Biology and Medical Modelling*, 15(1):1–11, 2018.
- [7] YA Terefe, H Gaff, Morgan Kamga, and Luther van der Mescht. Mathematics of a model for zika transmission dynamics. *Theory in Biosciences*, 137(2):209–218, 2018.
- [8] Xiaoyan Yuan, Yijun Lou, Daihai He, Jinliang Wang, and Daozhou Gao. A zika endemic model for the contribution of multiple transmission routes. *Bulletin of Mathematical Biology*, 83(11):1–28, 2021.
- [9] Pushendra Kumar, Norodin A Rangaig, Hamadjam Abboubakar, Anoop Kumar, and M. Manickam. Prediction studies of the epidemic peak of coronavirus disease in japan: from caputo derivatives to atangana-baleanu derivatives. *International Journal of Modeling, Simulation, and Scientific Computing*, 2021.
- [10] Abdon Atangana. Fractional discretization: the african’s tortoise walk. *Chaos, Solitons & Fractals*, 130:109399, 2020.

- [11] André Schmidt and Lothar Gaul. On the numerical evaluation of fractional derivatives in multi-degree-of-freedom systems. *Signal Processing*, 86(10):2592–2601, 2006.
- [12] Ebraheem Alzahrani, M.M. El-Dessoky, and Dumitru Baleanu. Mathematical modeling and analysis of the novel coronavirus using atangana-baleanu derivative. *Results in Physics*, 25:104240, 2021.
- [13] Stefano Boccaletti, William Ditto, Gabriel Mindlin, and Abdon Atangana. Modeling and forecasting of epidemic spreading: The case of covid-19 and beyond. *Chaos, Solitons, and Fractals*, 135:109794, 2020.
- [14] Aatif Ali, Saeed Islam, M Riaz Khan, Saim Rasheed, FM Allehiyani, Jamel Baili, Muhammad Altaf Khan, and Hijaz Ahmad. Dynamics of a fractional order zika virus model with mutant. *Alexandria Engineering Journal*, 2021.
- [15] Muhammad Altaf Khan, Saif Ullah, and Muhammad Farhan. The dynamics of zika virus with caputo fractional derivative. *AIMS Math*, 4(1):134–146, 2019.
- [16] Faiz Muhammad Khan, Amjad Ali, Zia Ullah Khan, MR Alharthi, and Abdel-Haleem Abdel-Aty. Qualitative and quantitative study of zika virus epidemic model under caputo’s fractional differential operator. *Physica Scripta*, 96(12):124030, 2021.
- [17] R Rakkiyappan, V Preethi Latha, and Fathalla A Rihan. A fractional-order model for zika virus infection with multiple delays. *Complexity*, 2019, 2019.
- [18] Hamadjam Abboubakar, Lausaire Kemayou Kombou, Adamou Dang Koko, Henri Paul Ekobena Fouda, and Anoop Kumar. Projections and fractional dynamics of the typhoid fever: A case study of mbandjock in the centre region of cameroon. *Chaos, Solitons & Fractals*, 150:111129, 2021.
- [19] Changpin Li and Fanhai Zeng. The finite difference methods for fractional ordinary differential equations. *Numerical Functional Analysis and Optimization*, 34(2):149–179, 2013.
- [20] Pauline Van den Driessche and James Watmough. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical biosciences*, 180(1-2):29–48, 2002.
- [21] Hong Li, Jun Cheng, Hou-Biao Li, and Shou-Ming Zhong. Stability analysis of a fractional-order linear system described by the caputo–fabrizio derivative. *Mathematics*, 7(2):200, 2019.
- [22] Weronika Wojtak, Cristiana J Silva, and Delfim FM Torres. Uniform asymptotic stability of a fractional tuberculosis model. *Mathematical Modelling of Natural Phenomena*, 13(1):9, 2018.
- [23] Kai Diethelm. An algorithm for the numerical solution of differential equations of fractional order. *Electron. Trans. Numer. Anal*, 5(1):1–6, 1997.

A ACKNOWLEDGEMENTS

The first author thanks the Zukunftscolleg of the University of Konstanz for their hospitality during his research stay which permits him to improve this work. The authors express their gratitude to the anonymous reviewers for their comments and suggestions which helped to improve the manuscript.