

Shipping and environmental pressure: Ship-whale interactions off the Guadeloupe archipelago

Jean-Luc Jung, Iwan Le Berre, Bénédicte Madon, Laurent Bouveret, Benjamin de Montgolfier, Damien Le Guyader, Eric Foulquier, Pascal Jean Lopez

▶ To cite this version:

Jean-Luc Jung, Iwan Le Berre, Bénédicte Madon, Laurent Bouveret, Benjamin de Montgolfier, et al.. Shipping and environmental pressure: Ship-whale interactions off the Guadeloupe archipelago. International Symposium of LabEx DRIIHM 2022, Jun 2022, Nantes, France. , 2022. hal-03704048

HAL Id: hal-03704048 https://hal.science/hal-03704048v1

Submitted on 24 Jun2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Shipping and environmental pressure: Ship-whale interactions off the Guadeloupe archipelago

J-L Jung^{1*}, I. Le Berre^{2*}, B. Madon³, L. Bouveret⁴, B. de Montgolfier⁵, D Le Guyader⁶, E. Foulquier², P.J. Lopez⁷

R6554 CNRS, IUEM UBO, Plouzané; ³AMARE, LIENSs UMR7266 CNRS, Université de La Rochelle; ⁴OMMAG, Port-Louis, Guadeloupe, France; ³Aquasearch & ISMER-Université du Québec à Rimouski, Sainte-Luce, Martinique; ⁶GEO4SEAS consulting agency, Brest; ⁷BOREA, UMR8067 CNRS-MNHN, Paris.

60° W

AGOA Sanctuary

The Guadeloupe archipelago is home to a rich and precious marine biodiversity that concerns all levels of food webs, including marine mammals. More than 20 species of cetaceans have been listed in the archipelago, which is part of the Agoa sanctuary, a marine protected area in the French Caribbean waters dedicated to marine mammals. Some species are well known, abundant throughout the year (bottlenose dolphins, sperm whales) or only during certain periods determined by their migratory routes, such as humpback whales. Other species are rarer, difficult to observe and data are lacking to estimate their status in the archipelago.

Caribbean Se

Maritime traffic is an activity growing locally in the context of the expansion of the Grand Port Maritime de Guadeloupe. Its potential and real impacts on marine ecosystems are particularly difficult to decipher. For cetaceans, considered here both as emblematic and sentinel species of the quality of the natural environment, the impacts can be direct (collisions, disturbances) or indirect (noise or chemical pollution).


4

0

We studied the interactions between cetaceans and maritime traffic in the Guadeloupe archipelago, using complementary approaches.


1 - The marine traffic was mapped in detail using AIS data.

400 million ship positions registered during the whole year 2019 were analysed



3 - The presence of cetaceans was studied over the long term through citizen science monitoring.


4704 observations of 21 species were gathered in an open access database named Kakila* *Kakila means "who is there?" in Guadeloupean creole



These comparative analyses highlight potentially specify negative and interactions between certain cetacean species and marine traffic in several areas of the Guadeloupe archipelago. e.g. the most plains-dependent species (such as Fraser's dolphins and short-finned pilot whales) have to face the heavy traffic between Pointe-à-Pitre and Marie Galante. Speed pressure impacts are high at the surface of sperm and beaked whale preferred habitats, highlighting an increased risk of direct collision for these species. Humpback whales and pantropical spotted dolphins face the greatest risk of disturbance from high average vessel traffic occupancy in the shallow continental shelf waters.


Shipping trajectories density around Guadeloupe and the Agoa sanctuary en 2019

2 - Cumulative pressure of the marine traffic was estimated in terms of intensity, occupancy and speed



Example of heat maps constructed from the *Kakila* database, showing sightings of a cetacean species, the sperm whale (P. macrocephalus).

4 - Cumulative pressure of the marine traffic and cetacean habitat preferences were compared to underwater topography



Such promising results encourage us to extend the study. In particular:

- by improving the cetacean presence data (including by using environmental DNAbased approaches),
- by integrating hydrological variables to the habitat characterization,

The shipping pressure expressed by the cumulative score of the intensity, occupancy and speed indices

> Underwater topography of the study area and marine ecological habitat features for cetaceans

by extending the AIS data analysis over several years to study the chronological variations (monthly and interannual). The identification of marine mammal conservation hotspots in Guadeloupe, as well as mitigation proposals, are the goals of our study.

Further readings

Coché L. et al. (2021) Kakila database: Towards a FAIR community approved database of cetacean presence in the waters of the Guadeloupe archipelago based on citizen science. Biodiversity Data Journal. https://doi.org/10.3897/BDJ.9.e69022 Madon B. et al. (2022) Pairing AIS data and underwater topography to assess maritime traffic pressures on cetaceans: Case study in the Guadeloupean waters of the Agoa Sanctuary. Marine Policy. In press





0.5

1.5

International Symposium of Labex DRIIHM - 2022 Inter-Disciplinary Research Facility on Human-Environment Interactions - ANR-11-LABX-0010 **20-22 june 2022 - Nantes (France)**