



HAL
open science

Global Change and Conservation of Solitary Mammals

Charlotte-Anaïs Olivier, Carsten Schradin, Lindelani Makuya

► **To cite this version:**

Charlotte-Anaïs Olivier, Carsten Schradin, Lindelani Makuya. Global Change and Conservation of Solitary Mammals. *Frontiers in Ecology and Evolution*, 2022, 10, 10.3389/fevo.2022.906446 . hal-03706078

HAL Id: hal-03706078

<https://hal.science/hal-03706078>

Submitted on 27 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Global Change and Conservation of Solitary Mammals

Charlotte-Anaïs Olivier^{1,2*}, Carsten Schradin^{1,2} and Lindelani Makuya¹

¹ School of Animal, Plant & Environmental Sciences, University of the Witwatersrand, Johannesburg, South Africa, ² IPHC, UNISTRA, CNRS, Strasbourg, France

Keywords: social organization, conservation, mammals, solitary species, group-living species

INTRODUCTION

Here we argue that understanding the social system of species can help us to conserve them. Many endangered mammals are solitary living (Clutton-Brock, 2021). Here we show how understanding the constraints and the adaptive value of solitary living is important to understand how solitary mammals will be affected and can respond to the different aspects of global change. For example, solitary species might be more vulnerable to global change than sociable species as they miss benefits of living in groups that could mitigate change.

GLOBAL CHANGE IMPACTS BIODIVERSITY

Global change is a major threat to biodiversity (Pimm, 2009; Harley, 2011; Thornton et al., 2014). Anthropogenic-induced global change includes habitat loss (Galbraith et al., 2002), fragmentation (Fahrig, 2003), and climate change (Chapin et al., 2000), all of which are accelerating at a rate unprecedented in earth's history (Etterson and Shaw, 2001). While the factors causing global change are known, the speed and strength at which it occurs is still unsure and we do not know the way in which species will be able to cope and adapt to these changes. Global change occurs at an unprecedented rate and is expected to have severe impacts on ecosystems (Friedlingstein, 2008), decreasing biodiversity (Vitousek et al., 1997; Sih et al., 2011).

UNDERSTANDING THE SOCIAL SYSTEM OF SPECIES, INCLUDING SOLITARY ONES, CAN HELP US TO CONSERVE THEM

In order to overcome and slow biodiversity loss, we need studies informing conservation programs, including those on mammals. There are many ways to prevent the decline of mammal species. One of them is through understanding and analyzing behavior, because changes in behavior can allow for resistance toward change, and constraints on behavioral flexibility restrict resilience (Rymer et al., 2016). Animal social systems have always been a key subject in behavioral ecology as it gives us information about the mating system, the care system, the social organization and the social structure of species (Kappeler, 2019). These four components of social systems are interrelated and influence each other. This has revealed stunning interspecific diversity in the size, composition, and cohesion of social units, as well as in the patterning of reproductive skew, cooperation, and competition among their members (for mammals see Clutton-Brock, 2021). Within mammals, species display a wide variety of social systems ranging from solitary living in many carnivores, to pair-living in some primates, to groups in some species, such as elephants (*Loxodonta africana*) and marmosets (*Callithrix jacchus*). By understanding the social system of a species we are able to predict how a species will respond to global change, and how its responses will be constrained, important information to decide on the best conservation efforts for each species.

OPEN ACCESS

Edited by:

Tasmin Rymer,
James Cook University, Australia

Reviewed by:

Sheng-Feng Shen,
Academia Sinica, Taiwan

*Correspondence:

Charlotte-Anaïs Olivier
charlotte-anaïs.olivier@iphc.cnrs.fr

Specialty section:

This article was submitted to
Behavioral and Evolutionary Ecology,
a section of the journal
Frontiers in Ecology and Evolution


Received: 28 March 2022

Accepted: 26 May 2022

Published: 24 June 2022

Citation:

Olivier C-A, Schradin C and Makuya L
(2022) Global Change and
Conservation of Solitary Mammals.
Front. Ecol. Evol. 10:906446.
doi: 10.3389/fevo.2022.906446



SOLITARY SPECIES MISS THE BENEFITS FROM GROUP-LIVING AND MIGHT BE MORE VULNERABLE TO GLOBAL CHANGE THAN SOCIABLE SPECIES

Conservation efforts for solitary species might differ from those of group-living species. The social system of a species induces costs and benefits that are reliant on the environment, especially with regards to competition for resources (mates, food shelter), and how the social system impacts on predation risk (Kappeler et al., 2013). The net outcome of these costs and benefits will change when the environment changes. Species that live in groups might be buffered from these changes due to the benefits of group living, especially improved foraging and predator avoidance (Krause and Ruxton, 2002). The extent that solitary species show behavioral flexibility to cope with changes in resource availability and predation pressure is unknown, i.e., the extent that individuals can alter their behavior reversibly in response to changing environmental conditions (Gordon, 1991). This is why we need to study the behavioral flexibility of solitary species under varying environmental conditions to assess their vulnerability to global change (Huey et al., 2012). As solitary species miss the benefits from group-living they might be more vulnerable than sociable species.

ALLEE EFFECT AND SOLITARY SPECIES

The Allee effect describes a positive correlation between individual fitness and population density, with individuals benefitting from the presence of conspecifics (Allee, 1927). When

living in groups, individuals can benefit from the Allee effect, by reducing predation pressure and gaining easier access to mating partners (Stephens and Sutherland, 2000). The Allee effect can thus be seen as a driving force behind animal spatial structure and sociality (Stephens and Sutherland, 2000). Due to climate change (obligate) sociable species might suffer from fitness costs induced by reduced group sizes. However, even solitary species might benefit from higher population density making it easier to find mating partners or reducing predation pressure (because predators have more prey to choose from), unless—as in sociable species—high population density increases intra-specific competition. Thus, so far it is unknown in how far the correlation between fitness and population density differs between sociable and solitary species, nor do we know at which population density positive Allee effects are overrun by negative effects of intra-specific competition. Therefore, in both sociable and solitary species the Allee effect should be taken into account when predicting how changes in population density induced by climate change affect extinction risk.

SOLITARY SPECIES MIGHT BE UNDERSTUDIED WHEN COMPARED TO SOCIABLE SPECIES

Solitary species might be understudied when compared to sociable species, especially with regards to their social behavior (Figure 1). First, for decades researchers rather studied sociable than solitary species as the focus of behavioral ecology was on the benefits of group-living and aspects like alloparental care (Clutton-Brock, 2021). Second, species that live solitary

are often more difficult to observe as they are nocturnal and shy, live in dense habitat (many small mammals), or have huge home ranges (solitary carnivores). In addition, the fact that they live alone makes them more difficult to monitor (Figure 1). The combination of a research focus on sociable species and the discrete nature of solitary species explains why they are understudied with regards to behavior, their trade-offs (energy, resources, time), and how these trade-offs might change and affect their resilience under conditions of global change.

TO CONSERVE MAMMALIAN BIODIVERSITY, WE NEED TO UNDERSTAND WHY SOLITARY LIVING IS AN ADAPTIVE TRAIT

Six hundred and seventy nine of the 5,500 mammalian species are endangered (IUCN redlist) and many of them are solitary living. The first mammal that became extinct as a direct cause of climate warming was the solitary Bramble Cay mosaic-tailed rat (*Melomys rubicola*; Fulton, 2017; Waller et al., 2017). It became extinct because of an ocean inundation, i.e., high water levels and storm surges caused by an increased frequency of extreme weather events, which were caused by anthropogenic climate change (Waller et al., 2017). Other well-known examples of endangered solitary

mammals that are threatened by climate change include polar bears (*Ursus maritimus*), snow leopards (*Panthera uncia*), tigers (*Panthera tigris*), and panda bears (*Ailuropoda melanoleuca*). It can be expected that many less well-monitored solitary small mammal species are threatened by climate change.

Many endangered species are solitary and thus understanding constraints and the adaptive value of solitary living is important to conserve them. The persistence of mammals being solitary living after 170 my of mammalian evolution (Rowe, 1988) indicates that it is an adaptive social system for many environments. To conserve mammalian biodiversity, we need to understand why solitary living is adaptive for many species. However, while for decades the costs and benefits of group living have been studied, it is not known what makes solitary living adaptive. Without knowing the costs and benefits of solitary living we will not know how the trade-offs that solitary species face will change with global change. At the moment, we do not even know which are their most restricting trade-offs, whether it is with regards to food, shelter, predation, or maybe finding mating partners.

AUTHOR CONTRIBUTIONS

C-AO, CS, and LM conceived of and wrote the manuscript. All authors approved the final manuscript.

REFERENCES

- Allee, W. C. (1927). Animal aggregations. *Q. Rev. Biol.* 2, 367–398. doi: 10.1086/394281
- Chapin, F., Zavaleta, E., Eviner, V., et al. (2000). Consequences of changing biodiversity. *Nature* 405, 234–242. doi: 10.1038/35012241
- Clutton-Brock, T. (2021). Social evolution in mammals. *Science* 373:eabc9699. doi: 10.1126/science.abc9699
- Etterson, J. R., and Shaw, R. G. (2001). Constraint to adaptive evolution in response to global warming. *Science* 294, 151–154. doi: 10.1126/science.1063656
- Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. *Annu. Rev. Ecol. Syst.* 34, 487–515. doi: 10.1146/annurev.ecolsys.34.011802.132419
- Friedlingstein, P. (2008). A steep road to climate stabilization. *Nature*. 451, 297–298. doi: 10.1038/nature06593
- Fulton, G. R. (2017). The Bramble Cay melomys: the first mammalian extinction due to human-induced climate change. *Pac. Conserv. Biol.* 23, 1–3. doi: 10.1071/PCv23n1_ED
- Galbraith, H., Jones, R., Park, R., Clough, J., Herrod-Julius, S., Harrington, B., et al. (2002). Global climate change and sea level rise: potential losses of intertidal habitat for shorebirds. *Waterbirds* 25, 173–183. doi: 10.1675/1524-4695(2002)0250173:GCCASL2.0.CO;2
- Gordon, D. M. (1991). Behavioral flexibility and the foraging ecology of seed-eating ants. *Am. Nat.* 138, 379–411. doi: 10.1086/285223
- Harley, C. D. G. (2011). Climate change, keystone predation, and biodiversity loss. *Science* 334, 1124–1127. doi: 10.1126/science.1210199
- Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M., and Williams, S. E. (2012). Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation. *Philos. Trans. R. Soc. B Biol. Sci.* 367, 1665–1679. doi: 10.1098/rstb.2012.0005
- Kappeler, P. M. (2019). A framework for studying social complexity. *Behav. Ecol. Sociobiol.* 73, 1–14. doi: 10.1007/s00265-018-2601-8
- Kappeler, P. M., Barrett, L., Blumstein, D. T., and Clutton-Brock, T. H. (2013). Constraints and flexibility in mammalian social behaviour: introduction and synthesis. *Philos. Trans. R. Soc. B Biol. Sci.* 368:20120337. doi: 10.1098/rstb.2012.0337
- Krause, J., and Ruxton, G. D. (2002). *Living in Groups*. Oxford: Oxford University Press.
- Pimm, S. L. (2009). Climate disruption and biodiversity. *Curr. Biol.* 19, 595–601. doi: 10.1016/j.cub.2009.05.055
- Rowe, T. (1988). Definition, diagnosis, and origin of Mammalia. *J. Vert. Paleontol.* 8, 241–264. doi: 10.1080/02724634.1988.10011708
- Rymer, T. L., Pillay, N., and Schradin, C. (2016). Resilience to droughts in mammals: a conceptual framework for estimating vulnerability of a single species. *Q. Rev. Biol.* 91, 133–176. doi: 10.1086/686810
- Sih, A., Ferrari, M. C., and Harris, D. J. (2011). Evolution and behavioural responses to human-induced rapid environmental change. *Evolut. Appl.* 4, 367–387. doi: 10.1111/j.1752-4571.2010.00166.x
- Stephens, P. A., and Sutherland, W. J. (2000). “Vertebrate mating systems, rarity and conservation,” in *Vertebrate Mating Systems*, eds M. Apollonio, M. Festa-Bianchet, and D. Mainardi (London: World Scientific Publishing), 186–213. doi: 10.1142/9789812793584_0009
- Thornton, P. K., Erickson, P. J., Herrero, M., and Challinor, A. J. (2014). Climate variability and vulnerability to climate change: a review. *Glob. Chang. Biol.* 20, 3313–3328. doi: 10.1111/gcb.12581
- Vitousek, P. M., Aber, J. D., Howarth, R. W., Likens, G. E., Matson, P. A., Schindler, D. W., et al. (1997). Human alteration of the global nitrogen cycle: sources and consequences. *Ecol. Appl.* 7, 737–750. doi: 10.1890/1051-0761(1997)007[0737:HAOTGN]2.0.CO;2

Waller, N. L., Gynther, I. C., Freeman, A. B., Lavery, T. H., and Leung, L. K.-P. (2017). The Bramble Cay melomys *Melomys rubicola* (Rodentia: Muridae): a first mammalian extinction caused by human-induced climate change? *Wildl. Res.* 44, 9–21. doi: 10.1071/WR16157

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of

the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Olivier, Schradin and Makuya. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.