



HAL
open science

Ambient BTEX Concentrations during the COVID-19 Lockdown in a Peri-Urban Environment (Orléans, France)

Min Cai, Yangang Ren, Rodrigo Gibilisco, Benoit Grosselin, Max McGillen,
Chaoyang Xue, Abdelwahid Mellouki, Véronique Daële

► **To cite this version:**

Min Cai, Yangang Ren, Rodrigo Gibilisco, Benoit Grosselin, Max McGillen, et al.. Ambient BTEX Concentrations during the COVID-19 Lockdown in a Peri-Urban Environment (Orléans, France). Atmosphere, 2021, 13 (1), pp.10. 10.3390/atmos13010010 . hal-03715748

HAL Id: hal-03715748

<https://hal.science/hal-03715748v1>

Submitted on 6 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 Article

2 Ambient BTEX Concentrations During the COVID-19 3 Lockdown in a Peri-urban Environment (Orléans, France)

4 Min Cai ¹, Yangang Ren ^{1,a}, Rodrigo G. Gibilisco ^{1,b}, Benoit Grosselin ¹, Max R. McGillen ¹, Chaoyang Xue ², Abdel-
5 wahid Mellouki ¹, Véronique Daële ¹ and *

6 ¹ Institut de Combustion, Aérothermique, Réactivité Environnement (ICARE), CNRS, 1C Avenue de la Re-
7 cherche Scientifique, 45071 Orléans Cedex 02, France; min.cai@cnrs-orleans.fr, benoit.grosselin@cnrs-or-
8 leans.fr, max.mcgillen@cnrs-orleans.fr, mellouki@cnrs-orleans.fr
9 ² Laboratoire de Physique et de Chimie de l'Environnement et de l'Espace (LPC2E), 3A Avenue de la Re-
10 cherche Scientifique, 45071 Orléans Cedex 2, France; chaoyang.xue@cnrs-orleans.fr
11 ^a present address: Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing
12 100085, China ; ygren@rcees.ac.cn
13 ^b present address: Laboratorio de Estudios Atmosféricos (LEA) - INQUINOA (CONICET-UNT), Facultad de
14 Bioquímica, Química y Farmacia, Universidad Nacional de Tucumán, San Lorenzo 456, T4000CAN, San
15 Miguel de Tucumán, Argentina ; rodrigogibilisco@gmail.com
16 * Correspondence: veronique.daele@cnrs-orleans.fr Tel.: +33-2-38-25-54-94

17 **Abstract:** During the period from March 17th to May 10th 2020, France saw dramatic shifts in domes-
18 tic, industrial and transport sectors as a national lockdown was introduced. So far, studies have
19 generally focused on urban settings, by contrast, this work reports data for a periurban location. Air
20 samples were collected and analyzed using a fully automated GC-MS-FID system in an air quality
21 monitoring station situated in the suburbs of Orléans, France. Average concentrations of BTEX (ben-
22 zene, toluene, ethylbenzene and xylenes) before, during and after lockdown, were 402 ± 143, 800 ±
23 378 and 851 ± 445 pptv, respectively. Diurnal variation in BTEX and correlations between each of its
24 components were analyzed to determine its various sources. The toluene/benzene (T/B) and m,p-
25 xylene/ethylbenzene (MP/E) ratios, photochemical ages were used to explore whether the BTEX
were from local or more distant sources. Together with a host of complementary measurements
including NO_x, O₃, black carbon, meteorological parameters and anthropogenic activities, we were
able to make some inferences on the sources of BTEX. The results suggest that although anomalous
local anthropogenic activity can lead to significant changes in BTEX concentrations, pollution levels
in Orléans are mostly dependent on meteorological conditions, specifically whether the winds are
coming from the Paris Region. It appears, based on these measurements, that the pollution in the
Orléans area is very much tied to the nearby megacity of Paris, this may be true for other peri-urban
sites with implications for city planning and pollution mitigation strategies.

26 **Citation:** Lastname, F.; Lastname, F.;
27 Lastname, F. Title. *Atmosphere* 2021,
28 12, x. <https://doi.org/10.3390/xxxxx>

29
30 Academic Editor: Firstname Last-
31 name

32 Received: date

33 Accepted: date

34 Published: date

35
36 **Publisher's Note:** MDPI stays neu-
tral with regard to jurisdictional
claims in published maps and institu-
tional affiliations.



37
38 **Copyright:** © 2021 by the author
39 Submitted for possible open access
40 publication under the terms and
41 conditions of the Creative Commons
42 Attribution (CC BY) license
43 ([https://creativecommons.org/licenses/](https://creativecommons.org/licenses/by/4.0/)
44 [by/4.0/](https://creativecommons.org/licenses/by/4.0/)).
45
46

Keywords: BTEX; COVID-19; lockdown; Orleans; peri-urban

1. Introduction

The COVID-19 pandemic has resulted in reduced emissions into the atmosphere worldwide as a result of the actions taken to limit or even stop certain human activities, which are generally linked to transportation and industrial activities. In some locations, these measures led to substantial reductions in the emissions of important atmospheric pollutants such as nitrogen oxides (NO_x), particulate matter (PM), and volatile organic compounds (VOCs) as reported by a number of studies around the globe. For example, Barré et al. [1] and Guevara et al. [2] reported emissions dropping by up to 60% for NO_x, and up to 15% for non-methane volatile organic compounds (NMVOC) during the March/April 2020 lockdown in Europe while Shi and Brasseur [3] showed that the surface concentrations of PM_{2.5} and NO₂ in China were reduced by 35% and 60%, respectively. It

is expected that the NO_x levels decreased in urban areas during lockdown periods since they are major combustion by-products from vehicle emissions. Many other regions have recorded significant reductions in air pollution globally [4–6].

While NO_x levels have generally decreased, NO_x concentration from other sources and the concentrations of other species may not follow the same trends, especially for those emitted into the atmosphere through plants, agricultural, biomass burning and indoor activities. In addition, atmospheric pollution is not a linear “chemistry” [7,8]. The reduction of NO_x alone does not necessitate a direct improvement of air quality, for example, lowering NO_x emissions may lead to an increase of ozone levels depending on a number of factors such as the types of VOCs and their concentrations as well as the solar irradiation. In fact, a number of studies have reported an increase of ozone concentrations in various areas during the lockdown despite the reduction in the NO_x levels [9,10].

Atmospheric chemistry is complex; a complete understanding of the processes occurring and the resulting impacts on air quality and health require the knowledge of many parameters, not only the concentrations of NO_x and PMs. More specifically, it requires the speciation of many other pollutants such as VOCs, which may play an important role in tropospheric ozone formation for example. Among the VOCs emitted into the atmosphere, the mono-aromatic hydrocarbons such as benzene, toluene, ethylbenzene, and xylene isomers (BTEX) are of special interest because of their role in the tropospheric chemistry and their adverse health effects such as asthma, dizziness, fatigue, and eye, nose, and throat irritation [11–13]. BTEX are emitted into the atmosphere from gasoline and diesel-powered automobiles, residential heating, industrial emissions and biogenic sources [14–17]. Once in the atmosphere, BTEX could react near the emission source or undergo transport depending on the atmospheric composition and meteorological conditions [18]. It is expected that the main degradation process in the gas phase during the day is the reaction with OH (hydroxyl radicals), resulting in a considerable contribution to photo-oxidant and secondary organic aerosol formation in the atmosphere [19]. The ratios of toluene/benzene (T/B) and m,p-xylene/ethylbenzene (MP/E) are considered to be useful indicators of atmospheric photochemical activity as well as the sources [20]. Assessing the impact of the lockdown on the BTEX atmospheric levels is important to the evaluation of the stay-at-home measures on the air quality during the pandemic period.

We report in the present work measurements of the concentrations of BTEX levels in a peri-urban area in France. The main objectives are: (1) to determine real time BTEX concentrations before, during and after the period of COVID-19 lockdown in Orléans, France; (2) to rationalize the obtained results based on the concentrations of secondary pollutants such as NO_x and O₃, particulate matter and black carbon and (3) to investigate the effect of meteorological parameters such as wind speed, wind direction, and temperature on the diurnal BTEX variations during the studied period.

2. Materials and Methods

2.1. Sampling Site

As shown in Fig. 1, the sampling site is positioned within the campus of the Centre National de la Recherche Scientifique (CNRS) about 10 km south from Orléans city center (47°50'17"N, 1°56'39"E). Orléans is the capital of the Centre-Val de Loire region, located in central France about 120 kilometers southwest of Paris, with a population of 300 thousand inhabitants. The site is close to a forest belt, and farms. There are no obstructing buildings around the sampling site within 50 m. The measurements reported in this study were made during the period from Feb. 26th to May 31st in 2020.




Figure 1. Location of the sampling site at the CNRS-campus in Orléans, France.

2.2. Instrumentation

2.2.1. BTEX Measurements

An automated online gas chromatographic system (GC-FID, AirmoVOC C6–C16 Chromatotec®) was used for in situ BTEX measurements with a time resolution of 40 min. Ambient air was sampled continuously from an inlet mounted at a height of 4 m above the ground through a Teflon tube (length 2.5 m, diameter 1/4 inch) at a 16 L min^{-1} bypass flow, and then entered the trap at a flow rate of 45 mL min^{-1} through a heated stainless-steel tube (length $\sim 1.5 \text{ m}$, diameter 1/8 inch). This sampling setup could ensure a short residence time in the sampling line and improve the sampling efficiency of the C6–C16 VOCs. The volume of air samples was 1350 ml for a 30 min sampling period. The trap was heated to 380°C within 4 min. VOCs were thermally desorbed and injected into capillary columns (analytical column: MXT 30 CE, film thickness: $1 \mu\text{m}$, id: 0.28 mm , length: 30 m) with H_2 around 3 to 4 mL min^{-1} for separation before detection. The temperature ramp of the capillary column (a plot of temperature vs time is provided in Figure. A1 in the supporting information) was set as follows: increased from 38°C to 50°C at a rate of 2°C min^{-1} ; increased to 80°C in 3 min; increased to 220°C at a rate of $15^\circ\text{C min}^{-1}$, then went to 230°C at a rate of 2°C min^{-1} ; finally reached 269°C at a rate of 9°C min^{-1} and held for 3 min. A flame ionization detector (FID) set at 200°C was used for quantification.

The accuracy of the system (quantification and identification) was controlled every 12h using an automatic calibration system, which is equipped with four internal permeation tubes containing four internal standards (benzene, n-butane, n-decane, and n-hexane). The detection limits for benzene, toluene, ethylbenzene, m, p-xylene, and o-xylene were $\sim 5 \text{ pptv}$.

2.2.2. NO_x, O₃, PM₁₀ and BC measurements

The concentrations of ozone (O₃), NO_x (include NO and NO₂), black carbon (BC), and particulate matter with a diameter less than 10 μm (PM₁₀) were provided by the air quality monitoring agency at the Centre-Val de Loire region by Lig'Air (<http://www.ligair.fr/>). These species were monitored at the same location as the BTEX measurements reported in the present work. Ambient air was sampled continuously from separate Teflon tubes (diameter 1/4 inch) at a height of 1 m above the roof of the container and about 4 m from the ground. BC was measured continuously at 1-min intervals using an aethalometer (AE33 model, Magee Scientific). The O₃ and NO–NO_x concentrations were continuously monitored by UV absorption (Thermo Scientific™ Model 49i) and chemiluminescence (Thermo Scientific™ Model 42i), respectively. A DIGITEL DHA-80 high volume aerosol sampler was also used for PM₁₀ sampling. Most of data processing and figures were performed using the R software environment [21] and particularly the Openair package, designed by Carslaw et al. [22].

2.2.3. Meteorological data

The meteorological data were used from Orléans-Bricy air base station (<https://www.infoclimat.fr/>) due to missing data for several weeks during lockdown.

3. Results and discussions

3.1. Data Overview

The measurement period (Feb. 26th to May 31st 2020) was divided in three regimes (a) before lockdown (Feb. 26th – Mar. 16th), (b) during lockdown (Mar. 17th – May 10th) and (c) after lockdown (May 11th – May 31st). Average BTEX mixing ratios along with their associated arithmetic mean and interquartile range (IQR) in each period are listed in Table 1. Benzene and ethylbenzene were the most abundant BTEX compounds throughout the study period. On average, the highest concentrations of BTEX were observed during and after the lockdown.


Table 1. Average mixing ratios (pptv) of BTEX for each period (before, during and after lockdown) in Orléans, France (SD: Standard deviation, IQR: interquartile range).

	Before lockdown		Lockdown		After lockdown	
	Feb. 26 th - Mar. 16 th		Mar. 17 th - May 10 th		May 11 th - May 31 st	
	Mean ± SD	IQR	Mean ± SD	IQR	Mean ± SD	IQR
N° samples	620		1860		595	
Benzene	83 ± 48	30	271 ± 143	233	280 ± 163	191
Toluene	56 ± 36	32	121 ± 80	112	118 ± 63	79
Ethylbenzene	99 ± 37	47	172 ± 80	125	205 ± 113	121
m,p-xylene	86 ± 29	37	124 ± 59	92	122 ± 58	76
o-xylene	79 ± 20	27	113 ± 49	65	128 ± 64	59
Total BTEX	402 ± 143	155	800 ± 378	626	851 ± 445	523

The temporal variations of BTEX, NO, NO₂, O₃, PM₁₀, BC, temperature at this site, and meteorological parameters (wind speed in m s⁻¹ and wind direction in a color-coded way) during the measurement period are illustrated in Figure. 2. Furthermore, the relative changes in daily regional mobility data from the Google Community Mobility Reports (<https://www.google.com/covid19/mobility/>) were also used for this analysis (the bottom of Figure. 2). The reports chart movement trends over time by geography, across different categories of places such as retail and recreation, groceries and pharmacies, parks, transit stations, workplaces, and residential. These reports are created with aggregated, anonymized sets of data from users who have turned on the Location History setting, which is off by default. This provides a breakdown by category (grocery stores, parks, residential, workplaces, transit stations, and retail) in changes to daily movement at a regional level.

159
160

The data “percent charge in the grocery and pharmacy” was used to perform the comparisons along the studied period.



161

162 **Figure 2.** Time series of hourly BTEX (pptv, y axis of BTEX 1k=1×10³), NO (ppbv), NO₂ (ppbv), O₃ (ppbv), PM₁₀ (µg m⁻³),
163 Black Carbon (µg m⁻³) and Temperature (°C) in Orléans during the studied period, with daily “percent change in the
164 grocery and pharmacy”, and meteorological parameters (wind speed in m s⁻¹, wind direction in a color-coded way).

165

3.2. Diurnal variation and correlations

166

167

168

169

170

171

The study of diurnal variations of air pollutants could provide valuable information about the sources, transport and chemical formation/destruction of such pollutants. The diurnal variations of the BTEX, BC, PM₁₀, NO_x and O₃ during the present study are shown in Figure. 3. (a series of diurnal variations in benzene, toluene, ethylbenzene, m,p-xylene and o-xylene mixing ratios before, during and after lockdown provided in the supporting information, Figure. A2)




Figure 3. Diurnal variations in BTEX, BC, PM₁₀, NO_x, O₃ and temperature before, during and after lockdown. The shaded area represent 95% confidence interval in mean.

BTEX concentrations have been observed to be generally higher during the night than daytime. The concentrations-time profiles show a decrease of the total BTEX from around 6 am to around 3 pm in the three periods. This is likely to result from a combination of effects such as photochemical reactions, an increase in the atmospheric mixing depth and/or a reduction in emission rates [23]. During the afternoon, an increase in concentration is observed around 3 pm. This increase is observed markedly during the period before the lockdown, which could be attributed to traffic emissions and home heating fuels consumed in winter in the late afternoon. High values were recorded also during rush hours in the morning and late afternoon before and after the lockdown. These feature was less clear during lockdown, revealing the possible contribution from the traffic emissions during periods before and after the lockdown.

Moreover, a strong correlation coefficient ($r = 0.83$) was obtained for ethylbenzene / xylene isomers before the lockdown (Figure. 4 a). This fact suggests that both compounds

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188
189
190
191
192
193
194

could be emitted from the gasoline vehicles and stations [24]. Toluene is also widely used as a solvent in paint manufacture, the production of adhesives and glues, while benzene emissions could be mainly due to wintertime domestic heating combustion in France [25]. During the lockdown (Figure. 4 b), the high correlations among all the BTEX suggests that the contributions are likely to be emitted by the same sources. The correlation becomes even stronger after the lockdown period as shown by Figure 4 c, which indicates that the source of BTEX were becoming increasingly homogenous.

195

196
197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213




Figure 4. Correlation coefficients in percentage between Benzene, Toluene, Ethylbenzene, m,p-Xylene and o-Xylene (a) before, (b) during and (c) after the lockdown.

3.3. BTEX ratios assessment

The Toluene/Benzene ratio (T/B) is often used in BTEX source apportionment studies [26]. It is a tool for characterizing the distance from (and age of) vehicular emission sources. While toluene emissions are associated with traffic and industries, benzene from these sources has been reduced as a consequence of gasoline regulations. The principal source of benzene in France is therefore likely to be residential heating [27].

It can be observed in Figure. 5 that the T/B ratio falls below 1 in general, with some higher values observed in the first week of lockdown. This suggests that there were no major changes in BTEX sources during our study period, which is surprising, given the decrease in transport activities and the change in the population's daily routines during lockdown. When the location of the measurement site in Orléans is considered in relation to potential BTEX sources in Paris, the T/B ratios < 1 that we observe are broadly consistent with the measurements of Salameh [28], who show that in suburban sites such as Melun and other locations on the outskirts of Paris, T/B ratios are close to 1. Nevertheless, it is possible that a significant contribution of the benzene concentration comes from residential heating as mentioned above..




Figure 5. Variation of toluene / benzene (T/B), photochemical age of the air (Δt), and m, p-xylene / ethylbenzene (MP/E) concentration ratios in the observation site during the entire measurement period.

In this study, the photochemical age was used to estimate the origin of the air masses. Its definition depends on the fact that the more reactive chemicals are consumed faster than the less reactive ones, so that as an air mass ages there are systematic changes in concentration ratios [29]. The T/B ratio in the sampled air as described by Roberts [30]:

$$\Delta t = \frac{1}{[\text{OH}](k_{\text{toluene}} - k_{\text{benzene}})} \times \left[\ln \left(\frac{[\text{toluene}]_t}{[\text{benzene}]_t} \right) - \ln \left(\frac{[\text{toluene}]_0}{[\text{benzene}]_0} \right) \right] \quad (1)$$

where k_{toluene} and k_{benzene} are the rate coefficients for the reaction with OH ($k_{\text{toluene}} = 5.63 \times 10^{-12}$ and $k_{\text{benzene}} = 1.22 \times 10^{-12}$ at 298 K in $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ [31]. $[\text{OH}]$ is the average concentration of the OH radical ($2.1 \times 10^6 \text{ molecule cm}^{-3}$, [32]). The initial emission ratio of T/B was set to 1.69 (Initial value of B/T = 0.59), it was taken from Airparif station, which is located on the roof of the main Airparif building, close to busy roads, in the Paris city center [33].

It can be seen from Figure. 5 that when the T/B ratio increases, the photochemical age decreases. For example, on the evening of Mar. 13th and the early morning of Mar. 16th, it is as low as 1.37 and 1.14 days, respectively. This suggests that the pollutants are fresher and the source of the pollution is relatively close to the site (100–300 km). In most of the observation period, the photochemical age is more than 3 days, indicating that long-distance air transport could contribute to the air pollution in this region (> 500 km).

The m,p-xylene to ethylbenzene (MP/E) ratio could also be used to evaluate the age of air parcels and as an indicator for the photochemical age of the VOCs in the atmosphere. The xylenes are more reactive towards the OH radical than ethylbenzene so the low ratio of MP/E could be used as an indication of an aged air parcel. Relatively constant MP/E ratios ranging from 2.8 to 4.6 with a mean value of 3.5 due to near traffic exhaust emissions have been previously reported [34]. MP/E ratio along the studied period is presented in Figure. 5. MP/E ratios average shows a quite small but noticeable difference before (0.87), during (0.72) and after (0.59) the lockdown. The values in the three periods suggest a rather aged air parcel, and the decreasing trend is an expected effect due to the increase in solar radiation as the spring season approaches. During and after lockdown the variation in the MP/E ratio reaches its maximum value before sunrise and its minimum at mid-afternoon (Figure. 5), in agreement with what is expected for a photochemically aged air parcel [35]. On the other hand, before the lockdown, this cyclical variation in the MP/E ratio between day and night is not clearly observed, this may be due to the different emission sources during winter (Figure. 6). Figure. 2 shows that the prevailing wind direction changed significantly after the first period. This is likely to have also a strong effect, transporting air from different locations with different distance.




Figure 6. Diurnal variations in MP/E ratio (a) before, (b) during and (c) after the lockdown.

3.4. Impact from community mobility and meteorological effects on the measured BTEX

In order to get a better understanding of the BTEX sources we made a series of supporting measurements including meteorological information, O_3 , NO , NO_2 , PM_{10} and BC . For this purpose, we investigated correlations over 4 episodes taken from the whole period of investigation: before lockdown (Feb. 26th - Mar. 12th), declared lockdown (Mar. 13th, Mar. 15th, Mar. 16th), lockdown (Mar. 17th - May 10th) and after the lockdown (May 11th - May 31st). Figure. 7 shows correlation coefficient between BTEX, O_3 , NO , NO_2 , PM_{10} , BC and temperature (T) for each period.

BTEX average concentrations were relatively low before lockdown, with an average mixing ratio around 402 ± 143 pptv. Nevertheless, on March 13th and 15th to 16th, two higher concentration peaks of BTEX were detected, with about 820 pptv and 1040 pptv, respectively. To better understand the possible causes and identify the main contribution sources for these particular events, correlation coefficients were calculated using the corresponding data of these particular days as shown in Figure. 7b. From this figure, it can be seen that a high positive correlation ($r = 0.90$) between BC and NO_2 , but BTEX didn't correlate very well with either of them ($r = 0.51$ with NO_2 and $r = 0.63$ with BC), suggesting that it could be another source besides fuels and/or biomass burning [36,37].

During the same period, the warm weather would suggest that it is less likely that the burning of wood for home heating represented a significant source, however, given the extremely different daily schedules of people during this timeframe, it remains possible that residential heating was a contributing factor. Considering these facts, BTEX concentration increase during these episodes could be related to transportation emissions rather than other sources. This may be linked to the announcement of official measures to be implemented in order to prevent the spread of COVID-19 on Mar. 12th. Similarly, for Mar. 15th and 16th with the order from the French Government to close all non-essential public places and businesses, including restaurants, cafes, shops, and entertainment venues from Mar. 14th in order to limit social movements. To support this hypothesis, the Google Community Mobility Reports data for Friday 13th, Sunday 15th and Monday 16th of March were also used. From these data, displayed in Figure. 2, it is possible to observe a substantial increase in the mobility of people (percent change) on the same dates where the maximum concentrations of BTEX were observed. Considering this, it could be speculated that most of the emissions detected during Mar. 13th, 15th and 16th were strongly related to the local use of transport vehicles on an elevated scale to visit pharmacies and grocery stores.




Figure 7. Correlation coefficient in percentage between BTEX, NO, NO₂, O₃, PM₁₀, BC and temperature (T) for each period (before, declared, during and after lockdown).

During the lockdown, social mobility decreased significantly as it can be seen from the relative changes in the Mobility Reports (Figure. 2). At the same time, NO concentrations also showed a progressive decrease, which is likely to be associated with the reduction of local traffic emissions. However, BTEX concentrations showed multiple elevated episodes during the lockdown period. From the correlation coefficients shown in Figure. 7, it is possible to observe that this increase does not correlate with the NO concentration ($r = 0.02$), weakly correlated with NO₂ ($r = 0.29$), and only shows a moderate correlation with black carbon ($r = 0.45$). Besides, from Figure. 2, it is possible to observe that wind direction changed markedly in the region, from southwest before lockdown to northeast during the lockdown.

To consider a possible effect of these variables, polar plots of each pollutant and temperature against wind direction were constructed in Figure. 8, where each factor is plotted as a function of wind speed and wind direction. The dominant wind direction came from the northeast 43.1% of the time, which was most frequent during the lockdown. Other wind directions were: north 15%, east 8.7%, southeast 5.2%, south 8.1%, southwest 12%, west 5.8% and northwest 2%.

286
287
288

289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304




Figure 8. Polar plots of BTEX, NO, NO₂, O₃, PM₁₀, BC and temperature (T) for lockdown (from Mar. 17th to May 10th); ws = wind speed (m s⁻¹).

For NO and NO₂, polar plot representations clearly show that maximum concentrations were observed for winds blowing from the southwest. The southwest of the site is located near a traffic intersection, gas station and provincial road, which would indicate very local sources.

Another area with high NO₂ concentration was found in the north, which has a similar situation for BC and PM₁₀. It is noted that the site is located in the south of Orléans (distance 6 km from city center) and south west of Paris (distance 90 km), and that the major source of volatile organic compounds during wintertime in the Paris region is wood burning [38]. Hence, impact of biomass or coal burning emissions, especially during the lockdown, was clearly evidenced on NO₂, BC and PM₁₀ atmospheric concentrations.

Concerning the BTEX mixing ratios, the probability to observe high concentrations decreased with the wind speed. Higher concentrations were associated with a speed ranging between 0 and 4 m s⁻¹, lower concentrations of all pollutants were observed when the wind speed was greater than 12 m s⁻¹, intuitively, this result indicates that the high wind speed plays a role in “cleaning up” the air over the measurements site area.

4. Conclusions

The ambient levels and possible sources of atmospheric BTEX during the lockdown period in Orléans are reported in this study. The variation of the mean BTEX concentration suggests that besides source strength, the seasonal and diurnal variations of atmospheric BTEX in periurban areas also strongly depend on meteorological conditions and photochemical activity.

T/B ratios are generally low (<1) at the site during all three periods. In contrast to the observations of Salameh [28], which focused on the Paris area, we find that temperature

is not a strong determinant of T/B, which suggests that evaporation rates are not the major control on this ratio in our study, and that photochemical age may be playing a more important role. Previous observations in Parisian suburbs such as Melun [28] find T/B ratios of approximately 1, which is similar to our own observations and suggests that we are observing a more aged air mass in line with the overall findings of the MEGAPOLI campaign [28].

Surprisingly, the overall BTEX mixing ratios were in general at their highest (~ 1 ppb) during the lockdown period, a time when we would expect to have the lowest volumes of traffic, which is supported by the Google Community mobility reports. Some of this effect is likely to be caused by meteorological conditions, where a change in the prevailing wind direction from SW to NE occurred near the onset of lockdown, suggesting that Orléans during this time could be more affected by plumes from the Paris megacity. It is noted that the overall concentrations are lower than those observed in Paris, as would be expected from photochemical losses as well as mixing and diffusion during the transport process. Changes to daily routines that could result in increases of residential heating emissions could also explain elevated BTEX mixing ratios during lockdown, although it is noted that the meteorological conditions during this time were generally warmer than the period that preceded lockdown. This hypothesis is supported on some days by the higher loading of BC and PM₁₀ observed during lockdown, which could be attributable to a wood smoke source.

In addition to the BTEX measurements, we present a suite of supporting measurements including NO_x, O₃, PM₁₀, BC and meteorological data. In general, we didn't observe strong correlations between these parameters and BTEX, suggesting that their respective sources could be different.

Even though we can be sure of changes in emission sources during the lockdown period, we find that the highest ambient pollution levels occurred during a time of national inactivity. Given that BTEX are considered to be dominated by traffic [25], and that local traffic volumes were reduced during the lockdown, we conclude therefore that ambient pollution levels in Orléans are strongly connected with the neighboring Paris region, which may be true of other suburban and peri-urban sites around the globe. Furthermore, if the higher BTEX episodes observed during lockdown are indeed related to more local sources of residential heating, then this implies that emission control of wood burning should be prioritized.

Author Contributions: M.C., Y.R., R.G. and B.G. conducted the measurements, M.C., Y.R., S.X. and B.G. performed the analysis of the data, M.C., M.M.G and A.M. wrote the manuscript, all coauthors commented on the manuscript, V.D. and A.M. supervised the work and provided the resources.

Funding: This research was funded through PIVOTS project by the Région Centre – Val de Loire (ARD 2020 program), CPER 2015 -2020 and European Union who invests in Centre-Val de Loire with the European Regional Development Fund. This work is also part of the "Investments d'Avenir" Programme overseen by the French National Research Agency (ANR) (LabEx BASC; ANR-11-LABEX-0034).

Institutional Review Board Statement: Not applicable

Informed Consent Statement: Not applicable.

Data Availability Statement: <https://zenodo.org/record/5617970#.YXwM4prP1PY>

Acknowledgments: We are grateful to Météo France and Lig'Air for the use of their meteorological and other pollutants data, respectively. M.C. is grateful to the China Scholarship Council for the financial support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 380 1. Barré, J.; Petetin, H.; Colette, A.; Guevara, M.; Peuch, V.-H.; Rouil, L.; Engelen, R.; Inness, A.; Flemming, J.; Pérez García-Pando,
381 C.; et al. *Estimating Lockdown Induced European NO₂ Changes*; Gases/Remote Sensing/Troposphere/Chemistry (chemical
382 composition and reactions), 2020;
- 383 2. Guevara, M.; Jorba, O.; Soret, A.; Petetin, H.; Bowdalo, D.; Serradell, K.; Tena, C.; Denier van der Gon, H.; Kuenen, J.; Peuch, V.-
384 H.; et al. Time-Resolved Emission Reductions for Atmospheric Chemistry Modelling in Europe during the COVID-19
385 Lockdowns. *Atmospheric Chem. Phys.* **2021**, *21*, 773–797, doi:https://doi.org/10.5194/acp-21-773-2021.
- 386 3. Shi, X.; Brasseur, G.P. The Response in Air Quality to the Reduction of Chinese Economic Activities During the COVID-19
387 Outbreak. *Geophys. Res. Lett.* **2020**, *47*, e2020GL088070, doi:https://doi.org/10.1029/2020GL088070.
- 388 4. Briz-Redón, Á.; Belenguer-Sapiña, C.; Serrano-Aroca, Á. Changes in Air Pollution during COVID-19 Lockdown in Spain: A
389 Multi-City Study. *J. Environ. Sci.* **2021**, *101*, 16–26, doi:10.1016/j.jes.2020.07.029.
- 390 5. Le, T.; Wang, Y.; Liu, L.; Yang, J.; Yung, Y.L.; Li, G.; Seinfeld, J.H. Unexpected Air Pollution with Marked Emission Reductions
391 during the COVID-19 Outbreak in China. *Science* **2020**, *369*, 702–706, doi:10.1126/science.abb7431.
- 392 6. Singh, V.; Singh, S.; Biswal, A.; Kesarkar, A.P.; Mor, S.; Ravindra, K. Diurnal and Temporal Changes in Air Pollution during
393 COVID-19 Strict Lockdown over Different Regions of India. *Environ. Pollut.* **2020**, *266*, 115368, doi:10.1016/j.envpol.2020.115368.
- 394 7. Kroll, J.H.; Heald, C.L.; Cappa, C.D.; Farmer, D.K.; Fry, J.L.; Murphy, J.G.; Steiner, A.L. The Complex Chemical Effects of
395 COVID-19 Shutdowns on Air Quality. *Nat. Chem.* **2020**, *12*, 777–779, doi:10.1038/s41557-020-0535-z.
- 396 8. Menut, L.; Bessagnet, B.; Siour, G.; Mailler, S.; Pennel, R.; Cholokian, A. Impact of Lockdown Measures to Combat Covid-19 on
397 Air Quality over Western Europe. *Sci. Total Environ.* **2020**, *741*, 140426, doi:10.1016/j.scitotenv.2020.140426.
- 398 9. Collivignarelli, M.C.; Abbà, A.; Bertanza, G.; Pedrazzani, R.; Ricciardi, P.; Carnevale Miino, M. Lockdown for CoViD-2019 in
399 Milan: What Are the Effects on Air Quality? *Sci. Total Environ.* **2020**, *732*, 139280, doi:10.1016/j.scitotenv.2020.139280.
- 400 10. Schripp, T.; Langer, S.; Salthammer, T. Interaction of Ozone with Wooden Building Products, Treated Wood Samples and Exotic
401 Wood Species. *Atmos. Environ.* **2012**, *54*, 365–372, doi:10.1016/j.atmosenv.2012.02.064.
- 402 11. Lee, S.C. Volatile Organic Compounds (VOCs) in Urban Atmosphere of Hong Kong., doi:https://doi.org/10.1016/S0045-
403 6535(02)00040-1.
- 404 12. Liu, J.; Mu, Y.; Zhang, Y.; Zhang, Z.; Wang, X.; Liu, Y.; Sun, Z. Atmospheric Levels of BTEX Compounds during the 2008
405 Olympic Games in the Urban Area of Beijing. *Sci. Total Environ.* **2009**, *408*, 109–116, doi:10.1016/j.scitotenv.2009.09.026.
- 406 13. Mehlman, Dangerous Properties of Petroleum-Refining Products: Carcinogenicity of Motor Fuels (Gasoline).,
407 doi:10.1002/tcm.1770100505.
- 408 14. Demir, S.; Saral, A.; Ertürk, F.; Kuzu, S.L.; Goncaloğlu, B.İ.; Demir, G. Effect of Diurnal Changes in VOC Source Strengths on
409 Performances of Receptor Models. *Environ. Sci. Pollut. Res.* **2012**, *19*, 1503–1514, doi:10.1007/s11356-011-0636-8.
- 410 15. Guo, H.; So, K.L.; Simpson, I.J.; Barletta, B.; Meinardi, S.; Blake, D.R. C1–C8 Volatile Organic Compounds in the Atmosphere of
411 Hong Kong: Overview of Atmospheric Processing and Source Apportionment. *Atmos. Environ.* **2007**, *17*.
- 412 16. Kansal, A. Sources and Reactivity of NMHCs and VOCs in the Atmosphere: A Review. *J. Hazard. Mater.* **2009**, *10*.
- 413 17. Lanz, V.A.; Buchmann, B.; Hueglin, C.; Locher, R.; Reimann, S.; Staehelin, J. Factor Analytical Modeling of
414 C₁–C₈ Hydrocarbon Sources at an Urban Background Site in Zurich
415 (Switzerland): Changes between 1993–1994 and 2005–2006. *Atmospheric Chem. Phys. Discuss.* **2008**, *8*, 907–955, doi:10.5194/acpd-
416 8-907-2008.
- 417 18. Finlayson-Pitts, B.J.; Pitts, J.N. Atmospheric Chemistry of Tropospheric Ozone Formation: Scientific and Regulatory
418 Implications. *Air Waste* **1993**, *43*, 1091–1100, doi:10.1080/1073161X.1993.10467187.
- 419 19. Calvert, J.; Mellouki, A.; Orlando, J.; Pilling, M.; Wallington, T. *Mechanisms of Atmospheric Oxidation of the Oxygenates*; Oxford
420 University Press, New York, USA, 2011;
- 421

- 422 20. Nelson, R.M.; Quigley, J.P. The m,p-Xylenes:Ethylbenzene Ratio. A Technique for Estimating Hydrocarbon Age in Ambient Atmospheres.
423 *Atmospheric Environ.* **1983**, *17*, 659–662, doi:10.1016/0004-6981(83)90141-5.
- 424 21. R Core Team R: A Language and Environment for Statistical Computing. *Found. Stat. Comput.* **2018**.
- 425 22. Carslaw, D.C. Openair - An R Package for Air Quality Data Analysis. *Environ. Model.* **2012**, *10*.
- 426 23. Jiang, Z.; Grosselin, B.; Daële, V.; Mellouki, A.; Mu, Y. Seasonal and Diurnal Variations of BTEX Compounds in the Semi-Urban
427 Environment of Orleans, France. *Sci. Total Environ.* **2017**, *574*, 1659–1664, doi:10.1016/j.scitotenv.2016.08.214.
- 428 24. Hajizadeh, Y.; Mokhtari, M.; Faraji, M.; Mohammadi, A.; Nemati, S.; Ghanbari, R.; Abdollahnejad, A.; Fard, R.F.; Nikoonahad,
429 A.; Jafari, N.; et al. Trends of BTEX in the Central Urban Area of Iran: A Preliminary Study of Photochemical Ozone Pollution
430 and Health Risk Assessment. *Atmospheric Pollut. Res.* **2018**, *9*, 220–229, doi:10.1016/j.apr.2017.09.005.
- 431 25. Borbon, A.; Boynard, A.; Salameh, T.; Baudic, A.; Gros, V.; Gauduin, J.; Perrussel, O.; Pallares, C. Is Traffic Still an Important
432 Emitter of Monoaromatic Organic Compounds in European Urban Areas? *Environ. Sci. Technol.* **2018**, *52*, 513–521,
433 doi:10.1021/acs.est.7b01408.
- 434 26. Gelencsér, A.; Siszler, K.; Hlavay, J. Toluene–Benzene Concentration Ratio as a Tool for Characterizing the Distance from
435 Vehicular Emission Sources. *Environ. Sci. Technol.* **1997**, *31*, 2869–2872, doi:10.1021/es970004c.
- 436 27. Sauvage, S.; Plaisance, H.; Locoge, N.; Wroblewski, A.; Coddeville, P.; Galloo, J.C. Long Term Measurement and Source
437 Apportionment of Non-Methane Hydrocarbons in Three French Rural Areas. *Atmos. Environ.* **2009**, *43*, 2430–2441,
438 doi:10.1016/j.atmosenv.2009.02.001.
- 439 28. Salameh, T.; Sauvage, S.; Locoge, N.; Gauduin, J.; Perrussel, O.; Borbon, A. Spatial and Temporal Variability of BTEX in Paris
440 Megacity: Two-Wheelers as a Major Driver. *Atmospheric Environ. X* **2019**, *1*, 100003, doi:10.1016/j.aeaoa.2018.100003.
- 441 29. Kleinman, L.I.; Daum, P.H.; Lee, Y.-N.; Nunnermacker, L.J.; Springston, S.R.; Weinstein-Lloyd, J.; Hyde, P.; Doskey, P.; Rudolph,
442 J.; Fast, J.; et al. Photochemical Age Determinations in the Phoenix Metropolitan Area. *J. Geophys. Res. Atmospheres* **2003**, *108*,
443 doi:https://doi.org/10.1029/2002JD002621.
- 444 30. Roberts, J.M.; Fehsenfeld, F.C.; Liu, S.C.; Bollinger, M.J.; Hahn, C.; Albritton, D.L.; Sievers, R.E. Measurements of Aromatic
445 Hydrocarbon Ratios and NO_x Concentrations in the Rural Troposphere: Observation of Air Mass Photochemical Aging and
446 NO_x Removal. *Atmospheric Environ.* **1984**, *18*, 2421–2432, doi:10.1016/0004-6981(84)90012-X.
- 447 31. Atkinson, R.; Baulch, D.L.; Cox, R.A.; Crowley, J.N.; Hampson, R.F. Summary of Evaluated Kinetic and Photochemical Data for
448 Atmospheric Chemistry. **2006**, *60*.
- 449 32. Warneke, C.; McKeen, S.A.; Gouw, J.A. de; Goldan, P.D.; Kuster, W.C.; Holloway, J.S.; Williams, E.J.; Lerner, B.M.; Parrish, D.D.;
450 Trainer, M.; et al. Determination of Urban Volatile Organic Compound Emission Ratios and Comparison with an Emissions
451 Database. *J. Geophys. Res. Atmospheres* **2007**, *112*, doi:10.1029/2006JD007930.
- 452 33. Languille, B.; Gros, V.; Petit, J.-E.; Honoré, C.; Baudic, A.; Perrussel, O.; Foret, G.; Michoud, V.; Truong, F.; Bonnaire, N.; et al.
453 Wood Burning: A Major Source of Volatile Organic Compounds during Wintertime in the Paris Region. *Sci. Total Environ.* **2020**,
454 *711*, 135055, doi:10.1016/j.scitotenv.2019.135055.
- 455 34. Monod, A.; Sive, B.C.; Avino, P.; Chen, T.; Blake, D.R.; Sherwood Rowland, F. Monoaromatic Compounds in Ambient Air of
456 Various Cities: A Focus on Correlations between the Xylenes and Ethylbenzene. *Atmos. Environ.* **2001**, *35*, 135–149,
457 doi:10.1016/S1352-2310(00)00274-0.
- 458 35. Rad, H.D.; Babaei, A.A.; Goudarzi, G.; Angali, K.A.; Ramezani, Z.; Mohammadi, M.M. Levels and Sources of BTEX in Ambient
459 Air of Ahvaz Metropolitan City. *Air Qual. Atmosphere Health* **2014**, *7*, 515–524, doi:10.1007/s11869-014-0254-y.
- 460 36. Kirchstetter, T.W.; Novakov, T.; Hobbs, P.V. Evidence That the Spectral Dependence of Light Absorption by Aerosols Is
461 Affected by Organic Carbon. *J. Geophys. Res. Atmospheres* **2004**, *109*, doi:https://doi.org/10.1029/2004JD004999.
- 462 37. Logan, J.A. Nitrogen Oxides in the Troposphere: Global and Regional Budgets. *J. Geophys. Res. Oceans* **1983**, *88*, 10785–10807,
463 doi:https://doi.org/10.1029/JC088iC15p10785.

- 464 38. Languille, B. Wood Burning: A Major Source of Volatile Organic Compounds during Wintertime in the Paris Region. *Sci. Total*
465 *Environ.* **2020**, *16*.
466