



HAL
open science

A Gaussian correlation inequality for plurisubharmonic functions

Franck Barthe, Dario Cordero-Erausquin

► **To cite this version:**

Franck Barthe, Dario Cordero-Erausquin. A Gaussian correlation inequality for plurisubharmonic functions. 2022. hal-03716857

HAL Id: hal-03716857

<https://hal.science/hal-03716857>

Preprint submitted on 7 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Gaussian correlation inequality for plurisubharmonic functions

F. Barthe and D. Cordero-Erausquin

July 7, 2022

Abstract

A positive correlation inequality is established for circular-invariant plurisubharmonic functions, with respect to complex Gaussian measures. The main ingredients of the proofs are the Ornstein-Uhlenbeck semigroup, and another natural semigroup associated to the Gaussian $\bar{\partial}$ -Laplacian.

1 Introduction

The motivation of the present work comes from a Gaussian moment inequality in \mathbb{C}^n due to Arias de Reyna [1]. We will show that his result is in fact a very particular case of a new correlation inequality, which can be seen as the complex analogue of the following correlation inequality for convex function in \mathbb{R}^n due to Hu [6]: if μ is a centered Gaussian measure on \mathbb{R}^n and if $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ are *convex* functions in $L^2(\mu)$ and f is *even*, then

$$\int f g d\mu \geq \int f d\mu \int g d\mu.$$

We will say that a function on \mathbb{C}^n is *circular-symmetric* if it is invariant under the action of S^1 (i.e. multiplication by complex numbers of modulus one) ; in other words a function f defined on \mathbb{C}^n is circular-symmetric if

$$f(e^{i\theta}w) = f(w) \quad \forall \theta \in \mathbb{R}, \quad \forall w \in \mathbb{C}^n.$$

A function $u : \mathbb{C}^n \rightarrow [-\infty, +\infty)$ is plurisubharmonic (psh) if it is upper semi-continuous and for all $a, b \in \mathbb{C}^n$ the function $z \in \mathbb{C} \mapsto u(a + zb)$ is subharmonic. Classically, a twice continuously differentiable function $u : \mathbb{C}^n \rightarrow \mathbb{R}$ is psh if and only if for all $w, z \in \mathbb{C}^n$

$$\sum_{j,k} \partial_{z_j \bar{z}_k}^2 u(z) w_j \bar{w}_k \geq 0,$$

where $\partial_{z_j} = \frac{1}{2}(\partial_{x_j} - i\partial_{y_j})$, $\partial_{\bar{z}_j} = \frac{1}{2}(\partial_{x_j} + i\partial_{y_j})$, $z = x + iy$ with $x, y \in \mathbb{R}^n$. The later condition means that the complex Hessian $D_{\mathbb{C}}^2 u$ is pointwise Hermitian semi-definite positive. We refer e.g. to the textbook [5] for more details.

We consider the standard complex Gaussian measure γ on \mathbb{C}^n ,

$$d\gamma(w) = d\gamma_n(w) = \pi^{-n} e^{-w \cdot \bar{w}} d\ell(w) = \pi^{-n} e^{-|w|^2} d\ell(w),$$

where ℓ denotes the Lebesgue measure on $\mathbb{C}^n \simeq \mathbb{R}^{2n}$ and $w \cdot w' = \sum w_j w'_j$ for $w, w' \in \mathbb{C}^n$. For convenience, let us introduce the following class of $L^2(\gamma)$ functions with controlled growth at infinity:

$$\mathcal{G} := \left\{ f : \mathbb{C}^n \rightarrow \mathbb{C}; f \in L^2_{loc}(\lambda) \text{ and } \exists \epsilon, c, C > 0 \text{ such that } |f(w)| \leq e^{c|w|^{2-\epsilon}}, \forall |w| \geq C \right\}.$$

In particular any function (locally L^2) dominated by a polynomial function on \mathbb{R}^{2n} belongs to \mathcal{G} . Our main result reads as follows:

Theorem 1 (Correlation for psh functions). *Let $f, g : \mathbb{C}^n \rightarrow [-\infty, \infty)$ be two plurisubharmonic functions belonging to \mathcal{G} . If f is circular-symmetric, then*

$$\int f g d\gamma \geq \int f d\gamma \int g d\gamma.$$

One can extend the result by approximation to more general psh functions in $L^2(\gamma)$. The inequality also extends to arbitrary centered complex Gaussian measure, which are images of γ by \mathbb{C} -linear maps. Indeed composing a psh function with a \mathbb{C} -linear map gives another psh function.

Let us give some direct consequences of this theorem. First, we see that when F, G are holomorphic functions, or simply complex polynomial functions, belonging to $L^2(\gamma)$ and F is homogeneous, then for any $\alpha, \beta \geq 0$ we have

$$\int |F|^\alpha |G|^\beta d\gamma \geq \int |F|^\alpha d\gamma \int |G|^\beta d\gamma.$$

Indeed, if F is holomorphic $f = |F|^\beta$ is psh, and if F is homogeneous, then f is also circular-symmetric. This argument can also be used for products of the form

$$f := |F_1|^{\alpha_1} \dots |F_k|^{\alpha_k}$$

where the F_j are holomorphic and the α_j 's are nonnegative real numbers, so that f is log-psh, in the sense that

$$\log f(w) = \sum_{\ell=1}^k \alpha_\ell \log |F_\ell(w)|$$

is psh. This implies that f is also psh, and if the holomorphic functions F_j are homogeneous then f is also circular-symmetric.

Theorem 2. *Let F_1, \dots, F_N be a family of homogeneous polynomial functions on \mathbb{C}^n . Then for any $\alpha, \dots, \alpha_N \geq 0$ and $k \leq N - 1$ we have*

$$\int \prod_{j=1}^N |F_j|^{\alpha_j} d\gamma \geq \left(\int \prod_{j=1}^k |F_j|^{\alpha_j} d\gamma \right) \left(\int \prod_{j=k+1}^N |F_j|^{\alpha_j} d\gamma \right) \geq \prod_{j=1}^N \int |F_j|^{\alpha_j} d\gamma.$$

A standard complex Gaussian vector in \mathbb{C}^n is a random vector taking values in \mathbb{C}^n according to the distribution $\gamma = \gamma_n$. A random vector $X = (X_1, \dots, X_N) \in \mathbb{C}^N$ is a centered complex Gaussian vector if there is an n , a standard complex Gaussian vector G in \mathbb{C}^n and a \mathbb{C} -linear map $A : \mathbb{C}^n \rightarrow \mathbb{C}^N$ such that $X = AG$. It turns out that the law for X is then characterized by its complex covariance matrix $(\mathbb{E}(X_k \overline{X_\ell}))_{1 \leq k, \ell \leq N}$. Denoting by $a_1, \dots, a_N \in \mathbb{C}^n$ the rows the matrix of A in the canonical basis, $X_j = G \cdot a_j$. Applying the later theorem to the complex linear forms $F_j(w) = w \cdot a_j$ yields the following result.

Theorem 3. *Let $(X_1, \dots, X_N) \in \mathbb{C}^N$ be a centered complex Gaussian vector, and let $\alpha_1, \dots, \alpha_N \in \mathbb{R}^+$. Then, for any $k \leq N - 1$*

$$\mathbb{E} \prod_{j=1}^N |X_j|^{\alpha_j} \geq \left(\mathbb{E} \prod_{j=1}^k |X_j|^{\alpha_j} \right) \left(\mathbb{E} \prod_{j=k+1}^N |X_j|^{\alpha_j} \right) \quad (1)$$

and in particular

$$\mathbb{E} \prod_{j=1}^N |X_j|^{\alpha_j} \geq \prod_{j=1}^N \mathbb{E} |X_j|^{\alpha_j}. \quad (2)$$

In other words, among centered complex Gaussian vectors $(X_1, \dots, X_N) \in \mathbb{C}^N$ with fixed diagonal covariance (i.e. $(\mathbb{E}|X_j|^2)_{j \leq N}$ fixed) the expectation of $\prod_{j=1}^N |X_j|^{\alpha_j}$ is minimal when the variables are independent.

Inequality (2) is an extension of an inequality of Arias de Reyna [1], who established the particular case where all the $\alpha_j = 2p_j$ are even integer by rewriting the left hand side in terms of a permanent of a $2m$ matrix ($m = \sum p_j$) and using an inequality for permanents due to Lieb. Actually, Inequality (1) in the case where the α_j are even integers is equivalent to Lieb's permanent inequality, so in particular we are giving a new proof of this inequality.

In the next section we will introduce the tools that will be used in the proof, that is two semi-groups: the usual Ornstein-Uhlenbeck semi-group and another natural semi-group associated to the $\bar{\partial}$ operator (the generator of which could be called, depending from the context, Landau or magnetic Laplacian). In the last section we give the proof of our correlation inequality.

2 Semi-groups

To get the result, we will let the circular-symmetric psh function evolve along the Ornstein-Uhlenbeck semi-group on $\mathbb{C}^n = \mathbb{R}^n + i\mathbb{R}^n \simeq \mathbb{R}^{2n}$ associated with the measure γ and the scalar product : $\langle w, w' \rangle = \Re(w \cdot \overline{w'})$. We recall that its generator is given, for smooth functions f , writing $w = x + iy$, $x, y \in \mathbb{R}^n$, by

$$\begin{aligned} L^{\text{ou}} f(w) &:= \frac{1}{4} \Delta f(w) - \frac{1}{2} \langle w, \nabla \rangle f(w) \\ &= \sum_{j=1}^n \frac{1}{4} \left(\partial_{x_j x_j}^2 f(w) + \partial_{y_j y_j}^2 f(w) \right) - \frac{1}{2} (x_j \partial_{x_j} f(w) + y_j \partial_{y_j} f(w)). \end{aligned}$$

Note that the normalization differs slightly from the usual one on \mathbb{R}^{2n} because our Gaussian measure has complex covariance Id_n but real covariance equal to $\frac{1}{2} \text{Id}_{2n}$. Accordingly, the spectrum of $-L^{\text{ou}}$ on $L^2(\gamma)$ is here $\frac{1}{2} \mathbb{N} = \{0, \frac{1}{2}, 1, \dots\}$. The Ornstein-Uhlenbeck semi-group $P_t^{\text{ou}} = e^{tL^{\text{ou}}}$ admits the representation, for suitable functions $f : \mathbb{C}^n \rightarrow \mathbb{C}$,

$$P_t^{\text{ou}} f(z) = \int f(e^{-t/2} z + \sqrt{1 - e^{-t}} w) d\gamma(w) \quad (3)$$

$$= \pi^{-n} (1 - e^{-t})^{-n} \int f(w) e^{-\frac{1}{1-e^{-t}} |w - e^{-t/2} z|^2} d\ell(w) \quad (4)$$

As usual in semi-group methods, it is convenient to work with a nice stable space of functions. Here, we can for instance consider

$$\mathcal{G}^\infty := \{f \in C^\infty(\mathbb{C}^n); f \text{ and all its derivatives belong to } \mathcal{G}\}.$$

Note that for $f \in \mathcal{G}$, we can define $P_t^{\text{ou}} f$ by (3) and then we have that $(t, z) \mapsto P_t^{\text{ou}} f(z)$ is smooth on $(0, \infty) \times \mathbb{C}^n$, with $P_t^{\text{ou}} f \in \mathcal{G}^\infty$ and $\partial_t P_t^{\text{ou}} f = L^{\text{ou}} P_t^{\text{ou}} f$ for every $t > 0$, and $P_t^{\text{ou}} f \rightarrow f$ in $L^2(\gamma)$ as $t \rightarrow 0$. We refer to [3] for details. Let us also mention here that with formula (3) it is readily checked that properties like convexity, subharmonicity, pluri-subharmonicity are preserved along P_t^{ou} .

Another natural operator will be used. Indeed, since pluri-subharmonicity is characterized through a “ $\partial_{\bar{z}\bar{z}}$ operation”, we shall also use the following differential operator on smooth functions f on \mathbb{C}^n :

$$Lf = \sum_{j=1}^n \left(\partial_{z_j \bar{z}_j}^2 f - \bar{z}_j \partial_{\bar{z}_j} f \right) = \sum_{j=1}^n e^{|z|^2} \partial_{z_j} (e^{-|z|^2} \partial_{\bar{z}_j} f)$$

Note that $Lf = 0$ if (and only if, see below) f is holomorphic. Formally $L = -\bar{\partial}^* \bar{\partial}$ on $L^2(\gamma)$ equipped with the Hermitian structure $(f, g) = \int f \bar{g} d\gamma$. More precisely, denoting for a differentiable function

$$\partial_{\bar{z}} f = (\partial_{\bar{z}_1} f, \dots, \partial_{\bar{z}_n} f) \quad \text{and} \quad \partial_z f := (\partial_{z_1} f, \dots, \partial_{z_n} f)$$

we have the following standard fact.

Fact 4 (Integration by parts). *For regular enough functions $f, g : \mathbb{C}^n \rightarrow \mathbb{C}$, for instance for functions in \mathcal{G}^∞ , we have*

$$\int (Lf) \bar{g} d\gamma = - \int \partial_{\bar{z}} f \cdot \overline{\partial_{\bar{z}} g} d\gamma \quad (5)$$

and in particular $\int (Lf) \bar{f} d\gamma = - \int |\partial_{\bar{z}} f|^2 d\gamma \leq 0$. We can also write

$$\int (Lf) g d\gamma = - \int \partial_{\bar{z}} f \cdot \partial_z g d\gamma = \int f(\bar{L}g) d\gamma \quad (6)$$

where $\bar{L}f := \sum_{j=1}^n \left(\partial_{z_j \bar{z}_j}^2 f - z_j \partial_{z_j} f \right)$.

Indeed, it suffices to sum over j the equations

$$\int [e^{|z|^2} \partial_{z_j} (e^{-|z|^2} \partial_{\bar{z}_j} f)] \bar{g} d\gamma = \pi^{-n} \int \partial_{z_j} (e^{-|z|^2} \partial_{\bar{z}_j} f) \bar{g} d\lambda(z) = - \int \partial_{\bar{z}_j} f \partial_{z_j} \bar{g} d\gamma = - \int \partial_{\bar{z}_j} f \overline{\partial_{z_j} g} d\gamma.$$

The assumption that $f, g \in \mathcal{G}^\infty$ guarantees that the boundary terms (at infinity) in the integration by parts vanish.

As a consequence, we see that the Gaussian measure γ is invariant for L , and actually that L is a symmetric nonpositive operator on $L^2(\gamma)$ with the above-mentioned Hermitian structure. The kernel of L is the Bargmann space \mathcal{H}_0 formed by the holomorphic functions on \mathbb{C}^n that belong to $L^2(\gamma)$.

We want to work with the semi-group $P_t = e^{tL}$ which is also Hermitian (formally):

$$\int (P_t f) \bar{g} d\gamma = \int f \overline{P_t g} d\gamma \quad (7)$$

Although we will not explicitly use it, let us discuss a bit the (well known) spectral analysis of L on the complex Hilbert space $L^2(\gamma)$. This analysis is indeed fairly standard using the ideas introduced by Landau. Following for instance the presentation given in [4, Section 4], consider the ‘‘annihilation’’ operators $a_j = \partial_{\bar{z}_j}$ and their adjoints, the ‘‘creation’’ operators $b_j := a_j^* = \bar{z}_j - \partial_{z_j}$. Then $L = - \sum_{j \leq n} b_j \circ a_j$, with $[a_j, b_j] = 1$, and all these operators commute for distinct indices j . Plainly, if a function f and a scalar $\lambda \in \mathbb{C}$ satisfy $-Lf = \lambda f$, then $-L(a_j f) = (\lambda - 1)a_j f$ and $-L(b_j f) = (\lambda + 1)b_j f$. This implies that the spectrum of $-L$ is \mathbb{N} and that the eigenspace associated to the eigenvalue $k \in \mathbb{N}$ is given by the sum of the spaces $b^m \mathcal{H}_0$ with $m = (m_1, \dots, m_n) \in \mathbb{N}^n$, $|m| := \sum_{j \leq n} m_j = k$ and the convention $b^m := b_1^{m_1} \circ \dots \circ b_n^{m_n}$. Moreover, if we introduce the classical projection $\Pi_0 : L^2(\gamma) \rightarrow \mathcal{H}_0$ onto holomorphic functions

$$\Pi_0 f(z) := \int f(w) e^{z \cdot \bar{w}} d\gamma(w) = \int f(z + w) e^{-\bar{z} \cdot w} d\gamma(w),$$

then the projector Π_k on the k -eigenspace can be expressed in terms of Π_0 and the creation and annihilation operators. This allows to compute the reproducing kernel of Π_k , in terms of classical families of orthogonal polynomials. Next, one can sum over k and obtain the kernel $K_t(z, w)$ for $e^{tL} = \sum_k e^{-kt} \Pi_k$. Only the formula for K_t will be useful in the sequel and we shall actually *check* below that this suggested formula is indeed the kernel of e^{tL} .

An explicit formula for K_t can be found in [2]: setting

$$K_t(z, w) := \frac{1}{\pi^n (1 - e^{-t})^n} \exp \left(z \cdot \bar{w} - \frac{e^{-t} |z - w|^2}{1 - e^{-t}} - |w|^2 \right), \quad (8)$$

then

$$\begin{aligned} P_t f(z) &= \int f(w) K_t(z, w) d\ell(w) \\ &= (1 - e^{-t})^{-n} \int f(w) e^{z \cdot \bar{w} - \frac{e^{-t} |z - w|^2}{1 - e^{-t}}} d\gamma(w) \end{aligned} \quad (9)$$

Next, let us note that by performing the change of variable $w = z + \sqrt{1 - e^{-t}} \xi$ for fixed z we find

$$P_t f(z) = \int f(z + \sqrt{1 - e^{-t}} \xi) e^{-\sqrt{1 - e^{-t}} \bar{z} \cdot \xi} d\gamma(\xi). \quad (10)$$

On this formula, we see immediately that $P_0 = \text{Id}$ and $P_\infty = \Pi_0$.

To avoid discussions regarding unbounded operators and existence of semi-groups, we will proceed in the opposite direction and use the previous formula to *define* $P_t f$. Actually, to be fair, we should mention that later, in the proof of our result, we only need to work with smooth functions $f \in \mathcal{G}^\infty$; these functions provide nice initial data, ensuring existence and uniqueness of strong solutions for the semi-group equation. Nevertheless, we feel it is of independent interest to start from the integral formula (9) or (10) and derive from it the semi-group properties; we will first check that (9) solves (11) indeed. The drawback is that some properties that are obvious (formally) for $e^{tL} f$ need to be checked thoroughly when using this kernel representation, in particular because the kernel (8) is not Markovian.

Formulas (9)-(10) make sense pointwise, for $z \in \mathbb{C}^n$ fixed, as long as $f \in L^2(\gamma)$; actually we have the pointwise estimate $|P_t f(z)| \leq C_t(z) \|f\|_{L^2(\gamma)}$ for some constant $C_t(z) > 0$ depending on t and z . In order to derive some properties of $P_t f$, a stronger integrability condition (as $f \in \mathcal{G}$) will be assumed.

Lemma 5. *Given $f \in \mathcal{G}$, let us define $P_t f$ using the formula (9) or (10). Then one has that $(t, z) \rightarrow P_t f(z)$ is smooth on $(0, \infty) \times \mathbb{C}^n$, with for any $t > 0$, $P_t f \in \mathcal{G}_\infty$ and*

$$\partial_t P_t f = L P_t f. \quad (11)$$

Moreover we have

$$\|P_t f\|_{L^2(\gamma)} \leq \|f\|_{L^2(\gamma)}$$

and $P_t f \rightarrow f$ in $L^2(\gamma)$ as $t \rightarrow 0$.

Proof. It is readily checked that, for any fixed $w \in \mathbb{C}^n$ one has

$$\partial_t K_t(\cdot, w) = L K_t(\cdot, w).$$

Moreover, for $T, R, k > 0$ fixed, there exists constants $c = c(T, R, k), C = C(T, R, k)$ such that for $F = \partial K_t$, or $F = K_t$ or else F being any of the first k th partial derivatives of K_t , it holds that $|F(z, w)| \leq C e^{-c|w|^2}$ for all $w \in \mathbb{C}^n$, all $t \in (0, T)$ and all $|z| \leq R$. From this and the definition of \mathcal{G} , we can call upon dominated convergence to conclude to the smoothness of $(t, z) \rightarrow P_t f(z)$ and to the fact that $P_t f \in \mathcal{G}^\infty$ with $\partial_t P_t f = L P_t f$.

Regarding the contraction property, we want to avoid direct computations or spectral arguments, and so we make a detour and use some obvious but important properties of P_t .

First, we will use the semi-group property: for $t, s > 0$ that $P_t \circ P_s = P_{t+s}$. This can be seen in two ways. One can invoke that for a *smooth* function $F := P_{t_0} f \in \mathcal{G}^\infty$ then there is existence and uniqueness for the equation $\partial_t F_t = L F_t$ with initial condition $F_0 = F$ and we have seen that $(P_{t_0+t} f)$ is a solution; from there one can conclude. Or else, in a more pedestrian way, one can check that

$$\int K_t(z, w) K_s(\xi, z) d\ell(z) = K_{t+s}(\xi, w).$$

For this, one may use that given $z, \xi \in \mathbb{C}^n$ and $c \in \mathbb{C}$ with $\Re(c) > 0$,

$$\int e^{z \cdot w + \xi \cdot \bar{w}} e^{-c|w|^2} d\ell(w) = \pi^n c^{-n} e^{z \cdot \xi / c}. \quad (12)$$

Next, recall that P_t is Hermitian, in the sense (7), on $\mathcal{G} \subset L^2(\gamma)$; this can be seen directly from the integral formula since $K_t(z, w) = \overline{K_t(w, z)}$. Finally, we will use that $\|P_t f\|_{L^2(\gamma)}$ decreases for $t \in (0, \infty)$. This is immediate from the non-positivity of L since for $t > 0$ we have $P_t f \in \mathcal{G}^\infty$ and

$$\frac{d}{dt} \int |P_t f|^2 d\gamma = 2 \int (L P_t f) \overline{P_t f} d\gamma = -2 \int |\partial_{\bar{z}} P_t f|^2 d\gamma \leq 0.$$

So for our $f \in \mathcal{G}$ and $t > 0$ we have

$$\|P_t f\|_{L^2(\gamma)}^2 = \int (P_{2t} f) \bar{f} d\gamma \leq \|P_{2t} f\|_{L^2(\gamma)} \|f\|_{L^2(\gamma)} \leq \|P_t f\|_{L^2(\gamma)} \|f\|_{L^2(\gamma)}$$

which implies the contraction property in $L^2(\gamma)$.

To prove the continuity at $t = 0$ in $L^2(\gamma)$ we first assume that f is smooth and compactly supported. Using (10) we see that $P_t f$ converge point-wise to f and that for $t \in (0, 1)$ we have $|P_t f(z)| \leq C e^{c|z|}$ for some constant $c, C > 0$; so we can conclude by dominated convergence. For $f \in \mathcal{G}$ and $\epsilon > 0$, introduce g smooth compactly supported such that $\|f - g\|_{L^2(\gamma)} \leq \epsilon$ and let $\delta > 0$ be such that $t \leq \delta$ ensures that $\|P_t g - g\|_{L^2(\gamma)} \leq \epsilon$ holds. For $t \leq \delta$,

$$\|P_t f - f\|_{L^2(\gamma)} \leq \|P_t f - P_t g\|_{L^2(\gamma)} + \|P_t g - g\|_{L^2(\gamma)} + \|g - f\|_{L^2(\gamma)} \leq 2\|f - g\|_{L^2(\gamma)} + \epsilon \leq 3\epsilon.$$

This establishes the desired continuity. \square

Remark 6 (Contraction property).

1. We observe that some results, which can be deduced from the spectral decomposition and Hilbertian analysis, may be derived in a soft way thanks to flexible semigroup techniques. We have proved that starting from formula (9) we have $\|P_t f\|_{L^2(\gamma)} \leq \|f\|_{L^2(\gamma)}$ on the dense subspace \mathcal{G} , which together with the pointwise estimate given just before the previous Lemma implies by density that

$$\|P_t\|_{L^2(\gamma) \rightarrow L^2(\gamma)} \leq 1.$$

Formally, by letting $t \rightarrow \infty$ in $\|P_t\|_{L^2(\gamma) \rightarrow L^2(\gamma)} \leq 1$ we recover that

$$\|\Pi_0\|_{L^2(\gamma) \rightarrow L^2(\gamma)} \leq 1.$$

Actually, the convergence of P_t towards Π_0 can be quantified rigorously through Hörmander's inequality,

$$\|\varphi - \Pi_0 \varphi\|_{L^2(\gamma)}^2 \leq \|\partial_{\bar{z}} \varphi\|_{L^2(\gamma)}^2 = \int (-L\varphi) \varphi d\gamma$$

valid for any suitable φ , for instance for $\varphi \in \mathcal{G}^\infty$. Note that from formula (10), P_t acts as the identity on holomorphic functions, so $P_t \Pi_0 = \Pi_0 P_t = \Pi_0$. A classical Grönwall type argument (using the previous Lemma to justify the computation of $\frac{d}{dt} \int |P_t(f - \Pi_0 f)|^2 d\gamma$) ensures that for $f \in \mathcal{G}$ and $t \geq 0$

$$\|P_t f - \Pi_0 f\|_{L^2(\gamma)}^2 \leq e^{-2t} \|f - \Pi_0 f\|_{L^2(\gamma)}^2.$$

2. In analogy with the Markovian case P_t^{ou} we may wonder if P_t is also a contraction on some $L^p(\gamma)$. However, for any $p \neq 2$ we have

$$\|P_t\|_{L^p(\gamma) \rightarrow L^p(\gamma)} = +\infty,$$

as it can be seen by taking, in dimension $n = 1$, for $a \in \mathbb{R}$,

$$f_a(w) := e^{aw + \bar{w}}, \quad w \in \mathbb{C}.$$

Indeed, repeated applications of (12) with $c = 1$ show, setting $s_t := \sqrt{1 - e^{-t}}$ and using (10), that $P_t f_a(z) = e^{s_t^2 a} e^{az + (1 - s_t^2) \bar{z}}$ and that

$$\frac{\|P_t f_a\|_{L^p(\gamma)}^p}{\|f_a\|_{L^p(\gamma)}^p} = C(t, p) e^{a s_t^2 (p - p^2/2)}.$$

The next result describes how derivatives and P_t commute, an important issue in semi-group methods.

Lemma 7 (Commutation relations). *For any suitable f , say $f \in \mathcal{G}^\infty$, and $t > 0$ we have for every $1 \leq j \leq n$ and $z \in \mathbb{C}^n$,*

$$\partial_{z_j}(P_t f(z)) = P_t(\partial_{z_j} f)(z) \quad \text{and} \quad \partial_{\bar{z}_j}(P_t f(z)) = e^{-t} P_t(\partial_{\bar{z}_j} f)(z).$$

Proof. We use (10). The first equality is obvious. For the second one, setting $s_t = \sqrt{1 - e^{-t}}$, we have

$$\partial_{\bar{z}_j}(P_t f)(z) = P_t(\partial_{\bar{z}_j} f)(z) - s_t \int f(z + s_t \xi) e^{-s_t \bar{z}_j \xi} \xi_j d\gamma(\xi),$$

and

$$\begin{aligned} \pi^{-n} \int f(z + s_t \xi) e^{-s_t \bar{z}_j \xi} \xi_j e^{-\xi \cdot \bar{\xi}} d\ell(\xi) &= -\pi^{-n} \int f(z + s_t \xi) e^{-s_t \bar{z}_j \xi} \partial_{\xi_j}(e^{-\xi \cdot \bar{\xi}}) d\ell(\xi) \\ &= s_t \int (\partial_{\bar{z}_j} f)(z + s_t \xi) e^{-s_t \bar{z}_j \xi} d\gamma(\xi) = s_t P_t(\partial_{\bar{z}_j} f)(z). \end{aligned}$$

□

Now, and for the rest of this section, we focus on the case of circular-symmetric functions. Given $\theta \in \mathbb{R}$ and a function f we denote f_θ the function $f_\theta(w) = f(e^{i\theta} w)$. Note that

$$P_t(f_\theta) = (P_t f)_\theta. \quad (13)$$

Recall that a function f is said to be circular-symmetric if $f_\theta = f$ for every θ . It is worth noting that a holomorphic function on \mathbb{C}^n is necessarily constant when circular-symmetric. Indeed if $h : \mathbb{C} \rightarrow \mathbb{C}$ has both properties then invariance and the Cauchy formula give $h(1) = \int_0^{2\pi} h(e^{i\theta}) d\theta / (2\pi) = h(0)$; next if $f : \mathbb{C}^n \rightarrow \mathbb{C}$ is holomorphic and circular symmetric, then for any $(z_1, \dots, z_n) \in \mathbb{C}^n$, the function $h : z \in \mathbb{C} \mapsto f(z z_1, \dots, z z_n)$ is also holomorphic and circular-symmetric, hence $f(z_1, \dots, z_n) = h(1) = h(0) = f(0)$. Accordingly if $f \in L^2(\gamma)$ is circular-symmetric then $\Pi_0 f \equiv \int f d\gamma$ since the gaussian density is also circular-symmetric. Actually, much more can be said, as we shall see below.

Let us first investigate the relation between L and L^{ou} . Note that one can write

$$L f = L^{\text{ou}} f + \frac{i}{2} \sum_{j=1}^n (y_j \partial_{x_j} f - x_j \partial_{y_j} f). \quad (14)$$

So we have $L^{\text{ou}} = \Re(L) = \frac{L + \bar{L}}{2}$ where $\bar{L} f = \sum_{j=1}^n \left(\partial_{z_j}^2 \bar{z}_j f - z_j \partial_{z_j} f \right)$ has a kernel formed by the anti-holomorphic functions. The operators L and \bar{L} are Hermitian symmetric, whereas L^{ou} is symmetric for the real and the Hermitian product, and preserves the subspace of real valued functions. As we said, its spectrum is $-\frac{1}{2}\mathbb{N}$, as can be seen also from the formula $L^{\text{ou}} = \frac{L + \bar{L}}{2}$. Let us illustrate this on two examples, obtained by applying the creation operator $b_1 = \bar{z}_1 - \partial_{z_1}$ to the holomorphic functions $z \mapsto 1$, and $z \mapsto z_1$. The function $z \mapsto \bar{z}_1$ is an eigenfunction for L with eigenvalue -1 , for \bar{L} with eigenvalue 0 , and for L^{ou} with eigenvalue $-1/2$. The function $z \mapsto |z_1|^2 - 1$ is an eigenfunction for L with eigenvalue -1 , for \bar{L} with eigenvalue -1 , and for L^{ou} with eigenvalue -1 .

The special role played by circular-symmetric functions is due to the fact that these operators, and the associated semi-groups, coincide for them.

Lemma 8 (Action of L and P_t on circular-symmetric functions). *If f is a smooth circular-symmetric function, then we have*

$$L f = \bar{L} f = L^{\text{ou}} f.$$

In particular we have, when $f, g \in \mathcal{G}^\infty$ and f is circular-symmetric,

$$\int (L f) g d\gamma = \int f L g d\gamma = - \int \partial_z f \cdot \partial_{\bar{z}} g d\gamma.$$

Also, if $f \in L^2(\gamma)$ is circular-symmetric then we have

$$P_t f = P_t^{\text{ou}} f$$

for every $t \geq 0$.

Proof. Writing $w = x + iy$, $x, y \in \mathbb{R}^n$, the symmetry rewrites as $f((\cos(\theta)x - \sin(\theta)y) + i(\cos(\theta)y + \sin(\theta)x)) = f(x + iy)$. Taking the derivative at $\theta = 0$ we find

$$\sum_{j=1}^n \left(-y_j \partial_{x_j} f(x + iy) + x_j \partial_{y_j} f(x + iy) \right) = 0,$$

and this for every $x, y \in \mathbb{R}^n$. This implies in view of (14) that $Lf = L^{\text{ou}} f = \bar{L}f$. Next, for any smooth function g we have, using that $Lf = \bar{L}f$ and (6),

$$\int (Lf)g \, d\gamma = \int (\bar{L}f)g \, d\gamma = \int fLg \, d\gamma = - \int \partial_z f \cdot \partial_{\bar{z}} g \, d\gamma.$$

Although it is formally trivial that equality of L and L^{ou} on circular-symmetric functions implies equality of the semi-groups P_t and P_t^{ou} , a bit more should be said since we defined the semi-group using the explicit formula (9). And it is anyway instructive to compute the kernels on circular-symmetric functions. Denote by K_t^{ou} the kernel of the Ornstein-Uhlenbeck semi-group that we recalled above: $K_t^{\text{ou}}(z, w) = \pi^{-n} (1 - e^{-t})^{-n} e^{-\frac{1}{1-e^{-t}}|w - e^{-t/2}z|^2}$. So we have, setting $c_t := e^{-t/2}$ and $s_t := \sqrt{1 - e^{-t}}$,

$$K_t^{\text{ou}}(z, w) = \pi^{-n} s_t^{-2n} e^{-s_t^{-2}|w|^2 - s_t^{-2}c_t^2|z|^2} e^{s_t^{-2}c_t(w \cdot \bar{z} + \bar{w} \cdot z)}$$

and

$$K_t(z, w) = \pi^{-n} s_t^{-2n} e^{-s_t^{-2}|w|^2 - s_t^{-2}c_t^2|z|^2} e^{s_t^{-2}(c_t^2 w \cdot \bar{z} + \bar{w} \cdot z)}.$$

Note that only the last exponentials differ in these two formulas. When f is circular-symmetric, in order to check that $P_t f = P_t^{\text{ou}} f$ it suffices to check that for fixed w, z, t one has

$$\frac{1}{2\pi} \int_0^{2\pi} K_t(z, e^{i\theta} w) \, d\theta = \frac{1}{2\pi} \int_0^{2\pi} K_t^{\text{ou}}(z, e^{i\theta} w) \, d\theta.$$

Observe that for $a, b \in \mathbb{C}$, we have

$$\frac{1}{2\pi} \int_0^{2\pi} e^{ae^{i\theta} + be^{-i\theta}} \, d\theta = \frac{1}{2\pi} \int_0^{2\pi} \sum_{p, q \in \mathbb{N}} \frac{a^p b^q}{p! q!} e^{i(p-q)\theta} \, d\theta = \sum_{n \geq 0} \frac{(ab)^n}{(n!)^2} = B(ab)$$

with $B(x) := \sum_{n \geq 0} \frac{x^n}{(n!)^2}$. Therefore, we have

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} K_t(z, e^{i\theta} w) \, d\theta &= \pi^{-n} s_t^{-2n} e^{-s_t^{-2}|w|^2 - s_t^{-2}c_t^2|z|^2} B(s_t^{-4}c_t^2|w \cdot \bar{z}|^2) \\ &= \frac{1}{2\pi} \int_0^{2\pi} K_t^{\text{ou}}(z, e^{i\theta} w) \, d\theta, \end{aligned}$$

as wanted. □

3 Proof of Theorem 1

First, let us note that we can assume that g is smooth, and actually that $g \in \mathcal{G}^\infty$. Indeed, if $g \in \mathcal{G}$ then $P_t^{\text{ou}} g \in \mathcal{G}^\infty$ and we mentioned that $P_t^{\text{ou}} g$ converges to g in $L^2(\gamma)$ and therefore also in $L^1(\gamma)$, as $t \rightarrow 0$. Consequently, if we know the conclusion for a function in \mathcal{G}^∞ , then

$$\int f P_t^{\text{ou}} g \, d\gamma \geq \int f \, d\gamma \int P_t^{\text{ou}} g \, d\gamma$$

and by passing to the limit when $t \rightarrow 0$ we know it also for $g \in \mathcal{G}$. For the same reason, we can assume that $f \in \mathcal{G}^\infty$, recalling that $P_t^{\text{ou}}f$ is circular-symmetric when f is.

So in the sequel, we are given two psh functions $f, g \in \mathcal{G}^\infty$, with f circular-symmetric.

As in the proof of the correlation for convex functions [6], we will compute some kind of second derivative in t for integrals involving $P_t^{\text{ou}}f$; recall that $\partial_t P_t^{\text{ou}}f = L^{\text{ou}}P_t^{\text{ou}}f$. The novelty is that, along the way, we will also use $P_t f$ which satisfies $\partial_t P_t f = L P_t f$ (Lemma 5).

Consider

$$\alpha(t) := \int (P_t^{\text{ou}}f) g d\gamma = \int (P_t f) g d\gamma \in \mathbb{R}.$$

The function α is, by construction, smooth on $(0, \infty)$ and continuous on $[0, \infty)$ (see Lemma 5 for the continuity at zero). Since $P_t^{\text{ou}}f$ tends to the constant $\int f d\gamma$ when $t \rightarrow \infty$, we have

$$\alpha(t) \rightarrow \int f d\gamma \int g d\gamma$$

In order to conclude, it suffices to show that α decreases. Actually we will prove that α is convex; which is enough, since a convex function with a bounded limit at $+\infty$ cannot increase. It holds

$$\alpha'(t) = \int L^{\text{ou}}(P_t^{\text{ou}}f) g d\gamma = \int L P_t f g d\gamma. \quad (15)$$

Since $P_t f$ is also circular-symmetric, we can invoke Lemma 8 and write

$$\alpha'(t) = - \int \partial_z P_t f \cdot \partial_{\bar{z}} g d\gamma.$$

Next, using the first commutation relation from Lemma 7 we get

$$\alpha'(t) = - \sum_{j=1}^n \int P_t(\partial_{z_j} f) \partial_{\bar{z}_j} g d\gamma$$

We stress that $\partial_z f$ is no longer circular-symmetric, so we cannot exchange P_t and P_t^{ou} . The second derivative of α is, using Fact 4,

$$\alpha''(t) = - \sum_{j=1}^n \int L(P_t(\partial_{z_j} f)) \partial_{\bar{z}_j} g d\gamma = \sum_{j=1}^n \int \partial_{\bar{z}}(P_t(\partial_{z_j} f)) \cdot \partial_z(\partial_{\bar{z}_j} g) d\gamma = \sum_{j,k=1}^n \int \partial_{\bar{z}_k} P_t(\partial_{z_j} f) \partial_{z_k \bar{z}_j}^2 g d\gamma.$$

Using the commutation relation from Lemma 7 we can write

$$\alpha''(t) = \sum_{j,k=1}^n \int \partial_{\bar{z}_k z_j}^2 (P_t f) \partial_{z_k \bar{z}_j}^2 g d\gamma = \int \text{Tr} \left((D_{\mathbb{C}}^2 P_t f)(z) (D_{\mathbb{C}}^2 g)(z) \right) d\gamma$$

where for a C^2 function h on \mathbb{C}^n the notation $D_{\mathbb{C}}^2 h(z)$ refers to the Hermitian matrix $\left(\partial_{z_j \bar{z}_k}^2 h(z) \right)_{j,k \leq n}$.

Since $P_t f = P_t^{\text{ou}}f$ and g are psh, the corresponding matrices are nonnegative Hermitian matrices, which means that the trace of their product is still nonnegative. This shows that $\alpha'' \geq 0$ and finishes the proof of the theorem. \square

We would like to conclude with a discussion of the differences between the real case and the complex case. After all, we are computing second derivatives of the same object

$$\alpha(t) = \int (P_t^{\text{ou}}f) g d\gamma$$

along the Ornstein-Uhlenbeck semi-group exactly as in the case of convex functions, so what is going on?

In both cases we prove that α decreases by showing that $\alpha' \leq 0$ using the next derivative somehow, but we compute these derivatives differently. The argument for convex function goes as follows. A direct computation and usual commutation properties show that, if $\int \nabla f d\gamma = 0$, which is the case when f is even, then

$$\alpha'(t) = -e^{-t/2} \int_t^\infty \left(\int \operatorname{Tr} \left((D^2 P_s^{\text{ou}} f)(z) (D^2 g)(z) \right) d\gamma(z) \right) e^{s/2} ds$$

where D^2 refers to the usual (real) Hessian on \mathbb{R}^{2n} ; from this we conclude to the correlation for convex functions. On the other hand, we have proved, when f is circular-symmetric, that

$$\alpha'(t) = - \int_t^\infty \left(\int \operatorname{Tr} \left((D_{\mathbb{C}}^2 P_s^{\text{ou}} f)(z) (D_{\mathbb{C}}^2 g)(z) \right) d\gamma(z) \right) ds.$$

Note that we have used here that $\alpha'(t)$ tends to 0 when $t \rightarrow +\infty$: this follows from the fact that α is convex and has a finite limit at $+\infty$, and can also be seen from (15) since $P_t^{\text{ou}} f$ tends to a constant when $t \rightarrow +\infty$. It is because we wanted to work with complex derivatives that we aimed at inserting L in place of L^{ou} ; recall that $\partial_{z_j} f$ need not be circular-symmetric even when f is, although the second derivatives are again circular-symmetric.

Finally, let us observe that if we consider in dimension 1 the circular-symmetric psh functions $f(w) = |w|^{1/3}$ and $g(w) = |w|^4$ on $\mathbb{C} \simeq \mathbb{R}^2$, then, as we already mentioned,

$$\operatorname{Tr} \left((D_{\mathbb{C}}^2 f) (D_{\mathbb{C}}^2 g) \right) = \Delta f \Delta g \geq 0,$$

but a direct computation shows that

$$\operatorname{Tr} \left((D^2 f)(z) (D^2 g)(z) \right) = -\frac{4}{3} |z|^{1/3} \leq 0 \quad \forall z \in \mathbb{C}.$$

Of course, this discrepancy cannot hold at all times for $P_t^{\text{ou}} f$ in place of f (and moreover f is not smooth at zero, although this is not really an issue). But it suggests that the two formulas above for $\alpha'(t)$ are indeed quite different.

Acknowledgements: We thank Laurent Charles, Vincent Guedj and Pascal Thomas for useful discussions, and for pointing out to us relevant references.

References

- [1] J. Arias-de-Reyna, *Gaussian variables, polynomials and permanents*, Linear Algebra Appl. 285 (1998), no. 1-3, 107–114.
- [2] N. Askour, A. Intissar and Z. Mouayn, *Explicit formulas for reproducing kernels of generalized Bargmann spaces of \mathbb{C}^n* , J. Math. Phys. 41 (2000), no. 5, 3057–3067.
- [3] D. Bakry, I. Gentil, Ivan and M. Ledoux, *Analysis and geometry of Markov diffusion operators*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 348. Springer, Cham, 2014.
- [4] L. Charles, *Landau levels on a compact manifold*, preprint, Arxiv 2021.
- [5] L. Hörmander. *Notions of convexity*. Progress in Mathematics, 127. Birkhäuser Boston, Inc., Boston, MA, 1994. viii+414 pp.
- [6] Y. Hu, *Itô-Wiener chaos expansion with exact residual and correlation, variance inequalities*, J. Theoret. Probab. 10 (1997), no. 4, 835–848.
- [7] D. Malicet, I. Nourdin, G. Peccati, G. Poly, *Squared chaotic random variables: new moment inequalities with applications*, J. Funct. Anal. 270 (2016), no. 2, 649–670.

Franck Barthe: Institut de Mathématiques de Toulouse (UMR 5219). Université de Toulouse & CNRS. UPS, F-31062 Toulouse Cedex 9, France.

Dario Cordero-Erausquin: Institut de Mathématiques de Jussieu (UMR 7586), Sorbonne Université & CNRS, 75252 Paris Cedex 5, France.