



**HAL**  
open science

## Synthesis and further use of SF<sub>5</sub> -alkynes as platforms for the design of more complex SF<sub>5</sub> -containing products

Lucas Popek, Thi Mo Nguyen, Nicolas Blanchard, Dominique Cahard,  
Vincent Bizet

### ► To cite this version:

Lucas Popek, Thi Mo Nguyen, Nicolas Blanchard, Dominique Cahard, Vincent Bizet. Synthesis and further use of SF<sub>5</sub> -alkynes as platforms for the design of more complex SF<sub>5</sub> -containing products. *Tetrahedron*, 2022, 117-118, pp.132814. 10.1016/j.tet.2022.132814 . hal-03717188

**HAL Id: hal-03717188**

**<https://hal.science/hal-03717188v1>**

Submitted on 8 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Synthesis and further use of SF<sub>5</sub>-alkynes as platforms for the design of more complex SF<sub>5</sub>-containing products.

Lucas Popek,<sup>a</sup> Thi Mo Nguyen,<sup>b</sup> Nicolas Blanchard,<sup>a</sup> Dominique Cahard,<sup>b</sup> Vincent Bizet<sup>a,\*</sup>

a) Université de Haute-Alsace, Université de Strasbourg, CNRS, LIMA, UMR 7042, 68000 Mulhouse, France

b) CNRS, UMR 6014 COBRA, Normandie Université, 76821 Mont Saint Aignan, France

## Contents

Introduction

Synthesis of SF<sub>5</sub>-alkynes

By means of SF<sub>5</sub>Cl

By means of SF<sub>5</sub>Br

Cycloaddition reactions with SF<sub>5</sub>-alkynes

SF<sub>5</sub>-alkynes in Diels-Alder cycloadditions

SF<sub>5</sub>-alkynes in Hetero-Diels-Alder reactions

SF<sub>5</sub>-alkynes in 1,3 dipolar cycloadditions

Nucleophilic additions onto SF<sub>5</sub>-alkynes

Synthesis of *cis*- and *trans*-1-SF<sub>5</sub>-2-methoxy-ethene

Synthesis of SF<sub>5</sub>-substituted thienothiophene


Synthesis of 2-SF<sub>5</sub>-(aza)indoles via intramolecular cyclization

Hydrofunctionalization of SF<sub>5</sub> alkynes

Conclusion

## 1 Introduction

Pentafluorosulfanyl group is known as an emerging fluorinated group, it has been reported for the first time in 1960, and then remained an exotic fluorinated group for years, but nowadays this group is rising in interest and many groups and companies start to investigate this fluorinated motif in more depth. Indeed, SF<sub>5</sub> is known as a "super CF<sub>3</sub>", gathering and magnifying the benefits of CF<sub>3</sub> that are a large volume, a high electronegativity and a high lipophilicity, which are highly valuable properties for drugs development. Several general reviews on the synthesis and use of SF<sub>5</sub> compounds have been published recently.[1–6] In parallel, a few applications of SF<sub>5</sub>-containing molecules has begun to appear in materials science,[7,8] drug design[9] or crop protection[10] (Scheme 1), but innovation remains highly dependent on the synthetic methods available to introduce SF<sub>5</sub> in organic molecules. In this report, we turned our attention more specifically to SF<sub>5</sub>-alkynes, from their synthesis to applications.




Scheme 1 : Selected examples of applications of SF<sub>5</sub> containing molecules

## 2 Synthesis of SF<sub>5</sub>-alkynes

### 2.1 By means of SF<sub>5</sub>Cl


Pentafluorosulfanyl chloride (SF<sub>5</sub>Cl) is a commercially available gas and the most frequently used reagent to functionalize aliphatic compounds with an SF<sub>5</sub> group.[11] Moreover, a convenient synthesis of this gas has been reported recently starting from molecular sulfur, potassium fluoride and trichloroisocyanuric acid as oxidant; SF<sub>5</sub>Cl can be stored and used in hexanes solution.[12,13] SF<sub>5</sub>Cl adds to C=C or C≡C bonds under thermal, photochemical, or chemical activation; this latter activation being essentially conducted with the aid of Et<sub>3</sub>B.

Thermal reaction of an acetylene with SF<sub>5</sub>Cl led regioselectively to the corresponding SF<sub>5</sub>-chloro-olefin in relatively low yield. The subsequent dehydrohalogenation reaction is of great interest because it yields a polarized SF<sub>5</sub>-alkyne. In 1961, Case and co-workers reported the first example of SF<sub>5</sub>-alkyne by HCl elimination in basic conditions, which produced the SF<sub>5</sub>-propyne **3a** in a low 6% overall yield (Scheme 2).[14] Unfortunately this sequence was not applicable to other substrates.



Scheme 2 : Addition reaction of SF<sub>5</sub>Cl onto propyne and subsequent dehydrochlorination


In 1964, Hoover and Coffman described a four-step strategy through initial thermal addition of SF<sub>5</sub>Cl onto acetylene (Scheme 3).[15] Next, the photochemical addition of bromine on the chloro SF<sub>5</sub>-olefin **2b**, followed by base-mediated dehydrobromination afforded olefin **5b** as a mixture of isomers. A final step of reduction with zinc led to SF<sub>5</sub>-acetylene **3b** in high yield albeit with a low 11% overall yield.



Scheme 3: Synthesis of SF<sub>5</sub>-acetylene.


At that time, the harsh thermal conditions appeared as a strong limitation to diversify easily the scope of SF<sub>5</sub>-alkynes in decent yields. A revolution appeared in 2002, when Dolbier reported an easy and practical radical activation procedure based on Et<sub>3</sub>B-catalyzed addition of SF<sub>5</sub>Cl to terminal alkynes in hexane (Scheme 4).[16–19] The reaction proceeded smoothly at low temperature and provided the *trans*-1,2-addition products **2c-h** in high yields for alkylacetylenes and moderate yields with arylacetylenes. When intermediates **2** were treated with an excess of lithium hydroxide in DMSO, HCl

elimination occurred and delivered the corresponding SF<sub>5</sub>-alkynes **3c-g** in moderate to high yields, depending on the substitution pattern of alkynes.




Scheme 4: Dolbier's procedure for the synthesis of SF<sub>5</sub>-alkynes

This two-step strategy was also used by Röschenthaler and co-workers to synthesize liquid crystalline compounds with a terminal SF<sub>5</sub>-substituted acetylene moiety **3i-k** (Scheme 5).[20,21] In this case, hexane was replaced by dichloromethane and the reaction mixture was maintained at -20 °C for the SF<sub>5</sub>-addition step. The subsequent elimination step with 5 equivalents of LiOH required higher temperature (50 °C) and longer reaction time (12 h) compared to Dolbier's procedure showed in Scheme 4. In 2019, the Dolbier's method was illustrated by Paquin in relatively low overall yields (Scheme 6).[22]




Scheme 5: Synthesis of liquid crystals containing a SF<sub>5</sub>-alkyne



Scheme 6: Synthesis of various SF<sub>5</sub>-alkynes using Dolbier's procedure


In 2021, Bizet and co-workers modified the original Dolbier's method with 2-ethynyl aniline substrates **6** by performing the radical addition of SF<sub>5</sub>Cl in ethyl acetate, which gave the corresponding adducts **7** in high yields. Moreover, they developed a new elimination protocol for the synthesis of SF<sub>5</sub>-alkynes by using an excess of LiHMDS at -78 °C, affording SF<sub>5</sub>-2-ethynyl aniline derivatives **8** in high yields. These conditions proved to be tolerant to sensitive functional groups such as ester, nitrile, or halogens (Scheme 7).[23]



Scheme 7 : Synthesis of SF<sub>5</sub>-alkynes derived from 2-ethynyl anilines


## 2.2 By means of SF<sub>5</sub>Br

SF<sub>5</sub>Br is much more reactive than SF<sub>5</sub>Cl. However, one big limitation is that this gas is not commercially available nor easily manufactured. Nevertheless, it has been used in reactions with alkenes or functionalized double bond systems providing vicinal bromo-pentafluorosulfanyl alkanes, which illustrated the compatibility of the reagent with several functional groups, although the reagent is less selective compared to SF<sub>5</sub>Cl. Like its chlorinated analogue, SF<sub>5</sub>Br was used for addition onto different alkynes. Gard, Shreeve and co-workers improved the previous works of Case and Hoover and develop a straightforward synthesis by using of SF<sub>5</sub>Br instead of SF<sub>5</sub>Cl. In that case, the dehydrobromination was more efficient and gave the SF<sub>5</sub>-acetylene **3b** in 39% overall yield (Scheme 8).[24,25]




Scheme 8 : Alternative synthesis of SF<sub>5</sub>-acetylene with SF<sub>5</sub>Br

This methodology is not limited to acetylene but could be extended to diversely substituted alkynes [26–31] or diacetylenes [32,33]. Firstly, when diacetylene **10** reacted neat with one equivalent of SF<sub>5</sub>Br at -78 °C, it gave the mono-SF<sub>5</sub> adduct **11** in 50% yield, while the di-SF<sub>5</sub> adduct **13** was formed with two equivalents of SF<sub>5</sub>Br at -45 °C. Treatment with KOH at 25 °C delivered the mono- or disubstituted SF<sub>5</sub> diacetylenes **12** & **14**, in approximatively 50% yield in both cases (Scheme 9).




Scheme 9 : Synthesis of  $SF_5$ -diacetylenes using  $SF_5Br$

Similarly, addition reaction of  $SF_5Br$  onto several substituted alkynes has been reported and it was shown that the  $SF_5Br$  can react with electron deficient alkynes (**1w-y**) thanks to thermal activation to give the corresponding 2-bromo-1- $SF_5$ -alkenes **9** in moderate to good yields. Few  $SF_5$ -alkynes **3** were obtained in good yields by dehydrobromination with KOH (Scheme 10).[26–31]



Scheme 10 : Synthesis of  $SF_5$ -alkynes by bromopentafluorosulfanylation reaction and HBr elimination

In the 80s and 90s, Seppelt and Berry used these alkynes for coordination chemistry of nickel [34] or cobalt [35–37] complexes. More recently, in 2006, polymer chemists became interested in  $SF_5$  alkynes as monomer precursor and  $SF_5$ -ethynylthiophene **17** was synthesized with an excellent 78% yield over two steps with the use of NaOH for the dehydrobromination step; thus expanding the scope of alkynes to those featuring heteroaromatic substituents (scheme 11).[38]




Scheme 11 : Synthesis of  $SF_5$ -ethynylthiophene

### 3 Cycloaddition reactions with $SF_5$ -alkynes

#### 3.1 $SF_5$ -alkynes in Diels-Alder cycloadditions


The common "normal" electron demand Diels-Alder reaction takes place by HOMO/LUMO interactions between an electron-rich diene and an electron-deficient dienophile. The  $SF_5$  motif, which is a strong electron-withdrawing group makes  $SF_5$ -alkynes good electron-deficient dienophiles for such reaction.[1,2] The group of Dolbier has largely contributed to the use of  $SF_5$  alkynes as dienophiles in [4+2] cycloaddition reactions. For instance,  $SF_5$ -acetylene **3b** reacted with butadiene **18a** or 2,3-dimethylbuta-1,3-diene **18b**, to give  $SF_5$ -substituted cyclohexa-1,4-dienes **19** in moderate to good

yields. These intermediates **19** could give SF<sub>5</sub>-substituted benzenes **20** after aromatizing dehydrogenation either over platinum gauze at 575 °C or by refluxing with chloranil in *p*-xylene.[15] When *n*-Bu substituted SF<sub>5</sub>-alkyne **3g** was heated with cyclopentadiene **21** at 120 °C for 5 h, the norbornadiene derivative **22** was obtained in 50% yield (Scheme 12).[39]



Scheme 12 : Diels-Alder reactions of SF<sub>5</sub>-alkynes with 1,3-dienes


The electron-deficient alkyne **3w** was successfully used as versatile dienophile in [4+2] Diels-Alder cycloaddition with a series of cyclic 1,3-dienes **23** (scheme 13). The reactions occurred in DCM at room temperature or in refluxing toluene and provided the corresponding cycloadducts **24** in high yields in most cases. Starting from pyranone **23d** as diene, the corresponding cycloadduct **24d** was not stable to be isolated and instead the reaction yielded 1-SF<sub>5</sub>-2-CF<sub>3</sub>-benzene **25** after elimination of CO<sub>2</sub> in a spontaneous retro-Diels-Alder reaction. Similarly, the diene **26** was reacted with the SF<sub>5</sub>-alkyne **3w** to yield the fluoranthene **27** in 75% yield.[31]



Scheme 13 : Diels-Alder reactions of F<sub>3</sub>C-C≡C-SF<sub>5</sub> with cyclic 1,3-dienes

### 3.2 SF<sub>5</sub>-alkynes in hetero-Diels-Alder reactions


The preparation of SF<sub>5</sub>-substituted heterocycles from SF<sub>5</sub>-alkynes is scarce in the literature and only one example of hetero-Diels-Alder reaction has been described by Dolbier. The 3-butyl-4-SF<sub>5</sub>-furan **30** was obtained in 58% yield following a tandem Diels-Alder/retro-Diels-Alder sequence between SF<sub>5</sub>-hexyne **3g** and 4-phenyloxazole **28** (scheme 14).[17,39]



Scheme 14 : Synthesis of 4-SF<sub>5</sub>-furan derivative **30**

### 3.3 SF<sub>5</sub>-alkyne in 1,3 dipolar cycloadditions


The group of Dolbier also studied the formation of nitrogen-containing heterocycles from SF<sub>5</sub>-alkynes using 1,3-dipolar cycloaddition reactions. Reaction of *tert*-butyl aziridine **31** at high temperature generated the intermediate azomethine ylide **33**, which reacted with SF<sub>5</sub>-alkyne **3** to yield SF<sub>5</sub>-pyrroles **32** after oxidation with DDQ (scheme 15a).[18] Few years later, a similar strategy was used using another azomethine ylide precursor **34**, which was activated in softer conditions at room temperature with TFA and afforded SF<sub>5</sub>-pyrroles **35** in high yields (scheme 15b).[40] This method has been extended to the synthesis of SF<sub>5</sub>-thiophenes **38**, using SO<sub>2</sub>Cl<sub>2</sub> in a second step to re-aromatize the dihydrothiophene to thiophene in moderate yields (scheme 15c).



Scheme 15 : Synthesis of SF<sub>5</sub>-substituted pyrroles and thiophenes by 1,3-dipolar cycloadditions


Aryl SF<sub>5</sub>-alkynes **3** proved to be suitable substrates in 1,3-dipolar cycloadditions for the synthesis of 4-SF<sub>5</sub>-isoxazoles **41** and isooxazolines **43**, by reaction with benzohydroxyiminoyl chlorides **40** and *N*-benzylidene aniline *N*-oxides **42**, respectively (scheme 16).[19,41]





Scheme 16 : Preparation of 4-SF<sub>5</sub>-isoxazoles and isoxazolines via 1,3-dipolar cycloadditions

The synthesis of SF<sub>5</sub> pyrazoles **44** and **45** and triazoles **46** was reported by Shreeve and co-workers, using diazomethane or “click” chemistry process with azide, for applications in materials science.[42,43] The addition of diazomethane to SF<sub>5</sub>-acetylenes led to a 3:2-mixture of 3- and 4-SF<sub>5</sub>-pyrazoles **44** and **45**,[15] while the single regioisomer **45** was obtained from the triisopropylsilyl (TIPS) derivative **3v** in excellent yields (scheme 17a-b).[44] The Huisgen cycloaddition was successfully applied to SF<sub>5</sub>-acetylene and delivered triazole **46** in good yields (scheme 17c). Interestingly, the 2,4-diazido-6-diazomethyl-1,3,5-triazine **47** was heated with SF<sub>5</sub>-acetylene **3b** in ethanol to give selectively the pyrazole **48** in 59% yield, while the two azido groups remained unreactive under these conditions (scheme 17d).




Scheme 17 : Synthesis of SF<sub>5</sub>-pyrazoles and triazoles**4 Nucleophilic additions onto SF<sub>5</sub>-alkynes**


SF<sub>5</sub>-Substituted alkynes are highly polarized as a result of the strong electron-withdrawing property of the SF<sub>5</sub> that renders the β-carbon of the alkyne a highly reactive site for nucleophilic additions. However, for intramolecular process, the reaction can take place of the α-carbon driven by the formation of 5-membered heteroaromatics.

**4.1 Synthesis of *cis*- and *trans*-1-SF<sub>5</sub>-2-methoxy-ethene**


In 1964, Hoover and co-workers described the first reaction involving a SF<sub>5</sub>-alkyne to form an enol-ether.[15] Interestingly, they observed that treatment of 2-chloro-1-SF<sub>5</sub>-ethene **2b** with MeOH and KOH gave exclusively *trans*-1-SF<sub>5</sub>-2-methoxyethene **49** in 40% yield by nucleophilic substitution of the chlorine atom by a methoxy group under basic conditions. On the other hand, the reaction of SF<sub>5</sub>-acetylene **3b** in similar conditions provided mainly the opposite *cis*-isomer **50** in 65% yield with small amount of the *trans*-isomer **49** (scheme 18).

Scheme 18 : Synthesis of *trans*- and *cis*-1-SF<sub>5</sub>-2-methoxy-ethenes**4.2 Synthesis of SF<sub>5</sub>-substituted thienothiophene**

The SF<sub>5</sub>-ethynylthiophene **17**, depicted earlier in the text (see scheme 11), was engaged in halogen-lithium exchange using *n*-BuLi at -78 °C. Then, treatment with elemental sulfur gave the SF<sub>5</sub>-substituted thienothiophene **51** in 43% yield (scheme 19).[38]

Scheme 19 : Synthesis of SF<sub>5</sub>-substituted thienothiophene**4.3 Synthesis of 2-SF<sub>5</sub>-(aza)indoles via intramolecular cyclization**


Recently, the synthesis of 2-SF<sub>5</sub>-indoles **53** was developed by Bizet and co-workers starting from 2-ethynylaniline derivatives following a one-pot telescoped procedure.[23] Interestingly, it was shown that by treatment of **7** in classical basic conditions using an excess of LiOH in DMSO, the HCl elimination took place to form **8** but this step was rapidly followed by 5-*endo*-dig cyclization (**52**) and *N*-tosyl deprotection affording 2-SF<sub>5</sub>-(aza)indoles **53** in moderate to good and reproducible yields (Scheme 20a). The synthesis was then decomposed stepwise and K<sub>3</sub>PO<sub>4</sub> was identified as a suitable base to transform selectively **8** into **52**. Noteworthy, with electron-withdrawing substituents R, some non-negligible amount of unprotected indole **53** was also formed (Scheme 20b).



Scheme 20 : Synthesis of 2-SF<sub>5</sub>-(aza)indoles via a base-mediated 5-endo-dig cyclization


#### 4.4 Hydrofunctionalization of SF<sub>5</sub> alkynes

The gold-catalyzed hydrofluorination of SF<sub>5</sub>-alkyne has been reported on a single example obtained in a low 15% yield (**55b**), which is much lower than the 49% yield obtained for the CF<sub>3</sub>-analog **55a**.<sup>[45]</sup> The bulkiness of the SF<sub>5</sub> compared to the CF<sub>3</sub> was proposed to explain the lower reactivity of **3d** in these conditions (Scheme 21).



Scheme 21 : Gold-catalyzed hydrofluorination of R<sub>f</sub>-alkynes

In contrast, gold-catalyzed hydration of SF<sub>5</sub>-alkynes was much more efficient and delivered  $\alpha$ -SF<sub>5</sub> ketones **56** in yields ranging from 60 to 78% both from aromatic and aliphatic substrates (Scheme 22).<sup>[22]</sup>



Scheme 22 : Gold-catalyzed hydration of SF<sub>5</sub>-alkynes

## 5 Conclusion

Over the past sixty years, the synthetic access to variously decorated SF<sub>5</sub>-containing compounds has remained a highly challenging task. Within the past few years, the synthesis of SF<sub>5</sub>-alkynes has been significantly improved making a strong new impulse to the development of small SF<sub>5</sub>-building blocks. A wider scope of SF<sub>5</sub>-alkynes was synthesized and the application of these alkynes in synthetic transformations has been discussed in this article. In the coming years, many applications of SF<sub>5</sub>-small molecules are expected in various fields such as crop science, drug discovery or materials science. Development of new synthetic methods is always timely and relevant to bring molecular complexity a step further. This topic of synthetic organofluorine chemistry is rising and many innovative discoveries are expected in near future.

## 6 Acknowledgment

This work was supported by the French National Research Agency – ANR SulFive project (Grant No. ANR-21-CE07-0042), Normandie Université, the Université de Haute-Alsace, the Université de Strasbourg and the CNRS.

## 7 References

- [1] S. Altomonte, M. Zanda, Synthetic chemistry and biological activity of pentafluorosulphanyl (SF<sub>5</sub>) organic molecules, *J. Fluorine Chem.* 143 (2012) 57–93. <https://doi.org/10.1016/j.jfluchem.2012.06.030>.
- [2] P. R. Savoie, J. T. Welch, Preparation and Utility of Organic Pentafluorosulfanyl-Containing Compounds, *Chem. Rev.* 115 (2015) 1130–1190. <https://doi.org/10.1021/cr500336u>.
- [3] G. Haufe, Pentafluorosulfanylation of Aliphatic Substrates, in: D. Cahard, J. Ma (Eds.), *Emerging Fluorinated Motifs*, 1st ed., Wiley, 2020: pp. 571–609. <https://doi.org/10.1002/9783527824342.ch19>.
- [4] P. Beier, Pentafluorosulfanylation of Aromatics and Heteroaromatics, in: *Emerging Fluorinated Motifs*, John Wiley & Sons, Ltd, 2020: pp. 551–570. <https://doi.org/10.1002/9783527824342.ch18>.
- [5] G. Haufe, Synthesis and application of pentafluorosulfanylation reagents and derived aliphatic SF<sub>5</sub>-containing building blocks, *Tetrahedron.* 109 (2022) 132656. <https://doi.org/10.1016/j.tet.2022.132656>.
- [6] M. Magre, S. Ni, J. Cornella, (Hetero)aryl–S(VI) Fluorides: Synthetic Development and Opportunities, *Angew. Chem. Int. Ed.* (2022). <https://doi.org/10.1002/anie.202200904>.
- [7] M. V. Ponomarenko, N. Kalinovich, Y. A. Serguchev, M. Bremer, G.-V. Rösenthaller, Synthesis of pentafluoro- $\lambda^6$ -sulfanyl substituted acetylenes for novel liquid crystals, *J. Fluorine Chem.* 135 (2012) 68–74. <https://doi.org/10.1016/j.jfluchem.2011.08.011>.
- [8] P. Kenyon, S. Mecking, Pentafluorosulfanyl Substituents in Polymerization Catalysis, *J. Am. Chem. Soc.* 139 (2017) 13786–13790. <https://doi.org/10.1021/jacs.7b06745>.
- [9] T. Mo, X. Mi, E.E. Milner, G.S. Dow, P. Wipf, Synthesis of an 8-pentafluorosulfanyl analog of the antimalarial agent mefloquine, *Tetrahedron Lett.* 51 (2010) 5137–5140. <https://doi.org/10.1016/j.tetlet.2010.07.113>.
- [10] O. S. Kanishchev, W. R. Dolbier Jr., Ni/Ir-Catalyzed Photoredox Decarboxylative Coupling of S-Substituted Thiolactic Acids with Heteroaryl Bromides: Short Synthesis of Sulfoxaflor and Its SF<sub>5</sub> Analog, *Chem. Eur. J.* 23 (2017) 7677–7681. <https://doi.org/10.1002/chem.201701627>.
- [11] D. J. Burton, Y. Wang, V. Bizet, D. Cahard, Pentafluorosulfanyl Chloride, in: *Encyclopedia of Reagents for Organic Synthesis*, John Wiley & Sons, Ltd, 2020: pp. 1–7. <https://doi.org/10.1002/047084289X.rn00413.pub2>.
- [12] J.-Y. Shou, X.-H. Xu, F.-L. Qing, Chemoselective Hydro(chloro)pentafluorosulfanylation of Diazo Compounds with Pentafluorosulfanyl Chloride, *Angew. Chem. Int. Ed.* 60 (2021) 15271–15275. <https://doi.org/10.1002/anie.202103606>.
- [13] C. R. Pitts, N. Santschi, A. Togni, Method for Preparing a Polyfluorinated Compound, WO2019229103 (A1), 2019. [https://worldwide.espacenet.com/publicationDetails/biblio?FT=D&date=20191205&DB=EPODOC&locale=fr\\_EP&CC=WO&NR=2019229103A1&KC=A1&ND=4](https://worldwide.espacenet.com/publicationDetails/biblio?FT=D&date=20191205&DB=EPODOC&locale=fr_EP&CC=WO&NR=2019229103A1&KC=A1&ND=4) (accessed March 24, 2022).
- [14] J. R. Case, N. H. Ray, H. L. Roberts, 393. Sulphur chloride pentafluoride: reaction with unsaturated hydrocarbons, *J. Chem. Soc.* (1961) 2066–2070. <https://doi.org/10.1039/JR9610002066>.
- [15] F. W. Hoover, D. D. Coffman, Synthesis and Chemistry of Ethynylsulfur Pentafluoride, *J. Org. Chem.* 29 (1964) 3567–3570. <https://doi.org/10.1021/jo01035a030>.
- [16] S. Ait-Mohand, W. R. Dolbier, New and Convenient Method for Incorporation of Pentafluorosulfanyl (SF<sub>5</sub>) Substituents Into Aliphatic Organic Compounds, *Org. Lett.* 4 (2002) 3013–3015. <https://doi.org/10.1021/ol026483o>.

- [17] W. R. Dolbier, S. Ait-Mohand, T. D. Schertz, T. A. Sergeeva, J. A. Cradlebaugh, A. Mitani, G. L. Gard, R. W. Winter, J. S. Thrasher, A convenient and efficient method for incorporation of pentafluorosulfanyl (SF<sub>5</sub>) substituents into aliphatic compounds, *J. Fluorine Chem.* 127 (2006) 1302–1310. <https://doi.org/10.1016/j.jfluchem.2006.05.003>.
- [18] W. R. Dolbier, Z. Zheng, Preparation of Pentafluorosulfanyl (SF<sub>5</sub>) Pyrrole Carboxylic Acid Esters, *J. Org. Chem.* 74 (2009) 5626–5628. <https://doi.org/10.1021/jo9007699>.
- [19] S. E. Lopez, A. Mitani, P. Pena, I. Ghiviriga, W. R. Dolbier, Use of 1-pentafluorosulfanyl-phenylacetylenes for the preparation of SF<sub>5</sub>-substituted five-membered ring heterocycles through 1,3-dipolar cycloadditions. Isoxazoles and isoxazolines, *J. Fluorine Chem.* 176 (2015) 121–126. <https://doi.org/10.1016/j.jfluchem.2015.06.006>.
- [20] M. V. Ponomarenko, N. Kalinovich, Y. A. Serguchev, M. Bremer, G.-V. Rösenthaller, Synthesis of pentafluoro-λ<sup>6</sup>-sulfanyl substituted acetylenes for novel liquid crystals, *J. Fluorine Chem.* 135 (2012) 68–74. <https://doi.org/10.1016/j.jfluchem.2011.08.011>.
- [21] O. Kazakova, M. Ponomarenko, G.-V. Rösenthaller, Synthesis of Pentafluoro-λ<sup>6</sup>-sulfanyl-Substituted Acetylenes for Liquid Crystals, in: *Efficient Preparations of Fluorine Compounds*, John Wiley & Sons, Ltd, 2013: pp. 214–219. <https://doi.org/10.1002/9781118409466.ch30>.
- [22] M. Cloutier, M. Roudias, J.-F. Paquin, Regioselective Gold-Catalyzed Hydration of CF<sub>3</sub>- and SF<sub>5</sub>-alkynes, *Org. Lett.* 21 (2019) 3866–3870. <https://doi.org/10.1021/acs.orglett.9b01379>.
- [23] V. Debrauwer, I. Leito, M. Lõkov, S. Tshepelevitsh, M. Parmentier, N. Blanchard, V. Bizet, Synthesis and Physicochemical Properties of 2-SF<sub>5</sub>-(Aza)Indoles, a New Family of SF<sub>5</sub> Heterocycles, *ACS Org. Inorg. Au.* 1 (2021) 43–50. <https://doi.org/10.1021/acsorginorgau.1c00010>.
- [24] J. M. Canich, M. M. Ludvig, G. L. Gard, J. M. Shreeve, Synthesis and spectroscopic properties of ethynylsulfur pentafluoride, *Inorg. Chem.* 24 (1985) 3668–3670. <https://doi.org/10.1021/ic00216a039>.
- [25] R. J. Terjeson, J. A. Canich, G.L. Gard, M.R. Colman, M.A. Uremovich, S.H. Strauss, Ethynylpentafluoro-λ<sup>6</sup>-Sulfane (SF<sub>5</sub>C≡CH), in: *Inorganic Syntheses*, John Wiley & Sons, Ltd, 1990: pp. 329–332. <https://doi.org/10.1002/9780470132586.ch66>.
- [26] R. W. Winter, R. Dodean, L. Holmes, G.L. Gard, New acrylate systems: derivatives of β-SF<sub>5</sub>-acrylic acid, *J. Fluorine Chem.* 125 (2004) 37–41. <https://doi.org/10.1016/j.jfluchem.2003.09.004>.
- [27] A. D. Berry, R. A. De Marco, W. B. Fox, Synthesis and characterization of (pentafluorosulfur)acetylenes, *J. Am. Chem. Soc.* 101 (1979) 737–738. <https://doi.org/10.1021/ja00497a042>.
- [28] Q. C. Wang, H. F. White, G. L. Gard, The addition of SF<sub>5</sub>Br to propyne and 3,3,3-trifluoropropyne, *J. Fluorine Chem.* 13 (1979) 455–461. [https://doi.org/10.1016/S0022-1139\(00\)81581-X](https://doi.org/10.1016/S0022-1139(00)81581-X).
- [29] G. S. Lal, K.E. Minnich, Sulfurpentafluoride compounds and methods for making and using same, US6479645B1, 2002. <https://patents.google.com/patent/US6479645B1/en> (accessed March 8, 2022).
- [30] J. Wessel, G. Kleemann, K. Seppelt, Sulfuranonium-Ionen, R-SF<sub>4</sub><sup>+</sup>, Darstellung, Nachweis und Stabilität, *Chem. Ber.* 116 (1983) 2399–2407. <https://doi.org/10.1002/cber.19831160702>.
- [31] B. Duda, D. Lentz, Simultaneous introduction of trifluoromethyl and λ<sup>6</sup>-pentafluorosulfanyl substituents using F<sub>5</sub>S–C≡C–CF<sub>3</sub> as a dienophile, *Org. Biomol. Chem.* 13 (2015) 5625–5628. <https://doi.org/10.1039/C5OB00610D>.
- [32] T. A. Kovacina, R. A. De Marco, A. W. Snow, Synthesis and characterization of mono- and bis(pentafluorosulfur)diacetylene, *J. Fluorine Chem.* 21 (1982) 261–264. [https://doi.org/10.1016/S0022-1139\(00\)81247-6](https://doi.org/10.1016/S0022-1139(00)81247-6).
- [33] T. A. Kovacina, R. A. De Marco, A. W. Snow, Syntheses and characterizations of poly(pentafluorosulfur diacetylenes), *Ind. Eng. Chem. Prod. Res. Dev.* 22 (1983) 170–172. <https://doi.org/10.1021/i300010a003>.
- [34] A. D. Berry, R. A. De Marco, Reaction of pentafluoro[(trifluoromethyl)acetylenyl]sulfur with nickel tetracarbonyl, *Inorg. Chem.* 21 (1982) 457–458. <https://doi.org/10.1021/ic00131a091>.
- [35] J. Wessel, H. Hartl, K. Seppelt, Derivate der Acetylene HC≡C-SF<sub>5</sub> und SF<sub>5</sub>-C≡C-SF<sub>5</sub>, *Chem. Ber.* 119 (1986) 453–463. <https://doi.org/10.1002/cber.19861190208>.

- [36] R. Damerius, D. Leopold, W. Schulze, K. Seppelt, Strukturen von SF<sub>5</sub>-substituierten Metallkomplexen, *Z. Anorg. Allg. Chem.* 578 (1989) 110–118. <https://doi.org/10.1002/zaac.19895780113>.
- [37] T. Henkel, A. Klauck, K. Seppelt, Pentafluoro- $\lambda^6$ -sulfanylacetylene complexes of cobalt, *J. Organomet. Chem.* 501 (1995) 1–6. [https://doi.org/10.1016/0022-328X\(94\)05271-C](https://doi.org/10.1016/0022-328X(94)05271-C).
- [38] S. Zahn, A. F. Nordquist, K. E. Minnich, G. S. Lal, W. F. J. Burgoyne, A. Klauck-Jacobs, Pentafluorosulfanyl-Substituted Thienothiophene Monomers and Conducting Polymers, US2006074250 (A1), 2006. [https://worldwide.espacenet.com/publicationDetails/biblio?FT=D&date=20060406&DB=&locale=en\\_EP&CC=US&NR=2006074250A1&KC=A1&ND=1](https://worldwide.espacenet.com/publicationDetails/biblio?FT=D&date=20060406&DB=&locale=en_EP&CC=US&NR=2006074250A1&KC=A1&ND=1) (accessed January 2, 2022).
- [39] A. Mitani, W.R. Dolbier, Synthesis of Pentafluorosulfanyl (SF<sub>5</sub>)-Substituted Heterocycles and Alkynes, WO2007106818 (A1), 2007. [https://worldwide.espacenet.com/publicationDetails/biblio?FT=D&date=20070920&DB=&locale=en\\_EP&CC=WO&NR=2007106818A1&KC=A1&ND=1](https://worldwide.espacenet.com/publicationDetails/biblio?FT=D&date=20070920&DB=&locale=en_EP&CC=WO&NR=2007106818A1&KC=A1&ND=1) (accessed January 2, 2022).
- [40] W. R. Dolbier, Z. Zheng, Use of 1,3-dipolar reactions for the preparation of SF<sub>5</sub>-substituted five-membered ring heterocycles. Pyrroles and thiophenes, *J. Fluorine Chem.* 132 (2011) 389–393. <https://doi.org/10.1016/j.jfluchem.2011.03.017>.
- [41] S. Emamian, Understanding the regioselectivity and molecular mechanism in the synthesis of isoxazoles containing pentafluorosulfanyl substitution via a [3+2] cycloaddition reaction: A DFT study, *J. Fluorine Chem.* 178 (2015) 165–172. <https://doi.org/10.1016/j.jfluchem.2015.07.019>.
- [42] T. Abe, G.-H. Tao, Y.-H. Joo, R. W. Winter, G. L. Gard, J. M. Shreeve, 5-(1,2,3-Triazol-1-yl)tetrazole Derivatives of an Azidotetrazole via Click Chemistry, *Chem. Eur. J.* 15 (2009) 9897–9904. <https://doi.org/10.1002/chem.200901029>.
- [43] S. Garg, J. M. Shreeve, Trifluoromethyl- or pentafluorosulfanyl-substituted poly-1,2,3-triazole compounds as dense stable energetic materials, *J. Mater. Chem.* 21 (2011) 4787–4795. <https://doi.org/10.1039/C0JM04229C>.
- [44] C. Ye, G. L. Gard, R. W. Winter, R. G. Syvret, B. Twamley, J. M. Shreeve, Synthesis of Pentafluorosulfanylpyrazole and Pentafluorosulfanyl-1,2,3-triazole and Their Derivatives as Energetic Materials by Click Chemistry, *Org. Lett.* 9 (2007) 3841–3844. <https://doi.org/10.1021/ol701602a>.
- [45] R. Gauthier, M. Mamone, J.-F. Paquin, Gold-Catalyzed Hydrofluorination of Internal Alkynes Using Aqueous HF, *Org. Lett.* 21 (2019) 9024–9027. <https://doi.org/10.1021/acs.orglett.9b03425>.