



HAL
open science

Goal-oriented error estimation based on equilibrated flux reconstruction for the approximation of elliptic and parabolic problems

Emmanuel Creusé, Serge Nicaise, Zuqi Tang

► **To cite this version:**

Emmanuel Creusé, Serge Nicaise, Zuqi Tang. Goal-oriented error estimation based on equilibrated flux reconstruction for the approximation of elliptic and parabolic problems. 2022. hal-03719266v1

HAL Id: hal-03719266

<https://hal.science/hal-03719266v1>

Preprint submitted on 11 Jul 2022 (v1), last revised 22 Nov 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Goal-oriented error estimation based on equilibrated flux reconstruction for the approximation of elliptic and parabolic problems

Emmanuel Creusé*, Serge Nicaise

Univ. Polytechnique Hauts-de-France, INSA Hauts-de-France,

CERAMATHS - Laboratoire de Matériaux Céramiques et de Mathématiques,

F-59313 Valenciennes, France

and

Zuqi Tang

Univ. Lille, Arts et Metiers Institute of Technology, Centrale Lille, Junia,

ULR 2697-L2EP, F-59000 Lille, France

July 5, 2022

Abstract

We present unified frameworks for goal-oriented estimates for elliptic and parabolic problems that combine the dual-weighted residual method with equilibrated flux reconstruction. These frameworks allow to analyze simultaneously different approximation schemes for the space discretization of the primal and the dual problems, as conforming or nonconforming finite element method, discontinuous Galerkin methods, or finite volume method. Our main contribution is the splitting of the error on the quantity of interest into a fully computable estimator and a remainder that is bounded, up to an explicit constant, by the product of the fully computable estimators of the primal and dual problems. Some illustrative numerical examples that validate our theoretical results are finally presented.

AMS (MOS) subject classification: 65N30, 65N15, 65M15.

Key Words: goal-oriented estimates, quantity of interest, elliptic and parabolic problems.

1 Introduction

Many engineering problems require computing some quantities of interest, like the local or global mean value of the solution. These quantities of interest are usually linear functionals on a vector space that contains the solution of the considered boundary value problem. Error estimations on such functionals are called goal-oriented error estimations. The main ingredient for such estimations relies on the resolution of the dual problem. Several approaches have been proposed for elliptic problems, let us quote: goal-oriented error estimates based on energy norm of the errors on the solutions of the primal and dual problems [23, 22, 2, 1], the dual weighted residual method [4, 3, 16], the constitutive relation error [13, 12, 24, 25], and the equilibrated flux reconstruction method [17, 15, 26]. To the best of our knowledge, there exist few papers concerning goal-oriented error estimations for parabolic problems [18, 19, 20], where the estimations are based on energy

*Corresponding author: emmanuel.creuse@uphf.fr

norm of the errors on the solutions of the primal and dual problems.

The goal of this paper is then to extend the equilibrated flux reconstruction method (combined with the dual-weighted residual method) for general reaction-diffusion problems and to parabolic problems. Indeed based on equilibrated flux reconstructions, we furnish a decomposition of the error \mathcal{E} on the quantity of interest in the form

$$\mathcal{E} = \eta_{QOI} + \mathcal{R}, \quad (1)$$

where $\eta_{QOI} = \sum_{j=1}^J \eta_j$, each term η_j being fully computable, with some positive integer J depending on the problem, is our proposed estimator on the quantity of interest, while the remainder term \mathcal{R} may depend on the solution of the primal and dual problems but for which we have the bound

$$|\mathcal{R}| \leq C\eta\eta^*, \quad (2)$$

where C is a positive constant independent of the meshsize (and time step size) as explicit as possible, and η (resp. η^*) is an estimator of the error of the solution of the primal (resp. dual) problem using an approximation in a finite dimensional space V_h (resp. V_h^*) not necessarily included into the energy space. These two properties lead to

$$|\mathcal{E}| \leq |\eta_{QOI}| + C\eta\eta^*, \quad (3)$$

and therefore one can chose as error estimator $|\eta_{QOI}| + C\eta\eta^*$ to implement an adaptive algorithm.

On the other hand, if we use a higher order approximation space V_h^* to approximate the dual problem, one may hope to obtain an efficient error estimator (effectivity index close to 1) [2, 17]. In other words, the remainder term is negligible, hence it may be disregarded and the sole term $|\eta_{QOI}|$ can be used as error estimator. More precisely since the estimator η_{QOI} and the right-hand side of (2) are fully computable, we can estimate the ratio

$$\frac{\mathcal{R}}{\eta_{QOI}},$$

during a refinement procedure based on the use of η_{QOI} and check if it tends to zero or not. In the positive case, due to the identity (1) this means that the ratio

$$\frac{\mathcal{E}}{\eta_{QOI}}$$

tends to 1 and will validate the asymptotic exactness of the estimator $|\eta_{QOI}|$.

For a purely diffusion problem with piecewise constant diffusion tensor and piecewise polynomial righ-hand sides, this approach was successfully developed in [15], we here extend their approach to reaction-diffusion problems and to parabolic problems. For a diffusion problem, the decomposition (1) is also obtained in [17], but with an a priori bound for the remainder, which is less interesting for adaptive purposes.

In many cases, the use of $|\eta_{QOI}|$ is sufficient to obtain an efficient error estimator, but it has been observed in [12, 13, 15] that it can overestimate the error. As we can compare all η_i and $\eta\eta^*$, we can improve the approximated value of the quantity of interest by adding the dominant term from η_{QOI} . For instance if η_1 reveals to be a dominant term, we can add η_1 to the approximated value of the quantity of interest and use $|\eta_{QOI} - \eta_1| + C\eta\eta^*$ as error estimator, see [13, Figure 10] and [15, Remark 4.7 and Theorem 4.8].

In summary, the developed approach is flexible and may be then adapted to the different situations to treat.

Let us introduce some notation used in the paper. In the whole paper $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$, $d \geq 2$, is an open, bounded and connected set with a Lipschitz and polyhedral boundary $\partial\Omega$. The usual norm and semi-norm of $H^s(\Omega)$ ($s \geq 0$) are denoted by $\|\cdot\|_{s,\Omega}$ and $|\cdot|_{s,\Omega}$, respectively. In this paper we consider real valued function spaces. For $s = 0$ we drop the index s . Similarly the inner product of $L^2(\Omega)$ will be denoted by (\cdot, \cdot) (resp. $\|\cdot\|$). The duality pairing between $H^{-1}(\Omega)$ (the dual space of $H_0^1(\Omega)$) and $H_0^1(\Omega)$ will be denoted by $\langle \cdot, \cdot \rangle$. By $a \lesssim b$, we mean that there exists a constant $C > 0$ independent of a, b , such that $a \leq Cb$.

The paper is organized as follows. Section 2 is devoted to the reaction-diffusion problem. The problem is defined in subsection 2.1 and the goal oriented functional as well as the adjoint problem are introduced in subsection 2.2. Then the discrete setting is defined in subsection 2.3 and the error representation given in subsection 2.4. The main results about the reaction-diffusion problem are obtained in Theorems 2.1 and 2.3. Similarly, section 3 is devoted to the parabolic problem and follows the same process from subsection 3.1 to subsection 3.4. The main results about the parabolic problem are obtained in Theorems 3.2 and 3.4. Finally, section 4 presents some numerical tests in order to illustrate the previous theoretical results.

2 The reaction-diffusion problem

2.1 Problem definition

In this section we consider the following reaction-diffusion problem

$$\begin{cases} -\operatorname{div}(D\nabla u) + ru = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (4)$$

where $D \in L^\infty(\Omega; \mathbb{R}^{d \times d})$ is a diffusion tensor which is supposed to be a symmetric matrix valued functions that is uniformly bounded from below, namely

$$D(x)\xi \cdot \xi \gtrsim |\xi|^2, \forall \xi \in \mathbb{R}^d, \text{ and a.a. } x \in \Omega,$$

while $r \in L^\infty(\Omega)$ is the reaction function supposed to be nonnegative. The source term f is supposed to be in $L^2(\Omega)$.

By setting

$$B(u, v) = \int_{\Omega} (D\nabla u \cdot \nabla v + ruv) dx, \forall u, v \in H_0^1(\Omega), \quad (5)$$

and

$$F(v) = \int_{\Omega} fv dx, \forall v \in H_0^1(\Omega),$$

the variational formulation of problem (4) is

$$B(u, v) = F(v), \forall v \in H_0^1(\Omega), \quad (6)$$

that has a unique (weak) solution u in $H_0^1(\Omega)$.

2.2 The goal oriented functional and the adjoint problem

We here consider the output functional that represents the physical quantity of interest [17, 15] given by

$$Q(v) = \int_{\Omega} qv dx, \forall v \in L^2(\Omega), \quad (7)$$

where $q \in L^2(\Omega)$ is a given function.

Accordingly the associated dual problem consists in looking for u^* solution of the adjoint equation

$$\begin{cases} -\operatorname{div}(D\nabla u^*) + ru^* = q & \text{in } \Omega, \\ u^* = 0 & \text{on } \partial\Omega. \end{cases} \quad (8)$$

As before, its weak formulation is

$$B(u^*, v) = Q(v), \forall v \in H_0^1(\Omega), \quad (9)$$

that has a unique (weak) solution u^* in $H_0^1(\Omega)$.

2.3 The discrete setting of the reaction-diffusion problem

To discretize problems (6) and (9), we suppose given a partition \mathcal{T} of Ω into polygonal elements T that covers exactly Ω . For simplicity, we assume that the mesh is simplicial and matching (extensions to general polygonal and nonmatching meshes are possible). On such a mesh we introduce the so-called broken Sobolev space

$$H^1(\mathcal{T}) = \{v \in L^2(\Omega) \mid v|_T \in H^1(T), \forall T \in \mathcal{T}\}.$$

As in [10, 15], in order to analyse simultaneously different approximation schemes, the primal problem is approximated in a finite dimensional subspace V_h of $H^1(\mathcal{T})$, while the dual problem will be approximated in a finite dimensional subspace V_h^* of $H^1(\mathcal{T})$, that may be different from V_h . In other words, we suppose given an approximation $u_h \in V_h$ of the solution u of (6) and $u_h^* \in V_h^*$ of the solution u^* of (9). We further assume that a flux reconstruction θ_h is available (using u_h and the datum f), belongs to $H(\operatorname{div}, \Omega)$ and satisfies the following conservation properties (compare with [10, identity (18)])

$$(\operatorname{div} \theta_h + ru_h - f, 1)_T = 0, \forall T \in \mathcal{T}. \quad (10)$$

In the same manner, we assume that a flux reconstruction θ_h^* can be constructed, belongs to $H(\operatorname{div}, \Omega)$ and satisfies the following conservation properties

$$(\operatorname{div} \theta_h^* + ru_h^* - q, 1)_T = 0, \forall T \in \mathcal{T}. \quad (11)$$

With the help of these objects, in the spirit of [17] (see also [15]), let us show that the error $\mathcal{E} = Q(u - u_h)$ on the quantity of interest can be expressed into a fully computable expression and a remainder that will be estimated by a fully computable quantity (but is usually of higher order and can then be disregarded).

2.4 The error representation of the reaction-diffusion problem

Theorem 2.1 *Let $s_h \in H_0^1(\Omega)$ be a potential reconstruction of u_h , then we have*

$$\mathcal{E} = Q(u - u_h) = \eta_{QOI} + \mathcal{R}, \quad (12)$$

where the estimator η_{QOI} is given by

$$\begin{aligned} \eta_{QOI} &= \int_{\Omega} q(s_h - u_h) dx \\ &+ \int_{\Omega} (f - \operatorname{div} \theta_h - ru_h) u_h^* dx \\ &+ \int_{\Omega} (\theta_h + D\nabla s_h) \cdot D^{-1} \theta_h^* dx \\ &- \int_{\Omega} ru_h^* (s_h - u_h) dx, \end{aligned} \quad (13)$$

while the remainder term \mathcal{R} is defined by

$$\begin{aligned}\mathcal{R} &= \int_{\Omega} (f - \operatorname{div} \theta_h - ru_h)(u^* - u_h^*) dx \\ &\quad - \int_{\Omega} (\theta_h + D\nabla s_h) \cdot (D^{-1}\theta_h^* + \nabla u^*) dx \\ &\quad - \int_{\Omega} r(u^* - u_h^*)(s_h - u_h) dx.\end{aligned}$$

Proof. By the definition of the error and Green's formula we have

$$\mathcal{E} = F(u^*) - Q(u_h).$$

Introducing artificially s_h and using (9), this is equivalent to

$$\begin{aligned}\mathcal{E} &= F(u^*) - Q(u_h - s_h) \\ &\quad - \int_{\Omega} (D\nabla u^* \cdot \nabla s_h + ru^* s_h) dx.\end{aligned}\tag{14}$$

Adding and subtracting the term

$$\int_{\Omega} \nabla u^* \cdot \theta_h dx$$

we find

$$\begin{aligned}\mathcal{E} &= F(u^*) - Q(u_h - s_h) \\ &\quad - \int_{\Omega} \nabla u^* \cdot (D\nabla s_h + \theta_h) dx \\ &\quad + \int_{\Omega} (\nabla u^* \cdot \theta_h - ru^* s_h) dx.\end{aligned}$$

Using Green's formula in the fourth term of this right-hand side we find

$$\begin{aligned}\mathcal{E} &= F(u^*) - Q(u_h - s_h) \\ &\quad - \int_{\Omega} \nabla u^* \cdot (D\nabla s_h + \theta_h) dx \\ &\quad + \int_{\Omega} u^* (-\operatorname{div} \theta_h - rs_h) dx.\end{aligned}$$

Replacing s_h by $s_h - u_h + u_h$ in this last term and recalling that $F(u^*) = \int_{\Omega} fu^* dx$, we obtain

$$\begin{aligned}\mathcal{E} &= -Q(u_h - s_h) \\ &\quad - \int_{\Omega} \nabla u^* \cdot (D\nabla s_h + \theta_h) dx \\ &\quad + \int_{\Omega} u^* (f - \operatorname{div} \theta_h - ru_h) dx \\ &\quad - \int_{\Omega} ru^* (s_h - u_h) dx.\end{aligned}$$

Writing $u^* = u_h^* + u^* - u_h^*$ and $D\nabla u^* = -\theta_h^* + D\nabla u^* + \theta_h^*$, we arrive at (12). ■

Note that we do not need the conservation properties (10) and (11) to obtain the splitting (12) from Theorem 2.1, but they are minimal assumptions to guarantee that θ_h (resp. θ_h^*) is a correct approximation of the continuous flux $-D\nabla u$ (resp. $-D\nabla u^*$).

Remark 2.2 (Comparison with earlier literature)

1. In the spirit of [17, Theorem 1 and Remark 1], our estimator η_{QOI} admits the decomposition

$$\eta_{QOI} = \eta_{\mathcal{O}} + \eta_{\nabla} + \eta_{\mathcal{H}},$$

where

$$\begin{aligned}\eta_{\mathcal{O}} &= \int_{\Omega} (f - \operatorname{div} \theta_h - ru_h) u_h^* dx, \\ \eta_{\nabla} &= \int_{\Omega} (\theta_h + D\nabla s_h) \cdot D^{-1} \theta_h^* dx \\ \eta_{\mathcal{H}} &= \int_{\Omega} q(s_h - u_h) dx - \int_{\Omega} ru_h^*(s_h - u_h) dx.\end{aligned}$$

Here $\eta_{\mathcal{O}}$ represents the data oscillation with respect to the primal problem weighted by the dual approximate solution (that slightly differs from the one from [17] because of our weaker assumption (10)), the flux estimator η_{∇} that measures the deviation of $-D\nabla s_h$ from the reconstructed flux θ_h and $\eta_{\mathcal{H}}$ that measures the deviation of u_h from $H_0^1(\Omega)$, these two last quantities also differ from the ones from [17] due to the use of the potential reconstruction s_h .

2. We recover the splitting (4.13) from [15, Remark 4.7] if $r = 0$, D is constant and the data f and q are piecewise polynomials simply by replacing in the expression of η_{∇} from (13) the factor θ_h^* by $-D\nabla s_h^*$, where $s_h^* \in H_0^1(\Omega)$ is a potential reconstruction of u_h^* and plugging the term

$$\int_{\Omega} (\theta_h + D\nabla s_h) \cdot D^{-1} (\theta_h^* + D\nabla s_h^*) dx,$$

in the remainder. More generally in the expression of η_{∇} , we may replace the factor θ_h^* by

$$\theta_h^{*,\alpha} = (1 - \alpha)\theta_h^* - \alpha D\nabla s_h^*,$$

with $\alpha \in \mathbb{R}$ and plug the term

$$\int_{\Omega} (\theta_h + D\nabla s_h) \cdot D^{-1} (\theta_h^* - \theta_h^{*,\alpha}) dx = \alpha \int_{\Omega} (\theta_h + D\nabla s_h) \cdot D^{-1} (\theta_h^* + D\nabla s_h^*) dx$$

in the remainder. For $\alpha = \frac{1}{2}$, we recover the splitting (4.17) from [15, Remark 4.7].

Let us now show that the remainder can be explicitly estimated using the error estimators for u and u^* obtained in [10, Theorem 7]. Namely denote by η (resp. η^*) the error estimator mainly obtained in this reference (for D diagonal and constant but easily extended to the case treated here) using the discrete spaces V_h (resp. V_h^*) to approximate the solution u of the primal problem (6) (resp. u^* of the dual problem (9)), so that

$$\|u - u_h\|_h \leq \eta, \tag{15}$$

$$\|u^* - u_h^*\|_h \leq \eta^*, \tag{16}$$

where the mesh-dependent norm $\|\cdot\|_h$ is defined by

$$\|w\|_h^2 = \|D^{\frac{1}{2}} \nabla_h w\|^2 + \|r^{\frac{1}{2}} w\|^2, \forall w \in H_0^1(\Omega) \cup V_h,$$

where $\nabla_h w$ means the piecewise gradient of w , namely

$$\nabla_h w = \nabla w \text{ on } T, \forall T \in \mathcal{T},$$

and

$$\eta^2 = (\eta_{NC,T}^2 + \eta_{R,T}^2 + \eta_{DF,T}^2),$$

with

$$\begin{aligned}\eta_{NC,T} &= \|u_h - s_h\|_{h,T}, \\ \eta_{R,T} &= m_T \|f - \operatorname{div} \theta_h + r u_h\|_T, \\ \eta_{DF,T} &= \|D^{-\frac{1}{2}}(\theta_h + D\nabla u_h)\|_T,\end{aligned}$$

recalling that

$$\begin{aligned}\|w\|_{h,T} &= \int_T (D\nabla w \cdot \nabla w + r w^2) dx, \forall w \in H^1(T), \\ m_T &:= \min\{C_{P,T}^{\frac{1}{2}} \|D^{-\frac{1}{2}}\|_{\infty,T} h_T, \|r^{-\frac{1}{2}}\|_{\infty,T}\},\end{aligned}$$

and that $C_{P,T}$ is the Poincaré constant, namely the smallest positive constant such that

$$\|\varphi - \mathcal{M}_T \varphi\|_T^2 \leq C_{P,T} h_T^2 \|\nabla \varphi\|_T^2, \forall \varphi \in H^1(T), \quad (17)$$

where $\mathcal{M}_T \varphi$ is the mean of φ in T , namely

$$\mathcal{M}_T \varphi = |T|^{-1} \int_T \varphi(x) dx.$$

We further notice that by convention we set $\|r^{-\frac{1}{2}}\|_{\infty,T} = \infty$ if $\inf_{x \in T} r(x) = 0$.

The expression of η^* is similar by replacing u_h by u_h^* , θ_h by θ_h^* , s_h by s_h^* , and f by q .

Theorem 2.3 *With η (resp. η^*) defined before, we have*

$$|\mathcal{R}| \leq 4\eta\eta^*. \quad (18)$$

Proof. We estimate each term of \mathcal{R} separately. For the first term, using the property (10), we have

$$\mathcal{R}_1 = \int_{\Omega} (f - \operatorname{div} \theta_h - r u_h)(u^* - u_h^*) dx = \int_T (f - \operatorname{div} \theta_h - r u_h)((u^* - u_h^*) - \mathcal{M}_T(u^* - u_h^*)) dx.$$

By Cauchy-Schwarz's inequality, we find

$$|\mathcal{R}_1| \leq \|f - \operatorname{div} \theta_h - r u_h\|_T \|(u^* - u_h^*) - \mathcal{M}_T(u^* - u_h^*)\|_T. \quad (19)$$

Let us now show that

$$\|(u^* - u_h^*) - \mathcal{M}_T(u^* - u_h^*)\|_T \leq m_T \|u^* - u_h^*\|_{h,T}. \quad (20)$$

Indeed first using the estimate (17), we find

$$\begin{aligned}\|(u^* - u_h^*) - \mathcal{M}_T(u^* - u_h^*)\|_T &\leq C_{P,T}^{\frac{1}{2}} h_T \|\nabla(u^* - u_h^*)\|_T \\ &\leq C_{P,T}^{\frac{1}{2}} h_T \|D^{-\frac{1}{2}}\|_{\infty,T} \|D^{\frac{1}{2}} \nabla(u^* - u_h^*)\|_T \\ &\leq C_{P,T}^{\frac{1}{2}} h_T \|D^{-\frac{1}{2}}\|_{\infty,T} \|u^* - u_h^*\|_{h,T}.\end{aligned}$$

This shows (20) if $\inf_{x \in T} r(x) = 0$. On the contrary if $\inf_{x \in T} r(x) > 0$, then we directly write

$$\|(u^* - u_h^*) - \mathcal{M}_T(u^* - u_h^*)\|_T \leq \|u^* - u_h^*\|_T \leq \|r^{-\frac{1}{2}}\|_{\infty,T} (r^{\frac{1}{2}}(u^* - u_h^*))\|_T.$$

In that case, this estimate combined with the previous one yield (20).

Coming back to (19) and using (20), we get

$$|\mathcal{R}_1| \leq \|f - \operatorname{div} \theta_h - ru_h\|_{TmT} \|u^* - u_h^*\|_{h,T}.$$

By discrete Cauchy-Schwarz's inequality, the definition of the estimator η and the estimate (16), we find

$$|\mathcal{R}_1| \leq \eta\eta^*. \quad (21)$$

For the second term, by Cauchy-Schwarz's inequality, we have

$$|\mathcal{R}_2| \leq \|D^{-\frac{1}{2}}(\theta_h + D\nabla s_h)\| \|D^{-\frac{1}{2}}(\theta_h^* + D\nabla u^*)\|.$$

Replacing ∇u^* by $\nabla_h(u^* - u_h^*) + \nabla_h u_h^*$ and using the triangular inequality, one gets

$$|\mathcal{R}_2| \leq \|D^{-\frac{1}{2}}(\theta_h + D\nabla s_h)\| (\|D^{-\frac{1}{2}}(\theta_h^* + D\nabla_h u_h^*)\| + \|D^{\frac{1}{2}}\nabla_h(u^* - u_h^*)\|).$$

Using (16) and the definition of η and η^* , we obtain

$$|\mathcal{R}_2| \leq 2\eta\eta^*. \quad (22)$$

For the last term, again Cauchy-Schwarz's inequality yields

$$\begin{aligned} \left| \int_{\Omega} r(u^* - u_h^*)(s_h - u_h) dx \right| &\leq \|r^{\frac{1}{2}}(u^* - u_h^*)\| \|r^{\frac{1}{2}}(s_h - u_h)\| \\ &\leq \eta\eta^*, \end{aligned}$$

again by (16) and the definition of η .

This estimate and the estimates (21), (22) lead to the conclusion. ■

Remark 2.4 According to the observations from [13, Figure 10] (see also [15]), the flux estimator may overestimate the error due the previous Theorem 2.3 and point 2 of Remark 2.2, therefore, if necessary, it may be added to the approximate quantity of interest.

Remark 2.5 Since the estimator η_{QOI} and the right-hand side of (18) are fully computable, we can estimate the ratio

$$\frac{\mathcal{R}}{\eta_{QOI}},$$

by computing $\frac{4\eta\eta^*}{\eta_{QOI}}$, during a refinement procedure based on the use of η_{QOI} and check if it tends to zero or not. In the positive case, due to the identity (12) this means that the ratio

$$\frac{\mathcal{E}}{\eta_{QOI}}$$

tends to 1 and will validate the asymptotic exactness of the estimator η_{QOI} .

In any case, we can use the estimate

$$|\mathcal{E}| \leq |\eta_{QOI}| + 4\eta\eta^*, \quad (23)$$

that follows from (12) and (18), and then chose as estimator $|\eta_{QOI}| + 4\eta\eta^*$ to implement an adaptive algorithm.

Remark 2.6 Our approach applies to the diffusion-reaction equation with mixed boundary conditions

$$\begin{cases} -\operatorname{div}(D\nabla u) + ru = f & \text{in } \Omega, \\ u = 0 & \text{on } \Gamma_D, \\ -D\nabla u \cdot \mathbf{n} = g_N & \text{on } \Gamma_N, \end{cases} \quad (24)$$

where Γ_D and Γ_N are two open parts of $\partial\Omega$ such that $\bar{\Gamma}_D \cap \bar{\Gamma}_N = \partial\Omega$, and $g_N \in L^2(\Gamma_N)$. Its variational formulation is (6) by replacing $H_0^1(\Omega)$, by

$$H_{\Gamma_D}^1(\Omega) = \{u \in H^1(\Omega) \mid u = 0 \text{ on } \Gamma_N\},$$

with B defined by (5) and

$$F(v) = \int_{\Omega} f v \, dx - \int_{\Gamma_N} g_N v \, d\sigma(x), \forall v \in H_{\Gamma_D}^1(\Omega),$$

while the weak formulation of the adjoint problem is still (9) by replacing $H_0^1(\Omega)$ by $H_{\Gamma_D}^1(\Omega)$.

For this problem, the flux reconstruction $\theta_h \in H(\text{div}, \Omega)$ has to satisfy (10) but furthermore

$$\int_e (\theta_h \cdot \mathbf{n} - g_N) \, d\sigma(x) = 0, \forall e \in \mathcal{E}_h \text{ s.t. } e \subset \bar{\Gamma}_N,$$

while the potential reconstruction s_h of u_h has to be in $H_{\Gamma_D}^1(\Omega) \cap V_h$ and to satisfy $\int_{\Omega} s_h \, dx = 0$ if $\text{meas } \Gamma_D = 0$ and $r = 0$ (a.e.). Similarly the flux reconstruction $\theta_h^* \in H(\text{div}, \Omega)$ has to satisfy (11) and

$$\int_e \theta_h^* \cdot \mathbf{n} \, d\sigma(x) = 0, \forall e \in \mathcal{E}_h \text{ s.t. } e \subset \bar{\Gamma}_N,$$

while s_h^* has to be in $H_{\Gamma_D}^1(\Omega)$ and $\int_{\Omega} s_h^* \, dx = 0$ if $\text{meas } \Gamma_D = 0$ and $r = 0$ (a.e.). With these slight modifications, Theorem 2.1 remains valid with the same definition for the estimator except for $\eta_{\mathcal{O}}$ which is here given by

$$\eta_{\mathcal{O}} = \int_{\Omega} (f - \text{div } \theta_h - r u_h) u_h^* \, dx + \int_{\Gamma_N} (\theta_h \cdot \mathbf{n} - g_N) u_h^* \, d\sigma(x),$$

the remainder being modified accordingly. Further using the results from [8, 10], estimates like (15)-(16) are available, allowing to prove a result like Theorem 2.3.

Remark 2.7 Our approach also applies to the convection-diffusion-reaction equation

$$\begin{cases} -\text{div}(D\nabla u - \mathbf{w}u) + ru = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (25)$$

where $\mathbf{w} \in L^\infty(\Omega)^d$ is divergence-free. Its variational formulation is (6) with

$$B(u, v) = \int_{\Omega} ((D\nabla u - \mathbf{w}u) \cdot \nabla v + ruv) \, dx, \forall u, v \in H_0^1(\Omega),$$

while the weak formulation of the adjoint problem is

$$B^*(u^*, v) = Q(v), \forall v \in H_0^1(\Omega), \forall u^* \in H_0^1(\Omega),$$

where

$$B^*(u, v) = \int_{\Omega} ((D\nabla u + \mathbf{w}u) \cdot \nabla v + ruv) \, dx, \forall u, v \in H_0^1(\Omega).$$

In this setting, Theorem 2.1 remains valid with a slight difference in the definition of η_{∇} that here takes the form

$$\eta_{\nabla} = \int_{\Omega} (\theta_h + D\nabla s_h - \mathbf{w}s_h) \cdot D^{-1}(\theta_h^* + \mathbf{w}u_h^*) \, dx.$$

Further using the results from [8, 10], estimates like (15)-(16) are available, allowing to prove a result like Theorem 2.3.

3 The parabolic problem

3.1 Problem definition

We here consider the following heat type equation:

$$\begin{cases} \frac{\partial u}{\partial t} - \operatorname{div}(D\nabla u) = f & \text{in } C := \Omega \times I, \\ u(\cdot, t) = 0 & \text{on } \partial\Omega \times I, \\ u(\cdot, 0) = u_0 & \text{in } \Omega, \end{cases} \quad (26)$$

where the time interval $I = (0, T)$, with a fixed time T , $D \in L^\infty(\Omega; \mathbb{R}^{d \times d})$ is a diffusion tensor satisfying the same assumptions than in section 2. The datum f is supposed to satisfy $f \in L^2(C)$ and the initial value $u_0 \in L^2(\Omega)$. Under these assumptions, problem (26) or equivalently

$$\langle \partial_t u(\cdot, t), v \rangle + (D\nabla u(\cdot, t), \nabla v) = (f(\cdot, t), v), \forall v \in H_0^1(\Omega), \forall \text{ a.e. } t \in I, \quad (27)$$

has a unique (weak) solution in $Y := \{y \in X := L^2(I; H_0^1(\Omega)) \mid \partial_t u \in X' = L^2(I; H^{-1}(\Omega))\}$. Note that Y is a Hilbert space with the norm

$$\|y\|_Y = \|y\|_X + \|\partial_t y\|_{X'},$$

where the space-time energy norm is given by

$$\|y\|_X^2 := \int_0^T \|D^{\frac{1}{2}} \nabla y\|^2 dt, \forall y \in X.$$

Let us further notice that by [14, Théorème 3.4.1], Y is continuously embedded into $C(\bar{I}; L^2(\Omega))$, which gives a meaning to the initial condition in (26). For further purposes, denote by C_Y the smallest positive constant such that

$$\|y\|_{C([0, T]; L^2(\Omega))} \leq C_Y \|y\|_Y, \forall y \in Y. \quad (28)$$

Note that an explicit upper bound of C_Y can be obtained, namely if we denote by C_Ω the smallest positive constant such that

$$\|u\|_\Omega \leq C_\Omega \|D^{\frac{1}{2}} \nabla u\|, \forall u \in H_0^1(\Omega),$$

then one can show that

$$C_Y \leq \frac{C_\Omega}{\sqrt{T}} + \sqrt{2},$$

see [7, p. 55]. Let us finally remark that $C_\Omega \leq \|D^{\frac{1}{2}}\|_{\infty, \Omega} \frac{\sqrt{d_\Omega}}{d_\Omega}$, where d_Ω is the diameter of Ω , see [6, Theorem VI-3 and p. 301]

3.2 The goal oriented functional and the adjoint problem

We here consider the output functional that represents the physical quantity of interest given by (see [19, 20] for a similar choice)

$$Q(v) = \int_I (g(\cdot, t), v) dt + (u_T^*, v(\cdot, T)), \forall v \in C(\bar{I}, L^2(\Omega)), \quad (29)$$

where $g \in L^2(C)$ and $u_T^* \in L^2(\Omega)$ are given functions.

Accordingly the associated dual problem consists in looking for u^* solution of the backward heat type equation

$$\begin{cases} -\frac{\partial u^*}{\partial t} - \operatorname{div}(D\nabla u^*) = g & \text{in } C, \\ u^*(\cdot, t) = 0 & \text{on } \partial\Omega \times I, \\ u^*(\cdot, T) = u_T^* & \text{in } \Omega. \end{cases} \quad (30)$$

As before, its weak formulation is

$$-\langle \partial_t u^*(\cdot, t), v \rangle + (D\nabla u^*(\cdot, t), \nabla v) = (g(\cdot, t), v), \forall v \in H_0^1(\Omega), \forall \text{ a.e. } t \in I, \quad (31)$$

whose unique (weak) solution u^* has the same regularity than before, namely $u^* \in L^2(I; H_0^1(\Omega)) \cap C(\bar{I}; L^2(\Omega))$ with $\partial_t u^* \in L^2(I; H^{-1}(\Omega))$.

3.3 The discrete setting of the parabolic problem

To discretize problem (26), we suppose given a sequence of discrete times $\{t^n\}_{n=0}^N$, with $N \in \mathbb{N}^*$ such that $t^0 = 0$, $t^N = T$ and $t^n < t^{n+1}$, for all $n = 0, \dots, N-1$. For all $n = 1, \dots, N$, we define $I_n = [t^{n-1}, t^n]$ and let $\tau^n = t^n - t^{n-1}$ be its length. For all $n = 0, \dots, N$, we also suppose given a partition \mathcal{T}^n of polygonal elements T that covers exactly Ω . As before for simplicity, we assume that the meshes are simplicial and matching. For all $n = 0, \dots, N$, we set $h^n = \max_{T \in \mathcal{T}^n} h_T$.

In a standard a posteriori error analysis of the primal problem (26), the initial mesh \mathcal{T}^0 is used to approximate the initial data u_0 , while for $n \geq 1$, \mathcal{T}^n corresponds to a refinement or a coarsening of \mathcal{T}^{n-1} as time evolves. Here the situation is more involved since our estimator is also based on the resolution of the dual problem. Anyway, since we use an implicit scheme in time (for the primal and the dual problems), we suppose that for all $n = 1, \dots, N$, \mathcal{T}^{n-1} and \mathcal{T}^n has a common refinement mesh $\mathcal{T}^{n-1, n}$.

For all $n = 0, \dots, N$, we finally suppose given two finite-dimensional spaces V_h^n and $V_h^{*,n}$ of $H^1(\mathcal{T}^n)$.

For a vector space W (of functions defined on Ω), and $k = 0$ or 1 , we denote by

$$P_\tau^k(W) := \{v \in L^2(0, T, W) \mid v|_{I_n} \in \mathbb{P}^k(I_n, W), \forall n = 1, \dots, N\},$$

where $\mathbb{P}^0(I_n, W)$ (resp. $\mathbb{P}^1(I_n, W)$) denotes the space of functions from I_n to W constant (resp. affine) in time. Note that any function $v \in P_\tau^1(W)$ is uniquely determined by its values $v^n := v(\cdot, t^n)$ at t^n , $n = 0, \dots, N$. On the contrary, any function $v \in P_\tau^0(W)$ is uniquely determined by its values $v^n = v(\cdot, t)|_{I_n}$, $n = 1, \dots, N$. Let us finally observe that the time derivative $\partial_t v$ of $v \in P_\tau^1(W)$ belongs to $P_\tau^0(W)$ and that

$$(\partial_t v)^n = \partial_t v|_{I_n} = \frac{v^n - v^{n-1}}{\tau^n}, \forall n = 1, \dots, N. \quad (32)$$

For shortness we set $\partial_t v^n = (\partial_t v)^n$.

Let us also introduce an approximation $\tilde{f} \in P_\tau^0(L^2(\Omega))$ of f defined by

$$\tilde{f}^n = \frac{1}{\tau_n} \int_{I_n} f(\cdot, t) dt, \forall n = 0, \dots, N.$$

As in [9], we suppose given an approximation $u_{h\tau} \in P_\tau^1(L^2(\Omega))$ of the solution u of (27) such that $u_{h\tau}^n = u_{h\tau}(t^n) \in V_h^n$, for all $n = 0, \dots, N$. Similarly, we suppose given an approximation $u_{h\tau}^* \in P_\tau^1(L^2(\Omega))$ of the solution u^* of (31) such that $u_{h\tau}^{*,n} = u_{h\tau}^*(t^n) \in V_h^{*,n}$, for all $n = 0, \dots, N$.

We also assume that a flux reconstruction $\theta_{h\tau}$ and a potential reconstruction $s_{h\tau}$ are available and satisfy the following conservation properties. $\theta_{h\tau} \in P_\tau^0(H(\text{div}, \Omega))$ and satisfies (compare with [9, identity (4.5)])

$$(\partial_t u_{h\tau}^n + \text{div } \theta_{h\tau}^n - \tilde{f}^n, 1)_T = 0, \forall T \in \mathcal{T}^n, n = 1, \dots, N. \quad (33)$$

On the other hand, $s_{h\tau} \in P_\tau^1(H_0^1(\Omega))$ and satisfies

$$(s_{h\tau}^n, 1)_{T'} = (u_{h\tau}^n, 1)_{T'}, \forall T' \in \mathcal{T}^{n, n+1}, n = 0, \dots, N. \quad (34)$$

A direct consequence of this property is that (see [9, Lemma 3.1])

$$(\partial_t s_{h\tau}^n, 1)_T = (\partial_t u_{h\tau}^n, 1)_T, \forall T \in \mathcal{T}^n, n = 1, \dots, N, \quad (35)$$

holds.

Note further that (35) and (33) directly leads to

$$(\partial_t s_{h\tau}^n + \operatorname{div} \theta_{h\tau}^n - \tilde{f}^n, 1)_T = 0, \forall T \in \mathcal{T}^n, n = 1, \dots, N. \quad (36)$$

In the same manner, we assume that a flux reconstruction $\theta_{h\tau}^*$ and a potential reconstruction $s_{h\tau}^*$ are available and satisfy the following conservation properties: $\theta_{h\tau}^* \in P_\tau^0(H(\operatorname{div}, \Omega))$ and satisfies

$$(-\partial_t u_{h\tau}^{*,n} + \operatorname{div} \theta_{h\tau}^{*,n} - \tilde{g}^n, 1)_T = 0, \forall T \in \mathcal{T}^n, n = 0, \dots, N-1. \quad (37)$$

As before, $s_{h\tau}^* \in P_\tau^1(H_0^1(\Omega))$ is such that

$$(s_{h\tau}^{*,n}, 1)_{T'} = (u_{h\tau}^{*,n}, 1)_{T'}, \forall T' \in \mathcal{T}^{n-1}, n = 0, \dots, N, \quad (38)$$

which yields

$$(\partial_t s_{h\tau}^{*,n}, 1)_T = (\partial_t u_{h\tau}^{*,n}, 1)_T, \forall T \in \mathcal{T}^n, n = 0, \dots, N-1. \quad (39)$$

3.4 The error representation of the parabolic problem

Let us first transform $Q(u)$ as a function of the solution of the adjoint problem.

Lemma 3.1 *It holds*

$$Q(u) = \int_I (f(\cdot, t), u^*(\cdot, t)) dt + (u_0, u^*(0)). \quad (40)$$

Proof. Taking in (31) a test function $v \in L^2(I, H_0^1(\Omega))$ and integrating in time yields

$$\int_I (-\langle \partial_t u^*(\cdot, t), v(\cdot, t) \rangle + (D\nabla u^*(\cdot, t), \nabla v(\cdot, t))) dt = \int_I (g(\cdot, t), v(\cdot, t)) dt, \forall v \in L^2(I, H_0^1(\Omega)). \quad (41)$$

Hence taking $v = u$, we get

$$\int_I (-\langle \partial_t u^*(\cdot, t), u(\cdot, t) \rangle + (D\nabla u^*(\cdot, t), \nabla u(\cdot, t))) dt = \int_I (g(\cdot, t), u(\cdot, t)) dt.$$

Due to the regularities of u and u^* , the following integration by parts formula is valid

$$\int_I \langle \partial_t u^*(\cdot, t), u(\cdot, t) \rangle dt = - \int_I \langle \partial_t u(\cdot, t), u^*(\cdot, t) \rangle dt + (u^*(\cdot, T), u(\cdot, T)) - (u^*(\cdot, 0), u(\cdot, 0)).$$

Inserting this identity into the previous one, and using the initial and final conditions on u and u^* , we find

$$\int_I (\langle \partial_t u(\cdot, t), u^*(\cdot, t) \rangle + (D\nabla u^*(\cdot, t), \nabla u(\cdot, t))) dt - (u_T^*, u(\cdot, T)) + (u^*(\cdot, 0), u_0) = \int_I (g(\cdot, t), u(\cdot, t)) dt.$$

As before as (26) implies that

$$\int_I (\langle \partial_t u(\cdot, t), u^*(\cdot, t) \rangle + (D\nabla u^*(\cdot, t), \nabla u(\cdot, t))) dt = \int_I (f(\cdot, t), u^*(\cdot, t)) dt,$$

we conclude by inserting this identity in the last one. ■

The error on the quantity of interest is defined by

$$\mathcal{E} = Q(u) - Q(u_{h\tau}). \quad (42)$$

Inspired by section 2.3, we show that it can be decomposed into a fully computable expression and a remainder that will be estimated by a fully computable quantity.

Theorem 3.2 *With the previous notation, we have*

$$\mathcal{E} = \eta_{QOI} + \mathcal{R}, \quad (43)$$

where the estimator η_{QOI} is given by

$$\begin{aligned} \eta_{QOI} &= (u_0 - s_{h\tau}(0), u_{h\tau}^*(0)) + (u_{h\tau}^{*,N}, s_{h\tau}(T) - u_{h\tau}^N) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt \\ &+ \int_I (f(\cdot, t) - \tilde{f}, u_{h\tau}^*(\cdot, t)) dt \\ &+ \int_I (\tilde{f}(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, u_{h\tau}^*(\cdot, t)) dt \\ &+ \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot D^{-1} \theta_{h\tau}^* dx dt, \end{aligned} \quad (44)$$

while the remainder term \mathcal{R} is defined by

$$\begin{aligned} \mathcal{R} &= (u_0 - s_{h\tau}(0), (u^* - u_{h\tau}^*)(0)) + (u_T^* - u_{h\tau}^{*,N}, s_{h\tau}(T) - u_{h\tau}^N) \\ &+ \int_I (f(\cdot, t) - \tilde{f}, (u^* - u_{h\tau}^*)(\cdot, t)) dt \\ &+ \int_I (\tilde{f}(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, (u^* - u_{h\tau}^*)(\cdot, t)) dt \\ &- \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot D^{-1} (D\nabla u^* + \theta_{h\tau}^*) dx dt. \end{aligned} \quad (45)$$

Proof. Using the definition (42) of the error and Lemma 3.1, we have

$$\mathcal{E} = \int_I (f(\cdot, t), u^*(\cdot, t)) dt + (u_0, u^*(0)) - \int_I (g(\cdot, t), u_{h\tau}) dt - (u_T^*, u_{h\tau}(\cdot, T)).$$

Introducing artificially $s_{h\tau}$, we get

$$\mathcal{E} = \int_I (f(\cdot, t), u^*(\cdot, t)) dt + (u_0, u^*(0)) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt - \int_I (g(\cdot, t), s_{h\tau}) dt - (u_T^*, u_{h\tau}(\cdot, T)).$$

Using (30), this is equivalent to

$$\begin{aligned} \mathcal{E} &= \int_I (f(\cdot, t), u^*(\cdot, t)) dt + (u_0, u^*(0)) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt \\ &+ \int_I (\partial_t u^* + \operatorname{div}(D\nabla u^*), s_{h\tau}) dt - (u_T^*, u_{h\tau}(\cdot, T)). \end{aligned} \quad (46)$$

As $s_{h\tau}$ is in $H^1(I, H_0^1(\Omega))$ an integration by parts in space and in time leads to

$$\begin{aligned} \int_I (\operatorname{div}(D\nabla u^*), s_{h\tau}) dt &= - \int_I \int_{\Omega} D\nabla u^* \cdot \nabla s_{h\tau} dx dt, \\ \int_I (\partial_t u^*, s_{h\tau}) dt &= - \int_I (u^*, \partial_t s_{h\tau}) dt + (u^*(T), s_{h\tau}(T)) - (u^*(0), s_{h\tau}(0)). \end{aligned}$$

Inserting these identities in (46) we find

$$\begin{aligned} \mathcal{E} &= (u_0 - s_{h\tau}(0), u^*(0)) + (u^*(T), s_{h\tau}(T) - u_{h\tau}^N) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt \\ &+ \int_I ((f(\cdot, t), u^*(\cdot, t)) - (u^*, \partial_t s_{h\tau})) dt \\ &- \int_I \int_{\Omega} D\nabla u^* \cdot \nabla s_{h\tau} dx dt. \end{aligned}$$

Adding and subtracting the term

$$\int_I \int_{\Omega} \nabla u^* \cdot \theta_{h\tau} dx dt,$$

we find

$$\begin{aligned} \mathcal{E} &= (u_0 - s_{h\tau}(0), u^*(0)) + (u^*(T), s_{h\tau}(T) - u_{h\tau}^N) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt \\ &+ \int_I (f(\cdot, t) - \partial_t s_{h\tau}, u^*(\cdot, t)) dt \\ &+ \int_I \int_{\Omega} \nabla u^* \cdot \theta_{h\tau} dx dt \\ &- \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot \nabla u^* dx dt. \end{aligned}$$

Using Green's formula in the fourth term of this right-hand side we find

$$\begin{aligned} \mathcal{E} &= (u_0 - s_{h\tau}(0), u^*(0)) + (u^*(T), s_{h\tau}(T) - u_{h\tau}^N) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt \\ &+ \int_I (f(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, u^*(\cdot, t)) dt \\ &- \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot \nabla u^* dx dt. \end{aligned}$$

Replacing f by $f - \tilde{f} + \tilde{f}$, we find

$$\begin{aligned} \mathcal{E} &= (u_0 - s_{h\tau}(0), u^*(0)) + (u^*(T), s_{h\tau}(T) - u_{h\tau}^N) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt \\ &+ \int_I (f(\cdot, t) - \tilde{f}, u^*(\cdot, t)) dt \\ &+ \int_I (\tilde{f}(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, u^*(\cdot, t)) dt \\ &- \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot \nabla u^* dx dt \end{aligned}$$

Writing $u^* = u_{h\tau}^* + u^* - u_{h\tau}^*$ and $D\nabla u^* = -\theta_{h\tau}^* + D\nabla u^* + \theta_{h\tau}^*$, we arrive at (43). ■

Remark 3.3 As in Remark 2.2, our estimator η_{QOI} admits the decomposition

$$\eta_{QOI} = \eta_{\mathcal{O}} + \eta_{\nabla} + \eta_{\mathcal{H}},$$

where

$$\begin{aligned} \eta_{\mathcal{O}} &= \int_I (f(\cdot, t) - \tilde{f}^n, u_{h\tau}^*(\cdot, t)) dt \\ &+ \int_I (\tilde{f}(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, u_{h\tau}^*(\cdot, t)) dt, \\ \eta_{\nabla} &= \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot D^{-1} \theta_{h\tau}^* dx dt, \\ \eta_{\mathcal{H}} &= (u_0 - s_{h\tau}(0), u_{h\tau}^*(0)) + (u_{h\tau}^{*,N}, s_{h\tau}(T) - u_{h\tau}^N) - \int_I (g(\cdot, t), u_{h\tau} - s_{h\tau}) dt. \end{aligned}$$

As before $\eta_{\mathcal{O}}$ represents the data oscillation with respect to the primal problem weighted by the dual approximate solution, the flux estimator η_{∇} that measures the deviation of $-D\nabla s_{h\tau}$ from the reconstructed flux $\theta_{h\tau}$ and $\eta_{\mathcal{H}}$ that measures the deviation of $u_{h\tau}$ from $H_0^1(\Omega)$.

Let us now show that the remainder can be explicitly estimated using the error estimators for u and u^* obtained in [9, Theorem 3.2]. Namely denote by η (resp. η^*) the estimator obtained in this reference using the discrete spaces V_h^n (resp. $V_h^{*,n}$) to approximate the solution u of the primal problem (27) (resp. u^* of the dual problem (31)), so that

$$\|u - u_{h\tau}\|_Y \leq \eta + 3\|f - \tilde{f}\|_{X'}, \quad (47)$$

$$\|u^* - u_{h\tau}^*\|_Y \leq \eta^* + 3\|g - \tilde{g}\|_{X'}, \quad (48)$$

where

$$\begin{aligned} \eta &= 3 \left\{ \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\eta_{R,T}^n(t) + \eta_{DF,T}^n(t))^2 dt \right\}^{\frac{1}{2}} + \eta_{IC} \\ &+ \left\{ \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\eta_{NC1,T}^n(t))^2 dt \right\}^{\frac{1}{2}} + \left\{ \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\eta_{NC2,T}^n(t))^2 dt \right\}^{\frac{1}{2}}, \end{aligned}$$

with

$$\eta_{IC} = \sqrt{2}\|s_{h\tau}^0 - u_0\|,$$

and for all $n = 1, \dots, N$, $t \in I_n$ and $T \in \mathcal{T}^n$,

$$\begin{aligned} \eta_{R,T}^n(t) &= \pi^{-1} \|D^{-\frac{1}{2}}\|_{\infty, T} h_T \|\tilde{f} - \partial_t s_{h\tau} - \operatorname{div} \theta_{h\tau}^n\|_T, \\ \eta_{DF,T}^n(t) &= \|D^{-\frac{1}{2}}(\theta_{h\tau} + D\nabla s_{h\tau})\|_T, \\ \eta_{NC1,T}^n(t) &= \|D^{\frac{1}{2}}\nabla(s_{h\tau} - u_{h\tau})\|_T, \\ \eta_{NC2,T}^n(t) &= \pi^{-1} \|D^{-\frac{1}{2}}\|_{\infty, T} h_T \|\partial_t(s_{h\tau} - u_{h\tau})\|_T. \end{aligned}$$

The estimator η^* of the dual problem is defined similarly by replacing $u_{h\tau}$ by $u_{h\tau}^*$, $\theta_{h\tau}$ by $\theta_{h\tau}^*$, $s_{h\tau}$ by $s_{h\tau}^*$, $s_{h\tau}^0$ by $s_{h\tau}^{*,0}$, u_0 by u_T^* , f by q , and ∂_t by $-\partial_t$.

Theorem 3.4 *With η (resp. η^*) defined before, we have*

$$\begin{aligned} |\mathcal{R}| &\leq D_Y(\eta + \|(s_{h\tau} - u_{h\tau})(T)\| + \|f - \tilde{f}\|_{X'}) (\eta^* + \|(s_{h\tau}^* - u_{h\tau}^*)(0)\| + \|g - \tilde{g}\|_{X'}) \\ &+ \pi^{-1} h^n \|D^{-\frac{1}{2}}\|_{\infty, \Omega} \|f(\cdot, t) - \tilde{f}\|_C \eta^*, \end{aligned} \quad (49)$$

with $D_Y = \sqrt{2} \max\{1, 3C_Y\} + 11$.

Proof. We estimate each term of \mathcal{R} separately. For the first term, by Cauchy-Schwarz's inequality we have

$$|(u_0 - s_{h\tau}(0), (u^* - u_{h\tau}^*)(0))| \leq \|u_0 - s_{h\tau}(0)\| \|(u^* - u_{h\tau}^*)(0)\|. \quad (50)$$

Note that

$$\|u_0 - s_{h\tau}(0)\| \leq 2^{-1/2} \eta, \quad (51)$$

therefore it remains to estimate the second factor of the right-hand side of (50). But since $u_{h\tau}^*$ does not belong to Y , we write $(u^* - u_{h\tau}^*)(0) = (u^* - s_{h\tau}^*)(0) + (s_{h\tau}^* - u_{h\tau}^*)(0)$, and use the triangular inequality to find

$$\|(u^* - u_{h\tau}^*)(0)\| \leq \|(u^* - s_{h\tau}^*)(0)\| + \|(s_{h\tau}^* - u_{h\tau}^*)(0)\|. \quad (52)$$

For the first term of this right-hand side we use the continuous embedding $Y \hookrightarrow C([0, T]; L^2(\Omega))$ to find (see (28))

$$\|(u^* - s_{h\tau}^*)(0)\| \leq C_Y \|u^* - s_{h\tau}^*\|_Y. \quad (53)$$

Then using the identity

$$\|u^* - s_{h\tau}^*\|_Y = \|\partial_t(u^* - s_{h\tau}^*)\|_{L^2(0,T,H^{-1}(\Omega))} + \left(\sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^{n-1,n}} \|D^{\frac{1}{2}} \nabla(u^* - s_{h\tau}^*)(t)\|_T^2 dt \right)^{\frac{1}{2}},$$

inserting artificially $u_{h\tau}^*$ and using the triangular inequality, we find

$$\begin{aligned} \|u^* - s_{h\tau}^*\|_Y &\leq \|\partial_t(u^* - u_{h\tau}^*)\|_{L^2(0,T,H^{-1}(\Omega))} + \|\partial_t(u_{h\tau}^* - s_{h\tau}^*)\|_{L^2(0,T,H^{-1}(\Omega))} \\ &+ \left(\sum_{n=0}^{N-1} \int_{I_n} \sum_{T \in \mathcal{T}^{n-1,n}} \|D^{\frac{1}{2}} \nabla(u^* - u_{h\tau}^*)(t)\|_T^2 dt \right)^{\frac{1}{2}} \\ &+ \left(\sum_{n=0}^{N-1} \int_{I_n} \sum_{T \in \mathcal{T}^{n-1,n}} \|D^{\frac{1}{2}} \nabla(u_{h\tau}^* - s_{h\tau}^*)(t)\|_T^2 dt \right)^{\frac{1}{2}}. \end{aligned}$$

According to the definition of the norm Y for $u^* - u_{h\tau}^*$ and to the definition of η^* , we find

$$\begin{aligned} \|u^* - s_{h\tau}^*\|_Y &\leq \|u^* - u_{h\tau}^*\|_Y + \eta^* \\ &+ \|\partial_t(u_{h\tau}^* - s_{h\tau}^*)\|_{L^2(0,T,H^{-1}(\Omega))}. \end{aligned} \quad (54)$$

For this last term, for all $t \in (0, T)$, we define the $H^{-1}(\Omega)$ -norm by¹

$$\|\partial_t(u_{h\tau}^* - s_{h\tau}^*)(t)\|_{H^{-1}(\Omega)} = \sup_{\varphi \in H_0^1(\Omega), \varphi \neq 0} \frac{|\int_{\Omega} \partial_t(u_{h\tau}^* - s_{h\tau}^*)(t) \varphi dx|}{\|D^{\frac{1}{2}} \nabla \varphi\|}. \quad (55)$$

Hence using (39), we may write

$$\int_{\Omega} \partial_t(u_{h\tau}^* - s_{h\tau}^*)(t) \varphi dx = \sum_{T \in \mathcal{T}^n} \int_T \partial_t(u_{h\tau}^* - s_{h\tau}^*)(t) (\varphi - \mathcal{M}_T \varphi) dx,$$

where $\mathcal{M}_T \varphi$ is the mean of φ on T . Therefore by the Poincaré inequality stating that

$$\|\varphi - \mathcal{M}_T \varphi\|_T \leq \pi^{-1} h_T |\varphi|_{1,T},$$

we find

$$\|\varphi - \mathcal{M}_T \varphi\|_T \leq \pi^{-1} h_T \|D^{-\frac{1}{2}}\|_{\infty,T} \|D^{\frac{1}{2}} \nabla \varphi\|_T,$$

and hence

$$\left| \int_{\Omega} \partial_t(u_{h\tau}^* - s_{h\tau}^*)(t) \varphi dx \right| \leq \pi^{-1} \sum_{T \in \mathcal{T}^n} h_T \|D^{-\frac{1}{2}}\|_{\infty,T} \|\partial_t(u_{h\tau}^* - s_{h\tau}^*)(t)\|_T \|D^{\frac{1}{2}} \nabla \varphi\|_T.$$

Using the discrete Cauchy-Schwarz's inequality and inserting this estimate in (55), we obtain

$$\|\partial_t(u_{h\tau}^* - s_{h\tau}^*)(t)\|_{H^{-1}(\Omega)} \leq \pi^{-1} \left(\sum_{T \in \mathcal{T}^n} h_T^2 \|D^{-\frac{1}{2}}\|_{\infty,T}^2 \|\partial_t(u_{h\tau}^* - s_{h\tau}^*)(t)\|_T^2 \right)^{\frac{1}{2}}.$$

Integrating the square of this estimate in $0, T$ and using the definition of η^* , we arrive at

$$\|\partial_t(u_{h\tau}^* - s_{h\tau}^*)(t)\|_{L^2(0,T,H^{-1}(\Omega))} \leq \eta^*,$$

and using it into (54) yields

$$\|u^* - s_{h\tau}^*\|_Y \leq \|u^* - u_{h\tau}^*\|_Y^2 + 2\eta^*.$$

¹as D is bounded and positive definite this norm is equivalent to the usual one.

The estimate (48) then leads to

$$\|u^* - s_{h\tau}^*\|_Y \leq 3(\eta^* + \|g - \tilde{g}\|_{X'}). \quad (56)$$

Combining this estimate with (53) leads to

$$\|(u^* - s_{h\tau}^*)(0)\| \leq 3C_Y(\eta^* + \|g - \tilde{g}\|_{X'}),$$

that can be used in (52) to find

$$\|(u^* - u_{h\tau}^*)(0)\| \leq \max\{1, 3C_Y\}(\eta^* + \|(s_{h\tau}^* - u_{h\tau}^*)(0)\| + \|g - \tilde{g}\|_{X'}). \quad (57)$$

Coming back to (50) and using this last estimate and (51), we finally find

$$|(u_0 - s_{h\tau}(0), (u^* - u_{h\tau}^*)(0))| \leq 2^{-\frac{1}{2}} \max\{1, 3C_Y\} \eta(\eta^* + \|(s_{h\tau}^* - u_{h\tau}^*)(0)\| + \|g - \tilde{g}\|_{X'}). \quad (58)$$

The second term is the symmetric of the first one by exchanging the role of u and u^* , hence

$$|(u_T^* - u_{h\tau}^{*,N}, s_{h\tau}(T) - u_{h\tau}^N)| \leq 2^{-\frac{1}{2}} \max\{1, 3C_Y\}(\eta + \|(s_{h\tau} - u_{h\tau})(T)\| + \|f - \tilde{f}\|_{X'})\eta^*. \quad (59)$$

Concerning the third term, we again introduce artificially $s_{h\tau}^*$ to get

$$\begin{aligned} \mathcal{R}_3 &:= \int_I (f(\cdot, t) - \tilde{f}, (u^* - u_{h\tau}^*)(\cdot, t)) dt = \int_I (f(\cdot, t) - \tilde{f}, (u^* - s_{h\tau}^*)(\cdot, t)) dt \\ &\quad + \sum_{n=1}^N \int_{I_n} (f(\cdot, t) - \tilde{f}^n, (s_{h\tau}^* - u_{h\tau}^*)(\cdot, t)) dt, \end{aligned}$$

and by using a duality argument for the first term and Cauchy-Schwarz's inequality for the second one, we obtain

$$\begin{aligned} |\mathcal{R}_3| &\leq \|f(\cdot, t) - \tilde{f}\|_{X'} \|u^* - s_{h\tau}^*\|_X \\ &\quad + \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^{n, n-1}} \|f(\cdot, t) - \tilde{f}^n\|_T \|(s_{h\tau}^* - u_{h\tau}^*)(\cdot, t)\|_T dt. \end{aligned}$$

Due to the property (38) and the Poincaré inequality, we find

$$\begin{aligned} |\mathcal{R}_3| &\leq \|f(\cdot, t) - \tilde{f}\|_{X'} \|u^* - s_{h\tau}^*\|_X \\ &\quad + \pi^{-1} \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^{n, n-1}} \|f(\cdot, t) - \tilde{f}^n\|_T h_T \|\nabla(s_{h\tau}^* - u_{h\tau}^*)(\cdot, t)\|_T dt. \end{aligned}$$

Using the definition of the broken gradient of $u_{h\tau}^*$ and the fact that $h_T \leq h^n$ for all $T \in \mathcal{T}^{n-1, n}$, we get

$$\begin{aligned} |\mathcal{R}_3| &\leq \|f(\cdot, t) - \tilde{f}\|_{X'} \|u^* - s_{h\tau}^*\|_X \\ &\quad + \pi^{-1} h^n \|D^{-\frac{1}{2}}\|_{\infty, \Omega} \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} \|f(\cdot, t) - \tilde{f}^n\|_T \|D^{\frac{1}{2}} \nabla^{n-1, n}(s_{h\tau}^* - u_{h\tau}^*)(\cdot, t)\|_T dt. \end{aligned}$$

Hence using Cauchy-Schwarz's inequality, we find

$$|\mathcal{R}_3| \leq \|f(\cdot, t) - \tilde{f}\|_{X'} \|u^* - s_{h\tau}^*\|_X + \pi^{-1} h^n \|D^{-\frac{1}{2}}\|_{\infty, \Omega} \|f(\cdot, t) - \tilde{f}\|_C \eta^*.$$

Finally as by definition $\|u^* - s_{h\tau}^*\|_X \leq \|u^* - s_{h\tau}^*\|_Y$, the estimate (56) yields

$$|\mathcal{R}_3| \leq 3\|f(\cdot, t) - \tilde{f}\|_{X'}(\eta^* + \|g - \tilde{g}\|_{X'}) + \pi^{-1} h^n \|D^{-\frac{1}{2}}\|_{\infty, \Omega} \|f(\cdot, t) - \tilde{f}\|_C \eta^*. \quad (60)$$

For the fourth term again inserting artificially $s_{h\tau}^*$ we get

$$\begin{aligned}\mathcal{R}_4 &:= \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, (u^* - u_{h\tau}^*)(\cdot, t))_T dt \\ &= \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, (u^* - s_{h\tau}^*)(\cdot, t))_T dt \\ &\quad + \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, (s_{h\tau}^* - u_{h\tau}^*)(\cdot, t))_T dt.\end{aligned}$$

For the first term using (36), we can replace $u^* - s_{h\tau}^*$ by $u^* - s_{h\tau}^* - \mathcal{M}_T(u^* - s_{h\tau}^*)$ on $T \in \mathcal{T}^n$, while for the second term we reorganize the summation on the triangulation $\mathcal{T}^{n-1, n}$ and then find

$$\begin{aligned}\mathcal{R}_4 &= \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} (\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, (u^* - s_{h\tau}^*)(\cdot, t) - \mathcal{M}_T(u^* - s_{h\tau}^*))_T dt \\ &\quad + \sum_{n=1}^N \int_{I_n} \sum_{T' \in \mathcal{T}^{n-1, n}} (\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}, (s_{h\tau}^* - u_{h\tau}^*)(\cdot, t))_{T'} dt.\end{aligned}$$

Using Poincaré inequality for each term of this right-hand side (recalling the property (38)), we find

$$\begin{aligned}|\mathcal{R}_4| &\leq \pi^{-1} \sum_{n=1}^N \int_{I_n} \sum_{T \in \mathcal{T}^n} h_T \|D^{-\frac{1}{2}}\|_{\infty, T} \|\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}\|_T \|D^{\frac{1}{2}} \nabla(u^* - s_{h\tau}^*)(\cdot, t)\|_T dt \\ &\quad + \pi^{-1} \sum_{n=1}^N \int_{I_n} \sum_{T' \in \mathcal{T}^{n-1, n}} h_{T'} \|D^{-\frac{1}{2}}\|_{\infty, T'} \|\tilde{f}^n(\cdot, t) - \operatorname{div} \theta_{h\tau}(\cdot, t) - \partial_t s_{h\tau}\|_{T'} \|D^{\frac{1}{2}} \nabla(s_{h\tau}^* - u_{h\tau}^*)\|_{T'} dt.\end{aligned}$$

Using Cauchy-Schwarz's inequality and reorganizing the summation in the second term (using the fact that $h_{T'} \leq h_T$, for all $T' \in \mathcal{T}^{n-1, n}$ such that $T' \subset T \in \mathcal{T}^n$), we get

$$|\mathcal{R}_4| \leq \eta(\|u^* - s_{h\tau}^*\|_Y + \eta^*).$$

As before the estimate (56) yields

$$|\mathcal{R}_4| \leq 3\eta(\eta^* + \|g - \tilde{g}\|_{X'}) + \eta\eta^*. \quad (61)$$

It remains the last term that can be transformed as

$$\begin{aligned}\mathcal{R}_5 &:= \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot D^{-1}(D\nabla u^* + \theta_{h\tau}^*) dx dt \\ &= \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot \nabla(u^* - s_{h\tau}^*) dx dt \\ &\quad + \int_I \int_{\Omega} (\theta_{h\tau} + D\nabla s_{h\tau}) \cdot D^{-1}(D\nabla s_{h\tau} + \theta_{h\tau}^*) dx dt.\end{aligned}$$

Hence Cauchy-Schwarz's inequality directly yields

$$|\mathcal{R}_5| \leq \eta(\|u^* - s_{h\tau}^*\|_Y + \eta^*).$$

With the help of the estimate (56) we again obtain

$$|\mathcal{R}_5| \leq 3\eta(\eta^* + \|g - \tilde{g}\|_{X'}) + \eta\eta^*. \quad (62)$$

The conclusion follows from the estimate (58), (59), (60) (61), and (62). ■

Note that Remark 2.5 remains valid for our parabolic problem (26).

4 Numerical examples

In this section, we present some illustrative numerical examples that validate our theoretical results. Before going on, let us notice that all numerical simulations are realized by the free scientific computing software FreeFem++ [11].

4.1 Diffusion problems

In this subsection, we aim to illustrate the theoretical results obtained for diffusion problems, and specifically Theorems 2.1 and 2.3.

4.1.1 Regular solutions

We first consider the same benchmark as the one proposed in [17], section 8 and [15], section 6.1. The primal problem (4) is defined with $d = 2$, $\Omega =]0, 1[^2$, $D = I_{\mathbb{R}^2}$ and $r = 0$. The exact solution is chosen equal to $u(x, y) = 10^4 x(1-x)y(1-y)e^{-100(\rho(x,y))^2}$ with $\rho(x, y) = ((x-0.75)^2 + (y-0.75)^2)^{1/2}$ (see Figure 1(a)), and the right-hand-side f is computed accordingly. The dual problem is defined by (8) with $q = \mathbb{1}_\omega$ and $\omega = \{(x, y) \in \Omega : 1.5 \leq x + y \leq 1.75\}$ (see Figure 1(b)).




Figure 1: Diffusion problem, regular solution. (a) : regular solution. (b) : q function.

The approximation u_h of u solution of the primal problem is made using standard conforming \mathbb{P}_1 finite elements on regular meshes made of triangles (see Figure 2 for the first three refinements), and the corresponding flux reconstruction θ_h is approximated using some standard \mathbb{RT}_1 finite elements (Raviart-Thomas finite elements of degree 2 on each triangle). The approximation u_h^* of u^* solution of the dual problem is made using standard conforming \mathbb{P}_2 finite elements on the same meshes, and the corresponding flux reconstruction θ_h^* is approximated using some standard \mathbb{RT}_2 finite elements (Raviart-Thomas finite elements of degree 3 on each triangle).

For each mesh, the value of $\mathcal{E} = Q(u - u_h)$ defined by (7) is computed, as well as the one of η_{QOI} defined by (13). We plot on Figure 3(a) the values of \mathcal{E} as well as the one of η_{QOI} and $4\eta\eta^*$ as a function of h in a log-log scale. First of all, we remark that \mathcal{E} goes towards zero at the order $O(h^2)$, what is the expected behaviour since $u \in H^2(\Omega)$ and u_h is approximated by \mathbb{P}_1 finite elements. Moreover, the estimator η_{QOI} converges towards zero at the same rate of convergence. Figure 3(b) displays the so-called effectivity index defined by $I_{eff} = \mathcal{E}/\eta_{QOI}$ as a function of h . This one is always nearly equal to one, and consequently the estimator η_{QOI} is asymptotically

²here and below, $I_{\mathbb{R}^2}$ means the 2×2 identity matrix




Figure 2: Unstructured regular meshes, first three refinements.

exact. Coming back to Figure 3(a), we see that the value of $4\eta\eta^*$ goes towards zero faster than \mathcal{E} (here at the order $O(h^3)$), what illustrates the fact that \mathcal{R} is clearly superconvergent (see (18)).




Figure 3: Diffusion problem, $D = I_{\mathbb{R}^2}$, regular solution. (a) : rates of convergence. (b) : effectivity index.

We have also performed a similar simulation in the same setting than before except that we take a variable diffusion tensor, namely $D(x, y) = (1 + x^2 + y^2)I_{\mathbb{R}^2}$, for all $(x, y) \in]0, 1[^2$. The results are presented in Figure 4 and are very similar to the case $D = I_{\mathbb{R}^2}$, leading to the same conclusion than before.

4.1.2 A singular solution

This time we consider a singular case. The primal problem (4) is defined with $d = 2$, $\Omega =]-1, 1[^2$ and $r = 0$. The diffusion coefficient D is piecewise constant in Ω and defined by $D = d_i I_{\mathbb{R}^2}$ in Ω_i , $1 \leq i \leq 4$, with $\Omega_1 = (0, 1) \times (0, 1)$, $\Omega_2 = (-1, 0) \times (0, 1)$, $\Omega_3 = (-1, 0) \times (-1, 0)$ and $\Omega_4 = (0, 1) \times (-1, 0)$, $0 < d_1 = d_3 = a < 1$ and $d_2 = d_4 = 1$. Defining the singular exponent $\alpha = \frac{4}{\pi} \arctan(\sqrt{a})$, the exact solution is given by

$$u(x, y) = p(x, y) S(x, y),$$

where $p(x, y) = (1 - x^4)(1 - y^4)$ is a truncation function and $S(x, y) = \rho^\alpha v(\theta)$ with

$$\rho = \sqrt{x^2 + y^2} \quad , \quad \theta = \begin{cases} \arctan(y/x) & \text{in } \Omega_1 \cup \Omega_4, \\ \pi + \arctan(y/x) & \text{in } \Omega_2 \cup \Omega_3, \end{cases}$$




Figure 4: Diffusion problem, $D(x, y) = (1+x^2+y^2)I_{\mathbb{R}^2}$ regular solution. (a) : rates of convergence. (b) : effectivity index.

and

$$v(\theta) = \begin{cases} \sin\left(\alpha\left(\theta - \frac{\pi}{4}\right)\right) & \text{in } \Omega_1, \\ \sqrt{a} \cos\left(\alpha\left(\frac{3\pi}{4} - \theta\right)\right) & \text{in } \Omega_2, \\ \sin\left(\alpha\left(\frac{5\pi}{4} - \theta\right)\right) & \text{in } \Omega_3, \\ -\sqrt{a} \cos\left(\alpha\left(\frac{\pi}{4} + \theta\right)\right) & \text{in } \Omega_4. \end{cases}$$

The right-hand-side f is computed accordingly. Let us note that in such a configuration, for any $\varepsilon > 0$ we have $u \in H^{1+\alpha-\varepsilon}(\Omega)$ (see e.g. the appendix of [5] for a similar configuration).

The dual problem is defined by (8) with $q = \mathbb{1}_\omega$ and $\omega = (0, 0.5) \times (-0.25, 0.25)$. All the spatial discretizations are done similarly to the regular case of section 4.1.1.


The obtained results are respectively displayed in Figures 5 and 6 for $a = 1/5$ and $a = 1/20$, respectively leading to the values of $\alpha \approx 0.53$ and $\alpha \approx 0.28$. From these figures, we may notice that the error, the estimator η_{QOI} and $4\eta\eta^*$ both converge towards zero at the order $O(h^{2\alpha})$. For $a = 1/5$, as the effectivity index is almost one for h small enough, we can conclude that the remainder \mathcal{R} is nevertheless superconvergent. For $a = 1/20$, this phenomenon is less clear but is probably due to the very low order of convergence. In any case, for such problems with singular solutions, an adaptive algorithm should be based on the estimator η_{QOI} and the product $4\eta\eta^*$, since they have the same rate of convergence.

4.2 Parabolic problems


In this subsection, we aim to illustrate the theoretical results obtained for parabolic problems, and specifically Theorems 3.2 and 3.4.

4.2.1 The heat equation

We start to propose a simple problem, defining the primal problem by (26) with $d = 2$, $\Omega =]0, 1[^2$, $T = 1.0$ and $D = I_{\mathbb{R}^2}$. We choose the exact solution being equal to $u(t, x, y) = x^2 y^2 (x-1)^2 (y-1)^2 \sin(t)$ and compute the right-hand-side f accordingly. The dual problem is defined by (30),




(a)




(b)

Figure 5: Diffusion problem, singular solution, $a = 1/5$. (a) : rates of convergence. (b) : effectivity index.



(a)



(b)

Figure 6: Diffusion problem, singular solution, $a = 1/20$. (a) : rates of convergence. (b) : effectivity index.

where we choose $g = 0$ and $u_T^* = e^{-\frac{\rho(x,y)^2}{32 \ln 10}}$ with $\rho(x, y) = ((x - 0.5)^2 + (y - 0.5)^2)^{1/2}$.

Concerning the time approximation, we use the definition of $(\partial_t u)^n$ given in (32) to compute an implicit-Euler time integration for the resolution of the primal problem, where u^n is the approximation of $u(t_n, \cdot)$ and solution of the elliptic problem:

$$\begin{cases} \frac{u^n - u^{n-1}}{\tau} - \operatorname{div}(D\nabla u^n) = \tilde{f}^n & \text{in } \Omega, \\ u^n = 0 & \text{on } \partial\Omega, \\ u^0 = u_0 & \text{in } \Omega. \end{cases} \quad (63)$$

Since the scheme is fully implicit in time, there is no need to impose any stability condition linking h and τ . For accuracy reasons, we choose $\tau = h$. Similarly to the stationary tests, the spatial approximation u_h^n of u^n is made using standard \mathbb{P}_1 finite elements and the corresponding flux reconstruction θ_h is approximated using some standard \mathbb{RT}_1 finite elements. The time integration of the dual problem is also done using an implicit-Euler scheme (with $\tau = h$), where $u^{*,n}$ is the approximation of $u^*(t_n, \cdot)$ and solution of:

$$\begin{cases} -\frac{u^{*,n} - u^{*,n-1}}{\tau} - \operatorname{div}(D\nabla u^{*,n-1}) = g & \text{in } C, \\ u^*(\cdot, t) = 0 & \text{on } \partial\Omega \times I, \\ u^*(\cdot, T) = u_T^* & \text{in } \Omega. \end{cases} \quad (64)$$

The spatial approximation $u_h^{*,n}$ of $u^{*,n}$ is made using standard \mathbb{P}_2 finite elements on the same meshes, and the corresponding flux reconstruction θ_h^* is approximated using some standard \mathbb{RT}_2 finite elements.


For each mesh, we compute the value of $\mathcal{E} = Q(u - u_h)$ defined by (42) as well as the one of η_{QOI} defined by (44). We plot on Figure 7(a) the values of \mathcal{E} as well as the one of η_{QOI} and $\eta\eta^*$ as a function of h in a log-log scale. First of all, we remark that the error \mathcal{E} goes towards zero at the order $O(h)$, what is the expected behaviour since even if u is regular enough and u_h is approximated by \mathbb{P}_1 finite elements, the time integration is at order one and the choice of $\tau = h$ leads to a global rate of convergence equal to one. Moreover, the estimator η_{QOI} converges towards zero at the same rate of convergence. Figure 7(b) displays the effectivity index $I_{eff} = \mathcal{E}/\eta_{QOI}$ as a function of h . As for the stationary tests with regular solutions, it is always nearly equal to one and converges towards one. Consequently the estimator η_{QOI} is asymptotically exact. Finally, coming back to Figure 7(a), we see that the value of $\eta\eta^*$ goes towards zero faster than \mathcal{E} (here around the order $O(h^{1.75})$), what illustrates the fact that \mathcal{R} is clearly superconvergent (see (49)).

4.2.2 A moving Gaussian


Finally we consider a more difficult test coming from [21]. This time, the primal problem is defined by (26) with $d = 2$, $\Omega =]0, 1]^2$, $T = 1.0$ and $D = I_{\mathbb{R}^2}$. We choose the exact solution being equal to $u(t, x, y) = \beta(t) e^{-50(r(x,y,t))^2}$ with $r(x, y, t) = ((x - 0.4t - 0.3)^2 + (y - 0.4t - 0.3)^2)^{1/2}$ and $\beta(t)$ defined by:

$$\beta(t) = \begin{cases} 1 - e^{-50(0.98t+0.01)^2} & \text{if } t \leq 0.5, \\ 1 - e^{-50(1-0.98t+0.01)^2} & \text{else.} \end{cases}$$

It consists in a moving Gaussian from the position (0.3, 0.3) to the position (0.7, 0.7) at a non constant speed. As usual, the right-hand-side f is computed accordingly. The dual problem is defined by (30), where we choose $g = 0$ and $u_T^* = e^{-\frac{\rho(x,y)^2}{32 \ln 10}}$ with $\rho(x, y) = ((x - 0.7)^2 + (y -$



(a)




(b)


Figure 7: Heat equation. (a) : rates of convergence. (b) : asymptotical exactness.

$0.7)^2)^{1/2}$.

For the numerical simulation, we set $\tau = h/10$, all the other numerical parameters are the same as the ones of section 4.2.1. We can see in Figure 8(a) that once again, the estimator η_{QOI} converges towards zero at the order $O(h)$ as theoretically expected, whereas the product $\eta\eta^*$ goes faster towards zero (in the order $O(h^{1.75})$). As expected by Theorems 3.2 and 3.4, Figure 8(b) illustrates the asymptotic exactness of the estimator since the effectivity index $I_{eff} = \mathcal{E}/\eta_{QOI}$ goes towards 1 when h goes towards zero.



(a)



(b)

Figure 8: Moving Gaussian. (a) : rates of convergence. (b) : asymptotical exactness.

References

- [1] M. Ainsworth and J. T. Oden. *A Posteriori Error Estimation in Finite Element Analysis*. Wiley, New York, 2000.
- [2] M. Ainsworth and R. Rankin. Guaranteed computable bounds on quantities of interest in finite element computations. *Internat. J. Numer. Methods Engrg.*, 89(13):1605–1634, 2012.
- [3] W. Bangerth and R. Rannacher. *Adaptive finite element methods for differential equations*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2003.
- [4] R. Becker and R. Rannacher. An optimal control approach to a posteriori error estimation in finite element methods. *Acta Numer.*, 10:1–102, 2001.
- [5] M. Costabel, M. Dauge, and S. Nicaise. Singularities of Maxwell interface problems. *M2AN Math. Model. Numer. Anal.*, 33(3):627–649, 1999.
- [6] R. Courant and D. Hilbert. *Methods of mathematical physics. Vol. I*. Interscience Publishers, Inc., New York, N.Y., 1953.
- [7] J. Droniou. Intégration et Espaces de Sobolev à Valeurs Vectorielles. working paper or preprint, Apr. 2001.
- [8] A. Ern, A. F. Stephansen, and M. Vohralík. Guaranteed and robust discontinuous Galerkin a posteriori error estimates for convection-diffusion-reaction problems. *J. Comput. Appl. Math.*, 234(1):114–130, 2010.
- [9] A. Ern and M. Vohralík. A posteriori error estimation based on potential and flux reconstruction for the heat equation. *SIAM J. Numer. Anal.*, 48(1):198–223, 2010.
- [10] A. Ern and M. Vohralík. A unified framework for a posteriori error estimation in elliptic and parabolic problems with application to finite volumes. In *Finite volumes for complex applications VI. Problems & perspectives. Volume 1, 2*, volume 4 of *Springer Proc. Math.*, pages 821–837. Springer, Heidelberg, 2011.
- [11] F. Hecht. New development in freefem++. *J. Numer. Math.*, 20(3-4):251–265, 2012.
- [12] P. Ladevèze and L. Chamoin. Calculation of strict error bounds for finite element approximations of non-linear pointwise quantities of interest. *Internat. J. Numer. Methods Engrg.*, 84(13):1638–1664, 2010.
- [13] P. Ladevèze, F. Pled, and L. Chamoin. New bounding techniques for goal-oriented error estimation applied to linear problems. *Internat. J. Numer. Methods Engrg.*, 93(13):1345–1380, 2013.
- [14] J.-L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications. Vol. 1*. Travaux et Recherches Mathématiques, No. 17. Dunod, Paris, 1968.
- [15] G. Mallik, M. Vohralík, and S. Yousef. Goal-oriented a posteriori error estimation for conforming and nonconforming approximations with inexact solvers. *J. Comput. Appl. Math.*, 366:112367, 20, 2020.
- [16] M. S. Mommer and R. Stevenson. A goal-oriented adaptive finite element method with convergence rates. *SIAM J. Numer. Anal.*, 47(2):861–886, 2009.
- [17] I. Mozolevski and S. Prudhomme. Goal-oriented error estimation based on equilibrated-flux reconstruction for finite element approximations of elliptic problems. *Comput. Methods Appl. Mech. Engrg.*, 288:127–145, 2015.

- [18] J. Muñoz Matute, E. Alberdi, D. Pardo, and V. M. Calo. Time-domain goal-oriented adaptivity using pseudo-dual error representations. *Comput. Methods Appl. Mech. Engrg.*, 325:395–415, 2017.
- [19] J. Muñoz Matute, V. M. Calo, D. Pardo, E. Alberdi, and K. G. van der Zee. Explicit-in-time goal-oriented adaptivity. *Comput. Methods Appl. Mech. Engrg.*, 347:176–200, 2019.
- [20] J. Muñoz Matute, D. Pardo, V. M. Calo, and E. Alberdi. Forward-in-time goal-oriented adaptivity. *Internat. J. Numer. Methods Engrg.*, 119(6):490–505, 2019.
- [21] S. Nicaise and N. Soualem. A posteriori error estimates for a nonconforming finite element discretization of the heat equation. *M2AN Math. Model. Numer. Anal.*, 39(2):319–348, 2005.
- [22] J. T. Oden and S. Prudhomme. Goal-oriented error estimation and adaptivity for the finite element method. *Comput. Math. Appl.*, 41(5-6):735–756, 2001.
- [23] S. Prudhomme and J. T. Oden. On goal-oriented error estimation for elliptic problems: application to the control of pointwise errors. volume 176, pages 313–331. 1999. *New advances in computational methods (Cachan, 1997)*.
- [24] V. Rey, P. Gosselet, and C. Rey. Strict bounding of quantities of interest in computations based on domain decomposition. *Comput. Methods Appl. Mech. Engrg.*, 287:212–228, 2015.
- [25] V. Rey, P. Gosselet, and C. Rey. Strict lower bounds with separation of sources of error in non-overlapping domain decomposition methods. *Internat. J. Numer. Methods Engrg.*, 108(9):1007–1029, 2016.
- [26] Z. Tang, S. Lou, A. Benabou, E. Creusé, S. Nicaise, J. Korecki, and J.-C. Mipo. Guaranteed quantity of interest error estimate based on equilibrated flux reconstruction. *IEEE Transactions on Magnetics*, 57(6):1–4, 2021.