

Supporting Information

New protective coatings against lampenflora growing in the Pommery Champagne cellar

Isabel Franco-Castillo^{1,2}, Archismita Misra³, Sébastien Laratte⁴, Maxime Gommeaux⁴, Robin Perarnau⁴, Nathalie Vaillant-Gaveau⁵, Clément Pierlot⁶, Carsten Streb³, Scott G. Mitchell^{1,2*}, Stéphanie Eyssautier-Chuine^{4*}

¹ Instituto de Nanociencia y Materiales de Aragón (INMA-CSIC), Consejo Superior de Investigaciones Científicas-Universidad de Zaragoza, c/ Pedro Cerbuna 12, 50009 Zaragoza, Spain.

² CIBER de Bioingeniería, Biomateriales y Nanomedicina, Instituto de Salud Carlos III, 28029 Madrid, Spain.

* Email: scott.mitchell@csic.es

³ Institute of Inorganic Chemistry I, Ulm University, Albert-Einstein-Allee 11, 89081 Ulm, Germany

⁴ Groupe d'Étude sur les Géomatériaux et les Environnements Naturels Anthropiques et Archéologiques 3795 (GEGENAA) - SFR Condorcet FR CNRS 3417 – 2, Esplanade Roland Garros, Université de Reims Champagne-Ardenne, 51100 cedex, Reims, France.

* Email: stephanie.eyssautier@univ-reims.fr

⁵ Unité de Recherche EA 4707 Résistance Induite et Bioprotection des Plantes (RIBP), SFR Condorcet FR CNRS 3417, UFR Sciences Exactes et Naturelles, Université de Reims Champagne-Ardenne, Reims, France.

⁶ Vranken-Pommery Monopole, 5 Place du Général Gouraud, BP1049, 51689, Reims, cedex 2, France

Supporting Information



Figure S1. Localisation of the sampling locations for the identification of the biofilm microorganisms.

Table S1. Applied quantity of liquid product on each slab and equivalent dry weight for the three biocidal assays.

Assay	Weight of liquid coating/slab (g)			Weight of dry coating/slab (g)		
	POM-IL1	POM-IL2	PR80	POM-IL1	POM-IL2	PR80
1	0.24 ± 0.01	0.21 ± 0.01	0.28 ± 0.03	0.04 ± 0.01	0.03 ± 0.01	0.15 ± 0.02
2	0.42 ± 0.01	0.40 ± 0.01	0.44 ± 0.01	0.08 ± 0.02	0.09 ± 0.01	0.24 ± 0.01
3	0.10 ± 0.03	0.12 ± 0.01	0.11 ± 0.01			
	0.20 ± 0.04	0.22 ± 0.03	0.23 ± 0.02			
	0.41 ± 0.02	0.40 ± 0.01	0.41 ± 0.01			

Supporting Information

Table S2. Identification of algae and cyanobacteria species by the closest affiliation in NBCI BlastN with primer, similarity percentage. The nucleotide sequence data have been deposited NCBI nucleotide sequence database under accession number. *CR_GenBank accession number.

Phylogenetic closest affiliation of isolates	Primer	Similarity (%)	Accession number*
<i>Chromochloris zofingiensis</i> strain ACSSI	18S rRNA	98.19	OK217227.1
<i>Pseudostichococcus monallantoides</i> strain UTEX 2249	rbcL	98.46	OK432519
<i>Diplosphaera</i> sp. cort12	rbcL	98.28	OK432520
<i>Timaviella</i> sp.	16S rRNA	98.00	OK422863

Table S3. Hoagland medium composition.

Hoagland - Component	Stock Solution	Stock Solution ml / L
2M KNO ₃	202 g/L	2.5
2M Ca(NO ₃) ₂ •4H ₂ O	472 g/L	2.5
2M MgSO ₄ •7H ₂ O	493 g/L	1
1M KH ₂ PO ₄	136 g/L	1
H ₃ BO ₃	2.86 g/L	1
MnCl ₂ •4H ₂ O	1.81 g/L	1
ZnSO ₄ •7H ₂ O	0.22 g/L	1
CuSO ₄ •5H ₂ O	0.08 g/L	1
Na ₂ MoO ₄ •2H ₂ O ☆	0.12 g/L	1
Iron(III)-EDTA	0.14 g/L	1.5

Supporting Information

Table S4. Δa^* and F_v/F_M average and standard deviation for Assay 1.

coating	Time (week)	<i>Chromochloris</i> sp. inoculation				<i>Pseudostichococcus</i> sp. inoculation			
		Δa^* average	Δa^* SD	F_v/F_M average	F_v/F_M SD	Δa^* average	Δa^* SD	F_v/F_M average	F_v/F_M SD
NO	0			0.47	0.03			0.28	0.07
	1	-3.3	0.8	0.61	0.03	-14.0	2.0	0.44	0.07
	2	-7.3	1.3	0.49	0.05	-20.2	1.2	0.62	0.01
	3	-12.8	2.6	0.47	0.05	-17.5	2.2	0.50	0.02
	4	-15.1	3.7	0.50	0.02	-17.5	2.2	0.48	0.01
	5	-17.6	2.1	0.41	0.03	-19.3	1.1	0.50	0.03
	7	-14.6	2.3	0.39	0.03	-21.1	1.1	0.44	0.04
POM- IL1	0			0.44	0.05			0.28	0.07
	1	0.3	0.4	0.56	0.04	0.0	0.2	0.23	0.05
	2	0.7	0.4	0.08	0.04	0.2	0.3	0.20	0.02
	3	0.7	0.4	0.16	0.04	-0.8	0.6	0.14	0.04
	4	0.7	0.4	0.05	0.01	-2.9	1.3	0.22	0.08
	5	0.5	0.4	0.13	0.02	-6.3	1.8	0.24	0.09
	7	-0.9	0.7	0.17	0.04	-10.0	2.9	0.35	0.08
POM- IL2	0			0.38	0.03			0.33	0.04
	1	-1.4	0.3	0.57	0.02	-9.6	2.0	0.30	0.04
	2	-3.0	0.4	0.35	0.09	-18.5	1.3	0.57	0.02
	3	-7.5	1.0	0.31	0.07	-16.0	2.4	0.42	0.01
	4	-10.3	1.9	0.36	0.06	-15.8	2.6	0.47	0.02
	5	-12.6	2.8	0.45	0.03	-16.8	1.6	0.43	0.03
	7	-15.7	3.4	0.40	0.03	-17.0	1.6	0.37	0.04
PR80	0			0.41	0.04			0.34	0.04
	1	-1.4	0.5	0.50	0.02	-1.2	0.8	0.33	0.06
	2	-3.4	1.2	0.36	0.08	-5.9	3.3	0.41	0.06
	3	-9.1	2.3	0.32	0.05	-14.4	2.9	0.46	0.02
	4	-14.2	2.0	0.42	0.04	-16.3	2.1	0.41	0.03
	5	-16.7	1.1	0.46	0.02	-17.1	2.4	0.38	0.03
	7	-17.9	2.1	0.40	0.04	-16.5	1.8	0.31	0.04

Supporting Information

Table S5. Δa^* and F_v/F_M average and standard deviation for Assay 2.

coating	Time (week)	Δa^* average	Δa^* SD	F_v/F_M average	F_v/F_M SD
NO	0			0.12	0.06
	1	-2.1	1.8	0.32	0.11
	2	-15.7	3.3	0.47	0.05
	3	-18.4	2.0	0.42	0.04
	4	-19.2	1.5	0.46	0.02
	5	-19.1	1.7	0.33	0.03
POM- IL1	0			0.15	0.06
	1	0.4	0.3	0.06	0.03
	2	0.5	0.3	0.08	0.02
	3	0.5	0.3	0.03	0.01
	4	0.5	0.3	0.04	0.01
	5	0.4	0.3	0.03	0.01
POM- IL2	0			0.12	0.05
	1	0.5	0.2	0.05	0.02
	2	0.6	0.2	0.02	0.02
	3	0.5	0.2	0.01	0.02
	4	0.5	0.2	0.04	0.02
	5	0.5	0.2	0.01	0.01
PR80	0			0.14	0.04
	1	0.2	0.2	0.06	0.04
	2	0.4	0.2	0.03	0.02
	3	0.2	0.5	0.02	0.02
	4	-0.1	1.0	0.08	0.06
	5	-3.3	3.0	0.17	0.12

Supporting Information

Table S6. Δa^* and F_v/F_M average and standard deviation for Assay 3.

coating	Time (week)	Δa^* average	Δa^* SD	F_v/F_M average	F_v/F_M SD
NO	0			0.21	0.06
	1	-7.0	3.9	0.39	0.09
	2	-17.2	4.3	0.51	0.04
	3	-19.0	2.9	0.40	0.02
	4	-19.7	2.6	0.41	0.03
	5	-19.2	2.6	0.33	0.04
POM-IL1 0.1g	0			0.06	0.04
	1	0.7	0.4	0.05	0.02
	2	0.6	0.5	0.16	0.07
	3	0.6	0.4	0.06	0.02
	4	0.5	0.5	0.18	0.04
	5	0.5	0.5	0.06	0.03
POM-IL2 0.1g	0			0.04	0.02
	1	0.4	0.2	0.04	0.02
	2	0.5	0.2	0.12	0.11
	3	0.4	0.2	0.07	0.03
	4	0.2	0.3	0.14	0.04
	5	0.3	0.3	0.05	0.02
POM-IL1 0.2g	0			0.04	0.04
	1	1.0	0.4	0.04	0.01
	2	0.7	0.4	0.11	0.03
	3	0.8	0.4	0.06	0.03
	4	0.7	0.4	0.17	0.04
	5	0.7	0.4	0.04	0.02
POM-IL2 0.2g	0			0.05	0.05
	1	0.9	0.5	0.04	0.02
	2	0.8	0.6	0.11	0.04
	3	0.8	0.5	0.05	0.03
	4	0.8	0.6	0.19	0.03
	5	0.8	0.6	0.06	0.03
PR80 0.2g	0			0.06	0.04
	1	0.6	0.7	0.06	0.06
	2	0.4	1.3	0.19	0.10
	3	0.4	1.1	0.08	0.08
	4	-0.8	2.3	0.16	0.07
	5	-2.7	4.6	0.20	0.11

Supporting Information

coating	Time (week)	Δa^* average	Δa^* SD	F_v/F_M average	F_v/F_M SD
POM- IL1 0.4g	0			0.09	0.04
	1	1.0	0.7	0.02	0.01
	2	0.8	0.8	0.11	0.04
	3	0.9	0.8	0.06	0.02
	4	0.8	0.8	0.12	0.03
	5	0.8	0.8	0.04	0.05
POM- IL2 0.4g	0			0.09	0.05
	1	0.8	0.5	0.03	0.02
	2	0.7	0.5	0.09	0.04
	3	0.8	0.5	0.06	0.02
	4	0.7	0.5	0.13	0.04
	5	0.7	0.5	0.02	0.01
PR80 0.4g	0			0.05	0.02
	1	1.0	0.2	0.04	0.02
	2	0.8	0.3	0.09	0.03
	3	0.9	0.2	0.06	0.02
	4	0.8	0.2	0.14	0.03
	5	0.8	0.3	0.05	0.02

Supporting Information

Table S7. Δa^* and F_v/F_M average and standard deviation for Assay 4.

coating	Time (week)	Δa^* average	Δa^* SD	F_v/F_M average	F_v/F_M SD
NO	0			0.14	0.04
	1	-4.8	1.8	0.35	0.07
	2	-18.5	3.5	0.40	0.04
	3	-20.1	2.3	0.58	0.02
	4	-19.6	1.9	0.32	0.04
	5	-18.7	1.5	0.45	0.04
POM-IL1 0.1g wet	0			0.13	0.04
	1	0.3	0.4	0.18	0.05
	2	0.5	0.5	0.20	0.06
	3	0.5	0.5	0.11	0.04
	4	0.5	0.4	0.11	0.05
	5	-0.3	1.0	0.15	0.06
POM-IL2 0.1g wet	0			0.13	0.03
	1	0.3	0.4	0.17	0.05
	2	-0.4	1.0	0.18	0.05
	3	-2.5	3.3	0.16	0.04
	4	-8.0	6.6	0.24	0.09
	5	-14.4	8.2	0.27	0.06
POM-IL1 0.2g wet	0			0.16	0.04
	1	0.1	0.5	0.15	0.04
	2	0.4	0.6	0.09	0.09
	3	0.4	0.6	0.01	0.04
	4	0.2	0.8	0.16	0.05
	5	-0.4	1.4	0.13	0.05
POM-IL2 0.2g wet	0			0.14	0.04
	1	0.5	0.5	0.15	0.05
	2	0.6	0.4	0.15	0.04
	3	0.3	0.7	0.06	0.09
	4	-0.4	2.0	0.14	0.08
	5	-2.0	4.9	0.12	0.05
POM-IL1 0.4g dry	0			0.13	0.04
	1	0.5	0.3	0.18	0.06
	2	0.7	0.3	0.15	0.09
	3	0.6	0.3	0.16	0.04
	4	0.6	0.4	0.18	0.06
	5	0.4	0.6	0.13	0.04
POM-IL2 0.4g dry	0			0.12	0.04
	1	0.4	0.3	0.18	0.04
	2	0.6	0.2	0.11	0.09
	3	0.5	0.3	0.16	0.05
	4	0.5	0.4	0.19	0.08
	5	0.2	0.6	0.18	0.07

Supporting Information

PR80 0.4g wet	0		0.17	0.06
	1	-3.9	2.5	0.39
	2	-16.2	5.5	0.43
	3	-18.4	4.4	0.42
	4	-18.2	3.3	0.57
	5	-16.2	3.2	0.56