



HAL
open science

Non parametric estimation of the jump coefficient of a diffusion with jumps

Émeline Schmisser

► **To cite this version:**

Émeline Schmisser. Non parametric estimation of the jump coefficient of a diffusion with jumps. 2024. hal-03721374v2

HAL Id: hal-03721374

<https://hal.science/hal-03721374v2>

Preprint submitted on 29 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non parametric estimation of the jump coefficient of a diffusion with jumps

Émeline Schmisser

Université de Lille, CNRS, UMR 8524 - Laboratoire Paul Painlevé
Villeneuve d'Ascq, France

email: emeline.schmisser@univ-lille.fr

This work was supported by the Labex CEMPI (ANR-11-LABX-0007-01)

Abstract

In this article, we consider a jump diffusion process $(X_t)_{t \geq 0}$ with drift function b , diffusion coefficient σ and jump coefficient ξ . This process is supposed to be ergodic, exponentially β -mixing and stationary. It is observed at discrete times $t = 0, \Delta, \dots, n\Delta$. The sampling interval Δ tends to 0 and the time interval $n\Delta$ tends to infinity. We construct a robust, adaptive non-parametric estimator of the function ξ^4 thanks to a penalized least-square approach. We provide bounds of the empirical and L^2 -risk of our estimator.

Keywords: jump diffusions, nonparametric estimation, model selection.

Subject classification: 62G05, 62M05.

The code of the simulation study is available on math.univ-lille1.fr/~schmisser/recherche.html

1 Introduction

We consider the jump diffusion given by the one-dimensional stochastic differential equation

$$dX_t = b(X_t)dt + \sigma(X_t)dW_t + \xi(X_{t-})dL_t, \quad X_0 = \eta \quad (1)$$

with η a random variable, $(W_t)_{t \geq 0}$ a Brownian motion independent of η and $(L_t)_{t \geq 0}$ a pure jump centred Lévy process independent of (W_t, η) . It can be written

$$L_t = \int_0^t \int_{\mathbb{R}} z(\mu(dz, dt) - \nu(dz)dt)$$

where μ is a Poisson measure of intensity $\nu(dz)dt$. We do not assume that the jumps are of finite intensity, only that L_t is centred and has moments of any order.

The diffusion is observed at discrete times $t = 0, \Delta, \dots, n\Delta$ under the asymptotic framework: $n\Delta \rightarrow \infty$ and $\Delta \rightarrow 0$. Our aim is to construct adaptive non-parametric estimator of the jump coefficient ξ^4 on a compact interval $A \subset \mathbb{R}$.

Jump diffusions are used to modelize dynamical systems where the noise is discontinuous or too intensive to be modeled by a Brownian motion, for instance polymerization phenomenons

26 (Berestycki (2004)), telephone noise (Protter and Talay (1997)), or finance (see Aït-Sahalia and
 27 Jacod (2009) or Protter and Talay (1997) for instance). There are many articles on the estimation
 28 for jumps diffusions, although quite few estimate the jump coefficient nonparametrically.

29 For the parametric estimation, we can cite Shimizu (2006) or Shimizu and Yoshida (2006) who
 30 estimate the three coefficients b , σ and ξ for stationary finite intensity processes thanks to a contrast
 31 function. Their estimators are consistent and asymptotically normal if the sampling interval Δ is
 32 sufficiently small. The estimators of the parameter of b and ξ^2 converge with rate $\sqrt{n\Delta}$, whereas
 33 the estimator of the parameter of σ^2 converges with rate \sqrt{n} .

34 For nonparametric estimation, Bandi and Nguyen (2003) use kernel estimators of the infinitesi-
 35 mal moments, $M_1 = b(x)$, $M_2 = \sigma^2(x) + \xi^2(x) \int z^2 \nu(dz)$ and $M_4 = \xi^4(x) \int z^4 \nu(dz)$. Their estima-
 36 tors are asymptotically normal and converge with rate $(n\Delta)^{-2/5}$. Their estimator do not depends of
 37 the regularity of the functions, and are not adaptive. Hanif et al. (2012) and Hanif (2016) construct
 38 estimators of the infinitesimal second moment $\sigma^2 + \xi^2 \int z^2 \nu(dz)$ with reweighted Nadaraya-Watson
 39 and Gamma Nadaraya-Watson estimators respectively. Their estimators converge almost surely,
 40 and are asymptotically normal.

41 To estimate σ^2 or ξ^2 non parametrically, a threshold is often used to suppress (or find) the
 42 jumps. For instance, Mancini (2009) estimate the integrated volatility. The estimator converge,
 43 and in the finite activity cas, the estimator is asymptotically normal. Mancini and Renò (2011)
 44 (for the finite intensity case) and Song et al. (2022) (for compound Poisson processes) estimate the
 45 diffusion coefficient σ^2 , using local time and kernel estimators. Schmisser (2019) construct a robust
 46 and adaptive estimator of σ^2 and of the second infinitesimal moment, even for infinite intensity
 47 Lévy processes.

48 Mancini and Renò (2011) for the finite activity case and Park and Wang (2021) for the finite
 49 intensity case both construct estimators for the jump coefficient ξ^2 , using local times and kernel
 50 estimator. The rate of convergence of the kernel estimator of ξ^2 is $(n\Delta)^{-2/5} + \Delta^{1/2}$.

51 There are two possibilities to construct an estimator of the jump component. Either take the
 52 second infinitesimal moment when the increments are big enough (when a jump is suspected):
 53 indeed, for finite intensity processes (see for instance Park and Wang (2021, Section 5.2)).

$$\frac{(X_{(k+1)\Delta} - X_{k\Delta})^2}{\Delta} \mathbf{1}_{|X_{(k+1)\Delta} - X_{k\Delta}| \geq \Delta^{1/2-\varepsilon}} = \xi^2(X_{k\Delta}) \int z^2 \nu(dz) + \text{centred terms} + \text{small terms}.$$

54 The problem of this method is that the small jumps will not be kept. It is not very important if
 55 the Lévy process is a compound Poisson process, but if the jumps have infinite activity or infinite
 56 intensity, the small terms will not be so small.

57 In this article, like Bandi and Nguyen (2003), we use the second possibility and consider an
 58 estimator based on the fourth moments of the increments:

$$T_{k\Delta} = \frac{(X_{(k+1)\Delta} - X_{k\Delta})^4}{\Delta} = \xi^4(X_{k\Delta}) \int z^4 \nu(dz) + \text{centred terms} + \text{small terms}.$$

59 The Brownian terms are small because of the power 4, and it only remains the jumps. We introduce
 60 a sequence of increasing subspaces S_m of $L^2(A)$ and construct a sequence of estimators $\hat{\xi}_m^4$ by
 61 minimizing on each S_m the contrast function $\gamma_n(t) = \frac{1}{n} \sum_{k=1}^n (T_{k\Delta} - t(X_{k\Delta}))^2$. When $\int z^4 \nu(dz)$ is
 62 known, its L^2 -risk is bounded by

$$\mathbb{E} \left(\left\| \hat{\xi}^4 - \xi^4 \right\|_{L^2(A)}^2 \right) \lesssim \underbrace{\left\| \xi^4 - \xi_m^4 \right\|_{L^2(A)}^2}_{\text{bias term}} + \underbrace{\frac{\Xi D_m}{n\Delta}}_{\text{variance term}} + \Delta$$

63 where D_m is the dimension of the subspace S_m . The bias term decreases when the dimension
64 D_m increases, whereas the variance term increases. To find a good bias-variance compromise, we
65 introduce a penalty, $\widehat{pen}(m)$, proportional to the variance term. To choose the dimension, we
66 minimize the quantity $\gamma_n(\widehat{\xi}_m^4 I_4) + \widehat{pen}(m)$ and deduce the "best" dimension \tilde{m} . Finally, we prove
67 that the risk of the robust estimator, $\widehat{\xi}_{\tilde{m}}^4$, satisfies an oracle inequality.

68 The paper is divided as follows: the model and its assumptions are stated in Section 2. In
69 Section 3, we construct the estimator and bound its risk. Section 4 is devoted to the simulations
70 and the proofs are gathered in Section 5.

71 We introduce some notations.

72 In all the paper, C means a constant that does not depend on n , Δ or the dimension D_m , but
73 may be different from one line to another. We note $A \lesssim B$ if $A \leq CB$ where C is a constant that
74 does not depend on n , Δ or D_m . \mathbb{R}_+^* is the set of positive numbers, and \mathcal{C}^r the set of fonctions t
75 that are r times continuously differentiable.

76 We set $I_k = \int z^k \nu(dz)$ if this quantity exists.

77 For a function t in $L^2(A)$, and π a density, $\|t\|_\pi^2 = \int_A t^2(x) \pi(dx)$ and the empirical norm is
78 $\|t\|_n^2 = \frac{1}{n} \sum_{i=1}^n t^2(X_{k\Delta})$. We consider three different projections: t_m is the orthogonal projection of
79 t on S_m (for the L^2 -norm), $t_{m,\pi}$ is the orthogonal projection of t on S_m for the $\|\cdot\|_\pi$ -norm and $\Pi_m t$
80 is the orthogonal projection on S_m for the empirical norm $\|\cdot\|_n$.

81 2 Model and assumptions

82 We consider the stochastic differential equation given by (1) and assume the following assumptions:

83 Assumption A1.

84 a. The functions b , σ and ξ are Lipschitz.

85 b. The drift function b is elastic:

$$\exists M > 0, \exists C > 0, \forall x, |x| \geq M, xb(x) \leq -Cx^2.$$

86 c. The diffusion and jump functions σ and ξ are bounded from below and above: there exists
87 $\sigma_0, \sigma_1, \xi_0, \xi_1 \in (\mathbb{R}_+^*)^4$ such that

$$\forall x, 0 < \xi_0 \leq \xi(x) \leq \xi_1 \quad \text{and} \quad 0 < \sigma_0 \leq \sigma(x) \leq \sigma_1.$$

88 d. The Lévy measure has moments of any order: $\forall k \geq 2, \int_{\mathbb{R}} |z|^k \nu(dz) < +\infty$.

89 e. $I_4 = \int_{\mathbb{R}} z^4 \nu(dz) = 1$.

90 Item a ensures that SDE (1) has a unique strong solution (see Applebaum (2009, Theorem 6.2.9)).
91 According to Masuda (2007, Theorem 2.2 and Lemma 2.4), under Assumptions A1a-d, the process
92 $(X_t)_{t \geq 0}$ has a unique stationary distribution, is ergodic and exponentially β -mixing. Assumption e
93 ensures that the model is identifiable: indeed, if we replace $\nu(\cdot)$ by $c\nu(\cdot)$ and ξ by ξ/c , we obtain
94 the same process $(X_t)_{t \geq 0}$. If this assumption is not fulfilled, our estimator will estimate $\xi^4(X_{k\Delta})I_4$,
95 wich enables us to see at least the shape of ξ .

96 We can now assume:

97 **Assumption A2.**

98 a. The process $(X_t)_{t \geq 0}$ is stationary.

99 b. Its stationary measure has a density π which is bounded on any compact set.

100 c. the density π is bounded from below and above on the compact of estimation A :

$$\exists \pi_0, \pi_1; \forall x \in A, \quad 0 < \pi_0 \leq \pi(x) \leq \pi_1.$$

101 By Masuda (2007, Theorem 2.2), it implies that X_t has also moments of any order.

102 Let us consider the σ -algebra $\mathcal{F}_t = \sigma(\eta, (W_s)_{0 \leq s \leq t}, (L_s)_{0 \leq s \leq t})$.

103 The following proposition helps us to control the increments of the process. It is proved for
104 instance in Applebaum (2009, Theorems 4.4.21 and 4.4.23)).

105 **Proposition 1.** According to the Burkholder-Davis-Gundy inequality, for any $p > 0$, there exists
106 a constant C_p such that:

$$\mathbb{E} \left(\sup_{s \in [t, t+h]} \left| \int_t^s \sigma(X_u) dW_u \right|^p \middle| \mathcal{F}_t \right) \leq C_p \mathbb{E} \left(\left| \int_t^{t+h} \sigma^2(X_u) du \right|^{p/2} \middle| \mathcal{F}_t \right)$$

107 and

$$\begin{aligned} \mathbb{E} \left(\sup_{s \in [t, t+h]} \left| \int_t^s \xi(X_{u-}) dL_u \right|^p \middle| \mathcal{F}_t \right) &\leq C_p \mathbb{E} \left(\left| \int_t^{t+h} \xi^2(X_u) du \right|^{p/2} \middle| \mathcal{F}_t \right) I_2^{p/2} \\ &+ C_p \mathbb{E} \left(\int_t^{t+h} |\xi^p(X_u)| du \middle| \mathcal{F}_t \right) I_p. \end{aligned}$$

108 To simplify the notations, let us set

$$B_{k\Delta} = \int_{k\Delta}^{(k+1)\Delta} b(X_s) ds, \quad Z_{k\Delta} = \int_{k\Delta}^{(k+1)\Delta} \sigma(X_s) dW_s \quad \text{and} \quad J_{k\Delta} = \int_{k\Delta}^{(k+1)\Delta} \xi(X_{s-}) dL_s.$$

109 The following lemma follows almost directly from the Burkholder-Davis-Gundy inequality (see for
110 instance Schmisser (2014, proof of Proposition 1)).

111 **Lemma 2.** For any integer $p > 0$:

$$\forall u > 0, \quad \mathbb{E} \left(\sup_{0 \leq s \leq \Delta} (X_{s+u})^{2p} \middle| \mathcal{F}_u \right) \lesssim 1 + |X_u|^{2p}$$

112 and

$$\forall u \geq 0, \quad \mathbb{E} \left(\sup_{0 \leq s \leq t} (X_{s+u} - X_u)^{2p} \middle| \mathcal{F}_u \right) \leq C_{2p} \left(t \xi_1^{2p} I_{2p} + t^p \left(\sigma_1^{2p} + \xi_1^{2p} I_2^p \right) \right).$$

113 Then, as b , σ^2 and ξ^2 are Lipschitz: $\mathbb{E}(|B_{k\Delta}|^{2p}) \lesssim \Delta^{2p}$, $\mathbb{E}(J_{k\Delta}^{2p}) \lesssim \Delta^p$ and $\mathbb{E}(J_{k\Delta})^{2p} =$
114 $\Delta \xi^{2p}(X_{k\Delta}) I_{2p} + O(\Delta^{3/2})$.

115 We need to control the variance term, more precisely the moments of the jumps. The Lévy-
 116 Khintchine formula states that $\mathbb{E}(e^{iuL_\Delta}) = e^{-\Delta \int_{\mathbb{R}} (e^{iuz} - 1 - iuz)\nu(dz)}$. The following lemma is exactly
 117 Lemma 13 of [Carpentier et al. \(2021\)](#).

118 **Lemma 3.** *We get the following moments:*

$$\begin{aligned} \mathbb{E}(L_\Delta^2) &= I_2\Delta \\ \mathbb{E}(L_\Delta^4) &= I_4\Delta + 3I_2^2\Delta^2 \\ \mathbb{E}(L_\Delta^6) &= I_6\Delta + (15I_2I_4 + 10I_3^2)\Delta^2 + 15I_2^3\Delta^3 \\ \mathbb{E}(L_\Delta^8) &= I_8\Delta + (35I_4^2 + 56I_3I_5 + 28I_2I_6)\Delta^2 + (210I_2^2I_4 + 280I_2I_3^2)\Delta^3 + 105I_2^4\Delta^4. \end{aligned}$$

119 To bound the risk the adaptive estimator, we apply a Talagrand's inequality. To this end, we
 120 need independent and bounded random variables. As we have an exponentially β -mixing process,
 121 we can construct independent random variables thanks to the Berbee's coupling lemma. But to
 122 work with nearly bounded variables, we need the following assumption:

123 **Assumption A3.**

- 124 a. *The jumps are sub-exponential:* $\exists M, C, \lambda \in (\mathbb{R}_+^*)^3, \forall z, |z| \geq M \nu([-z, z]^c) \leq Ce^{-\lambda|z|}$.
 125 b. $\exists \eta, \eta > 1$, such that $\Delta^\eta = O(n^{-1})$.
 126 c. *The Blumenthal-Gettoor index is strictly less than 2:* there exists $\beta < 2$, $\int_{-1}^1 z^\beta \nu(dz) < \infty$.
 127 *This is not a very strong assumption, as $\int_{-1}^1 z^2 \nu(dz)$ is already finite.*

128 Under Assumption A1 and 3, for any $c \leq \lambda/\xi_1$, any $t > 0$, $\mathbb{E}(e^{cX_t}) < \infty$.

129 *Proof.* Apply Theorem 2.2 of [Masuda \(2007\)](#) with

$$f^*(x) = e^{-cx} \mathbf{1}_{|x| \geq 1} + e^c \left(1 + \frac{c^2}{2} - \frac{5c}{8} + \frac{3c - c^2}{4} x^2 + \frac{c^2 - c}{8} x^4 \right) \mathbf{1}_{|x| < 1}. \quad \square$$

130 Under this assumption, the increments can be bounded. The small jumps (smaller than $\sqrt{\Delta}$)
 131 have the same comportement than the Brownian motion terms. As we have only sub-exponential-
 132 jumps, the size of the bigger jumps is also nearly bounded. The following lemma is proved in
 133 [Schmisser \(2019, Lemma 2\)](#):

134 **Lemma 4.** *Under Assumptions A1-A3, we have:*

- 135 • $\forall \varepsilon > 0, \forall r > 0, \mathbb{P}(|B_{k\Delta}| \geq \Delta^{1-\alpha}) \lesssim n^{-r}$.
 136 • $\forall r > 0, \mathbb{P}(|Z_{k\Delta}| \geq r\sigma_1 \Delta^{1/2} \ln(n)) \leq 2n^{-r}$.
 137 • $\forall p > 0, \forall r > 0$, for any C_J and $c(p)$ large enough:

$$\mathbb{P}(|J_{k\Delta}| \geq C_J r^2 \ln(n)/\lambda) \lesssim n^{-r} \quad \text{and} \quad \mathbb{P}\left(\frac{1}{q_n} \sum_{j=1}^{q_n} J_{k\Delta}^{2p} \geq c(p)\xi_1^{2p}(r+1)^2 \Delta \ln^{2p}(n)\right) \lesssim n^{-r}$$

138 where $q_n \geq 2 \ln(n)/\beta$, and β the β -mixing coefficient.

3 Estimation of the jump coefficient

Our aim is to construct an adaptive nonparametric estimator of the jump function ξ on the interval A . To make the diffusion terms small, we consider the fourth moments of the increments:

$$T_{k\Delta} = \frac{(X_{(k+1)\Delta} - X_{k\Delta})^4}{\Delta} = \frac{(B_{k\Delta} + Z_{k\Delta} + J_{k\Delta})^4}{\Delta}.$$

We can write $T_{k\Delta} = \xi^4(X_{k\Delta})I_4 + E_{k\Delta} + F_{k\Delta}$ where

$$\begin{aligned} \Delta E_{k\Delta} &:= (B_{k\Delta} + Z_{k\Delta} + J_{k\Delta})^4 - (Z_{k\Delta} + J_{k\Delta})^4 + \mathbb{E}((Z_{k\Delta} + J_{k\Delta})^4 | \mathcal{F}_{k\Delta}) - \xi^4(X_{k\Delta})I_4\Delta \\ \Delta F_{k\Delta} &:= (Z_{k\Delta} + J_{k\Delta})^4 - \mathbb{E}((Z_{k\Delta} + J_{k\Delta})^4 | \mathcal{F}_{k\Delta}). \end{aligned}$$

The terms $E_{k\Delta}$ are small, whereas $F_{k\Delta}$ are centred. The following Lemma is proved in Section 5, using Lemma 2.

Lemma 5.

- $\mathbb{E}(E_{k\Delta}^2) \lesssim \Delta$, $\mathbb{E}(E_{k\Delta}^4) \lesssim \Delta$ and $\mathbb{E}(E_{k\Delta}^8) \lesssim \Delta$.
- $\mathbb{E}(F_{k\Delta} | \mathcal{F}_{k\Delta}) = 0$, $\mathbb{E}(F_{k\Delta}^4) \lesssim 1/\Delta^3$ and

$$\text{Var}(F_{k\Delta} | \mathcal{F}_{k\Delta}) = \frac{\xi^8(X_{k\Delta})I_8}{\Delta} + O\left(\frac{1}{\sqrt{\Delta}}\right).$$

3.1 Subspaces of approximation

To construct an adaptive estimator of ξ^4 , we consider a sequence of increasing subspaces $S_0 \subseteq \dots \subseteq S_m, \dots$ such that $\cup_m S_m$ is dense in $L^2(A)$. We minimize a contrast function $\gamma_n(t)$ on each S_m and then choose the best estimator by introducing a penalty function (see for instance Barron et al. (1999)). The rate of convergence of our estimator will depend on the regularity of the coefficient ξ , i.e. its modulus of smoothness.

Definition (Modulus of smoothness). *The modulus of continuity of a function f is defined by*

$$\omega(f, t) = \sup_{|x-y| \leq t} |f(x) - f(y)|.$$

If f is Lipschitz, the modulus of continuity is proportional to t . If $\omega(f, t) = o(t)$, then f is constant: the modulus of continuity cannot measure higher smoothness.

We define the modulus of smoothness by

$$\omega_r(f, t)_p = \sup_{0 < h \leq t} \|\Delta_h^r(f, \cdot)\|_{L^p} \quad \text{where} \quad \Delta_h^r(f, x) = \sum_{k=0}^r (-1)^k \binom{r}{k} f(x + kh).$$

If $f \in \mathcal{C}^r$ (that is if f is r -times continuously differentiable), then for $1 \leq p \leq \infty$:

$$\omega_r(f, t)_p \leq t^r \omega(f^{(r)}, t)_p.$$

159 **Definition** (Besov space). *The Besov space $B_{2,\infty}^\alpha$ is the set of functions:*

$$B_{2,\infty}^\alpha = \{f \in L^2, \sup_{t>0} t^{-\alpha} \omega_r(f, t)_2 < \infty\}$$

160 *where $r = \lfloor \alpha + 1 \rfloor$. The norm on a Besov space is defined by:*

$$\|f\|_{B_{2,\infty}^\alpha} := \sup_{t>0} t^{-\alpha} \omega_r(f, t)_2 + \|f\|_{L^2}.$$

161 For more details see [DeVore and Lorentz \(1993\)](#).

162 **Assumption A4.** *We consider a series of vectorial subspaces of $L^2(A)$, $(S_m)_{m \in \mathcal{M}_n}$ satisfying the*
 163 *assumptions*

164 a. *The subspaces S_m have finite dimension D_m and are increasing: $\forall m, S_m \subseteq S_{m+1}$.*

165 b. *The $\|\cdot\|_{L^2}$ and $\|\cdot\|_\infty$ norms are connected:*

$$\exists \phi_1, \forall m, \forall t \in S_m, \quad \|t\|_\infty^2 \leq \phi_1 D_m \|t\|_{L^2}^2 (A)$$

166 *with $\|t\|_{L^2(A)}^2 = \int_A t^2(x) dx$ and $\|t\|_\infty = \sup_{x \in A} |t(x)|$. This implies that, for an orthonormal*
 167 *basis φ_λ of S_m , $\left\| \sum_{\lambda=1}^{D_m} \varphi_\lambda^2 \right\|_\infty \leq \phi_1 D_m$.*

168 c. *There exists a constant ϕ_2 such that for any $m \in \mathbb{N}$, there exists an orthonormal basis $(\psi_\lambda)_{\lambda \in \Lambda}$*
 169 *of S_m such that*

$$\forall \lambda, \text{card}(\lambda', \|\psi_\lambda \psi_{\lambda'}\|_\infty \neq 0) \leq \phi_2.$$

170 d. *For any function $t \in \mathcal{B}_{2,\infty}^\alpha$,*

$$\exists c, \forall m, \|t - t_m\|_{L^2}^2 \leq c D_m^{-2\alpha}$$

171 *where t_m is the orthogonal projection L^2 of t on S_m .*

172 Assumptions **A4 a, b** and **d** are quite standard (see for instance [Comte et al. \(2007\)](#)). Assumption
 173 **A4 c** is an additional assumption, which is not satisfied by the trigonometric basis, for instance.
 174 However, it is fulfilled by subspaces generated by piecewise polynomials of degree r , spline functions
 175 of degree r , or compactly supported wavelets.

176 3.2 Estimator with fixed m

177 We then define the contrast function

$$\gamma_n(t) = \frac{1}{n} \sum_{k=1}^n (T_{k\Delta} - t(X_{k\Delta}))^2 \quad (2)$$

178 and for any $m \in \mathcal{M}_n = \{m, D_m^2 \leq (n\Delta)/\ln(n)\}$, we consider the estimator

$$\hat{\xi}_m^4 = \arg \min_{t \in S_m} \gamma_n(t). \quad (3)$$

179 The easiest bound is obtained for the empirical risk.

180 **Proposition 6.** Under Assumptions A1-A4, for any m such as $D_m \in \mathcal{M}_n$:

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 \right) \leq \|\xi^4 - \xi_{m,\pi}^4\|_\pi^2 + 12\Xi \frac{D_m}{n\Delta} + C\Delta + \frac{C'}{n}$$

181 where $\xi_A(x) = \xi(x)\mathbf{1}_{x \in A}$ and

$$\Xi = \min \left(\xi_1^8 I_8, \frac{\phi_1}{\pi_0} \mathbb{E}_\pi (\xi^8(X_\Delta)) I_8 \right).$$

182 The empirical risk is not very intuitive: indeed, the empirical norm is random, and the stationary
183 measure π is unknown. Nevertheless, the L^2 -risk can also be bounded, even if the bound is less
184 sharp.

185 **Corollary 7.** Under Assumptions A1-A4, for any m such as $D_m \leq \mathcal{D}_n = \sqrt{n\Delta/\ln(n)}$:

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi^4 \right\|_\pi^2 \right) \leq 5 \|\xi^4 - \xi_{m,\pi}^4\|_\pi^2 + 48\Xi \frac{D_m}{n\Delta} + 4C\Delta.$$

186 Moreover,

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi^4 \right\|_{L^2(A)}^2 \right) \leq 5 \frac{\pi_1}{\pi_0} \|\xi^4 - \xi_m^4\|_{L^2(A)}^2 + 48 \frac{\Xi}{\pi_0} \frac{D_m}{n\Delta} + \frac{4C}{\pi_0} \Delta.$$

187 3.3 Adaptive estimator

188 We obtain a sequence of estimators of ξ^4 , and have to choose the best. If ξ^4 belongs to the Besov
189 space $\mathcal{B}_{2,\infty}^\alpha$, then the bias term $\|\xi^4 - \xi_m^4\|_{L^2}^2$ is smaller than $D_m^{-2\alpha}$ and the risk is optimal for
190 $D_m = (n\Delta)^{-1/(1+2\alpha)}$. The optimal estimator satisfies:

$$\mathbb{E} \left(\left\| \hat{\xi}_{m_{opt}}^4 - \xi_A^4 \right\|_n^2 \right) \lesssim (n\Delta)^{-2\alpha/(2\alpha+1)} + \Delta.$$

191 The rate $(n\Delta)^{-2\alpha/(2\alpha+1)}$ is standard for nonparametric estimation of ξ . As the regularity of ξ^4
192 is unknown, we have to choose the best estimator automatically. To this aim, we introduce the
193 penalty function $pen(m) = \kappa\Xi D_m/(n\Delta)$ and choose the best dimension m :

$$\hat{m} = \arg \min_{m \in \mathcal{M}_n} \{ \gamma_n(\hat{\xi}_m^4) + pen(m) \}$$

194 Our estimator achieves the best rate of convergence (up to a multiplicative constant).

195 **Theorem 8.** Under Assumptions A1-A3, for any $\kappa \geq \kappa_0$ with κ_0 a universal constant,

$$\mathbb{E} \left(\left\| \hat{\xi}_{\hat{m}}^4 - \xi_A^4 \right\|_n^2 \right) \lesssim \inf_{m \in \mathcal{M}_n} \left(\|\xi^4 - \xi_{m,\pi}^4\|_\pi^2 + \Xi \frac{D_m}{n\Delta} \right) + \frac{1}{n\Delta} + \Delta.$$

196 The bound of the L^2 -risk of the estimator is the same:

Corollary 9.

$$\mathbb{E} \left(\left\| \hat{\xi}_{\hat{m}}^4 - \xi^4 \right\|_{L^2(A)}^2 \right) \lesssim \inf_{m \in \mathcal{M}_n} \left(\|\xi^4 - \xi_m^4\|_{L^2(A)}^2 + \Xi \frac{D_m}{n\Delta} \right) + \frac{1}{n\Delta} + \Delta.$$

3.4 Robust estimator

The constant $\Xi = \min(\xi_1^8 I_8, \frac{\phi_1}{\pi_0} \mathbb{E}_\pi (\xi^8(X_\Delta)) I_8)$ is unknown, but can be estimated to construct a robust estimator. Let us first estimate $h(x) = \xi^8(x) I_8$. Let us note

$$V_{k\Delta} := \frac{(X_{(k+1)\Delta} - X_{k\Delta})^8}{\Delta} = \xi^8(X_{k\Delta}) I_8 + G_{k\Delta} + H_{k\Delta},$$

where $G_{k\Delta}$ is small and $H_{k\Delta}$ is centred:

$$\begin{aligned} \Delta G_{k\Delta} &= (B_{k\Delta} + Z_{k\Delta} + J_{k\Delta})^8 - (Z_{k\Delta} + J_{k\Delta})^8 + \mathbb{E}((Z_{k\Delta} + J_{k\Delta})^8 | \mathcal{F}_{k\Delta}) - \xi^8(X_{k\Delta}) I_8 \\ \Delta H_{k\Delta} &= (Z_{k\Delta} + J_{k\Delta})^8 - \mathbb{E}((Z_{k\Delta} + J_{k\Delta})^8 | \mathcal{F}_{k\Delta}). \end{aligned}$$

The following lemma bounds the moments of G and H :

Lemma 10.

$$\begin{aligned} \mathbb{E}(G_{k\Delta}^2 | \mathcal{F}_{k\Delta}) &\lesssim \Delta, \quad \mathbb{E}(G_{k\Delta}^4 | \mathcal{F}_{k\Delta}) \lesssim \Delta, \\ \mathbb{E}(H_{k\Delta} | \mathcal{F}_{k\Delta}) &= 0, \quad \mathbb{E}(H_{k\Delta}^2 | \mathcal{F}_{k\Delta}) \lesssim \frac{1}{\Delta}, \quad \mathbb{E}(H_{k\Delta}^4 | \mathcal{F}_{k\Delta}) \lesssim \frac{1}{\Delta^3}. \end{aligned}$$

As previously, we consider the contrast function $\gamma_{2,n}(t) = \frac{1}{n} \sum_{k=1}^n (V_{k\Delta} - t(X_{k\Delta}))^2$ and the estimator $\hat{h}_m = \arg \min_{t \in S_m} \gamma_{2,n}(t)$. This estimator is consistent:

Lemma 11.

$$\mathbb{E} \left(\left\| h_A - \hat{h}_m \right\|_n^2 \right) \lesssim \|h - h_m\|_{L^2(A)}^2 + \frac{D_m}{n\Delta} + \Delta$$

where $h_A(x) = h(x) \mathbf{1}_{x \in A}$.

Let us set $\hat{h}_1 = \sup_{x \in A} |\hat{h}_{\ln(n)}(x)|$.

Let us now estimate the second term, $\phi_1/\pi_0 \mathbb{E}_\pi (\xi^8(X_\Delta)) I_8$. We consider the rectangular kernel $K(x) := \mathbf{1}_{|x| \leq 1/2}$ and set

$$\hat{\pi}_h(x) = \frac{1}{n} \sum_{k=1}^n K_h(X_i - x) \quad \text{with} \quad K_h(x) = \frac{1}{h} K(x/h).$$

We take a grid $(x_1, \dots, x_{\ln^2(n)})$ of equally spaced points of A , and compute the minimum of $\hat{\pi}_{\sqrt{n\Delta}}$ on this grid:

$$\hat{\pi}_0 := \min_{1 \leq j \leq \ln^2(n)} \hat{\pi}_{\sqrt{n\Delta}}(x_j).$$

Finally, we obtain an estimator of the constant Ξ :

$$\hat{\Xi} := \min \left(\hat{h}_1, \frac{\phi_1}{\hat{\pi}_0} \bar{V}_n \right) + \Delta^{1/4} + 1/\ln(n)$$

with $\bar{V}_n = \frac{1}{n} \sum_{k=1}^n V_{k\Delta}$. The estimator $\hat{\Xi}$ is consistent:

Lemma 12. *Under Assumptions A1-3, we have that*

$$\mathbb{P} \left(|\hat{\Xi} - \Xi| \geq C/\ln(n) + \Delta^{1/4} \right) \lesssim n^{-4}.$$

213 Let us set

$$\tilde{m} = \arg \inf_{m \in \mathcal{M}_n} \gamma_n(\hat{\xi}_m) + \kappa \hat{\Xi} \frac{D_m}{n\Delta}.$$

214 The resulting robust estimator $\hat{\xi}_{\tilde{m}}$ is also optimal (up to a multiplicative constant):

215 **Corollary 13.** *The robust estimator satisfies the oracle inequality: Under Assumptions A1-A3,*

$$\mathbb{E} \left(\left\| \hat{\xi}_{\tilde{m}}^4 - \xi_A^4 \right\|_n^2 \right) \lesssim \inf_{m \in \mathcal{M}_n} \left(\|\xi^4 - \xi_{m,\pi}^4\|_\pi^2 + \Xi \frac{D_m}{n\Delta} \right) + \frac{1}{n\Delta} + \Delta.$$

216 Moreover,

$$\mathbb{E} \left(\left\| \hat{\xi}_{\tilde{m}}^4 - \xi^4 \right\|_{L^2(A)}^2 \right) \lesssim \inf_{m \in \mathcal{M}_n} \left(\|\xi^4 - \xi_m^4\|_{L^2(A)}^2 + \Xi \frac{D_m}{n\Delta} \right) + \frac{1}{n\Delta} + \Delta.$$

217 3.5 Comparison with the estimator obtained by thresholding

218 Park and Wang (2021) and Mancini and Renò (2011) construct an estimator of ξ^2 by taking the
219 second infinitesimal moments and keeping only the increments big enough: they consider

$$\frac{(X_{(k+1)\Delta} - X_{k\Delta})^2}{\Delta} \mathbf{1}_{|X_{(k+1)\Delta} - X_{k\Delta}| \geq \Delta^{1/2-\varepsilon}}.$$

220 Then, they construct a kernel estimator of ξ^2 , in the case where the jumps have finite intensity and
221 ξ^2 belongs to \mathcal{C}^2 . Their estimator is asymptotically normal, and the risk for the optimal bandwidth
222 is bounded by

$$\mathbb{E} \left((\hat{\xi}^2(x) - \xi^2(x))^2 \right) \lesssim (n\Delta)^{-4/5} + \Delta.$$

223 Similarly, Schmisser (2019) estimate first the second infinitesimal moment $g = \sigma^2 + \xi^2$, then σ^2 by
224 thresholding. An estimator of ξ^2 can be obtained by subtraction. If g and σ^2 belong to a Besov
225 spaces of regularity α_g and α_{σ^2} , the risk of the adaptive estimator $\tilde{\xi}^2$ is bounded by:

$$\mathbb{E} \left(\left\| \tilde{\xi}^2 - \xi^2 \right\|_{L^2(A)}^2 \right) \lesssim (n\Delta)^{2\alpha_g/(2\alpha_g+1)} + n^{-2\alpha_\sigma/(2\alpha_\sigma+1)} + \Delta^{1-\beta/2}$$

226 where β is the Blumenthal-Gettoor index (that is, $\beta = \inf_{\alpha \geq 0} \{ \int (|z|^\alpha \wedge 1) \nu(dz) < \infty \}$). If the Lévy
227 process has finite activity, then $\beta = 0$. If the intensity is finite, $\beta \leq 1$. For both estimators, it is
228 assumed than $I_2 = \int z^2 \nu(dz) = 1$.

229 The problem is, the first estimator can only be calculated for finite intensity processes, and is
230 not adaptive. Whereas the rate of convergence of the second depends on the Blumenthal-Gettoor
231 index, and on the regularity of σ^2 .

232 In order to compare those bounds to the risk of our estimator $\hat{\xi}_{\tilde{m}}$, we can remark that

$$\left\| \sqrt{\hat{\xi}_{\tilde{m}}^4} - \xi^2 \right\|_{L^2(A)}^2 \leq \frac{\left\| \hat{\xi}_{\tilde{m}}^4 - \xi_A^4 \right\|_{L^2(A)}^2}{\min_A \xi^2(x)}.$$

233 Therefore, we get that, if I_4 is known:

$$\mathbb{E} \left(\left\| \sqrt{\hat{\xi}_{\tilde{m}}^4} - \xi^2 \right\|_{L^2(A)}^2 \right) \lesssim (n\Delta)^{-2\alpha/(2\alpha+1)} + \Delta.$$

234 This bound is more general than the rate obtained by [Park and Wang \(2021\)](#), and sharper than
 235 the one obtained by [Schmisser \(2019\)](#). In particular, if the jump coefficient ξ is more regular than
 236 the diffusion coefficient σ^2 , the rate is better. Moreover, if the jumps have infinite activity, the
 237 remainder term is smaller.

238 However, we assume in this paper than the fourth moment of the Levy measure, $I_4 = \int z^4 \nu(dz)$,
 239 is equal to 1, which is not the standard assumption. If no assumptions are made on the moments
 240 of ν , the function ξ is not identifiable. So it can only be estimated up to a multiplicative constant.
 241 Even if we want to impose $I_2 = 1$, it is interessant to know the shape of ξ .

242 4 Simulations

243 We choose to consider the linear subspaces S_m generated by the spline functions of degree r (that
 244 is, piecewise polynomials of degree r and \mathcal{C}^{r-1}). It is also possible to consider piecewise polynomials
 245 of degree r or subspaces generated by wavelets. We choose the spline functions because they are
 246 explicits, contrary to wavelets functions, and because the resulting estimator is continuous if $r \geq 1$,
 247 which is not the case for piecewise polynomials.

248 We compute the series of estimators $\hat{\xi}_{m,r}^4$ for r between 0 and 4 and m between 0 and $\max(\sqrt{n\Delta/\ln(n)}, 7)$
 249 (for $m = 7$, we already have $D_m = 128$, for m bigger, the computations become too slow). To obtain
 250 the adaptive estimator, we minimise first with regard to m , then to r .

251 We choose $\kappa = 8$ for the simulations. We consider 3 different models. For each model, we let n
 252 and Δ vary and realize 200 simulations for each value of (n, Δ) . We compute the robust estimator
 253 $\hat{\xi}_{\hat{m},\hat{r}}^4$, the selected dimension $D_{\hat{m},\hat{r}}$, the estimated penalty constant $\hat{\Xi}$ and the L^2 -error

$$\text{risk} = \sum_{j=0}^{N-1} \left(\hat{\xi}_{\hat{m},\hat{r}}^4(x_j) - \xi^4(x_j) \right)^2 (x_{j+1} - x_j) \simeq \int_A \left(\hat{\xi}_{\hat{m},\hat{r}}^4(x) - \xi^4(x) \right)^2 dx$$

254 for (x_0, x_1, \dots, x_N) a regular subdivision of A . We also compute the empirical error for each $\hat{\xi}_{m,r}$.
 255 Then we deduce the dimension (m_{opt}, r_{opt}) that minimizes the empirical error (denoted by err_{min}).
 256 To compare this oracle estimator to the robust one, we compute an oracle or which is the mean
 257 of the logarithmic ratio between the two errors. To compare our estimator to the one obtained
 258 by substraction, we also compute the error obtained by the square root of our estimator, $\text{risk1} =$
 259 $\left\| \sqrt{\hat{\xi}_{\hat{m},\hat{r}}^4} - \xi_A^2 \right\|_{L^2}^2$ and the risk of the previous estimator obtained by thresholding and substraction
 260 in [Schmisser \(2019\)](#): $\text{risk2} := \left\| \tilde{\xi}^2 - \xi_A^2 \right\|_{L^2}^2$. We add the empirical risks of the estimators of the
 261 diffusion coefficient σ^2 and the second moment $g(x) = \sigma^2(x) + \xi^2(x)$, also computed in [Schmisser](#)
 262 [\(2019\)](#).

263 For each model, we provide two tables. The first one show some results for the estimator $\hat{\xi}_{\hat{m},\hat{r}}^4$:
 264 the mean and standard deviation of the risk of the robust estimator $\hat{\xi}_{\hat{m},\hat{r}}^4$ (in fact, the square root of
 265 these quantity, to have a norm), a criteria of comparison between the risks of the robust estimator
 266 and the 'oracle' estimator; or = $\text{mean}(\ln(\text{risk}/\text{err}))$, the mean and the standard deviation of the
 267 estimated penalty constant $\hat{\Xi}$ and the mean and the standard deviation of the selected dimension
 268 $D_{\hat{m},\hat{r}}$.

269 On the second table, we write the mean and standard deviations of risk1 and risk2 (more
 270 precisely, the square roots of those quantities) in order to compare these two estimators. We also

271 add the means and standard deviation for the risk of the estimators of g and σ^2 .

272 4.1 Models

273 In the first two models, L_t is simply a compound Poisson Process:

$$\int_0^t \xi(X_{s-}) dL_s = \sum_{k=1}^{N_t} \xi(X_{t_k^-}) \zeta_k$$

274 where N_t is a Poisson Process of intensity λ , t_1, \dots, t_{N_t} the times of the jumps, and $(\zeta_k)_{k \geq 0}$ are
 275 independent, identically distributed of binomial law: $\mathbb{P}(\zeta_k = \pm 1) = 1/2$.

276 **Model 1: Ornstein-Uhlenbeck** Let us consider the process given by the equation:

$$dX_t = -2X_t dt + dW_t + dL_t$$

277 with binomial jumps: $\mathbb{P}(\zeta = 1) = \mathbb{P}(\zeta = -1) = 0.5$ and the intensity of the Poisson process is 1.
 278 We have that $I_2 = 1$, $I_4 = 1$, $I_6 = 1$, $I_8 = 1$.

Model 2: non-constant coefficient

$$dX_t = -2X_t dt + \sqrt{2 + 0.5 \sin(\pi X_t)} dW_t + \sqrt{\frac{X_{t-}^2 + 3}{X_{t-}^2 + 1}} dL_t$$

279 with binomial jumps: $\mathbb{P}(\zeta = 1) = \mathbb{P}(\zeta = -1) = 0.5$.

Model 3: nearly stable processes

$$dX_t = -2X_t + \sqrt{\frac{3 + X_t^2}{1 + X_t^2}} dW_t + \sqrt{2 + 0.5 \sin(X_{t-})} dL_t$$

280 with L_t a nearly stable process: $\nu(dz) = \frac{2-\beta}{2|z|^{1+\beta}} \mathbf{1}_{|z| \leq 1}$. β is the Blumenthal-Gettoor index of the
 281 process. We take $\beta = 1/2$ and we get that $I_2 = 1$, $I_4 = 3/7$, $I_6 = 3/11$ and $I_8 = 3/15$.


282 4.2 Results

283 The risk of the adaptive estimator $\hat{\xi}_{\hat{m}, \hat{r}}^4$ decreases when n increases (for fixed Δ). When the period
 284 of observation ($n\Delta$) is fixed, the risk seems to be either stable, either decreasing with Δ , especially
 285 when Δ is not small enough. When $n\Delta \leq 100$, our estimator nearly always select the smallest
 286 dimension, whereas for $n\Delta \geq 10^3$, the algorithm can select more complex models. Our estimator
 287 is truly adaptive, and the oracle seems to be quite small.

288 For the estimated penalty constant $\hat{\Xi}$, it can be noted that when n increases, the variance
 289 decreases as well: $\hat{\Xi}$ converges to a constant depending on Δ . When Δ decreases, this constant
 290 decreases as well. This is because

$$\bar{V}_n \rightarrow \mathbb{E}(V_\Delta) \simeq \Delta^{-1} \mathbb{E}((J_\Delta + Z_{k\Delta})^8) \geq \mathbb{E}(\xi^8(X_\Delta)) I_8.$$

Figure 1: Model 1
 $dX_t = -2X_t dt + dW_t + dL_t$ with binomial jumps.



— estimator $\sqrt{\hat{\xi}_{\bar{m}, \bar{r}}^4}$... estimator $\tilde{\xi}^2 = \hat{g}_{\bar{m}} - \hat{\sigma}_{\bar{m}}^2$.
 $n = 10^5, \Delta = 10^{-2}$.
 Risk of $\hat{\xi}_{\bar{m}, \bar{r}}^4$


n	Δ	risk			$\hat{\Xi}$		$D_{\bar{m}, \bar{r}}$	
		mean	sd	or	mean	sd	mean	sd
10^3	10^{-1}	0.67	0.75	0	27.5	43.3	1	0
10^4	10^{-1}	0.61	0.36	0	27.7	24.0	1	0
10^5	10^{-1}	0.60	0.21	0.0014	22.0	11.7	1.06	0.42
10^3	10^{-2}	0.50	0.92	0	4.54	7.99	1	0
10^4	10^{-2}	0.16	0.21	0	4.44	6.22	1	0
10^5	10^{-2}	0.087	0.089	0	3.86	2.49	1	0
10^4	10^{-3}	0.37	0.54	0	2.76	3.50	1	0
10^5	10^{-3}	0.11	0.14	0	2.13	2.42	1	0

Comparison of the estimators $\sqrt{\hat{\xi}_{\bar{m}, \bar{r}}^4}$ and $\tilde{\xi}^2$.

n	Δ	risk1		risk2		risk g		risk σ^2	
		mean	sd	mean	sd	mean	sd	mean	sd
10^3	10^{-1}	0.28	0.29	0.23	0.22	0.34	0.24	0.42	0.15
10^4	10^{-1}	0.27	0.15	0.20	0.12	0.33	0.14	0.42	0.091
10^5	10^{-1}	0.27	0.12	0.20	0.084	0.25	0.083	0.41	0.049
10^3	10^{-2}	0.22	0.32	0.37	0.43	0.40	0.42	0.12	0.075
10^4	10^{-2}	0.075	0.094	0.15	0.14	0.23	0.14	0.11	0.039
10^5	10^{-2}	0.042	0.043	0.14	0.11	0.21	0.050	0.093	0.067
10^4	10^{-3}	0.18	0.23	0.36	0.43	0.40	0.42	0.10	0.043
10^5	10^{-3}	0.056	0.069	0.15	0.15	0.23	0.15	0.099	0.039

Figure 2: Model 2

$$dX_t = -2X_t dt + \sqrt{2 + 0.5 \sin(\pi X_t)} dW_t + \sqrt{\frac{X_t^2 + 3}{X_t^2 + 1}} dL_t \text{ with binomial jumps.}$$



-- estimator $\sqrt{\hat{\xi}_{\tilde{m},\tilde{r}}^4}$... estimator $\tilde{\xi}^2 = \hat{g}_{\tilde{m}} - \hat{\sigma}_{\tilde{m}}^2$.

$n = 10^5, \Delta = 10^{-2}$.

Risk of the estimator $\hat{\xi}_{\tilde{m},\tilde{r}}^4$.


n	Δ	risk			$\hat{\Xi}$		$D_{\tilde{m},\tilde{r}}$	
		mean	sd	or	mean	sd	mean	sd
10^3	10^{-1}	2.87	3.20	0.19	614	870	1	0
10^4	10^{-1}	2.29	1.33	0.33	394	240	1.01	0.07
10^5	10^{-1}	2.10	1.02	0.16	338	141	2.35	1.53
10^3	10^{-2}	3.63	4.92	0.11	128	189	1	0
10^4	10^{-2}	1.86	1.25	0.70	131	174	1	0
10^5	10^{-2}	0.82	0.92	0.44	101	24	3.55	1.07
10^4	10^{-3}	3.25	3.51	0.10	91	50	1	0
10^5	10^{-3}	1.83	1.07	0.94	89	72	1.01	0.07

Comparison of the estimators $\sqrt{\hat{\xi}_{\tilde{m},\tilde{r}}^4}$ and $\tilde{\xi}^2$.

n	Δ	risk1		risk2		risk g		risk σ^2	
		mean	sd	mean	sd	risk	sd	risk	sd
10^3	10^{-1}	0.46	0.54	0.84	0.73	0.99	0.63	1.44	0.29
10^4	10^{-1}	0.37	0.26	0.77	0.38	0.95	0.39	1.43	0.18
10^5	10^{-1}	0.46	0.30	0.78	0.29	0.85	0.32	1.42	0.10
10^3	10^{-2}	0.58	0.75	1.24	1.59	1.25	1.43	0.55	0.31
10^4	10^{-2}	0.21	0.25	0.67	0.55	0.71	0.39	0.456	0.19
10^5	10^{-2}	0.48	0.38	0.54	0.35	0.48	0.38	0.43	0.084
10^4	10^{-3}	0.53	0.60	1.21	1.28	1.21	1.28	0.11	0.13
10^5	10^{-3}	0.20	0.22	0.71	0.42	0.71	0.42	0.033	0.026

Figure 3: Model 3:

$$dX_t = -2X_t dt + \sqrt{\frac{X_{t-}^2 + 3}{X_{t-}^2 + 1}} dW_t + \sqrt{2 + 0.5 \sin(\pi X_{t-})} dL_t \quad \nu(z) = \frac{1.5}{2|z|^{1.5}} \mathbf{1}_{|z| \leq 1}$$



--- estimator $\sqrt{\hat{\xi}_{\tilde{m}, \tilde{r}}^4}$... estimator $\tilde{\xi}^2 = \hat{g}_{\tilde{m}} - \hat{\sigma}_{\tilde{m}}^2$.
 $n = 10^5, \Delta = 10^{-2}$.
 Estimation of $\hat{\xi}_{\tilde{m}, \tilde{r}}^4$.

n	Δ	risk			$\hat{\Xi}$		$D_{\tilde{m}, \tilde{r}}$	
		mean	sd	or	mean	sd	mean	sd
10^3	10^{-1}	3.91	2.83	0.0098	182	222	1	0
10^4	10^{-1}	3.75	1.57	0.015	135	84	1.04	0.18
10^5	10^{-1}	3.73	0.88	0.0081	117	45	1.58	0.55
10^3	10^{-2}	1.41	2.24	0.23	23.0	61.9	1	0
10^4	10^{-2}	0.86	0.66	0.67	17.3	18.4	1.03	0.17
10^5	10^{-2}	0.61	0.35	0.20	15.1	8.1	2.11	0.97
10^4	10^{-3}	1.07	1.33	0.29	12.8	27.9	1	0
10^5	10^{-3}	0.68	0.45	1.16	10.2	17	1.14	0.35

Comparison of the estimators $\sqrt{\hat{\xi}_{\tilde{m}, \tilde{r}}^4}$ and $\tilde{\xi}^2$.

n	Δ	risk1		risk2		risk g		risk σ^2	
		mean	sd	mean	sd	mean	sd	mean	sd
10^3	10^{-1}	1.60	1.01	1.33	0.77	0.97	0.58	1.99	0.34
10^4	10^{-1}	1.58	0.61	1.30	0.47	0.94	0.36	1.99	0.21
10^5	10^{-1}	1.66	0.58	1.28	0.26	0.86	0.32	1.99	0.13
10^3	10^{-2}	0.59	0.86	0.93	0.97	0.84	0.72	0.82	0.37
10^4	10^{-2}	0.36	0.42	0.83	0.49	0.65	0.23	0.78	0.23
10^5	10^{-2}	0.59	0.36	0.71	0.31	0.24	0.28	0.76	0.11
10^4	10^{-3}	0.48	0.62	0.84	0.69	0.84	0.70	0.11	0.11
10^5	10^{-3}	0.33	0.50	0.64	0.30	0.65	0.30	0.057	0.038

291 This is not a drawback, as the penalty should be the variance of the centred terms, more precisely

$$\Xi^\Delta = \Delta \min \left(\|\text{Var}(F_{k\Delta} | \mathcal{F}_{k\Delta})\|_\infty; \frac{\phi_1}{\pi_0} \text{Var}_\pi(F_\Delta) \right).$$

292 Ξ is an estimator of this quantity, that converges when $\Delta \rightarrow 0$ and $n\Delta \rightarrow \infty$: by Proposition 6,
293 $\Xi^\Delta = \Xi + C\Delta^{1/2}$. In the simple case where the jump and diffusion coefficients are constants, then

$$\begin{aligned} \Xi^\Delta &= \Delta^{-1} \text{Var}(F_{k\Delta}) = \Delta^{-1} \text{Var}((\xi L_{k\Delta} + \sigma Z_{k\Delta})^4) \\ &= \Delta^{-1} (\xi^8 \mathbb{E}(L_{k\Delta}^8) + 28\sigma^2 \xi^6 \mathbb{E}(L_{k\Delta}^6 Z_{k\Delta}^2) - \xi^8 (\mathbb{E}(L_{k\Delta}^4))^2) + O(\Delta^2) \\ &= \xi^8 I_8 + \Delta (\xi^8 (34I_4^2 + 28I_2 I_6) + \sigma^2 \xi^6 28I_6) + O(\Delta^2). \end{aligned}$$

294 When Δ is not small enough, the term in Δ is not negligible, neither is the term in $O(\Delta^2)$.

295 Our estimator $\sqrt{\hat{\xi}_m^4}$ is better than $\hat{\xi}^2$ as soon as the path of discretisation Δ is small enough
296 ($\Delta \leq 10^{-2}$), even for compound Poisson processes.

297 5 Proof

298 5.1 Proof of Lemma 5 and 10

299 According to Proposition 1, we have that

$$\mathbb{E}(B_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) \lesssim \Delta^{2p}, \quad \mathbb{E}(Z_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) \lesssim \Delta^p, \quad \text{and} \quad \mathbb{E}(J_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) \lesssim \Delta I_{2p} + O(\Delta^2). \quad (4)$$

300 Let us set

$$\begin{aligned} \tilde{B}_{k\Delta} &= b(X_{k\Delta})\Delta, & \bar{B}_{k\Delta} &= B_{k\Delta} - \tilde{B}_{k\Delta}, \\ \tilde{Z}_{k\Delta} &= \sigma(X_{k\Delta})(W_{(k+1)\Delta} - W_{k\Delta}), & \bar{Z}_{k\Delta} &= Z_{k\Delta} - \tilde{Z}_{k\Delta}, \\ \tilde{J}_{k\Delta} &= \xi(X_{k\Delta})(L_{(k+1)\Delta} - L_{k\Delta}), & \bar{J}_{k\Delta} &= J_{k\Delta} - \tilde{J}_{k\Delta}. \end{aligned}$$

301 Conditionnally to $\mathcal{F}_{k\Delta}$, $\tilde{B}_{k\Delta}$, $\tilde{J}_{k\Delta}$ and $\tilde{Z}_{k\Delta}$ are independent. Moreover, we have that

$$\mathbb{E}(\bar{B}_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) \lesssim \Delta^{2p+1}, \quad \mathbb{E}(\bar{Z}_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) \lesssim \Delta^{p+1}, \quad \mathbb{E}(\bar{J}_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) \lesssim \Delta^2 I_{2p}. \quad (5)$$

302 Indeed, by Hölder inequality and Lemma 2, as the drift function b is Lipschitz,

$$\begin{aligned} \mathbb{E}(\bar{B}_{k\Delta}^{2p} | \mathcal{F}_{k\Delta}) &= \mathbb{E} \left(\left(\int_{k\Delta}^{(k+1)\Delta} (b(X_s) - b(X_{k\Delta})) ds \right)^{2p} \middle| \mathcal{F}_{k\Delta} \right) \\ &\leq \Delta^{2p-1} \int_{k\Delta}^{(k+1)\Delta} \mathbb{E}((b(X_s) - b(X_{k\Delta}))^{2p} | \mathcal{F}_{k\Delta}) ds \\ &\lesssim \Delta^{2p-1} \int_{k\Delta}^{(k+1)\Delta} \mathbb{E}(|X_s - X_{k\Delta}|^{2p} | \mathcal{F}_{k\Delta}) ds \lesssim \Delta^{2p+1}. \end{aligned}$$

303 By Proposition 1 and the same arguments, we get:

$$\begin{aligned}\mathbb{E} \left(\bar{Z}_{k\Delta}^{2p} \right) &\lesssim \mathbb{E} \left(\left(\int_{k\Delta}^{(k+1)\Delta} (\sigma(X_s) - \sigma(X_{k\Delta}))^2 ds \right)^p \right) \\ &\lesssim \Delta^{p-1} \int_{k\Delta}^{(k+1)\Delta} \mathbb{E} \left((\sigma(X_s) - \sigma(X_{k\Delta}))^{2p} \right) ds \lesssim \Delta^{p+1}\end{aligned}$$

304 and

$$\begin{aligned}\mathbb{E} \left(\bar{J}_{k\Delta}^{2p} \middle| \mathcal{F}_{k\Delta} \right) &\lesssim \mathbb{E} \left(\int_{k\Delta}^{(k+1)\Delta} (\xi(X_{s-}) - \xi(X_{k\Delta}))^{2p} ds \middle| \mathcal{F}_{k\Delta} \right) I_{2p} \\ &\quad + \mathbb{E} \left(\left(\int_{k\Delta}^{(k+1)\Delta} (\xi(X_{s-}) - \xi(X_{k\Delta}))^2 ds \right)^p \middle| \mathcal{F}_{k\Delta} \right) I_2^p \\ &\lesssim \Delta^2 I_{2p} + \Delta^{p+1} I_2^p.\end{aligned}$$

305 We first bound the moments of the small terms, $E_{k\Delta}$ in Lemma 5 and $G_{k\Delta}$ in Lemma 10.

306 **Bound of $E_{k\Delta}$.** Let us decompose the bias term

$$\Delta E_{k\Delta} = (B_{k\Delta} + Z_{k\Delta} + J_{k\Delta})^4 - (Z_{k\Delta} + J_{k\Delta})^4 + \mathbb{E} \left((Z_{k\Delta} + J_{k\Delta})^4 \middle| \mathcal{F}_{k\Delta} \right) - \Delta \xi^4(X_{k\Delta} I_4).$$

307 We can write $E_{k\Delta} = E_{k\Delta}^{(1)} + E_{k\Delta}^{(2)} + E_{k\Delta}^{(3)} + E_{k\Delta}^{(4)}$ where

$$\begin{aligned}\Delta E_{k\Delta}^{(1)} &= \mathbb{E} \left((J_{k\Delta} + Z_{k\Delta})^4 - (\tilde{J}_{k\Delta} + \tilde{Z}_{k\Delta})^4 \middle| \mathcal{F}_{k\Delta} \right) \\ \Delta E_{k\Delta}^{(2)} &= \mathbb{E} \left((\tilde{J}_{k\Delta} + \tilde{Z}_{k\Delta})^4 - \tilde{J}_{k\Delta}^4 \middle| \mathcal{F}_{k\Delta} \right) \\ \Delta E_{k\Delta}^{(3)} &= \mathbb{E} \left(\tilde{J}_{k\Delta}^4 \middle| \mathcal{F}_{k\Delta} \right) - \Delta \xi^4(X_{k\Delta} I_4) \\ \Delta E_{k\Delta}^{(4)} &= (J_{k\Delta} + Z_{k\Delta} + B_{k\Delta})^4 - (J_{k\Delta} + Z_{k\Delta})^4.\end{aligned}$$

308 The first three terms are conditional expectations, the last is random. For the first term, we factorize
309 the difference of two squares: indeed, for any $x, y \in \mathbb{R}^2$, $x^4 - y^4 = (x - y)(x + y)(x^2 + y^2)$. Then

$$\Delta E_{k\Delta}^{(1)} = \mathbb{E} \left((\bar{J}_{k\Delta} + \bar{Z}_{k\Delta}) \left(J_{k\Delta} + \tilde{J}_{k\Delta} + Z_{k\Delta} + \tilde{Z}_{k\Delta} \right) \left((J_{k\Delta} + Z_{k\Delta})^2 + (\tilde{J}_{k\Delta} + \tilde{Z}_{k\Delta})^2 \right) \middle| \mathcal{F}_{k\Delta} \right).$$

310 Then, as $\mathbb{E} \left((\bar{J}_{k\Delta} + \bar{Z}_{k\Delta})^2 \right) \lesssim \Delta^2$, we obtain by Cauchy-Schwarz inequality and (4):

$$\Delta E_{k\Delta}^{(1)} \lesssim (\Delta^2)^{1/2} \Delta^{1/4} \Delta^{1/4} \simeq \Delta^{3/2}.$$

311 Moreover, $E_{k\Delta}^{(2)}$ can be written $\Delta E_{k\Delta}^{(2)} = \sum_{j=1}^4 C_4^j \mathbb{E} \left(\tilde{Z}_{k\Delta}^j \tilde{J}_{k\Delta}^{4-j} \middle| \mathcal{F}_{k\Delta} \right)$. As $\tilde{Z}_{k\Delta}$ and $\tilde{J}_{k\Delta}$ are centred
312 and conditionnally independent, by (4), we obtain:

$$\Delta E_{k\Delta}^{(2)} \lesssim \sum_{j=1}^4 \Delta^{j/2} (\Delta \mathbf{1}_{j < 4} + \mathbf{1}_{j=4}) \lesssim \Delta^{3/2}.$$

313 Furthermore, by Lemma 3, $\mathbb{E}(L_{k\Delta}^4) = \Delta I_4 + O(\Delta^2)$ and therefore:

$$\Delta E_{k\Delta}^{(3)} = \xi^4(X_{k\Delta})(\mathbb{E}(L_{k\Delta}^4) - \Delta I_4) \lesssim \Delta^2.$$

314 As $\Delta E_{k\Delta}^{(4)} = \sum_{j=1}^4 C_4^j B_{k\Delta}^j (Z_{k\Delta} + J_{k\Delta})^{4-j}$, we get that

$$\mathbb{E}\left(\left(\Delta E_{k\Delta}^{(4)}\right)^{2p}\right) \lesssim \sum_{j=1}^4 \mathbb{E}\left(B_{k\Delta}^{2jp} (Z_{k\Delta} + J_{k\Delta})^{8p-2pj}\right). \quad (6)$$

315 By Cauchy-Schwartz inequality and (4), for $2 \leq j \leq 4$,

$$\begin{aligned} \mathbb{E}\left(B_{k\Delta}^{2jp} (Z_{k\Delta} + J_{k\Delta})^{8p-2pj}\right) &\leq \left(\mathbb{E}\left(B_{k\Delta}^{4jp}\right) \mathbb{E}\left((Z_{k\Delta} + J_{k\Delta})^{16p-4pj}\right)\right)^{1/2} \\ &\leq \Delta^{2jp} \left(\mathbf{1}_{j=4} + \Delta^{1/2} \mathbf{1}_{j<4}\right) \leq \Delta^{4p+1/2}. \end{aligned} \quad (7)$$

316 Moreover, as $B_{k\Delta} = \Delta b(X_{k\Delta}) + \bar{B}_{k\Delta}$, by Equations (4) and (5), for $j = 1$ we obtain:

$$\begin{aligned} \mathbb{E}\left(B_{k\Delta}^{2p} (Z_{k\Delta} + J_{k\Delta})^{6p}\right) &\lesssim \Delta^{2p} \mathbb{E}\left(b(X_{k\Delta})^{2p} \mathbb{E}\left((Z_{k\Delta} + J_{k\Delta})^{6p} \mid \mathcal{F}_{k\Delta}\right)\right) + \mathbb{E}\left(\bar{B}_{k\Delta}^{2p} (Z_{k\Delta} + J_{k\Delta})^{6p}\right) \\ &\lesssim \Delta^{2p+1} + \left(\mathbb{E}\left(\bar{B}_{k\Delta}^{4p}\right) \mathbb{E}\left((Z_{k\Delta} + J_{k\Delta})^{12p}\right)\right)^{1/2} \\ &\lesssim \Delta^{2p+1} + \Delta^{2p+1/2} \Delta^{1/2} = \Delta^{2p+1}. \end{aligned}$$

317 Then $\mathbb{E}\left(\left(\Delta E_{k\Delta}^{(4)}\right)^{2p}\right) \lesssim \Delta^{2p+1}$. As $\mathbb{E}\left(E_{k\Delta}^{2p}\right) \lesssim \mathbb{E}\left(\left(E_{k\Delta}^{(1)} + E_{k\Delta}^{(2)} + E_{k\Delta}^{(3)}\right)^{2p}\right) + \mathbb{E}\left(\left(E_{k\Delta}^{(4)}\right)^{2p}\right)$, we
318 get that, for $p \geq 1$,

$$\mathbb{E}\left(E_{k\Delta}^{2p}\right) \lesssim \Delta.$$

319 **Bound of $G_{k\Delta}$** We recall that

$$\Delta G_{k\Delta} = (B_{k\Delta} + Z_{k\Delta} + J_{k\Delta})^8 - (Z_{k\Delta} + J_{k\Delta})^8 + \mathbb{E}\left((Z_{k\Delta} + J_{k\Delta})^8 \mid \mathcal{F}_{k\Delta}\right) - \Delta \xi^8(X_{k\Delta}) I_8.$$

320 As for the bound of $E_{k\Delta}$, we can write $G_{k\Delta} = G_{k\Delta}^{(1)} + G_{k\Delta}^{(2)} + G_{k\Delta}^{(3)} + G_{k\Delta}^{(4)}$ with

$$\begin{aligned} \Delta G_{k\Delta}^{(1)} &:= \mathbb{E}\left((Z_{k\Delta} + J_{k\Delta})^8 \mid \mathcal{F}_{k\Delta}\right) - \mathbb{E}\left((\tilde{Z}_{k\Delta} + \tilde{J}_{k\Delta})^8 \mid \mathcal{F}_{k\Delta}\right) \\ \Delta G_{k\Delta}^{(2)} &:= \mathbb{E}\left((\tilde{Z}_{k\Delta} + \tilde{J}_{k\Delta})^8 \mid \mathcal{F}_{k\Delta}\right) - \mathbb{E}\left(\tilde{J}_{k\Delta}^8 \mid \mathcal{F}_{k\Delta}\right) \\ \Delta G_{k\Delta}^{(3)} &:= \mathbb{E}\left(\tilde{J}_{k\Delta}^8 \mid \mathcal{F}_{k\Delta}\right) - \Delta \xi^8(X_{k\Delta}) I_8 \\ \Delta G_{k\Delta}^{(4)} &:= (B_{k\Delta} + Z_{k\Delta} + J_{k\Delta})^8 - (Z_{k\Delta} + J_{k\Delta})^8. \end{aligned}$$

321 Let us note $JZ := J_{k\Delta} + Z_{k\Delta}$ and $\tilde{J}\tilde{Z} := \tilde{J}_{k\Delta} + \tilde{Z}_{k\Delta}$. Then by Cauchy-Schwarz, Equations (4) and
322 (5),

$$\begin{aligned} \Delta G_{k\Delta}^{(1)} &= \mathbb{E}\left((\bar{J}_{k\Delta} + \bar{Z}_{k\Delta})(JZ + \tilde{J}\tilde{Z})\left(JZ^2 + \tilde{J}\tilde{Z}^2\right)\left(JZ^4 + \tilde{J}\tilde{Z}^4\right) \mid \mathcal{F}_{k\Delta}\right) \\ &\lesssim (\Delta^2)^{1/2} \Delta^{1/4} \Delta^{1/8} \Delta^{1/8} \lesssim \Delta^{3/2}. \end{aligned}$$

323 Moreover, as $\tilde{Z}_{k\Delta}$ and $\tilde{J}_{k\Delta}$ are conditionnaly independant, we get:

$$\Delta G_{k\Delta}^{(2)} = \sum_{j=1}^8 C_8^j \mathbb{E} \left(\tilde{Z}_{k\Delta}^j \tilde{J}_{k\Delta}^{8-j} \middle| \mathcal{F}_{k\Delta} \right) \lesssim \sum_{j=1}^8 \Delta^{j/2} (\Delta \mathbf{1}_{j < 8} + \mathbf{1}_{j=8}) \lesssim \Delta^{3/2}.$$

324 By Lemma 3:

$$\Delta G_{k\Delta}^{(3)} = \xi^8 (X_{k\Delta}) (\mathbb{E} (L_{\Delta}^8) - \Delta I_8) \lesssim C \Delta^2.$$

325 Finally, by (5)

$$\begin{aligned} \mathbb{E} \left(\left(\Delta G_{k\Delta}^{(4)} \right)^{2p} \middle| \mathcal{F}_{k\Delta} \right) &\lesssim \sum_{j=2}^8 \mathbb{E} \left(B_{k\Delta}^{2pj} (Z_{k\Delta} + J_{k\Delta})^{16p-2pj} \middle| \mathcal{F}_{k\Delta} \right) \\ &\quad + \tilde{B}_{k\Delta}^{2p} \mathbb{E} \left((Z_{k\Delta} + J_{k\Delta})^{14p} \middle| \mathcal{F}_{k\Delta} \right) + \mathbb{E} \left(\bar{B}_{k\Delta}^{2p} (Z_{k\Delta} + J_{k\Delta})^{14p} \middle| \mathcal{F}_{k\Delta} \right) \\ &\lesssim \Delta^{2p+1}. \end{aligned}$$

326 Therefore

$$\mathbb{E} (G_{k\Delta}^2 | \mathcal{F}_{k\Delta}) \lesssim \Delta \quad \text{and} \quad \mathbb{E} (G_{k\Delta}^4 | \mathcal{F}_{k\Delta}) \lesssim \Delta.$$

327 Let us bound the centred terms, $F_{k\Delta}$ in Lemma 5 and $H_{k\Delta}$ in Lemma 10.

328 **Moments of $F_{k\Delta}$.** The term $F_{k\Delta} = \frac{1}{\Delta} ((Z_{k\Delta} + J_{k\Delta})^4 - \mathbb{E}((Z_{k\Delta} + J_{k\Delta})^4 | \mathcal{F}_{k\Delta}))$ is centred.

329 Moreover, by Equation (4), for any $p > 0$,

$$\mathbb{E} (F_{k\Delta}^{2p}) \lesssim \frac{\mathbb{E} (J_{k\Delta}^{8p}) + \mathbb{E} (Z_{k\Delta}^{8p})}{\Delta^{2p}} \lesssim \frac{1}{\Delta^{2p-1}}.$$

330 Then $\mathbb{E} (F_{k\Delta}^4) \lesssim \Delta^{-3}$. It remains to bound the quantity $\text{Var} (F_{k\Delta} | \mathcal{F}_{k\Delta}) - \frac{\xi^8 (X_{k\Delta}) I_8}{\Delta}$. By Equation
331 (4),

$$\begin{aligned} \Delta^2 \text{Var} (F_{k\Delta} | \mathcal{F}_{k\Delta}) &= \text{Var} ((Z_{k\Delta} + J_{k\Delta})^4 | \mathcal{F}_{k\Delta}) \\ &= \mathbb{E} ((Z_{k\Delta} + J_{k\Delta})^8 | \mathcal{F}_{k\Delta}) - (\mathbb{E} ((Z_{k\Delta} + J_{k\Delta})^4 | \mathcal{F}_{k\Delta}))^2 \\ &= \mathbb{E} ((Z_{k\Delta} + J_{k\Delta})^8 | \mathcal{F}_{k\Delta}) + O(\Delta^2). \end{aligned}$$

332 As

$$\mathbb{E} ((Z_{k\Delta} + J_{k\Delta})^8 | \mathcal{F}_{k\Delta}) - \Delta \xi^8 (X_{k\Delta}) I_8 = \Delta G_{k\Delta}^{(1)} + \Delta G_{k\Delta}^{(2)} + \Delta G_{k\Delta}^{(3)} \lesssim \Delta^{3/2},$$

333 we get:

$$\left| \text{Var} (F_{k\Delta} | \mathcal{F}_{k\Delta}) - \frac{\xi^8 (X_{k\Delta}) I_8}{\Delta} \right| \lesssim \Delta^{-1/2}.$$

334 **Bound of $H_{k\Delta}$.** We only need an upper bound for the moments of $H_{k\Delta}$. We have that

$$\mathbb{E} (H_{k\Delta}^2 | \mathcal{F}_{k\Delta}) \leq \frac{1}{\Delta^2} \mathbb{E} ((J_{k\Delta} + Z_{k\Delta})^{16} | \mathcal{F}_{k\Delta}) \lesssim \frac{\Delta^8 + \Delta}{\Delta^2} = \frac{1}{\Delta}$$

335 and

$$\mathbb{E} (H_{k\Delta}^4 | \mathcal{F}_{k\Delta}) \leq \frac{1}{\Delta^4} \mathbb{E} ((J_{k\Delta} + Z_{k\Delta})^{32} | \mathcal{F}_{k\Delta}) \lesssim \frac{\Delta^{16} + \Delta}{\Delta^4} = \frac{1}{\Delta^3}.$$

5.2 Proof of Proposition 6

We recall that the empirical norm is defined by $\|t\|_n^2 = \frac{1}{n} \sum_{k=1}^n t^2(X_{k\Delta})$. Then for any $t \in S_m$,

$$\begin{aligned} \gamma_n(t) &= \frac{1}{n} \sum_{k=1}^n (T_{k\Delta} - \xi^4(X_{k\Delta}) + \xi^4(X_{k\Delta}) - t(X_{k\Delta}))^2 \\ &= \frac{1}{n} \sum_{k=1}^n (T_{k\Delta} - \xi^4(X_{k\Delta}))^2 + \|t - \xi^4\|_n^2 + \frac{2}{n} \sum_{k=1}^n (T_{k\Delta} - \xi^4(X_{k\Delta}))(\xi^4(X_{k\Delta}) - t(X_{k\Delta})). \end{aligned} \quad (8)$$

As $\hat{\xi}_m^4$ minimizes $\gamma_n(t)$, $\gamma_n(\hat{\xi}_m^4) \leq \gamma_n(\Pi_m \xi^4)$ where Π_m is the orthogonal projection on S_m for the $\|\cdot\|_n$ -norm. Then by (8),

$$\begin{aligned} \left\| \xi^4 - \hat{\xi}_m^4 \right\|_n^2 + \frac{2}{n} \sum_{k=1}^n (T_{k\Delta} - \xi^4(X_{k\Delta}))(\xi^4(X_{k\Delta}) - \hat{\xi}_m^4(X_{k\Delta})) &\leq \\ \left\| \Pi_m \xi^4 - \xi^4 \right\|_n^2 + \frac{2}{n} \sum_{k=1}^n (T_{k\Delta} - \xi^4(X_{k\Delta}))(\xi^4(X_{k\Delta}) - \Pi_m \xi^4(X_{k\Delta})). \end{aligned}$$

Moreover, as $T_{k\Delta} - \xi^4(X_{k\Delta}) = E_{k\Delta} + F_{k\Delta}$, we get:

$$\left\| \hat{\xi}_m^4 - \xi^4 \right\|_n^2 \leq \left\| \Pi_m \xi^4 - \xi^4 \right\|_n^2 + \frac{2}{n} \sum_{k=1}^n (E_{k\Delta} + F_{k\Delta})(\hat{\xi}_m^4(X_{k\Delta}) - \Pi_m \xi^4(X_{k\Delta})).$$

We have

$$\left\| \hat{\xi}_m^4 - \xi^4 \right\|_n^2 = \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 + \left\| \Pi_m \xi^4 - \xi^4 \right\|_n^2 \text{ and } \left\| \Pi_m \xi^4 - \xi^4 \right\|_n^2 = \left\| \Pi_m \xi^4 - \xi_A^4 \right\|_n^2 + \left\| \xi_A^4 - \xi^4 \right\|_n^2. \quad (9)$$

By subtracting $\left\| \Pi_m \xi^4 - \xi^4 \right\|_n^2$ in both sides of the inequality, we obtain:

$$\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 \leq \frac{2}{n} \sum_{k=1}^n (E_{k\Delta} + F_{k\Delta})(\hat{\xi}_m^4(X_{k\Delta}) - \Pi_m \xi^4(X_{k\Delta})).$$

By geometric-arithmetic means inequality,

$$\frac{2}{n} \sum_{k=1}^n E_{k\Delta}(\hat{\xi}_m^4(X_{k\Delta}) - \Pi_m \xi^4(X_{k\Delta})) \leq \frac{1}{6} \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 + \frac{6}{n} \sum_{k=1}^n E_{k\Delta}^2.$$

We introduce the unit ball for the π -norm $\mathcal{B}_m = \{t \in S_m, \|t\|_\pi^2 \leq 1\}$ and the contrast function $\nu_n(t) = \frac{1}{n} \sum_{k=1}^n F_{k\Delta} t(X_{k\Delta})$. Then by geometric-arithmetic means inequality,

$$\begin{aligned} \frac{2}{n} \sum_{k=1}^n F_{k\Delta}(\hat{\xi}_m^4(X_{k\Delta}) - \Pi_m \xi^4(X_{k\Delta})) &\leq \left(\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 \sup_{t \in \mathcal{B}_m} \nu_n^2(t) \right)^{1/2} \\ &\leq \frac{1}{6} \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 + 6 \sup_{t \in \mathcal{B}_m} \nu_n^2(t). \end{aligned}$$

346 Collecting terms, we get:

$$\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 \leq 6 \sup_{t \in \mathcal{B}_m} \nu_n^2(t) + \frac{6}{n} \sum_{k=1}^n E_{k\Delta}^2 + \frac{1}{6} \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 + \frac{1}{6} \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2.$$

347 We have two different norms in the bound of the risk. For any deterministic function t , $\|t\|_\pi^2 =$
 348 $\mathbb{E} \left(\|t\|_n^2 \right)$. As $\hat{\xi}^4$ is random, we introduce the space Ω_n on which the norms $\|\cdot\|_n$ and $\|\cdot\|_\pi$ are
 349 equivalent:

$$\Omega_n = \left\{ \omega, \forall m \in \mathcal{M}_n, \forall t \in S_m, \left| \frac{\|t\|_n^2}{\|t\|_\pi^2} - 1 \right| \leq \frac{1}{2} \right\}.$$

350 [Comte et al. \(2007, Lemma 6.1\)](#) prove the following lemma for diffusion processes thanks to con-
 351 centration inequalities. However, its proof only relies on the boundedness of the stationary density
 352 π and of the mixing properties of the process $(X_t)_{t \geq 0}$.

353 **Lemma 14.** *As*

- 354 *i) $(X_t)_{t \geq 0}$ is exponentially β -mixing,*
- 355 *ii) $(X_t)_{t \geq 0}$ is stationary and its stationary density π is bounded from below and above on A ,*
- 356 *iii) the vectorial subspaces S_m satisfy Assumption A4,*

357 *then*

$$\mathbb{P}(\Omega_n^c) \leq c/n^8.$$

358 We first bound the risk on Ω_n , where the $\|\cdot\|_\pi^2$ is controlled. On Ω_n , $\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 \leq$
 359 $2 \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2$. Therefore

$$\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 \mathbf{1}_{\Omega_n} \leq \frac{12}{n} \sum_{k=1}^n E_{k\Delta}^2 + 12 \sup_{t \in \mathcal{B}_m} \nu_n^2(t). \quad (10)$$

360 The term $\sup_{t \in \mathcal{B}_m} \nu_n^2(t)$ can be bounded in two ways. First, let us consider $(\varphi_\lambda)_{\lambda \in \Lambda_m}$ an orthonor-
 361 mal basis (for the $\|\cdot\|_\pi$ -norm) of S_m . As $t \in \mathcal{B}_m$, it can be written $t = \sum_\lambda a_\lambda \varphi_\lambda$ where $\sum_\lambda a_\lambda^2 = 1$.
 362 Then, as ν is a linear function

$$\begin{aligned} \sup_{t \in \mathcal{B}_m} \nu_n^2(t) &= \sup_{\sum_\lambda a_\lambda^2 = 1} \nu_n^2 \left(\sum_\lambda a_\lambda \varphi_\lambda \right) = \sup_{\sum_\lambda a_\lambda^2 = 1} \left(\sum_\lambda a_\lambda \nu_n(\varphi_\lambda) \right)^2 \\ &\leq \sup_{\sum_\lambda a_\lambda^2 = 1} \left(\sum_\lambda a_\lambda^2 \right) \left(\sum_\lambda \nu_n^2(\varphi_\lambda) \right) \leq \sum_\lambda \nu_n^2(\varphi_\lambda). \end{aligned}$$

363 By Lemma 5, $\mathbb{E}(F_{k\Delta} | \mathcal{F}_{k\Delta}) = 0$ and therefore:

$$\begin{aligned} \mathbb{E}(\nu_n^2(\varphi_\lambda)) &= \mathbb{E}\left(\left(\frac{1}{n}\sum_{k=1}^n F_{k\Delta}\varphi_\lambda(X_{k\Delta})\right)^2\right) \\ &= \frac{1}{n^2}\sum_{k=1}^n \mathbb{E}(\varphi_\lambda^2(X_{k\Delta})\mathbb{E}(F_{k\Delta}^2 | \mathcal{F}_{k\Delta})) \\ &\quad + \frac{2}{n(n-1)}\sum_{j < k} \mathbb{E}(\varphi_\lambda(X_{k\Delta})\varphi_\lambda(X_{j\Delta})F_{j\Delta}\mathbb{E}(F_{k\Delta} | \mathcal{F}_{k\Delta})) \\ &= \frac{1}{n^2}\sum_{k=1}^n \mathbb{E}(\varphi_\lambda^2(X_{k\Delta})\mathbb{E}(F_{k\Delta}^2 | \mathcal{F}_{k\Delta})). \end{aligned}$$

364 By Lemma 5, we get that

$$\mathbb{E}(\nu_n^2(\varphi_\lambda)) \leq \frac{1}{n^2}\sum_{k=1}^n \mathbb{E}\left(\varphi_\lambda^2(X_{k\Delta})\left(\frac{\xi^8(X_{k\Delta})I_8}{\Delta} + \frac{C}{\sqrt{\Delta}}\right)\right).$$

365 Using that $\mathbb{E}(\varphi_\lambda^2(X_{k\Delta})) = \int_{\mathbb{R}} \varphi_\lambda^2(x)\pi(x)dx = 1$, and the boundness of ξ , we obtain

$$\mathbb{E}(\nu_n^2(\varphi_\lambda)) \leq \frac{\xi_1^8 I_8}{n\Delta} + \frac{C}{n\sqrt{\Delta}}.$$

366 Therefore,

$$\mathbb{E}\left(\sup_{t \in \mathcal{B}_m} \nu_n^2(t)\right) \leq \sum_{\lambda} \mathbb{E}(\nu_n^2(\varphi_\lambda)) \leq \xi_1^8 I_8 \frac{D_m}{n\Delta} + \frac{CD_m}{n\sqrt{\Delta}}.$$

367 The second way of bounding $\sup_{t \in \mathcal{B}_m} \nu_n^2(t)$ nearly use the same arguments. Let us remark that if
368 $t \in \mathcal{B}_m$, then $\|t\|_{L^2}^2 \leq \frac{1}{\pi_0} \|t\|_{\pi}^2 = \frac{1}{\pi_0}$. Then, if we consider $(\psi_\lambda)_{\lambda \in \Lambda_m}$ an orthonormal basis of S_m
369 for the L^2 -norm.

$$\sup_{t \in \mathcal{B}_m} \nu_n^2(t) \leq \sup_{t \in S_m: \|t\|_{L^2}^2 \leq \frac{1}{\pi_0}} \nu_n^2(t) \leq \frac{1}{\pi_0} \sum_{\lambda} \nu_n^2(\psi_\lambda).$$

370 By Assumption 4 b, $\sum_{\lambda \in \Lambda_m} \psi_\lambda^2(X_{k\Delta}) \leq \phi_1 D_m$, and as the process is stationary:

$$\mathbb{E}\left(\sup_{t \in \mathcal{B}_m} \nu_n^2(t)\right) \leq \frac{1}{\pi_0} \frac{1}{n^2} \sum_{k=1}^n \mathbb{E}\left(F_{k\Delta}^2 \sum_{\lambda \in \Lambda_m} \psi_\lambda^2(X_{k\Delta})\right) \leq \frac{\phi_1 D_m \mathbb{E}(F_{\Delta}^2)}{\pi_0 n}.$$

371 Therefore, as $\mathbb{E}(F_{\Delta}^2) = \mathbb{E}(\xi^8(X_{\Delta})) I_8 / \Delta + O\left(\frac{1}{\sqrt{\Delta}}\right)$:

$$\mathbb{E}\left(\sup_{t \in \mathcal{B}_m} \nu_n^2(t)\right) \leq \Xi \frac{D_m}{n\Delta} \tag{11}$$

372 where $\Xi = \min\left(\xi_1^8 I_8, \frac{\phi_1}{\pi_0} \mathbb{E}_{\pi}(\xi^8(X_{\Delta})) I_8\right)$. By Lemma 5, $\mathbb{E}(E_{k\Delta}^2) \lesssim \Delta$. By (10) and (11), we
373 obtain:

$$\mathbb{E}\left(\left\|\hat{\xi}_m^4 - \Pi_m \xi^4\right\|_n^2 \mathbf{1}_{\Omega_n}\right) \leq 12\Xi \frac{D_m}{n\Delta} + c\Delta. \tag{12}$$

374 As $\Pi_m \xi^4$ is the orthogonal projection for the $\|\cdot\|_n$ -norm, $\|\Pi_m \xi^4 - \xi_A^4\|_n^4 \leq \|\xi_{m,\pi}^4 - \xi_A^4\|_n^2$. And as
 375 $\xi_{m,\pi}$ and ξ are deterministic, $\mathbb{E} \left(\|\xi_{m,\pi}^4 - \xi_A^4\|_n^2 \right) = \|\xi_{m,\pi}^4 - \xi_A^4\|_\pi^2$. Then by (9)

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 \mathbf{1}_{\Omega_n} \right) \leq \|\xi_{m,\pi}^4 - \xi^4\|_\pi^2 + \Xi \frac{D_m}{n\Delta} + c\Delta.$$

376 It remains to bound the risk on Ω_n^c . The function $\hat{\xi}_m^4$ is the orthogonal projection (for the $\|\cdot\|_n$
 377 norm) of $(T_\Delta, \dots, T_{n\Delta})$ on the vectorial subspace $\{(t(X_\Delta), \dots, t(X_{n\Delta}))\}$, $t \in S_m$.

378 We have:

$$\begin{aligned} \left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 &= \|\Pi_m T - \xi_A^4\|_n^2 = \|\Pi_m \xi^4 - \xi_A^4\|_n^2 + \|\Pi_m E + \Pi_m F\|_n^2 \\ &\leq \|\xi_A^4\|_n^2 + \|E + F\|_n^2. \end{aligned}$$

379 By stationarity and Cauchy-Schwarz:

$$\begin{aligned} \mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 \mathbf{1}_{\Omega_n^c} \right) &\lesssim \mathbb{E} \left(\|\xi_A^4\|_n^2 \mathbf{1}_{\Omega_n^c} \right) + \mathbb{E} \left[\left(\frac{1}{n} \sum_{k=1}^n E_{k\Delta}^2 + F_{k\Delta}^2 \right) \mathbf{1}_{\Omega_n^c} \right] \\ &\lesssim \left[\mathbb{E} \left[\|\xi_A^4\|_n^4 \right] + \mathbb{E} \left[E_{k\Delta}^4 + F_{k\Delta}^4 \right] \right] \mathbb{P}(\Omega_n^c)^{1/2}. \end{aligned}$$

380 We obtain by Lemma 5:

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 \mathbf{1}_{\Omega_n^c} \right) \lesssim \frac{1}{\Delta^{3/2} n^4} \leq \frac{1}{n}.$$

381 5.3 Proof of Corollary 7

382 We again separate the risk on Ω_n (where the empirical norm $\|\cdot\|_n$ is equivalent to the $\|\cdot\|_\pi$ -norm)
 383 and Ω_n^c . As $\hat{\xi}_{m,\pi}^4$ is the orthogonal projection of ξ^4 on S_m for the $\|\cdot\|_\pi$ - norm:

$$\left\| \hat{\xi}_m^4 - \xi^4 \right\|_\pi^2 = \left\| \hat{\xi}_m^4 - \xi_{m,\pi}^4 \right\|_\pi^2 + \left\| \xi_{m,\pi}^4 - \xi^4 \right\|_\pi^2.$$

384 As $\hat{\xi}_m^4$ and $\xi_{m,\pi}^4$ both belong to the subspace S_m , on Ω_n :

$$\left\| \hat{\xi}_m^4 - \xi_{m,\pi}^4 \right\|_\pi^2 \mathbf{1}_{\Omega_n} \leq 2 \left\| \hat{\xi}_m^4 - \xi_{m,\pi}^4 \right\|_n^2 \leq 4 \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 + 4 \left\| \Pi_m \xi^4 - \xi_{m,\pi}^4 \right\|_n^2$$

385 and as $\Pi_m \xi^4$ is the orthogonal projection of ξ^4 on S_m for the empirical risk, $\left\| \Pi_m \xi^4 - \xi_{m,\pi}^4 \right\|_n^2 \leq$
 386 $\left\| \xi_A^4 - \xi_{m,\pi}^4 \right\|_n^2$. By (12), we obtain that

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi^4 \right\|_\pi^2 \mathbf{1}_{\Omega_n} \right) \leq 5 \|\xi^4 - \xi_{m,\pi}^4\|_\pi^2 + 48\Xi \frac{D_m}{n\Delta} + C\Delta.$$

387 In addition, by Lemma 5 and Lemma 14:

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi_{m,\pi}^4 \right\|_\pi^2 \mathbf{1}_{\Omega_n^c} \right) \leq \left(\mathbb{E} \left(\|\Pi_m T\|_\pi^4 + \|\xi^4\|_\pi^4 \right) \right)^{1/2} (\mathbb{P}(\Omega_n^c))^{1/2} \lesssim \frac{1}{n}$$

388 which give the bound for the $\|\cdot\|_\pi$ -norm.

389 To bound the L^2 -risk, we simply remark that $\|t\|_{L^2(A)}^2 \leq \frac{1}{\pi_0} \|t\|_\pi^2$ and that $\|\xi^4 - \xi_{\pi,m}^4\|_\pi^2 \leq$
 390 $\|\xi^4 - \xi_m^4\|_\pi^2 \leq \pi_1 \|\xi^4 - \xi_m^4\|_{L^2(A)}^2$. Then

$$\mathbb{E} \left(\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_{L^2(A)}^2 \right) \leq \frac{5\pi_1}{\pi_0} \|\xi^4 - \xi_m^4\|_{L^2(A)}^2 + \frac{48\Xi}{\pi_0} \frac{D_m}{n\Delta} + C\Delta.$$

391 5.4 Proof of Theorem 8

392 For any $m \in \mathcal{M}_n$,

$$\gamma_n(\hat{\xi}_{\hat{m}}^4) + \text{pen}(\hat{m}) \leq \gamma_n(\Pi_m \xi^4) + \text{pen}(m).$$

393 As previously, we decompose the risk on Ω_n and Ω_n^c . On Ω_n^c , the bound is the same as in the
 394 previous proof. On Ω_n , as in the proof of Proposition 6, we have:

$$\begin{aligned} \left\| \hat{\xi}_{\hat{m}}^4 - \xi_A^4 \right\|_n^2 &\leq \left\| \Pi_m \xi^4 - \xi_A^4 \right\|_n^2 + \frac{1}{12} \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^2 + \frac{1}{12} \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 + 12 \sum_{k=1}^n E_{k\Delta}^2 + 12 \sup_{t \in \mathcal{B}_{m, \hat{m}}} \nu_n^2(t) \\ &\quad + \text{pen}(m) - \text{pen}(\hat{m}) \end{aligned}$$

395 where $\mathcal{B}_{m, m'}$ is the unit ball for the $\|\cdot\|_\pi$ -norm of the vectorial subspace $S_m + S_{m'}$. On Ω_n ,
 396 $\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 \leq 2 \left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_n^4$. By the triangular inequality, $\left\| \hat{\xi}_m^4 - \Pi_m \xi^4 \right\|_\pi^2 \leq 2 \left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 +$
 397 $2 \left\| \Pi_m \xi^4 - \xi_A^4 \right\|_n^2$. Then

$$\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 \leq 3 \left\| \Pi_m \xi^4 - \xi_A^4 \right\|_n^2 + 2\text{pen}(m) - 2\text{pen}(\hat{m}) + 24 \sum_{k=1}^n E_{k\Delta}^2 + 24 \sup_{t \in \mathcal{B}_{m, \hat{m}}} \nu_n^2(t).$$

398 Let us set $p(m, m') = (\text{pen}(m) + \text{pen}(m'))/12$. Then

$$\left\| \hat{\xi}_m^4 - \xi_A^4 \right\|_n^2 \mathbf{1}_{\Omega_n} \leq 3 \left\| \Pi_m \xi^4 - \xi_A^4 \right\|_n^2 + 2\text{pen}(m) + 24 \sum_{k=1}^n E_{k\Delta}^2 + 24 \left(\sup_{t \in \mathcal{B}_{m, \hat{m}}} \nu_n^2(t) - p(m, \hat{m}) \right). \quad (13)$$

399 It remains to bound

$$\mathbb{E} \left(\sup_{t \in \mathcal{B}_{m, \hat{m}}} \nu_n^2(t) - p(m, \hat{m}) \right) \leq \sum_{m \in \mathcal{M}_n} \mathbb{E} \left(\sup_{t \in \mathcal{B}_{m, m'}} \nu_n^2(t) - p(m, m') \right).$$

400 We can do so using concentration inequalities. The following lemma is deduced from the Berbee's
 401 coupling lemma and a Talagrand inequality. It is proved for instance in [Schmisser \(2019, Appendix](#)
 402 [A\)](#).

403 **Lemma 15** (Talagrand inequality for β -mixing variables). *Let us consider Z_1, \dots, Z_n some random*
 404 *variables exponentially β -mixing. Let us set $b_0 \geq 1/\beta$ where β is the β -mixing coefficient). We define*
 405 *$q_n := 2b_0 \ln(n)$, $p_n = n/(2q_n)$. We have that $\beta(q_n) \leq ce^{-2\beta b_0 \ln(n)} \lesssim n^{-2}$. Let us consider*

$$I_n(t) = \frac{1}{n} \sum_{k=1}^n F_t(Z_k) - \mathbb{E} [F_t(Z_k)].$$

406 If we can find a triplet $(M_2, V$ and $H)$ such that:

$$\forall i, \sup_{t \in \mathcal{F}} \text{Var} \left(\frac{1}{q_n} \sum_{k=iq_n}^{(i+1)q_n} F_t(Z_k) \right) \leq \frac{V}{q_n},$$

$$\sup_{t \in \mathcal{F}} \left\| \frac{1}{q_n} \sum_{k=cq_n}^{(c+1)q_n} F_t(Z_k) \right\|_{\infty} \leq M_2 \quad \text{and} \quad \mathbb{E} \left[\sup_{t \in \mathcal{F}} |I_n(t)| \right] \leq \frac{H}{\sqrt{n}},$$

407 then we have:

$$\mathbb{E} \left[\sup_{t \in \mathcal{F}} \left| I_n^2(t) - \frac{6H^2}{n} \right| \right]_+ \leq C \frac{V}{n} \exp \left(-k_1 \frac{H^2}{12V} \right) + C' \frac{M_2^2}{p_n^2} \exp \left(-k_2 \frac{\sqrt{p_n} H}{\sqrt{q_n} M_2} \right) + 2 \frac{M_2^2}{n^2}$$

408 where C, C', k_1 and k_2 are universal constants.

409 The random variables $(F_{k\Delta}, X_{k\Delta})$ are exponentially β -mixing, with β -mixing coefficient pro-
410 portional to Δ . Let us set $F_t(F_{k\Delta}, X_{k\Delta}) := F_{k\Delta} t(X_{k\Delta})$. We want to apply Lemma 15 to
411 $\nu_n(t) = \frac{1}{n} \sum_{k=1}^n F_{k\Delta} t(X_{k\Delta})$. By Equation (11),

$$\mathbb{E} \left(\sup_{t \in \mathcal{B}_{m,m'}} \nu_n^2(t) \right) \leq \frac{\Xi(D_m + D_{m'})}{n\Delta},$$

412 so we can take $H^2 = \frac{\Xi(D_m + D_{m'})}{\Delta}$. Let us set $q_n = c \ln(n)\Delta$ with c such that $\beta(q_n\Delta) \lesssim n^{-2}$. By
413 Lemma 5, as $\mathbb{E}(F_{k\Delta} | \mathcal{F}_{k\Delta}) = 0$, we have that, for any $t \in \mathcal{B}_{m,m'}$,

$$\text{Var} \left(\frac{1}{q_n} \sum_{k=iq_n}^{(i+1)q_n} F_{k\Delta} t(X_{k\Delta}) \right) = \frac{1}{q_n^2} \sum_{k=iq_n}^{(i+1)q_n} \mathbb{E} \left(\mathbb{E}(F_{k\Delta}^2 | \mathcal{F}_{k\Delta}) t^2(X_{k\Delta}) \right) \leq \frac{\xi_1^8 I_8}{q_n \Delta}.$$

414 Then we can choose $V = \xi_1 I_8 / \Delta$. It remains to bound the variables $F_{k\Delta}$ with high probability. We
415 have the following decomposition:

$$F_{k\Delta} \lesssim \frac{J_{k\Delta}^4 + Z_{k\Delta}^4}{\Delta}.$$

416 Let us introduce a set where this quantity is bounded: for any $0 < \alpha < 1$,

$$\Omega_{B,\alpha} = \left\{ \omega, \forall k, \forall p \in \mathbb{N}, p \leq 8, |B_{k\Delta}| \leq \Delta^{1-\alpha}; \quad |Z_{k\Delta}| \leq 6\sigma_1 \Delta^{1/2} \ln(n); \quad |J_{k\Delta}| \leq 36\Delta^{-\alpha} \ln(n); \right.$$

$$\left. \frac{1}{q_n} \sum_{k=cq_n}^{(c+1)q_n} J_{k\Delta}^p \leq 49\Delta^{1-\alpha} \ln^p(n) \right\}.$$

417 By Lemma 4:

$$\mathbb{P}(\Omega_{B,\alpha}) \lesssim \frac{1}{n^5}. \tag{14}$$

418 On $\Omega_n \cap \Omega_{B,\alpha}$, if $t \in S_m + S_{m'}$, $t(X_{k\Delta}) \leq (D_m + D_{m'})^{1/2} \sqrt{\phi_1/\pi_0}$, and $F_{k\Delta} \lesssim (J_{k\Delta}^4 + Z_{k\Delta}^4)/\Delta \lesssim$
419 $C \ln^4(n) \Delta^{-\alpha}$. Then we apply Talagrand's inequality (lemma 15) to $\nu_n(t) = \frac{1}{n} \sum_{k=1}^n t(X_{k\Delta}) F_{k\Delta}$
420 with

$$V = \frac{\xi_1^8 I_8}{\Delta}, \quad H^2 = \frac{\Xi D}{\Delta}, \quad M_2 = C \frac{D^{1/2} \ln^4(n)}{\Delta^\alpha}$$

421 with $D = D_m + D_{m'}$. By Lemma 15, we obtain:

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in \mathcal{B}_{m,m'}} \nu_n^2(t) - 12H^2 \right)_+ &\lesssim \frac{1}{n\Delta} \exp \left(-c \frac{D/\Delta}{1/\Delta} \right) \\ &+ \frac{D \ln^8(n)}{p_n^2 \Delta^{2\alpha}} \exp \left(-c' \frac{\sqrt{p_n} D^{1/2} / \sqrt{\Delta}}{\sqrt{q_n} D^{1/2} \ln^4(n) / \Delta^\alpha} \right) + \frac{D \ln^8(n)}{n^2 \Delta^{2\alpha}} \\ &\lesssim \frac{1}{n\Delta} \exp(-c_1 D) + \frac{D \ln^{10}(n)}{n^2 \Delta^{2+2\alpha}} \exp \left(-c_2 \frac{\sqrt{n\Delta^{1+2\alpha}}}{\ln^5(n)} \right) + \frac{D \ln^8(n)}{n^2 \Delta^{2\alpha}}. \end{aligned}$$

422 We obtain that

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in \mathcal{B}_{m,\hat{m}}} \nu_n^2(t) - p(m, \hat{m}) \right) &\leq \sum_{D_m \in \mathcal{M}_n} \mathbb{E} \left(\sup_{t \in \mathcal{B}_{m,m'}} \nu_n^2(t) - H^2 \right) \\ &\leq \sum_{D=1}^{\sqrt{n\Delta}} \mathbb{E} \left(\sup_{t \in \mathcal{B}_{m,m'}} \nu_n^2(t) - H^2 \right) \\ &\lesssim \frac{1}{n\Delta} + \frac{1}{n\Delta^{1+2\alpha}} e^{-\sqrt{\frac{n\Delta^{1+2\alpha}}{\ln^5(n)}}} + \frac{\Delta^{1-2\alpha}}{n}. \end{aligned}$$

423 As we can take α as small as we want, we get that

$$\mathbb{E} \left(\sup_{t \in \mathcal{B}_{m,\hat{m}}} \nu_n^2(t) - p(m, \hat{m}) \mathbf{1}_{\Omega_n \cap \Omega_{B,\alpha}} \right) \leq \frac{C'}{n\Delta}. \quad (15)$$

424 The bound of the risk on $(\Omega_n \cap \Omega_{B,\alpha})^c$ is done as in the proof of Proposition 6.

425 5.5 Proof of Corollary 9

426 The proof is nearly the same as for the risk of the estimator with m fixed. We have that, for any
427 m :

$$\left\| \hat{\xi}_{\hat{m}}^4 - \xi_A^4 \right\|_\pi^2 \leq 2 \left\| \hat{\xi}_{\hat{m}}^4 - \xi_{m,\pi}^4 \right\|_\pi^2 + 2 \left\| \xi_{m,\pi}^4 - \xi_A^4 \right\|_\pi^2. \quad (16)$$

428 And on $\Omega_n \cap \Omega_{B,\alpha}$:

$$\begin{aligned} \left\| \hat{\xi}_{\hat{m}}^4 - \xi_{m,\pi}^4 \right\|_\pi^2 \mathbf{1}_{\Omega_n} &\leq 2 \left\| \hat{\xi}_{\hat{m}}^4 - \xi_{m,\pi}^4 \right\|_n^2 \leq 4 \left\| \hat{\xi}_{\hat{m}}^4 - \Pi_m \xi^4 \right\|_n^2 + 4 \left\| \Pi_m \xi^4 - \xi_{m,\pi}^4 \right\|_n^2 \\ &\leq 4 \left\| \hat{\xi}_{\hat{m}}^4 - \xi_A^4 \right\|_n^2 + 4 \left\| \xi_A^4 - \xi_{m,\pi}^4 \right\|_n^2. \end{aligned}$$

429 As $\left\| \Pi_m \xi^4 - \xi_A^4 \right\|_n^2 \leq \left\| \xi_{m,\pi}^4 - \xi_A^4 \right\|_n^2$, by Equations (13) and (16), we obtain:

$$\begin{aligned} \left\| \hat{\xi}_{\hat{m}}^4 - \xi_{m,\pi}^4 \right\|_\pi^2 \mathbf{1}_{\Omega_n \cap \Omega_{B,\alpha}} &\leq 8 \left\| \hat{\xi}_{\hat{m}}^4 - \xi_A^4 \right\|_n^2 + 8 \left\| \xi_A^4 - \xi_{m,\pi}^4 \right\|_n^2 + 2 \left\| \xi_{m,\pi}^4 - \xi_A^4 \right\|_\pi^2 \\ &\leq 32 \left\| \xi_{m,\pi}^4 - \xi_A^4 \right\|_n^2 + 2 \left\| \xi_{m,\pi}^4 - \xi_A^4 \right\|_\pi^2 + 16pen(m) \\ &+ 192 \sum_{k=1}^n E_{k\Delta}^2 + 192 \sup_{t \in \mathcal{B}_{m,\hat{m}}} (\nu_n^2(t) - p(m, \hat{m})). \end{aligned}$$

430 The risk on $(\Omega_n \cap \Omega_{B,\alpha})^c$ is the same as for m fixed, so we get by (15):

$$\mathbb{E} \left(\left\| \hat{\xi}_{\hat{m}}^4 - \xi^4 \right\|_{\pi}^2 \right) \lesssim \inf_{m \in \mathcal{M}_n} \left(\left\| \xi_{m,\pi}^4 - \xi^4 \right\|_{\pi}^2 + \text{pen}(m) \right) + \frac{1}{n\Delta} + \Delta$$

431 and

$$\mathbb{E} \left(\left\| \hat{\xi}_{\hat{m}}^4 - \xi^4 \right\|_{L^2(A)}^2 \right) \lesssim \inf_{m \in \mathcal{M}_n} \left(\left\| \xi_m^4 - \xi^4 \right\|_{L^2(A)}^2 + \text{pen}(m) \right) + \frac{1}{n\Delta} + \Delta.$$

432 5.6 Proof of Lemma 11

433 As in the proof of Proposition 6, we decompose the risk on Ω_n and Ω_n^c and get that, on Ω_n ,

$$\left\| \hat{h}_m - \Pi_m h \right\|_n^2 \mathbf{1}_{\Omega_n} \leq 12 \sup_{t \in \mathcal{B}_m} \nu_{n,2}^2(t) + \frac{12}{n} \sum_{k=1}^n G_{k\Delta}^2$$

434 where $\nu_{n,2}(t) = \frac{1}{n} \sum_{k=1}^n t(X_{k\Delta}) H_{k\Delta}$. By Lemma 10, we have that $\mathbb{E}(G_{k\Delta}^2) \lesssim \Delta$. Moreover, for
435 $(\varphi_\lambda)_{1 \leq \lambda \leq D_m}$ an orthonormal basis for the L_{π}^2 - norm of S_m , we have

$$\mathbb{E} \left(\sup_{t \in S_m} \nu_{n,2}^2(t) \right) \leq \sum_{\lambda} \mathbb{E}(\nu_{n,2}^2(\varphi_\lambda)) = \sum_{\lambda} \frac{1}{n^2} \sum_{k=1}^n \mathbb{E}(H_{k\Delta}^2 \varphi_\lambda^2(X_{k\Delta})) \lesssim \frac{D_m}{n\Delta}.$$

436 As $\left\| \hat{h}_m - h_A \right\|_n^2 = \left\| \hat{h}_m - \Pi_m h \right\|_n^2 + \left\| \Pi_m h - h_A \right\|_n^2$, and $\mathbb{E} \left(\left\| \Pi_m h - h_A \right\|_n^2 \right) \leq \mathbb{E} \left(\left\| h_{m,\pi} - h_A \right\|_n^2 \right) =$
437 $\left\| h_{m,\pi} - h_A \right\|_{\pi}^2$, we get that

$$\mathbb{E} \left(\left\| \hat{h}_m - h_A \right\|_n^2 \mathbf{1}_{\Omega_n} \right) \leq \left\| h_{m,\pi} - h \right\|_{\pi}^2 + \frac{CD_m}{n\Delta} + c\Delta.$$

438 And on Ω_n^c , we get that

$$\mathbb{E} \left(\left\| \hat{h}_m - h_A \right\|_n^2 \right) \lesssim \left[\mathbb{E} \left(\left\| h_A \right\|_n^4 \right) + \mathbb{E} \left(G_{k\Delta}^4 + H_{k\Delta}^4 \right) \right] \mathbb{P}(\Omega_n^c)^{1/2} \lesssim \frac{1}{\Delta^{3/2} n^4} \leq \frac{1}{n}$$

439 which ends the proof.

440 5.7 Proof of lemma 12

441 Let us set $h_1 = \sup_{x \in A} h(x)$ and $\hat{h}_1 = \sup_A \hat{h}_{\ln(n)}(x)$.

442 **Bound of $h_1 - \hat{h}_1$.** We have that

$$|\hat{h}_1 - h_1| \leq \left\| \hat{h}_{\ln(n)} - h \right\|_{L^\infty(A)} \leq \left\| \hat{h}_{\ln(n)} - h_{\ln(n)} \right\|_{L^\infty(A)} + \left\| h_{\ln(n)} - h \right\|_{L^\infty(A)}.$$

443 By DeVore and Lorentz (1993, 13, p182) and Barron et al. (1999, 4, Lemma 12), if h belongs to the
444 Besov space $\mathcal{B}_{2,\infty}^\alpha(A)$, then

$$\left\| h_m - h \right\|_{L^\infty(A)}^2 \leq CD_m^{1-2\alpha}. \quad (17)$$

445 As ξ is Lipschitz, $h = \xi^8 I_8$ belongs to $\mathcal{B}_{2,\infty}^1(A)$ and $\|h_m - h_A\|_{L^\infty(A)}^2 \lesssim D_m^{-1}$. It remains to bound
 446 $\|\hat{h}_m - h_m\|_{L^\infty(A)}$. As \hat{h}_m and h_m belongs to S_m ,

$$\|\hat{h}_m - h_m\|_{L^\infty(A)}^2 \leq \phi_1 D_m \|\hat{h}_m - h_m\|_{L^2}^2 \leq \frac{\phi_1 D_m}{\pi_0} \|\hat{h}_m - h_m\|_\pi^2.$$

447 On Ω_n ,

$$\|\hat{h}_m - h_m\|_\pi^2 \leq 2 \|\hat{h}_m - h_m\|_n^2 \leq 2 \|\hat{h}_m - \Pi_m h\|_n^2 + 2 \|\Pi_m h - h_m\|_n^2.$$

448 Those two terms are bounded thanks to a concentration inequality.

449 **Bound of $\|\Pi_m h - h_m\|_n^2$.** As $\Pi_m h$ is the orthogonal projection of h on S_m , $\|\Pi_m h - h_m\|_n^2 \leq$
 450 $\|h_m - h_A\|_n^2$. Its expectation is $\|h_m - h_A\|_\pi^2 \lesssim D_m^{-2}$. By (17), $\|h_m - h\|_{L^\infty(A)}^2 \lesssim D_m^{-1}$. Moreover,

$$\begin{aligned} \text{Var} \left(\sum_{j q_n + 1}^{(j+1)q_n} (h_m - h_A)^2(X_{k\Delta}) \right) &\leq q_n^2 \mathbb{E}((h_m - h_A)^4(X_\Delta)) \leq q_n^2 \|h_m - h_A\|_{L^\infty}^2 \|h_m - h_A\|_\pi^2 \\ &\lesssim q_n^2 D_m^{-3}. \end{aligned}$$

451 The following lemma is very useful to control the difference between a mean and its expectation.
 452 It is proved for instance in Schmisser (2019, Result 24).

453 **Lemma 16** (Bennet inequality for β -mixing variables). *Let us consider Y_1, \dots, Y_n some exponen-*
 454 *tially β -mixing random variables. Let us set $b_0 \geq 1/\beta$ where β is the β -mixing coefficient. We*
 455 *define $q_n := 2b_0 \ln(n)$, $p_n = n/(2q_n)$. We have that $\beta(q_n) \leq ce^{-2\beta b_0 \ln(n)} \lesssim n^{-2}$. Let us set*
 456 *$\bar{Y}_n = \frac{1}{n} \sum_{k=1}^n Y_k$. If there exists two constants V' and M'_2 such that, for any j ,*

$$\text{Var} \left(\sum_{k=j q_n + 1}^{(j+1)q_n} Y_k \right) \leq q_n V' \quad \text{and} \quad \left| \sum_{k=j q_n + 1}^{(j+1)q_n} Y_k - \mathbb{E}(Y_k) \right| \leq q_n M'_2$$

457 then, for any $x > 0$,

$$\mathbb{P}(|\bar{Y}_n - \mathbb{E}(\bar{Y}_n)| \geq x) \leq 2 \exp\left(-\frac{nx^2}{2(V' + xM'_2 q_n)}\right).$$

458 By Bennett inequality with $V' = q_n D_m^{-3}$ and $M'_2 = C D_m^{-1}$, as $D_m = \ln(n)$,

$$\begin{aligned} \mathbb{P}\left(\left|\|h_m - h_A\|_n^2 - \|h_m - h_A\|_\pi^2\right| \geq D_m^{-2}\right) &\lesssim \exp\left(-\frac{cn D_m^{-4}}{q_n D_m^{-3} + q_n D_m^{-3}}\right) = \exp(-cp_n D_m^{-1}) \\ &\lesssim \exp\left(-c \frac{n\Delta}{\ln^2(n)}\right) \lesssim n^{-5}. \end{aligned}$$

459 Then

$$\mathbb{P}\left(\|h_m - \Pi_m h\|_n^2 \geq D_m^{-2}\right) \lesssim n^{-5}. \quad (18)$$

460 **Bound of $\left\|\hat{h}_m - \Pi_m h\right\|_n$.** We know that, on Ω_n ,

$$\left\|\hat{h}_m - \Pi_m h\right\|_n^2 \leq 12 \sup_{t \in \mathcal{B}_m} \nu_{n,2}^2(t) + \frac{12}{n} \sum_{k=1}^n G_{k\Delta}^2. \quad (19)$$

461 Moreover, for $(\varphi_\lambda)_{\lambda \in \Lambda_m}$ an orthonormal basis for the L_π^2 -norm of S_m ,

$$\sup_{t \in \mathcal{B}_m} \nu_{n,2}^2(t) \leq \sum_\lambda \nu_{n,2}^2(\varphi_\lambda) = \sum_\lambda \left(\frac{1}{n} \sum_{k=1}^n H_{k\Delta} \varphi_\lambda(X_{k\Delta}) \right)^2$$

462 and we use again Bennett's inequality. We recall that $|H_{k\Delta}| \lesssim \Delta^{-1}(Z_{k\Delta}^8 + J_{k\Delta}^8)$. For any $t \in \mathcal{B}_m$,
463 this term is centred: $\mathbb{E}(\nu_{n,2}(t)) = 0$. In addition, as $\mathbb{E}(H_{k\Delta} | \mathcal{F}_{k\Delta}) = 0$, its variance is bounded by:

$$\text{Var} \left(\sum_{k=jq_n+1}^{(j+1)q_n} H_{k\Delta} \varphi_\lambda(X_{k\Delta}) \right) = q_n \text{Var}(H_\Delta \varphi_\lambda(X_\Delta)) \lesssim \frac{q_n}{\Delta}.$$

464 On $\Omega_{B,\alpha}$, as $\|\varphi_\lambda\|_{L^\infty}^2 \leq \frac{\phi_1}{\pi_0} D_m \|\varphi\|_{L^2}^2 = \frac{\phi_1}{\pi_0} D_m$, we have:

$$\left| \sum_{k=jq_n+1}^{(j+1)q_n} H_{k\Delta} \varphi_\lambda(X_{k\Delta}) \right| \leq D_m^{1/2} \sqrt{\frac{\phi_1}{\pi_0}} \sum_{k=jq_n+1}^{(j+1)q_n} |H_{k\Delta}| \lesssim q_n \sqrt{D_m} \ln^8(n) \Delta^{-\alpha}.$$

465 And by Bennett's inequality (Lemma 16) with $V' = C/\Delta$ and $M'_2 = C' \sqrt{D_m} \ln^8(n) \Delta^{-\alpha}$, we obtain:

$$\mathbb{P} \left(|\nu_{n,2}(\varphi_\lambda)| \geq \frac{1}{\sqrt{D_m} \ln^9(n)} \right) \lesssim \mathbb{P}(\Omega_{B,\alpha}^c) + \exp \left(-c \frac{n \Delta^{1+\alpha}}{D_m \ln^{18}(n)} \right) \lesssim n^{-5} + \exp \left(-\frac{n \Delta^{1+\alpha}}{\ln^{19}(n)} \right) \lesssim n^{-5}.$$

466 Then

$$\begin{aligned} \mathbb{P} \left(\sup_{t \in \mathcal{B}_m} \nu_{n,2}^2(t) \geq \frac{1}{\ln^{18}(n)} \right) &\leq \mathbb{P} \left(\sum_\lambda \nu_{n,2}^2(\varphi_\lambda) \geq \frac{1}{\ln^{18}(n)} \right) \\ &\leq \sum_\lambda \mathbb{P} \left(|\nu_{n,2}(\varphi_\lambda)| \geq \frac{1}{D_m \ln^{18}(n)} \right) \lesssim D_m n^{-5}. \end{aligned} \quad (20)$$

467 Let us now bound $G_{k\Delta}$. We have that $G_{k\Delta} = G_{k\Delta}^{(1)} + G_{k\Delta}^{(2)} + G_{k\Delta}^{(3)} + G_{k\Delta}^{(4)}$, with $|G_{k\Delta}^{(1)} + G_{k\Delta}^{(2)} + G_{k\Delta}^{(3)}| \lesssim$
468 $\Delta^{1/2}$. Moreover, on $\Omega_{B,\alpha}$, by Lemma 4,

$$\begin{aligned} \left| \frac{1}{n} \sum_{k=1}^n \left(G_{k\Delta}^{(4)} \right)^2 \right| &\lesssim \frac{\Delta^{-2}}{n} \sum_{k=1}^n \sum_{j=1}^8 B_{k\Delta}^{2j} (Z_{k\Delta}^{16-2j} + J_{k\Delta}^{16-2j}) \\ &\lesssim \Delta^{-2} \sum_{j=1}^8 \Delta^{2j-2j\alpha} \left(\Delta^{8-j} \ln^{16-2j}(n) + \frac{1}{n} \sum_{k=1}^n J_{k\Delta}^{16-2j} \right) \\ &\lesssim \sum_{j=1}^8 \Delta^{2j-2j\alpha-2} \left(\Delta^{8-j} \ln^{16-2j}(n) + \Delta^{1-\alpha} \mathbf{1}_{j < 8} \ln^{16-2j}(n) + \mathbf{1}_{j=8} \right) \\ &\lesssim \Delta^{1-3\alpha} \ln^{16}(n). \end{aligned}$$

469 Then, for n large enough:

$$\mathbb{P} \left(\frac{1}{n} \sum_{k=1}^n G_{k\Delta}^2 \geq \Delta \ln(n) + \Delta^{1-3\alpha} \ln^{17}(n) \right) \leq \mathbb{P}(\Omega_{B,\alpha}^c) \lesssim n^{-5}. \quad (21)$$

470 Therefore, by Equations (19), (21) and (20), we obtain:

$$\mathbb{P} \left(\left\| \hat{h}_m - \Pi_m h \right\|_n^2 \geq \Delta^{1-3\alpha} \ln^{17}(n) + \frac{1}{\ln^9(n)} \right) \leq \mathbb{P}(\Omega_n^c) + D_m n^{-5} + n^{-5} \lesssim D_m n^{-5}. \quad (22)$$

471 We recall that, on Ω_n :

$$|\hat{h}_1 - h_1| \lesssim \|h_{\ln(n)} + h\|_{L^\infty(A)} + \ln(n) \|h_{\ln(n)} - \Pi_{\ln(n)} h\|_n^2 + \ln(n) \left\| \hat{h}_{\ln(n)} - \Pi_{\ln(n)} h \right\|_n^2.$$

472 By Equations (17), (18) and (22), we obtain:

$$\mathbb{P} \left(|\hat{h}_1 - h_1| \gtrsim \frac{1}{\ln(n)} + \frac{\ln(n)}{\ln^9(n)} + \ln^{18}(n) \Delta^{1-3\alpha} \right) = \mathbb{P} \left(|\hat{h}_1 - h_1| \gtrsim \frac{1}{\ln(n)} \right) \lesssim \mathbb{P}(\Omega_n^c) + n^{-4} \lesssim n^{-4}.$$

473 **Bound of $\bar{V}_n - \mathbb{E}(\xi^8(X_\Delta)) I_8$.** To bound this quantity, we use Bennett's inequality.

474 By Lemma 10,

$$\mathbb{E}(\bar{V}_n) = \mathbb{E}(\xi^8(X_{k\Delta}) I_8 + G_{k\Delta} + H_{k\Delta}) = \mathbb{E}(\xi^8(X_{k\Delta})) I_8 + C' \Delta^{1/2}.$$

475 Moreover,

$$\begin{aligned} \text{Var} \left(\sum_{j=1}^{q_n} V_{j\Delta} \right) &\lesssim \text{Var} \left(\sum_{j=1}^{q_n} \xi^8(X_{j\Delta}) \right) + \text{Var} \left(\sum_{j=1}^{q_n} G_{j\Delta} \right) + \text{Var} \left(\sum_{j=1}^{q_n} H_{j\Delta} \right) \\ &\lesssim q_n^2 + q_n^2 \Delta + \frac{q_n}{\Delta} = q_n^2 + \frac{q_n}{\Delta}. \end{aligned}$$

476 And on $\Omega_{B,\alpha}$:

$$\left| \sum_{j=1}^{q_n} V_{j\Delta} \right| \lesssim \sum_{j=1}^{q_n} \frac{B_{j\Delta}^8 + Z_{j\Delta}^8 + J_{j\Delta}^8}{\Delta} \lesssim q_n \ln^8(n) \Delta^{-\alpha}.$$

477 Then, with $V' = q_n + 1/\Delta$ and $M'_2 = \ln^8(n) \Delta^{-\alpha}$, by Bennett's inequality, we get on $\Omega_{B,\alpha}$:

$$\begin{aligned} \mathbb{P} \left(|\bar{V}_n - \mathbb{E}(\xi^8(X_\Delta)) I_8| \geq \Delta^{1/4} + x \right) &\leq \exp \left(-\frac{nx^2}{2(V' + M'_2 x q_n)} \right) \\ &= \exp \left(-\frac{Cnx^2}{\frac{1}{\Delta} + \frac{\ln(n)}{\Delta} + \frac{\ln^9(n)x\Delta^{-\alpha}}{\Delta}} \right) \end{aligned}$$

478 Taking $x = 1/\ln(n)$, as $\Delta = O(n^{-\delta})$ and $\ln(n) = o(1/\Delta)$, we get:

$$\mathbb{P} \left(|\bar{V}_n - \mathbb{E}(\xi^8(X_\Delta) I_8)| \geq \frac{1}{\ln(n)} + \Delta^{1/4} \right) \lesssim \exp \left(-\frac{cn\Delta}{\ln^2(n) (1 + \ln(n) + \ln^8(n) \Delta^{-\alpha})} \right) \lesssim n^{-5}.$$

479 **Bound of $\pi_0 - \hat{\pi}_0$.** We have that $\hat{\pi}_h(x) = \frac{1}{n} \sum_{k=1}^n K_h(X_{k\Delta}) - x$. The expectation of this term is
 480 $\mathbb{E}(\hat{\pi}_h(x)) = \pi_h(x) := \int K_h(y-x)\pi(y)dy$. We have the following bounds:

$$\text{Var} \left(\sum_{j=kq_n+1}^{(k+1)q_n} K_h(X_{j\Delta}) \right) \lesssim \frac{q_n^2}{h} \quad \text{and} \quad \left| \sum_{j=kq_n+1}^{(k+1)q_n} K_h(X_{j\Delta}) \right| \lesssim \frac{q_n}{h}.$$

481 Then by Bennett's inequality (lemma 16) with $V' = Cq_n/h$ and $M'_2 = C'/h$, we get, for any x :

$$\mathbb{P} \left(|\hat{\pi}_h(x) - \pi_h(x)| \geq c \frac{\ln^2(n)}{(n\Delta)^{1/4}} \right) \lesssim \exp \left(- \frac{nc^2 \ln^4(n)/(n\Delta)^{1/2}}{\frac{Cq_n}{h} + \frac{C'cq_n \ln^2(n)}{(n\Delta)^{1/4}h}} \right) \lesssim \exp \left(- \frac{C''nh \ln^4(n)}{(n\Delta)^{1/2}q_n} \right).$$

482 Let us take $h = (n\Delta)^{-1/2}$, as q_n is proportional to $\frac{\ln(n)}{\Delta}$, we obtain:

$$\mathbb{P} \left(|\hat{\pi}_h(x) - \pi_h(x)| \geq c \frac{\ln^2(n)}{(n\Delta)^{1/4}} \right) \lesssim \exp(-C \ln^3(n)).$$

483 Moreover, as π is continuous, by [Tsybakov \(2004, Proposition 1.2 with \$\beta = 1\$ \)](#): $|\pi_h(x) - \pi(x)| \lesssim$
 484 $C'h$. Let $(x_1, \dots, x_{\ln^2(n)})$ be equally spaced points on the interval A . Then

$$\begin{aligned} |\hat{\pi}_0 - \pi_0| &\leq \min_{1 \leq j \leq \ln^2(n)} |\pi(x_j) - \pi_0| + \max_{1 \leq j \leq \ln^2(n)} |\hat{\pi}_{(n\Delta)^{-1/2}}(x_j) - \pi(x_j)| \\ &\lesssim \frac{1}{\ln^2(n)} + \frac{\ln^2(n)}{(n\Delta)^{1/2}} + \max_{1 \leq j \leq \ln^2(n)} |\hat{\pi}_{(n\Delta)^{-1/2}}(x_j) - \pi_{(n\Delta)^{-1/2}}(x_j)| \end{aligned}$$

485 and therefore

$$\mathbb{P} \left(|\hat{\pi}_0 - \pi_0| \geq \frac{1}{\ln(n)} + \frac{\ln^2(n)}{(n\Delta)^{1/4}} \right) \lesssim n^{-5}$$

486 which ends the proof.

487 Conflict of interest

488 On behalf of all authors, the corresponding author states that there is no conflict of interest.

489 References

- 490 Y. Aït-Sahalia and J. Jacod. Estimating the degree of activity of jumps in high frequency data.
 491 *Ann. Statist.*, 37(5A):2202–2244, 2009. ISSN 0090-5364,2168-8966. doi: 10.1214/08-AOS640.
 492 URL <https://doi.org/10.1214/08-AOS640>.
- 493 D. Applebaum. *Lévy processes and stochastic calculus*, volume 116 of *Cambridge Studies*
 494 *in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2009.
 495 ISBN 978-0-521-73865-1. doi: 10.1017/CBO9780511809781. URL [https://doi.org/10.1017/](https://doi.org/10.1017/CBO9780511809781)
 496 [CBO9780511809781](https://doi.org/10.1017/CBO9780511809781).

- 497 F. M. Bandi and T. H. Nguyen. On the functional estimation of jump-diffusion models. volume 116,
498 pages 293–328. 2003. doi: 10.1016/S0304-4076(03)00110-6. URL [https://doi.org/10.1016/
499 S0304-4076\(03\)00110-6](https://doi.org/10.1016/S0304-4076(03)00110-6). Frontiers of financial econometrics and financial engineering.
- 500 A. Barron, L. Birgé, and P. Massart. Risk bounds for model selection via penalization. *Probab.
501 Theory Related Fields*, 113(3):301–413, 1999. ISSN 0178-8051,1432-2064. doi: 10.1007/
502 s004400050210. URL <https://doi.org/10.1007/s004400050210>.
- 503 J. Berestycki. Exchangeable fragmentation-coalescence processes and their equilibrium measures.
504 *Electron. J. Probab.*, 9:no. 25, 770–824, 2004. ISSN 1083-6489. doi: 10.1214/EJP.v9-227. URL
505 <https://doi.org/10.1214/EJP.v9-227>.
- 506 A. Carpentier, C. Duval, and E. Mariucci. Total variation distance for discretely observed Lévy
507 processes: a Gaussian approximation of the small jumps. *Ann. Inst. Henri Poincaré Probab.
508 Stat.*, 57(2):901–939, 2021. ISSN 0246-0203,1778-7017. doi: 10.1214/20-aihp1102. URL [https:
509 //doi.org/10.1214/20-aihp1102](https://doi.org/10.1214/20-aihp1102).
- 510 F. Comte, V. Genon-Catalot, and Y. Rozenholc. Penalized nonparametric mean square estimation
511 of the coefficients of diffusion processes. *Bernoulli*, 13(2):514–543, 2007. ISSN 1350-7265,1573-
512 9759. doi: 10.3150/07-BEJ5173. URL <https://doi.org/10.3150/07-BEJ5173>.
- 513 R. A. DeVore and G. G. Lorentz. *Constructive approximation*, volume 303 of *Grundlehren der
514 mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-
515 Verlag, Berlin, 1993. ISBN 3-540-50627-6.
- 516 M. Hanif. A nonparametric approach to the estimation of jump-diffusion models with asymmetric
517 kernels. *Cogent Math.*, 3:Art. ID 1179247, 15, 2016. ISSN 2331-1835. doi: 10.1080/23311835.
518 2016.1179247. URL <https://doi.org/10.1080/23311835.2016.1179247>.
- 519 M. Hanif, H. Wang, and Z. Lin. Reweighted Nadaraya-Watson estimation of jump-diffusion models.
520 *Sci. China Math.*, 55(5):1005–1016, 2012.
- 521 C. Mancini. Non-parametric threshold estimation for models with stochastic diffusion coefficient
522 and jumps. *Scand. J. Stat.*, 36(2):270–296, 2009. ISSN 0303-6898,1467-9469. doi: 10.1111/j.
523 1467-9469.2008.00622.x. URL <https://doi.org/10.1111/j.1467-9469.2008.00622.x>.
- 524 C. Mancini and R. Renò. Threshold estimation of Markov models with jumps and interest rate
525 modeling. *J. Econometrics*, 160(1):77–92, 2011. ISSN 0304-4076,1872-6895. doi: 10.1016/j.
526 jeconom.2010.03.019. URL <https://doi.org/10.1016/j.jeconom.2010.03.019>.
- 527 H. Masuda. Ergodicity and exponential β -mixing bounds for multidimensional diffusions with
528 jumps. *Stochastic Process. Appl.*, 117(1):35–56, 2007. ISSN 0304-4149,1879-209X. doi: 10.1016/
529 j.spa.2006.04.010. URL <https://doi.org/10.1016/j.spa.2006.04.010>.
- 530 J. Y. Park and B. Wang. Nonparametric estimation of jump diffusion models. *J. Econometrics*,
531 222(1):688–715, 2021. ISSN 0304-4076,1872-6895. doi: 10.1016/j.jeconom.2020.07.020. URL
532 <https://doi.org/10.1016/j.jeconom.2020.07.020>.
- 533 P. Protter and D. Talay. The Euler scheme for Lévy driven stochastic differential equations. *Ann.
534 Probab.*, 25(1):393–423, 1997. ISSN 0091-1798,2168-894X. doi: 10.1214/aop/1024404293. URL
535 <https://doi.org/10.1214/aop/1024404293>.

- 536 E. Schmisser. Non-parametric adaptive estimation of the drift for a jump diffusion process. *Stochas-*
537 *tic Process. Appl.*, 124(1):883–914, 2014. ISSN 0304-4149,1879-209X. doi: 10.1016/j.spa.2013.09.
538 012. URL <https://doi.org/10.1016/j.spa.2013.09.012>.
- 539 E. Schmisser. Non parametric estimation of the diffusion coefficients of a diffusion with jumps.
540 *Stochastic Process. Appl.*, 129(12):5364–5405, 2019. ISSN 0304-4149,1879-209X. doi: 10.1016/j.
541 spa.2019.03.003. URL <https://doi.org/10.1016/j.spa.2019.03.003>.
- 542 Y. Shimizu. M -estimation for discretely observed ergodic diffusion processes with infinitely many
543 jumps. *Stat. Inference Stoch. Process.*, 9(2):179–225, 2006. ISSN 1387-0874,1572-9311. doi:
544 10.1007/s11203-005-8113-y. URL <https://doi.org/10.1007/s11203-005-8113-y>.
- 545 Y. Shimizu and N. Yoshida. Estimation of parameters for diffusion processes with jumps from
546 discrete observations. *Stat. Inference Stoch. Process.*, 9(3):227–277, 2006. ISSN 1387-0874,1572-
547 9311. doi: 10.1007/s11203-005-8114-x. URL <https://doi.org/10.1007/s11203-005-8114-x>.
- 548 K. Song, Y. Song, and H. Wang. Threshold reweighted Nadaraya-Watson estimation of jump-
549 diffusion models. *Probab. Uncertain. Quant. Risk*, 7(1):31–44, 2022. ISSN 2095-9672,2367-0126.
550 doi: 10.3934/puqr.2022003. URL <https://doi.org/10.3934/puqr.2022003>.
- 551 A. B. Tsybakov. *Introduction à l'estimation non-paramétrique*, volume 41 of *Mathématiques &*
552 *Applications (Berlin) [Mathematics & Applications]*. Springer-Verlag, Berlin, 2004. ISBN 3-540-
553 40592-5.