



Blasticidin S Deaminase: A New Efficient Selectable Marker for *Chlamydomonas reinhardtii*

Félix de Carpentier, Jeanne Le Peillet, Nicolas D Boisset, Pierre Crozet, Stéphane D Lemaire, Antoine Danon

► To cite this version:

Félix de Carpentier, Jeanne Le Peillet, Nicolas D Boisset, Pierre Crozet, Stéphane D Lemaire, et al.. Blasticidin S Deaminase: A New Efficient Selectable Marker for *Chlamydomonas reinhardtii*. *Frontiers in Plant Science*, 2020, 11, pp.242. 10.3389/fpls.2020.00242 . hal-03721478

HAL Id: hal-03721478

<https://hal.science/hal-03721478>

Submitted on 12 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Blasticidin S Deaminase: A New Efficient Selectable Marker for *Chlamydomonas reinhardtii*

Félix de Carpentier^{1,2†}, Jeanne Le Peillet^{1†}, Nicolas D. Boisset^{1,2}, Pierre Crozet¹, Stéphane D. Lemaire¹ and Antoine Danon^{1*}

¹ Institut de Biologie Physico-Chimique, UMR 8226, CNRS, Sorbonne Université, Paris, France, ² Université Paris-Saclay, Saint-Aubin, France

OPEN ACCESS

Edited by:

Dimitris Petroustos,
UMR 5168 Laboratoire de Physiologie
Cellulaire Végétale (LPCV), France

Reviewed by:

Xenie Johnson,
Commissariat à l'Energie Atomique et
aux Energies Alternatives (CEA),
France
Véronique Larosa,
University of Liège, Belgium
Ryutaro Tokutsu,
National Institute for Basic Biology,
Japan

*Correspondence:

Antoine Danon
antoine.danon@ibpc.fr

[†] These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Plant Biotechnology,
a section of the journal
Frontiers in Plant Science

Received: 13 November 2019

Accepted: 17 February 2020

Published: 05 March 2020

Citation:

de Carpentier F, Le Peillet J,
Boisset ND, Crozet P, Lemaire SD
and Danon A (2020) Blasticidin S
Deaminase: A New Efficient
Selectable Marker
for *Chlamydomonas reinhardtii*.
Front. Plant Sci. 11:242.
doi: 10.3389/fpls.2020.00242

Chlamydomonas reinhardtii is a model unicellular organism for basic or biotechnological research, such as the production of high-value molecules or biofuels thanks to its photosynthetic ability. To enable rapid construction and optimization of multiple designs and strains, our team and collaborators have developed a versatile *Chlamydomonas* Modular Cloning toolkit comprising 119 biobricks. Having the ability to use a wide range of selectable markers is an important benefit for forward and reverse genetics in *Chlamydomonas*. We report here the development of a new selectable marker based on the resistance to the antibiotic blasticidin S, using the *Bacillus cereus* blasticidin S deaminase (*BSR*) gene. The optimal concentration of blasticidin S for effective selection was determined in both liquid and solid media and tested for multiple laboratory strains. In addition, we have shown that our new selectable marker does not interfere with other common antibiotic resistances: zeocin, hygromycin, kanamycin, paromomycin, and spectinomycin. The blasticidin resistance biobrick has been added to the *Chlamydomonas* Modular Cloning toolkit and is now available to the entire scientific community.

Keywords: blasticidin, *Chlamydomonas reinhardtii*, antibiotic resistance, selectable marker, synthetic biology, modular cloning, algal biotechnology

INTRODUCTION

Chlamydomonas reinhardtii is a model microalga widely used for basic and biotechnological research such as photosynthesis, cilia/flagella, production of biofuels, or other molecules of interest (Georgianna and Mayfield, 2012; Barahimipour et al., 2016; Salomé and Merchant, 2019). In the last decades *Chlamydomonas* has been shown to be amenable to powerful genetic approaches including CRISPR-Cas9 gene editing (Jiang et al., 2014; Greiner et al., 2017; Kao and Ng, 2017). The creation of the *Chlamydomonas* Library Project (CLiP), which contains more than 62,000 mutants covering roughly 83% of *Chlamydomonas* genes (Li et al., 2019), has also greatly contributed to the development of *Chlamydomonas* reverse genetics. A wide range of molecular tools for engineering of the nuclear genome are also available in *Chlamydomonas*, most of which have been grouped in a Modular Cloning toolkit (*Chlamy* MoClo toolkit). This collection contains 119 biobricks (promoters, terminators, reporter genes, selectable markers, targeting peptides, antibiotic resistance genes, riboswitch, miRNA backbone, etc.), which can be easily assembled through Golden Gate cloning (Crozet et al., 2018). Most applications related to the manipulation

of nuclear gene expression require transformation of *Chlamydomonas* cells followed by selection of transformants on plates using a selectable marker, which is in most cases an antibiotic resistance gene enabling selection on a medium containing the appropriate antibiotic. In the case of multiple consecutive transformations several selectable markers are required. Therefore, it is essential for advanced genetic engineering to have as many selectable markers as possible. At the moment, six antibiotic resistances are commonly used in *Chlamydomonas* as selectable markers: zeocin (Stevens et al., 1996), hygromycin (Berthold et al., 2002), kanamycin (Barahimipour et al., 2016), paromomycin (Sizova et al., 2001), sulfadiazine (Tabatabaei et al., 2019), and spectinomycin (Meslet-Cladière and Vallon, 2011).

Blasticidin S (hereafter referred to as “blasticidin”) is an antibiotic that inhibits cytosolic protein synthesis by blocking ribosomal translation termination (Svidritskiy et al., 2013). Blasticidin S deaminase (BSR) detoxifies blasticidin by catalyzing its deamination (Seto et al., 1966; Endo et al., 1987). The BSR gene has been successfully used as a selectable marker in mammals (Izumi et al., 1991), plants (Kamakura et al., 1990), yeasts (Kimura et al., 1994; Fukuda and Kizaki, 1999), and algae, including the diatom *Phaeodactylum tricornutum* (Buck et al., 2018) and the *Chlamydomonas*-related volvocine alga *Volvox carteri* (Ortega-Escalante et al., 2018).

We report here that blasticidin can be used as an antibiotic in *Chlamydomonas*, for all six laboratory strains tested. We show that BSR is functional in *Chlamydomonas* and can be used as a selectable marker, without conferring resistance to the other commonly used antibiotics. We also show that BSR can be used in combination with all other antibiotic resistance genes available in *Chlamydomonas*.

MATERIALS AND METHODS

Strains, Media, and Growth Conditions

If not otherwise specified, the reference strain used in the present study is CC-4533 (CMJ030), the CLiP library recipient strain used to generate the insertional mutants (Li et al., 2019). We also used other common laboratory strains CC-4051 (4A+) (Kim et al., 2005), CC-400 (cw15) (Roy Davies, 1972), CC-4425 (D66) (Schnell and Lefebvre, 1993), UVM4 (Neupert et al., 2009), and CC-124 (137c) (Pröschold et al., 2005). *Chlamydomonas* cells were grown on agar plates or liquid medium, using Tris-acetate-phosphate (TAP) medium (Gorman and Levine, 1965) at 25°C, under continuous light (40–60 $\mu\text{mol photon}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and shaking (130 rpm). Growth analysis were performed in the Algem® lab scale double photobioreactor systems (Algenuity, Stewartby, United Kingdom) under continuous light (200 $\mu\text{mol photons}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and 120 rpm agitation in TAP or high salt medium (HSM) (Sueoka, 1960), with bubbling air. The absorbance at 740 nm was recorded every 10 min using the built-in sensor. The maximal growth rate was determined as the maximal slope of the growth curve ($\Delta\text{abs}/\Delta\text{time}$). Linearity of the growth curve was estimated through linear regression every 200 min over a period of 13 h for TAP or 300 min over a period

of 40 h for HSM condition. The slopes of these linear regressions were selected only for regressions displaying an $R^2 < 99.5\%$. Finally, the maximal slope was chosen as the maximal growth rate (Supplementary Figure S2B). Antibiotics used were blasticidin S (Sigma-Aldrich: SBR00022), zeocin (Invitrogen: R25005), hygromycin B (Sigma-Aldrich: H9773), kanamycin (Sigma-Aldrich: K1377), paromomycin (Sigma-Aldrich: P8692), or spectinomycin (Sigma-Aldrich: S4014). *Chlamydomonas* multi-well plates and Petri dishes were scanned using a Perfection V800 scanner (Epson).

Cell Death Quantification

Dead cells were detected using Evans blue (Sigma-Aldrich: E2129) at a final concentration of 0.2% w/v. Dead cells appear in blue whereas living cells that are impermeable to Evans blue remain green (Gaff and Okong'o-Ogola, 1971). Cells were observed with a microscope (Olympus BX43, Tokyo, Japan). For each sample multiple microscopic fields were analyzed and Evans blue-positive cells scored. For each value the percentage of dead cells was calculated on a minimum of 100 individuals.

Plasmid Construction

Protein and nucleic acid designs were performed *in silico* on Serial Cloner 2.6.1 software. *Bacillus cereus* BSR protein sequence (NCBI accession number: WP_076871832.1) was reverse translated using *Chlamydomonas* nuclear genome codon usage table¹. The resulting BSR coding sequence was domesticated by removing *BbsI* restriction sites, designed for the position B3–B5 of the common Plant MoClo syntax (Patron et al., 2015), synthesized (Twist Bioscience), and cloned by Golden Gate reaction with *BsaI*-HFv2 (New England Biolabs) in pICH41308 (Weber et al., 2011) yielding plasmid pCM0-120, numbered according to the *Chlamydomonas* MoClo toolkit nomenclature (Crozet et al., 2018). Two other parts from the toolkit were used to build the BSR module (pCM1-029), the promoter $P_{A/R}$ coupled to the 5'UTR of RBCS2 (pCM0-020) and the 3'UTR of RBCS2 coupled to terminator T_{RBCS2} (pCM0-115) (Schroda et al., 2002; Crozet et al., 2018). The other antibiotic resistance genes were built using these same regulatory sequences and the coding sequence (CDS) from pCM0-077 (zeocin), pCM0-073 (hygromycin), pCM0-074 (paromomycin), pCM0-075 (kanamycin) and pCM0-076 (spectinomycin) generating the plasmids pCM1-030 (zeocin), pCM1-031 (hygromycin), pCM1-032 (kanamycin), pCM1-027 (paromomycin), and pCM1-033 (spectinomycin), respectively. All plasmid sequences are available in Supplementary Files S1–S7.

Chlamydomonas Transformation

Transformations were performed as previously described (Crozet et al., 2018). Briefly, *Chlamydomonas* cells were grown in TAP to early exponential phase ($1\text{--}2 \times 10^6$ cells/mL), concentrated 100 times in TAP + 60 mM sucrose. After incubation of 250 μL of cells with DNA (55 fmol of purified resistance module excised with *BbsI*-HF; New England Biolabs) at 4°C for 20 min, they were electroporated (2000 V/cm, 25 μF , no shunt resistance)

¹www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=3055

and incubated for 16–20 h in 10 mL of TAP + 60 mM sucrose prior to be plated on TAP-agar complemented with appropriate antibiotic(s). Transformants were selected on TAP-agar medium containing blasticidin S (50 mg/L, unless otherwise specified), zeocin (15 mg/L), hygromycin B (20 mg/L), kanamycin (50 mg/L), paromomycin (20 mg/L), and/or spectinomycin (100 mg/L). Plates and transformants were analyzed after 5–7 days of growth in continuous light ($50 \mu\text{mol photon m}^{-2} \text{s}^{-1}$) at 25°C.

Insert Detection

Cells were pelleted (5 min, $2500 \times g$, room temperature) and lysed in 400 μL of extraction buffer (200 mM Tris-HCl pH 7.5; 200 mM NaCl; 25 mM EDTA; 0.5% SDS) for 10 min at 37°C under agitation (1400 rpm). After centrifugation (3 min, $17,000 \times g$, room temperature), the supernatant was harvested and the genomic DNA was precipitated with one volume of isopropanol for 10 min at room temperature, washed with 70% ethanol, dried and resuspended in water. PCR was performed using the Quick-Load® Taq 2 \times Master Mix (New England Biolabs) according to the manufacturer recommendations with the primers BSR.5 (5'-GCTGTACGAGGACAACAAGC-3'), TRBCS2.3 (5'-ACGGAGGATCGTTACAACC-3'), CBLP.5 (5'-GACGTCATCCACTGCCTGTG-3'), and CBLP.3 (5'-CGACGCATCCTCAACACACC-3').

RESULTS

Chlamydomonas Is Sensitive to Blasticidin

To test the sensitivity of *Chlamydomonas* to blasticidin, the cells were grown in the presence of increasing blasticidin concentrations (25, 50, or 75 mg/L) in both solid and liquid cultures. Blasticidin was very effective to kill *Chlamydomonas* cells and the minimum efficient concentration was 50 mg/L in both conditions (Figure 1A). Many reference strains are used within the community and they present an important genetic diversity and substantial phenotypic differences (Gallaher et al., 2015). Compared to the reference strain of this study (CC-4533), some strain specific phenotypes, such as the absence of the cell wall, could alter blasticidin resistance. To assess whether this genetic diversity among common reference strains affects their sensitivity to blasticidin, CC-4051 (4A+), CC-400 (cw15), CC-4425 (D66+), UVM4, and CC-124 (137c) were cultivated on a solid medium supplemented or not with blasticidin (50 mg/L). In all cases, none of the strains survived in the presence of blasticidin, regardless of the initial number of cells tested (Figure 1B).

Blasticidin S Deaminase Can Be Used as a Selectable Marker in *Chlamydomonas*

To test whether the BSR module could be used as a selectable marker in *Chlamydomonas*, we engineered *B. cereus* BSR coding sequence to fit the optimal codon bias of *Chlamydomonas* nuclear genome. This has been already shown to improve

transgene expression efficiency, including selection markers (Barahimipour et al., 2015, 2016). The engineered BSR coding sequence was then domesticated by removing *BbsI* and *BsaI* recognition motif and adding appropriate fusion sites of the Plant MoClo syntax (Patron et al., 2015). It was finally assembled with the parts $P_{A/R}$ (the hybrid promoter of *HSP70A* coupled to the 5'UTR of *RBCS2*) and T_{RBCS2} (coupling the 3'UTR and terminator of *RBCS2*) to form a functional module. $P_{A/R}$ is a chimeric constitutive promoter made up of *HSP70A* and *RBCS2* promoters that was proven to be very efficient in *Chlamydomonas* by significantly reducing gene silencing (Schroda et al., 2002). We also chose this promoter/terminator combination because it allows successful expression of the same BSR gene in *Volvox* (Ortega-Escalante et al., 2018). The resulting construct pCM1-029 (pCM stands for plasmid *Chlamydomonas* MoClo) is represented in Figure 2A using the MoClo nomenclature (Crozet et al., 2018). pCM0-120 and pCM1-029 plasmid sequences are available in Supplementary Files S1, S2, respectively. Cells transformed with pCM1-029 or an empty vector were incubated on plates containing blasticidin. Transformants appeared only when the cells were transformed with pCM1-029 (Blasticidin resistant cells are hereafter referred to as "Blast^R"), as shown in Figure 2B, indicating that BSR can be used as a selectable marker in *Chlamydomonas*. The insertion of the BSR module was confirmed by PCR in four independent Blast^R strains (Supplementary Figure S1).

To precisely evaluate the effect of blasticidin on the mortality of wild-type and Blast^R cells, we used Evans blue as a death marker (Gaff and Okong'o-Ogola, 1971). To avoid the heterogeneity that comes from the position of the transgene insertion locus, dozens of Blast^R colonies were collected from a Petri dish and then mixed in liquid culture (Blast^R culture). Wild-type and Blast^R cultures were grown in a 24-well plate with or without blasticidin, and the percentage of dead cells was evaluated several days after treatment. In the wild-type strain treated with blasticidin, the percentage of dead cells increased after 3 days and all the cells died after 5 days, while no significant levels of death was detected for Blast^R and untreated cells (Figure 2C). To assess a potential detrimental effect of BSR on *Chlamydomonas*, wild-type and Blast^R growth were quantified in a photobioreactor, in both mixotrophic (TAP) and autotrophic (HSM) conditions. No differences in the growth rate were observed suggesting that BSR does not affect growth or photosynthesis (Supplementary Figure S2).

Blasticidin Can Be Used in Combination With Other Antibiotics

To test the possible interactions of blasticidin with the other most commonly used selectable markers in *Chlamydomonas*, we first generated strains resistant to these antibiotics by transformation of wild-type cells with plasmids containing modules conferring resistance to zeocin (pCM1-030), hygromycin (pCM1-031), kanamycin (pCM1-032), paromomycin (pCM1-027), and spectinomycin (pCM1-033) under the control of the same regulatory sequences used for BSR in pCM1-029. Transformants were selected on plates containing the appropriate antibiotic




FIGURE 1 | Sensitivity of *Chlamydomonas* to blasticidin. **(A)** *Chlamydomonas* CC-4533 (CMJ030) was grown with increasing concentrations of blasticidin: 25, 50, or 75 mg/L, in both liquid and 24-well agar plates. For both liquid and solid media, cultures were photographed after 3 days of growth. **(B)** Spot tests were performed for six standard laboratory strains with or without blasticidin (50 mg/L). The results shown are representative of three biological replicates.

and pooled to take into account position effect, as for Blast^R culture. These cultures were called Zeo^R, Hygro^R, Kana^R, Paro^R, and Spec^R. The wild-type strain and the six antibiotic resistant cultures were cultivated in 96-well plates until exponential phase (5×10^6 cells/mL) prior to treatment with the different antibiotics. Five days after the addition of antibiotics, only the cells carrying the corresponding resistance gene survived the treatment, and importantly, only the Blast^R culture had survived upon blasticidin treatment (**Figure 3A**). It is to be noticed that Kana^R strains are also resistant to paromomycin (**Figure 3A**), as previously reported (Barahimipour et al., 2016).

If the resistance modules tested display no cross reactivity to one another, they can be combined to allow double selection. To use the BSR module and blasticidin in combination with another selectable marker, there must then be no interference between the different antibiotics and resistance modules. To verify possible interference, a single Blast^R strain was transformed with either pCM1-030 (zeocin resistance), pCM1-031 (hygromycin resistance), pCM1-032 (kanamycin resistance), pCM1-027 (paromomycin resistance), or pCM1-033 (spectinomycin resistance) (**Figure 3B**). Transformants were selected on plates containing both blasticidin and the appropriate antibiotic. For each combination tested, transformants resistant




FIGURE 2 | Blastcidin S deaminase expression in *Chlamydomonas*. **(A)** The blastcidin resistance module pCM1-029 is built from a constitutive promoter ($P_{A/R}$ + 5'UTR of RBCS2), the BSR coding sequence and a terminator (3'UTR of RBCS2 + T_{RBCS2}) and was assembled using Golden Gate cloning. The numbers 1, 6, 9, and 11 stand for the standard fusion sites used for cloning (Crozet et al., 2018). SBOL2.0 visual syntax is shown above the module (Roehner et al., 2016). **(B)** CC-4533 cells were transformed with pCM1-029 or an empty vector (pICH47742) (Weber et al., 2011) and plated on TAP-agar supplemented with blastcidin (50 mg/L). Results are representative of three biological replicates. **(C)** The percentage of dead cells was evaluated using Evans blue in wild-type and Blast^R cultures with or without blastcidin, several days after treatment. Values represent the means and standard deviations of three independent experiments, symbol (*) shows the samples found to be significantly different doing an ANOVA analysis for each time points ($p < 0.001$).

to both antibiotics could be obtained (**Figure 3B**), indicating that no interference exists between the selectable markers tested.


DISCUSSION

Here we report the development of a new selectable marker for *C. reinhardtii*. Our data show that using blastcidin at a concentration of 50 mg/L ensures proper selection for all common laboratory strains of *Chlamydomonas* (**Figure 2C**). This concentration is slightly higher than the efficient concentration reported for *V. carteri* (Ortega-Escalante et al., 2018) or diatoms (Buck et al., 2018), but remains comparable with other antibiotics used in *Chlamydomonas* (Crozet et al., 2018).

We successfully engineered BSR coding sequence to adapt it to *Chlamydomonas* and no addition of intron was necessary for efficient expression, contrary to what was reported for *V. carteri* (Ortega-Escalante et al., 2018). Now that we have developed the blastcidin resistance module, seven antibiotic-based selectable markers are available for *Chlamydomonas*. This new tool can become important for advanced synthetic biology

strategies requiring successive transformations of the same strain in combination with new engineering tools including the CLiP library (Li et al., 2019), the CRISPR/Cas technology (Jiang et al., 2014; Greiner et al., 2017; Kao and Ng, 2017) and the MoClo toolkit (Crozet et al., 2018). It is also important to increase the number of selectable markers available because the use of certain antibiotics should be taken with care. For instance zeocin is not always recommended since it can potentially cause DNA damages (Chankova et al., 2007; Čížková et al., 2019) and subsequent unwanted mutations. The kanamycin selectable marker *NPTII* is also conferring resistance to paromomycin (Barahimipour et al., 2015; **Figure 3A**) which makes it impossible to use in CLiP strains that are paromomycin resistant (Li et al., 2019). We show here that it is possible to use BSR and blastcidin in combination with all the other commonly used selectable markers in *Chlamydomonas*. BSR gene has been integrated as a new biobrick into the *Chlamydomonas* MoClo toolkit, and is now available to the entire community through the *Chlamydomonas* Resource Center².

²www.chlamycollection.org/



DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/Supplementary Material.

AUTHOR CONTRIBUTIONS

FC, JL, PC, SL, and AD designed the study and analyzed the data. FC, JL, NB, PC, SL, and AD wrote the manuscript. FC, JL, NB, and AD performed the experiments.

FUNDING

This work was supported in part by the CNRS, Sorbonne Université, and Université Paris-Saclay, by Agence Nationale de la Recherche Grant 17-CE05-0001 CalvinDesign and by LABEX DYNAMO ANR-LABX-011.

ACKNOWLEDGMENTS

We thank Sandrine Bujaldon for providing the *Chlamydomonas reinhardtii* strains CC-4051, CC-400, and CC-124; Kevin Sarkissian for his help in the cloning; and Cyrielle Durand, Théo Le Moigne, Dr. Christophe Marchand, and Dr. Julien Henri for stimulating discussions and suggestions.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2020.00242/full#supplementary-material>

FIGURE S1 | Detection of the *BSR* module. pCM1-029 plasmid and the genomic DNA of the wild type (CC-4533) and four independent transformants Blast^R were used to amplify by PCR a fragment of the *BSR* resistance module. A product at the expected size (546 bp) was amplified only for pCM1-029 (positive control) and

the Blast^R transformants. Amplification of CBLP (Cre06.g278222) was used as a positive control for the experiment.

FIGURE S2 | Growth of wild-type (CC-4533) and Blast^R strains in a controlled photobioreactor. **(A)** Measurement of growth as detected by absorbance at 740 nm every 10 min, in TAP (mixotrophy) and HSM medium (autotrophy), light was set to 200 $\mu\text{mol photons}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ with temperature and shaking set at 25°C and 120 rpm, respectively. **(B)** Quantification of the average growth rate of wild-type and Blast^R cells, corresponding to the growth curves shown in Graph **(A)**. Error bars represent the standard error to the mean of two biological replicates.

FIGURE S3 | Transformation of a blasticidin resistant strain with zeocin, hygromycin, kanamycin, paromomycin and spectinomycin resistance plasmids. A blasticidin resistant strain (Blast^R) carrying pCM1-029 was independently transformed with the resistance modules to zeocin (pCM1-030), hygromycin (pCM1-031), kanamycin (pCM1-032), paromomycin (pCM1-027), and spectinomycin (pCM1-033) or an empty vector (pICH50881). The numbers 1, 6, 9, and 11 stand for the standard fusion sites used for cloning (Crozet et al., 2018). Blast^R transformed cells were selected on plate containing both blasticidin and the appropriate antibiotic. CC-4533 was also transformed with the same five resistance modules or the empty vector and selected on the corresponding antibiotic with or without addition of blasticidin. The results showed are representative of three biological replicates.

FILE S1 | Annotated sequence of pCM0-120, the level 0 plasmid containing *BSR* coding sequence engineered for *Chlamydomonas* nuclear genome and designed for the position B3–B5 of the *Chlamydomonas* MoClo toolkit (Crozet et al., 2018).

FILE S2 | Annotated sequence of pCM1-029, the level 1 plasmid made up of the promoter AR ($P_{A/R}$ + 5'UTR of *RBCS2*), *BSR* coding sequence and the terminator of *RBCS2* (3'UTR of *RBCS2* + T_{RBCS2}) conferring blasticidin resistance in *Chlamydomonas* (Crozet et al., 2018).

FILE S3 | Annotated sequence of pCM1-027, the level 1 plasmid made up of the promoter AR ($P_{A/R}$ + 5'UTR of *RBCS2*), *aphVIII* coding sequence and the terminator of *RBCS2* (3'UTR of *RBCS2* + T_{RBCS2}) conferring paromomycin resistance in *Chlamydomonas* (Sizova et al., 2001; Crozet et al., 2018).

FILE S4 | Annotated sequence of pCM1-030, the level 1 plasmid made up of the promoter AR ($P_{A/R}$ + 5'UTR of *RBCS2*), *Ble* coding sequence and the terminator *RBCS2* (3'UTR of *RBCS2* + T_{RBCS2}) conferring zeocin resistance in *Chlamydomonas* (Stevens et al., 1996; Crozet et al., 2018).

FILE S5 | Annotated sequence of pCM1-031, the level 1 plasmid made up of the promoter AR ($P_{A/R}$ + 5'UTR of *RBCS2*), *aph7''* coding sequence and the terminator *RBCS2* (3'UTR of *RBCS2* + T_{RBCS2}) conferring hygromycin resistance in *Chlamydomonas* (Berthold et al., 2002; Crozet et al., 2018).

FILE S6 | Annotated sequence of pCM1-032, the level 1 plasmid made up of the promoter AR ($P_{A/R}$ + 5'UTR of *RBCS2*), *NPTII* coding sequence and the terminator *RBCS2* (3'UTR of *RBCS2* + T_{RBCS2}) conferring kanamycin resistance in *Chlamydomonas* (Barahimipour et al., 2016; Crozet et al., 2018).

FILE S7 | Annotated sequence of pCM1-029, the level 1 plasmid made up of the promoter AR ($P_{A/R}$ + 5'UTR of *RBCS2*), *aadA* coding sequence and the terminator *RBCS2* (3'UTR of *RBCS2* + T_{RBCS2}) conferring spectinomycin resistance in *Chlamydomonas* (Meslet-Cladière and Vallon, 2011; Crozet et al., 2018).

REFERENCES

- Barahimipour, R., Neupert, J., and Bock, R. (2016). Efficient expression of nuclear transgenes in the green alga *Chlamydomonas*: synthesis of an HIV antigen and development of a new selectable marker. *Plant Mol. Biol.* 90, 403–418. doi: 10.1007/s11103-015-0425-8
- Barahimipour, R., Strenkert, D., Neupert, J., Schroda, M., Merchant, S. S., and Bock, R. (2015). Dissecting the contributions of GC content and codon usage to gene expression in the model alga *Chlamydomonas reinhardtii*. *Plant J.* 84, 704–717. doi: 10.1111/tpj.13033
- Berthold, P., Schmitt, R., and Mages, W. (2002). An engineered *Streptomyces hygroscopicus aph 7* Gene Mediates dominant resistance against hygromycin B in *Chlamydomonas reinhardtii*. *Protist* 153, 401–412. doi: 10.1078/14344610260450136
- Buck, J. M., Río Bártulos, C., Gruber, A., and Kroth, P. G. (2018). Blasticidin-S deaminase, a new selection marker for genetic transformation of the diatom *Phaeodactylum tricornutum*. *PeerJ* 6:e5884. doi: 10.7717/peerj.5884
- Chankova, S. G., Dimova, E., Dimitrova, M., and Bryant, P. E. (2007). Induction of DNA double-strand breaks by zeocin in *Chlamydomonas reinhardtii* and the role of increased DNA double-strand breaks rejoining in the formation of an adaptive response. *Radiat. Environ. Biophys.* 46, 409–416. doi: 10.1007/s00411-007-0123-2
- Čížková, M., Slavková, M., Vitová, M., Zachleder, V., and Bišová, K. (2019). Response of the green alga *Chlamydomonas reinhardtii* to the DNA damaging agent zeocin. *Cells* 8:E735. doi: 10.3390/cells8070735
- Crozet, P., Navarro, F. J., Willmund, F., Mehrshahi, P., Bakowski, K., Lauersen, K. J., et al. (2018). Birth of a photosynthetic chassis: a MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas reinhardtii*. *ACS Synth. Biol.* 7, 2074–2086. doi: 10.1021/acssynbio.8b00251
- Endo, T., Furuta, K., Kaneko, A., Katsuki, T., Kobayashi, K., Azuma, A., et al. (1987). Inactivation of blasticidin S by *Bacillus cereus*. I. inactivation mechanism. *J. Antibiot.* 40, 1791–1793. doi: 10.7164/antibiotics.40.1791
- Fukuda, H., and Kizaki, Y. (1999). A new transformation system of *Saccharomyces cerevisiae* with blasticidin S deaminase gene. *Biotechnol. Lett.* 21, 969–971. doi: 10.1023/A:1005613206138
- Gaff, D. F., and Okong'o-Ogola, O. (1971). The Use of non-permeating pigments for testing the survival of cells. *J. Exp. Bot.* 22, 756–758. doi: 10.1093/jxb/22.3.756
- Gallagher, S. D., Fitz-Gibbon, S. T., Glaesener, A. G., Pellegrini, M., and Merchant, S. S. (2015). *Chlamydomonas* genome resource for laboratory strains reveals a mosaic of sequence variation, identifies true strain histories, and enables strain-specific studies. *Plant Cell* 27, 2335–2352. doi: 10.1105/tpc.15.00508
- Georgianna, D. R., and Mayfield, S. P. (2012). Exploiting diversity and synthetic biology for the production of algal biofuels. *Nature* 488, 329–335. doi: 10.1038/nature11479
- Gorman, D. S., and Levine, R. P. (1965). Cytochrome f and plastocyanin: their sequence in the photosynthetic electron transport chain of *Chlamydomonas reinhardtii*. *Proc. Natl. Acad. Sci. U.S.A.* 54, 1665–1669. doi: 10.1073/pnas.54.6.1665
- Greiner, A., Kelterborn, S., Evers, H., Kreimer, G., Sizova, I., and Hegemann, P. (2017). Targeting of photoreceptor genes in *Chlamydomonas reinhardtii* via zinc-finger nucleases and CRISPR/Cas9. *Plant Cell* 29, 2498–2518. doi: 10.1105/tpc.17.00659
- Izumi, M., Miyazawa, H., Kamakura, T., Yamaguchi, I., Endo, T., and Hanaoka, F. (1991). Blasticidin S-resistance gene (*bsr*): a novel selectable marker for mammalian cells. *Exp. Cell Res.* 197, 229–233. doi: 10.1016/0014-4827(91)90427-V
- Jiang, W., Brueggeman, A. J., Horken, K. M., Plucinak, T. M., and Weeks, D. P. (2014). Successful transient expression of Cas9 and Single Guide RNA Genes in *Chlamydomonas reinhardtii*. *Eukaryot Cell* 13, 1465–1469. doi: 10.1128/EC.00213-14
- Kamakura, T., Yoneyama, K., and Yamaguchi, I. (1990). Expression of the blasticidin S deaminase gene (*bsr*) in tobacco: fungicide tolerance and a new selective marker for transgenic plants. *Mol. Gen. Genet.* 223, 332–334. doi: 10.1007/bf00265072
- Kao, P.-H., and Ng, I.-S. (2017). CRISPRi mediated phosphoenolpyruvate carboxylase regulation to enhance the production of lipid in *Chlamydomonas reinhardtii*. *Bioresour. Technol.* 245, 1527–1537. doi: 10.1016/j.biortech.2017.04.111
- Kim, K.-S., Feild, E., King, N., Yaoi, T., Kustu, S., and Inwood, W. (2005). Spontaneous mutations in the ammonium transport gene *AMT4* of *Chlamydomonas reinhardtii*. *Genetics* 170, 631–644. doi: 10.1534/genetics.105.041574
- Kimura, M., Kamakura, T., Zhou Tao, Q., Kaneko, I., and Yamaguchi, I. (1994). Cloning of the blasticidin S deaminase gene (*BSD*) from *Aspergillus terreus* and its use as a selectable marker for *Schizosaccharomyces pombe*

- and *Pyricularia oryzae*. *Mol. Gen. Genet.* 242, 121–129. doi: 10.1007/BF00391004
- Li, X., Patena, W., Fauser, F., Jinkerson, R. E., Saroussi, S., Meyer, M. T., et al. (2019). A genome-wide algal mutant library and functional screen identifies genes required for eukaryotic photosynthesis. *Nat. Genet.* 51, 627–635. doi: 10.1038/s41588-019-0370-376
- Meslet-Cladière, L., and Vallon, O. (2011). Novel shuttle markers for nuclear transformation of the green alga *Chlamydomonas reinhardtii*. *Eukaryotic Cell* 10, 1670–1678. doi: 10.1128/EC.05043-11
- Neupert, J., Karcher, D., and Bock, R. (2009). Generation of *Chlamydomonas* strains that efficiently express nuclear transgenes. *Plant J.* 57, 1140–1150. doi: 10.1111/j.1365-3113X.2008.03746.x
- Ortega-Escalante, J. A., Kwok, O., and Miller, S. M. (2018). New Selectable Markers for *Volvox carteri* Transformation. *Protist* 170, 52–63. doi: 10.1016/j.protis.2018.11.002
- Patron, N. J., Orzaez, D., Marillonnet, S., Warzecha, H., Matthewman, C., Youles, M., et al. (2015). Standards for plant synthetic biology: a common syntax for exchange of DNA parts. *New Phytol.* 208, 13–19. doi: 10.1111/nph.13532
- Pröschold, T., Harris, E. H., and Coleman, A. W. (2005). Portrait of a species: *Chlamydomonas reinhardtii*. *Genetics* 170, 1601–1610. doi: 10.1534/genetics.105.044503
- Roehner, N., Beal, J., Clancy, K., Bartley, B., Misirli, G., Grünberg, R., et al. (2016). Sharing structure and function in biological design with SBOL 2.0. *ACS Synth. Biol.* 5, 498–506. doi: 10.1021/acssynbio.5b00215
- Roy Davies, D. (1972). Cell wall organisation in *Chlamydomonas reinhardtii*. *Mol. Gen. Genet.* 115, 334–348. doi: 10.1007/BF00333172
- Salomé, P. A., and Merchant, S. S. (2019). A series of fortunate events: introducing *Chlamydomonas* as a reference organism. *Plant Cell* 31, 1682–1707. doi: 10.1105/tpc.18.00952
- Schnell, R. A., and Lefebvre, P. A. (1993). Isolation of the *Chlamydomonas* regulatory gene NIT2 by transposon tagging. *Genetics* 134, 737–747.
- Schroda, M., Beck, C. F., and Vallon, O. (2002). Sequence elements within an HSP70 promoter counteract transcriptional transgene silencing in *Chlamydomonas*. *Plant J.* 31, 445–455. doi: 10.1046/j.1365-3113X.2002.01371.x
- Seto, H., Ôtake, N., and Yonehara, H. (1966). Biological transformation of Blastidicin S by *Aspergillus fumigatus* sp. *Agric. Biol. Chem.* 30, 877–886. doi: 10.1080/00021369.1966.10858693
- Sizova, I., Fuhrmann, M., and Hegemann, P. (2001). A *Streptomyces rimosus aphVIII* gene coding for a new type phosphotransferase provides stable antibiotic resistance to *Chlamydomonas reinhardtii*. *Gene* 277, 221–229. doi: 10.1016/S0378-1119(01)00616-3
- Stevens, D. R., Purton, S., and Rochaix, J.-D. (1996). The bacterial phleomycin resistance gene *ble* as a dominant selectable marker in *Chlamydomonas*. *Mol. Gen. Genet.* 251, 23–30. doi: 10.1007/BF02174340
- Sueoka, N. (1960). Mitotic replication of deoxyribonucleic acid in *Chlamydomonas Reinhardtii*. *Proc. Natl. Acad. Sci. U.S.A.* 46, 83–91. doi: 10.1073/pnas.46.1.83
- Svidritskiy, E., Ling, C., Ermolenko, D. N., and Korostelev, A. A. (2013). Blastidicin S inhibits translation by trapping deformed tRNA on the ribosome. *Proc. Natl. Acad. Sci. U.S.A.* 110, 12283–12288. doi: 10.1073/pnas.1304922110
- Tabatabaei, I., Dal Bosco, C., Bednarska, M., Ruf, S., Meurer, J., and Bock, R. (2019). A highly efficient sulfadiazine selection system for the generation of transgenic plants and algae. *Plant Biotechnol. J.* 17, 638–649. doi: 10.1111/pbi.13004
- Weber, E., Engler, C., Gruetzner, R., Werner, S., and Marillonnet, S. (2011). A modular cloning system for standardized assembly of multigene constructs. *PLoS One* 6:e16765. doi: 10.1371/journal.pone.0016765

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 de Carpentier, Le Peillet, Boisset, Crozet, Lemaire and Danon. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.