

# The taxonomic structure but not the functioning of riparian herbaceous communities varies with hydrological conditions on a large, highly regulated river: Evidence from a 2-year replicated study

Philippe Janssen, Hervé Piégay, André Evette

# ▶ To cite this version:

Philippe Janssen, Hervé Piégay, André Evette. The taxonomic structure but not the functioning of riparian herbaceous communities varies with hydrological conditions on a large, highly regulated river: Evidence from a 2-year replicated study. Ecohydrology, 2022, 15 (3), pp.1-36. 10.1002/eco.2405. hal-03721511

# HAL Id: hal-03721511 https://hal.science/hal-03721511

Submitted on 30 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1	The taxonomic structure but not the functioning of riparian herbaceous communities varies with
2	hydrological conditions on a large, highly regulated river: evidence from a two-year replicated study
3	
4	Philippe Janssen <sup>1,2</sup> *, Hervé Piégay <sup>2</sup> , André Evette <sup>1</sup>
5	
6	<sup>1</sup> Univ. Grenoble Alpes, INRAE, LESSEM, F-38402 St-Martin-d'Hères, France
7	<sup>2</sup> Univ. Lyon, UMR 5600 EVS, CNRS, ENS de Lyon, F-69342, Lyon, France
8	
9	* Corresponding author, email philippe.janssen@inrae.fr, phone +33 476762879
10	
11	Email addresses of other authors:
12	H. Piégay (herve.piegay@ens-lyon.fr), A. Evette (andre.evette@inrae.fr)
13	
14	Short title: Spatiotemporal variations of riparian communities in a regulated river
15	Keywords: Gravel bar, Inundation, Plant traits, Rhône River, Riverine vegetation

#### 16 ABSTRACT

17 Riparian vegetation encompasses dynamic communities structured by strong environmental gradients. Due to 18 anthropogenic changes, a greater uniformity of environmental conditions is observed along local gradients. To 19 maintain diverse and functional communities in regulated rivers, there is an urgent need to finely characterize 20 the factors controlling the spatial and temporal dynamics of riparian vegetation. In the summer of 2017 and 21 2019, we repeatedly sampled 8 gravel bars along the regulated Rhône River. Using a taxonomic and functional 22 approach, we assessed whether the diversity and composition of species and traits varied between the two 23 years at the corridor and site scales. Taxonomic diversity decreased and composition changed between years, 24 while functional diversity and composition did not. Specifically, we showed that variation in taxonomic diversity 25 was largely limited to species sharing the same trait values, suggesting some functional resilience of riparian 26 communities to environmental changes. In addition, we found that changes between the two years concerned 27 primarily the communities established on low-lying bars made up of coarse-grained sediments, more 28 frequently and intensely inundated. Beyond being the most responsive to environmental changes, these 29 riparian communities were also the most diverse in species and trait values. However, by inducing a strong 30 decrease in diversity, fine sedimentation seems to be a major threat to the functional and ecological integrity 31 of gravel bars. Overall, these results highlight the conservation priority of low-lying gravel surfaces and the 32 need to promote more morphologically complex riparian habitats by renaturalizing flow variability and 33 sediment regimes.

34

#### 35 1. INTRODUCTION

36 River ecosystems are intrinsically dynamic and exhibit strong environmental gradients that shape the structure 37 of communities over space and time (Allan and Castillo, 2007). In free-flowing, i.e., unregulated and 38 unchannelled rivers, complex hydro-geomorphic processes promote water and sediment exchanges in several 39 dimensions of the hydrosystem (Ward, 1989). These exchanges allow the creation and renewal of a mosaic of complex habitats (Junk et al., 1989) that benefit a large number of species (Dudgeon et al., 2006). This is 40 41 particularly the case in riparian zones, i.e., in the transitional area between terrestrial and aquatic habitats, where plant communities face wide variations in disturbance and stress intensities along the flood gradient 42 43 (Naiman and Decamps, 1997). These rapid changes in environmental conditions induce a strong turnover in 44 plant strategies, i.e., from ruderal-anoxia-tolerant species at the bottom of riverbanks to competitive-45 desiccation-tolerant species at the top of riverbanks (Janssen et al., 2020a; Kyle and Leishman, 2009; McCoy-46 Sulentic et al., 2017). River-floodplain connectivity, conceptualized within the flood pulse framework (Junk et 47 al., 1989), is thus a key property of hydrosystems capable of ensuring the functionality and ecological integrity 48 of riparian zones (Naiman and Decamps, 1997). 49 Around the world, river connectivity has been profoundly affected by human infrastructures such as dams and 50 dikes (Grill et al., 2019; Nilsson et al., 2005). These developments have dramatically altered the flow regime, 51 i.e., by decreasing the intensity and magnitude of inundations (Poff et al., 2007), as well as the sediment 52 regime, i.e., by trapping sediment load and favoring fine overbank depositions (Wohl et al., 2015), of most 53 rivers. These anthropogenic changes have in turn altered the structure and dynamic of riparian communities, 54 by reducing native species diversity and increasing invasion by exotic species (Catford et al., 2011; Richardson 55 et al., 2007). Specifically, modifications to the flow and sediment regimes due to regulation/channelization 56 works has promoted greater uniformity of environmental conditions along local gradients (Poff et al., 2007) 57 and has lead to a simplification of riparian communities (Aguiar et al., 2018; Bejarano et al., 2018; Tonkin et al., 58 2018). This can compromise the ability of degraded rivers to support functional and diverse communities and, 59 as a result, the wide range of ecosystem services that riparian vegetation delivers to societies (Riis et al., 2020). 60 Although many studies have documented the impacts of anthropogenic environmental changes on riparian 61 vegetation, many have focused on woody species and forested habitats established on riverbanks (Aguiar et al., 62 2018; Bejarano et al., 2018; Janssen et al., 2020b; Johnson et al., 2012). In regulated and channelized rivers,

63 knowledge of the factors that structure herbaceous communities at fine spatial and temporal scales is thus still 64 scarce (Su et al., 2020), particularly for riparian habitats of high conservation value, such as gravel bars (Gilvear 65 et al., 2008). Composed of accumulations of alluvial sediments, these dynamic and morphologically complex 66 habitats are nevertheless a key element of the river landscape and shelter numerous threatened plant and animal species (Kalníková et al., 2018; Rottenborn et al., 2018; Zeng et al., 2015). However, due to 67 68 anthropogenic changes in flow and sediment regimes, the area of these habitats has been reduced to the 69 extent that unvegetated or sparsely vegetated bars with mobile sediments are classified as vulnerable in the 70 European Red List of Habitats (Janssen et al., 2016). To preserve and restore these species-rich riparian habitats 71 along regulated rivers, it is therefore urgent to better understand the factors that control the structure and 72 dynamics of herbaceous plant communities on gravel bars over space, i.e., by contrasting the effects of local 73 and regional gradients at different scales, and time, i.e., by resampling vegetation over several years. 74 In riparian ecology, several recent studies have mobilized a functional trait-based approach to understand the 75 effects of hydraulic alterations on vegetation (Aguiar et al., 2018; Bejarano et al., 2018; Janssen et al., 2020b; 76 Su et al., 2020; Tabacchi et al., 2019). Combined with a taxonomic approach, this allows us to focus both on 77 conservation issues, by studying variations in species richness and composition, but also to overcome the limits 78 of taxonomic studies, under the control of the local species pool, by studying the underlying ecological 79 mechanisms captured by functional traits (Lavorel et al., 2008). Building on these conceptual foundations, we 80 used a replicated sampling design spaced two years apart to study the structure and dynamic of plant 81 communities on gravel bars, at the corridor and site scales, along the highly degraded Rhône River. As shifts in 82 assemblage patterns are likely to result from a combination of different environmental factors related to the 83 flow or sediment regimes (e.g., Gilvear et al., 2008; Škornik et al., 2017), we tested the relative influence of a 84 set of topographic, pedological and hydrological variables on riparian plants, at the community scale. 85 Topographic variables were used to assess the influence of the shape of the bar and thus better guide bank reprofiling operations. Pedological variables were used to assess the influence of fine sediment deposition and 86 87 thus the need to develop specific management plans. Hydrological variables were used to assess the influence of temporal variations in flows and thus the potential for managing riparian vegetation through environmental 88 89 flows. Based on this scheme, we addressed the following two questions: (a) In regulated and channelized 90 rivers, how and to what extent do environmental conditions and taxonomic and functional 91 diversity/composition of gravel bar plants vary over a two-year period at the corridor and gravel bar scales? (b)

- 92 At the community scale, what is the relative influence of topographic, pedological and hydrological variables in
- 93 explaining variations in taxonomic and functional diversity/composition over time and space?

94

### 95 2. MATERIALS & METHODS

# 96 2.1. Study area and experimental design

97 The study was carried out along the Rhône River (total length = 810 km, catchment area = 96,500 km<sup>2</sup>, mean annual discharge =  $1,700 \text{ m}^3/\text{s}$ ) on a 250 km long stretch of its middle reach (Figure 1). This area is 98 characterized by a temperate climate with mean annual temperatures and precipitation of 13.6°C and 755 mm 99 100 in its southern part and 11.6°C and 815 mm in its northern part. Two historical development phases have greatly altered the functioning of this river: (i) during the 19<sup>th</sup> century, a correction phase was carried out to 101 facilitate navigation; (ii) in the second half of the 20<sup>th</sup> century, a derivation phase was carried out to produce 102 103 hydropower and facilitate irrigation and navigation (for details see, Bravard and Gaydou, 2015). These two 104 development phases induced significant impacts in the bypassed reaches (i.e., part of the historical Rhône River 105 channel), including a channel dewatering due to flow by-passing, reduction in the frequency and magnitude of 106 the peak flows (Vázquez-Tarrío et al., 2019), and channel stabilization due to engineering infrastructures. These 107 reaches also underwent a bedload supply interruption due to sediment trapping and extractions at tributary 108 confluences inducing bed armouring within the main channel. Thus, with the exception of the Miribel bypassed 109 reach, which is located in the northernmost part of the study area, present-day bedload transport is limited to 110 fine-grained sediments (Vázquez-Tarrío et al., 2019).

111 During the summer of 2017, we sampled 8 gravel bars (GB1-8) belonging to 7 bypassed reaches, along a north-112 south section of about 250 km (Figure 1). The same 8 gravel bars were resampled, at the same location during 113 the summer of 2019. The studied gravel bars are fixed in the bypassed reaches, i.e., coarse sediments are no 114 longer remobilized during flood events and are subject to fine-grained overbank sedimentation and colonization by trees. These large fixed bars correspond to a part of the river bottom before derivation and 115 116 represent the only terrestrial habitats available for species within the Rhône riverbed. They are colonized by different plant communities, dominated by hygrophilic perennial (e.g., Carex acutiformis, Lysimachia vulgaris, 117 118 Phalaris arundinacea) or annual species (e.g., Persicaria lapathifolia, Persicaria mitis, Rorippa sylvestris) near 119 the water line and xero-nitrophilous species (e.g., Chenopodium album, Bromus sterilis, Elymus pungens) at

higher elevations. Many exotic species (e.g., *Ambrosia artemisiifolia, Aster x salignus, Cyperus eragrostis*) have
also taken advantage of these surfaces, forming locally dense patches.

To prevent encroachment of gravel bars by woody species and the associated channel narrowing, river managers periodically rejuvenate the vegetation from the waterline to the edge of the alluvial forest, resulting in a blockage of plant ecological successions. Specifically, in the study area, the four northern gravel bars (i.e., GB1, GB2, GB3 and GB4) are managed by vegetation clearing using a tractor-drawn grinder, while the four southern gravel bars (i.e., GB5, GB6, GB7 and GB8) are managed by plowing using a bulldozer. These vegetation maintenance works were performed in the fall, annually for brush clearing and every two years for plowing (in 2016 and 2018 for the southern banks studied).

#### 129 2.2. Vegetation data

130 On each gravel bar, the vegetation was surveyed along three transects positioned perpendicular to the river in 131 areas with clear elevation gradients relative to the water surface. The gradients started at the water line and 132 ended before the alluvial floodplain. Along each transect, four 5 x 5 m quadrats (25 m<sup>2</sup>) were positioned in homogenous areas while avoiding the aquatic parts of the river margin (i.e., 12 quadrats per gravel bar). Within 133 134 each quadrat ( $n_{2017}$  = 96,  $n_{2019}$  = 96), all vascular plants were surveyed following the Braun-Blanquet 135 abundance/dominance methodology (Braun-Blanquet, 1932). Because maintenance works rejuvenated the 136 bars every year or two, tree and shrub layers were absent and vegetation surveys were conducted by 137 considering the herbaceous layer only. To account for the time lag in vegetation development between high 138 and low level quadrat along the topographic gradient, two complete surveys were conducted per year (end of 139 June and end of July). Then, to characterize the plant community at the scale of each quadrat, the two surveys 140 were combined by listing all species encountered at least once during the two inventories and associating them 141 with the highest Braun-Blanquet abundance/dominance coefficient noted. Finally, to limit observer biases, all 142 surveys were performed by the first author (PJ).

### 143 2.3. Functional trait data

Three morphological traits – specific leaf area (SLA; leaf area per dry mass), plant height at maturity and seed mass – were extracted from the TRY database (Kattge et al., 2011). Within Westoby's plant ecology strategy scheme (Westoby, 1998), these three easily measurable traits, represent three axes of functional trade-offs to explain plant distribution. SLA is related to resource acquisition and conservation, contrasting "conservative" species with long-lived leaves to "acquisitive" species with rapid turnover of plant leaves. Plant height at

maturity is related to competitive ability, contrasting taller dominant-species with a high ability to pre-empt
resources but with high construction and maintenance costs to smaller dominated-species but with low
construction and maintenance costs. Seed mass is related to dispersal ability, contrasting light-seeded species
but with high seedling mortality to large-seeded species but with high long-term seedling survival.

153 For each trait, we calculated a mean value per species after removing all trait values with an error risk > 3, i.e., 154 having a maximum distance of 3 standard deviations from at least one of the mean trait values, and applied a 155 log<sub>10</sub>-transformation to comply with normality and to limit extreme values. Then, within each quadrat, we 156 computed (i) the community-weighted means (CWMs) and (ii) the functional dispersion (FDis) for each of the 157 three traits (FD package, Laliberté et al., 2014). CWM is defined as the mean of the trait values weighted by the 158 relative cover of each species bearing each value (Lavorel et al., 2008) and was used as a measure of functional 159 composition. FDis is defined as the mean distance of individual species to the weighted centroid of all species 160 in the assemblage (Laliberté and Legendre, 2010) and was used as a measure of functional diversity. Since the 161 trees/shrubs were present only at the seedling stage and the height values of these species strongly influenced the calculation of CWM for this trait, we arbitrarily set a height value of 0.10 m for all phanerophyte species (n 162 163 = 18, see Appendix S1 for details).

In addition, we studied the response of annual species, i.e., therophytes based on plant life-forms (Raunkiaer,
1934) extracted from the Baseflor database (Julve, 1998), which constitute communities often with high
conservation stakes on gravel bars (e.g., Rottenborn et al., 2018), and exotic species, based on a regional
hierarchical list of invasive species (Debay et al., 2020), which represent potential threats to native species
(Richardson et al., 2007) (see detailed information for each species in Appendix S1).

#### 169 **2.4. Environmental variables**

170 At the center of each quadrat, the elevation difference and distance to the water line were characterized using 171 topographical surveys. Topographic measurements were conducted using a laser rangefinder (TruPulse 200X) 172 positioned on a tripod at a constant height of 1.2 m and a target at a height of 2.0 m. The horizontal and 173 vertical distances between the tripod and the target were measured between the water line at the beginning of each transect and the center of the four successive quadrats. For soil properties, the proportion of subsoil 174 175 fine sediment was derived from two soil core samples (30-cm thickness) collected in representative areas 176 within each quadrat and pooled together. Each composite soil sample was then dried, weighed and sieved in the laboratory to estimate the proportion of fine sediments (i.e., <250-µm mesh). Moreover, fine sediment 177

thickness was estimated at the center of each quadrat using a soil corer inserted into the sediment up to the
interface with the gravel, and measuring the distance from the surface. All of these measurements were made
on each gravel bar in 2017 and 2019 during the survey carried out in June (for illustration of the variations in
topographic and pedological variables, within and between gravel bars, see Appendix S2).

For hydrological variables, the water elevation (m) of the Rhône was calculated for each transect using a 1D 182 183 hydrodynamic model based on the 1D shallow water equations (for details, see Dugué et al., 2015; Launay et al., 2019). This numerical model covers the entire length of the French Rhône, including its main tributaries and 184 185 operation rules of hydropower structures. It has been calibrated and validated on each reach using longitudinal water profiles, provided by the Compagnie Nationale du Rhône, over a wide range of discharges for frequent 186 187 hydrological events. The model correctly reproduces discharge time-series, especially during floods. For low 188 flow periods, results are not as accurate due to hydropower scheme subdaily operations that are not included 189 in the model. The output of the model provides a detailed elevation of the water, due to the real geometry of 190 the river bed as a series of cross-sections every 200-500 meters and the high temporal resolution of the hourly 191 results at any point in the model. Combined with centimetric measures of the quadrat elevation, we were able 192 to calculate two hydrological variables: the inundation duration, i.e., the sum of hourly measurements below 193 the water level expressed in days, and the inundation intensity, i.e., the highest water level expressed in 194 meters. For each quadrat, these variables were calculated considering a time extension from September 1 to 195 August 31 for the 2016/2017 and 2018/2019 periods (Appendix S3).

#### 196 **2.6. Statistical analysis**

197 The continuous independent variables were the elevation difference (denoted "Elevation" in tables and figures) 198 and distance to water line ("Distance"), the proportion ("Prop Sedim") and thickness ("Thick Sedim") of fine 199 sediments, the duration ("Dura\_Inund") and intensity ("Inte\_Inund") of inundation (Appendix S4). The 200 independent factors were the year of sampling ("Year", two-level factor), the gravel bar ("GB", eight-level 201 factor) and maintenance measures ("Manag", two-level factor). The skewed variables "Thick\_Sedim" and 202 "Dura\_Inund" were log<sub>10</sub>-transformed while the "Prop\_Sedim" (i.e., proportional data) was logit transformed. Finally, all continuous variables were standardized before performing the analyses. 203 204 To determine whether environmental conditions, taxonomic diversity, i.e., the richness of total, annual and

exotic species, and functional diversity, i.e., the functional dispersion (i.e., FDis) of SLA, height and seed trait

values, varied between 2017 and 2019, between gravel bars managed by brush clearing or plowing, as well as

207 with the interaction between both factors, we used two-way ANOVAs (stats package). To determine whether 208 environmental conditions, taxonomic diversity and functional diversity varied over time within each bar we 209 used paired t-test (stats package). We also used paired t-test to determine how inundation duration and 210 intensity varied along the elevation gradient, i.e. in each quadrat position, between years. As multiple comparison tests were performed, we adjusted the p-values with the Benjamini and Hochberg correction. 211 212 To determine how environmental conditions explained differences in taxonomic and functional diversity 213 between years, at the community scale, we used a modeling approach. We considered a set of 12 a priori 214 models, testing additive and interactive effects between the six environmental variables and year, plus a null 215 model (Appendix S5). Since observations within each bar were not independent, we used mixed models with 216 "GB" as a random effect (Ime4 package, Bates et al., 2015). Also, because maintenance measures influenced significantly the taxonomic diversity, we included "Manag" as a covariate in all a priori models. We then fitted 217 218 mixed models with a Poisson distribution to study variations in species richness and with a Normal distribution 219 to study variations in trait dispersion. To identify the most parsimonious model, we used Akaike's information criterion corrected for small sample sizes and performed model averaging, based on all a priori models, to 220 221 estimate the parameters and associated unconditional standard errors (MuMIn package, Barton, 2015). 222 To determine whether the taxonomic composition and functional composition varied between 2017 and 2019, 223 between gravel bars managed by brush clearing or plowing, as well as with the interaction between both 224 factors, we used two-way PERMANOVAs (Anderson and Walsh, 2013) with 999 permutations (vegan package, 225 Oksanen et al., 2013). For the taxonomic composition, we used a Hellinger distance matrix based on the cover 226 of each species in each of the quadrats, while for the functional composition, we used a Euclidean distance 227 matrix based on the mean values of each trait (i.e., CWM) in each of the quadrats. As multiple comparison tests 228 were performed, we adjusted the p-values with the Benjamini and Hochberg correction. To determine whether 229 the taxonomic composition and functional composition varied over time within each bar we used one-way PERMANOVAs and adjusted the p-values with the Benjamini and Hochberg correction. 230 231 To determine how variations in environmental conditions explained differences in taxonomic and functional

composition, at the community scale, we used canonical analysis of principal coordinates (CAP, Anderson and
Willis, 2003), with 999 permutations (vegan package, Oksanen et al., 2013). We then calculated the marginal
contribution of all independent variables to total constrained inertia and tested for their individual significance
(after all other variables were partialled out).

Analyses were performed with R version 3.5.1 (R Core Team, 2020) (see a summary of the statistical analyses
performed in Appendix S6).

238

239 3. RESULTS

#### 240 **3.1. Changes in environmental conditions between years**

241 Two-way ANOVAs showed that, on average, inundation intensity within quadrats was significantly lower in

242 2019 than in 2017, and that this effect was more pronounced on gravel bars managed by plowing, i.e. a

significant interaction effect (Table 1). In addition, compared to quadrats located on gravel bars managed by

244 brush clearing, quadrats on bars managed by plowing were at a shorter average distance and higher average

- elevation from the water line. In the same quadrats, i.e. managed by plowing, the average thickness of fine
- sediments, as well as the average duration of inundation, were significantly lower.
- Paired *t*-test showed that within gravel bars, the elevation increased significantly between 2017 and 2019 on

GB3 but decreased on GB7 and GB8, while the distance to water increased on GB5 but decreased on GB8. The

- 249 proportion of fine sediments increased significantly between years on GB2, GB4 and GB8 but decreased on
- 250 GB3, while the thickness of fine sediments decreased only on GB7. Finally, inundation intensity decreased
- significantly between years on all bars, except on GB1 where it increased, while inundation duration increased
- on GB1 and GB8 but decreased on GB3, GB5, GB6 and GB7 (Figure 2 & detailed results in Appendix S7).

253 However, while inundations were more intense in 2017 regardless of quadrat position along the elevation

254 gradient, they were longer in 2019 on low-level quadrats composed of coarse-grained sediments (Appendix

255 S8).

#### **3.2. Changes in taxonomic and functional diversity between years**

A total of 205 species (mean ±SD = 21 ±12) were recorded in the 192 quadrats along the Rhône River. Among
 these, 38 were exotic and 87 were annual.

Two-way ANOVAs showed that, on average, the richness of total, annual and exotic species within the quadrats decreased significantly between 2017 and 2019, while for functional diversity, trait dispersion values (i.e., FDis) remained stable (Table 1). In addition, the average richness of total, annual and exotic species, as well as the functional dispersion values of seed mass within the quadrats decreased significantly on gravel bars managed by brush clearing as compared to gravel bars managed by plowing. However, none of the measures of

taxonomic and functional diversity was significantly influenced by the interaction between the years andmanagement factors.

266 Paired *t*-test showed that within gravel bars, species richness decreased significantly between 2017 and 2019

267 on GB1, GB2, GB6 and GB8; functional dispersion values for plant height, seed mass and specific leaf area

268 decreased significantly between years on GB2; functional dispersion value for plant height increased

significantly between years on GB8 (Figure 3 & detailed results in Appendix S9).

#### 270 3.3. Changes in taxonomic and functional diversity with environmental conditions

271 Mixed models showed that models including pedological variables ranked first for the richness of annual and exotic species and the functional dispersion values (i.e., FDis) of height and seed mass values, while models 272 273 including hydrological variables ranked first for total richness and the functional dispersion value of SLA (Table 274 2 & detailed results in Appendix S10). Model averaging revealed that total, annual and exotic species richness, 275 as well as the functional dispersion value of seed mass were higher on bars maintained by plowing than on bars 276 maintained by brush clearing (Table 3). Also, the proportion and thickness of fine sediments had a significant 277 negative effect on species richness and the functional dispersion values of SLA and height. However, for species 278 richness, the interaction between fine sediments proportion and years was significant, revealing a large 279 variation in richness over time on the surfaces made up of coarse-grained sediments (see interaction plots in 280 Appendix S11). Total and exotic species richness, as well as the functional dispersion value of SLA, also 281 decreased with increasing elevation, while all richness measures varied more between years on the surfaces 282 closest to water. Conversely, total species richness and the functional dispersion value of SLA increased on 283 more intensely inundated surfaces, while all richness measures as well as the functional dispersion value of SLA 284 increased with inundation duration in 2017 but decreased in 2019.

# 285 **3.4. Changes in taxonomic and functional composition between years**

Two-way PERMANOVAs showed that the taxonomic composition differed significantly between 2017 and 2019 ( $F_{1,188} = 2.880, p = 0.002$ ), while the functional composition (i.e., CWM) did not ( $F_{1,188} = 1.201, p = 0.297$ ) (Figure 4). In addition, the taxonomic composition differed significantly between gravel bars managed by brush clearing and gravel bars managed by plowing ( $F_{1,188} = 15.395, p = 0.001$ ), while the functional composition did not ( $F_{1,188} = 2.573, p = 0.106$ ). However, both the taxonomic ( $F_{1,188} = 1.348, p = 0.130$ ) and the functional ( $F_{1,188} = 1.016, p = 0.343$ ) composition were not significantly influenced by the interaction between the years and management factors.

Within gravel bars, one-way PERMANOVAS showed that the taxonomic composition changed between 2017 and 2019 on GB7 ( $F_{1,22} = 5.032$ , *p*-adjust = 0.004) and GB8 ( $F_{1,22} = 5.027$ , *p*-adjust = 0.004), while the functional composition changed only on GB7 ( $F_{1,22} = 17.789$ , *p*-adjust = 0.016).

#### 296 **3.5. Changes in taxonomic and functional composition with environmental conditions**

297 Canonical analysis of principal coordinates showed that 16.8% ( $F_{6,185}$  = 7.415, p = 0.001) of the variation in 298 taxonomic composition and 22.9% ( $F_{6,185}$  = 10.475, p = 0.001) of the variation in functional composition were 299 explained by environmental variables (Figure 5). For the taxonomic composition, the first CAP axis was 300 positively related to elevation level and distance to water, which respectively explained 33.6% (p = 0.001) and 301 10.1% (p = 0.001) of the constrained inertia, and negatively related to the duration (8.4%, p = 0.001) and 302 intensity (12.2%, p = 0.001) of inundations. The second CAP axis was negatively related to the proportion 303 (20.6%, p = 0.001) and thickness (14.9%, p = 0.001) of fine sediments. For the functional composition, the first 304 CAP axis was positively related to elevation level (26.8%, p = 0.001) and distance to water (2.2%, p = 0.001), but 305 negatively to the duration (6.0%, p = 0.001) and intensity (1.7%, p = 0.008) of inundations. The second CAP axis 306 was positively related to the proportion (22.6%, p = 0.001) and thickness (40.7%, p = 0.001) of fine sediments. 307

## 308 4. DISCUSSION

309 Understanding the interactions between fluvial landforms and vegetation dynamics is essential for success in 310 restoring biodiversity and functionality to riparian communities on corridors heavily modified by human 311 activities (Palmer and Ruhi, 2019). By studying variations of abiotic and biotic conditions at the corridor and 312 gravel bar scales along the highly modified Rhône River between two seasons in 2017 and 2019, we showed 313 that (i) environmental conditions varied a little from year to year; (ii) taxonomic measures clearly tracked 314 environmental changes; (iii) functional measures were stable over time. Specifically, we showed that variations 315 in richness and composition of herbaceous plants between the two years and sites were largely limited to 316 species sharing the same trait values, underscoring the functional resilience of riparian communities to 317 environmental changes. Also, we highlighted that the strongest variations in taxonomic diversity between the years 2017 and 2019 were spatially restricted to low elevation surfaces, made up of coarse-grained sediments 318 319 and more frequently flooded. Beyond being the most vulnerable, riparian communities established under these 320 conditions were also the most taxonomically and functionally diverse. To improve both the biodiversity and

321 functionality of riparian herbaceous communities along the Rhône River, the conservation and restorations of 322 these gravel bar habitats should be a priority.

323 4.1. River regulation and bank stabilization lead to relatively stable riparian habitats

324 At the corridor scale there was no variation in topographic and pedological conditions between the two years. 325 Within the gravel bars, changes of low magnitudes were found between 2017 and 2019, mostly limited to 326 variations in the proportion of fine sediments. Although it cannot be ruled out that these small variations are 327 the results of methodological biases in the field measurements, this tends to confirm that hydro-geomorphic 328 processes along the Rhône River are limited to fine-grained sediments (Tena et al., 2020; Vauclin et al., 2020). Conversely, even if 2017 and 2019 were "calm" hydrological years, i.e., none of the peak flows recorded 329 330 exceeded the two-year return flood of the Rhône, at the corridor scale, a significant decrease in the intensity of 331 inundation could still be measured between the two years. Within the gravel bars and with the exception of 332 GB1, our results confirmed that inundations were more intense in 2017 but that their duration were either 333 longer or shorter depending on the site. However, additional analyses of floods duration showed heterogeneous variations along the elevation gradient and with the extent of fine sediment deposits (Appendix 334 335 S8). Thus, while mid- and high-level quadrats were inundated on average one day longer in 2017, low-level 336 quadrats, especially if they were composed of coarse-grained sediments, were inundated up to 20 days longer 337 in 2019. This shows that in this regulated and channelized river system, where the bypassed reaches receive 338 only a minimum baseflow (i.e., the residual flow is 10-30% of the mean annual flow arriving at the upstream 339 diversion dams) and are subject to a sharp reduction in the frequency and magnitude of peak flows (Vázquez-340 Tarrío et al., 2019), environmental factors indirectly related to the characteristics of the gravel bars are the 341 main sources of variation that can shape riparian communities. Moreover, this tends to confirm that water and 342 sediment exchanges, as well as scouring, in the absence of important floods, as was the case during this short 343 study period, are very limited in the bypassed reaches (Vázquez-Tarrío et al., 2019) and do not allow, under 344 current hydrological conditions, the creation or renewal of riparian habitats in the Rhône riverbed. 345 4.2. Taxonomic diversity and composition respond to environmental changes in space and time Species richness decreased between 2017 and 2019 at the corridor scale and followed a similar trend within 346 347 each of the bar. Specifically, we showed that maintenance measures, which are confused with the longitudinal 348 and climatic gradient, strongly influenced richness, i.e., northern bars maintained by brush clearing were species poor than southern bars maintained by plowing. By disturbing the soil surface, plowing artificially

13

350 increases microtopographic variations on bars (sensu, Pollock et al., 1998), allowing greater local co-occurrence 351 of species with contrasting habitat requirements (Janssen et al., 2019). After controlling for this confounding 352 spatial effect, which was constant between the two years studied, we demonstrated that temporal changes in 353 richness were independent of management practices and involved communities established under specific 354 environmental conditions. Indeed, plant richness varied more between years at low elevations, on more 355 frequently flooded surfaces consisting of coarse-grained sediments. Since flooding on the Rhone was shown to 356 be more intense in 2017 regardless of quadrat position along the elevation gradient, but longer in 2019 at the 357 low-level quadrats composed of coarse-grained sediments, we infer that these variations in hydrological 358 conditions were the primary cause of species loss. Indeed, it was shown that flooded areas received a more 359 diverse pool of colonizing species through hydrochory (Gurnell et al., 2008; Jansson et al., 2005) but also that 360 intermediate-intensity floods supported high species diversity (Pollock et al., 1998; Renofalt et al., 2005). 361 However, over the two years studied, the magnitude of flooding varied within a narrow range, and the peak 362 flows recorded cannot be characterized as intermediate disturbances (Appendix S3). At the same time, it has been shown that frequently flooded sites with low spatial variation in flood frequencies are species-poor 363 364 (Pollock et al., 1998) but also that in dam-regulated rivers, long unnatural inundations have a negatively effect 365 on species richness (e.g., Su et al., 2020; Zheng et al., 2021). On the Rhone, the spatial structuring of 366 inundations, longer in 2019 at low levels on pebble-dominated bars, and of species richness, lower in 2019 in 367 low-lying quadrats composed of coarse-grained sediments, supports these previous finding. Specifically, our 368 results suggest that a prolonged submergence of about 20 days can induce a strong decrease in the taxonomic 369 diversity of riparian herbaceous communities and highlight, in this highly regulated river system, the greater 370 vulnerability of plant species established on bar surfaces made up of coarse-grained sediments to variations in 371 hydrological conditions. Since along the gravel bars of the Rhone, as in other rivers (Kalníková et al., 2018; 372 Škornik et al., 2017), species richness is highest in these riparian habitats, we believe that their conservation 373 should be a priority (Gilvear et al., 2008; Rottenborn et al., 2018; Zeng et al., 2015). 374 Species composition also varied between the two years studied at the corridor scale. Within bars, changes in composition between years were however restricted to only two sites, which may be due to the small temporal 375 376 extent of this study but also to management practices that periodically reset succession. Furthermore,

377 community-wide analyses revealed that compositional changes were primarily explained by bar characteristics.

378 Indeed, elevation difference explained most of the variation (Gilvear et al., 2008; Kalníková et al., 2018; Škornik

379 et al., 2017), with a shift from flood-tolerant species at low elevation (e.g., Phalaris arundinacea and 380 Phragmites australis) to drought-tolerant species at high elevation (e.g., Lolium perenne and Elymus repens). 381 Also, the proportion of fine sediments on the bars profoundly influenced species composition (Gilvear et al., 382 2008; Kalníková et al., 2018). Thus, at low elevation, Phalaris arundinacea cover increased when there was little 383 fine sediment while Phragmites australis cover increased when there was a lot of fine sediment, and, at high 384 elevation, Lolium perenne cover increased when there was little fine sediment while Elymus repens cover 385 increased when there was a lot of fine sediment. Beyond bar characteristics, hydrological conditions 386 during the year preceding observations were of little inference value to explain changes in herbaceous plant 387 composition in our study system. This is likely related to the fact that the Rhone floods were of too low 388 intensity in 2017 and 2019 to trigger the scouring process necessary to initiate a new succession (Steiger et al., 389 2005). Overall, these results show that, even in a regulated river system, the taxonomic composition of riparian 390 habitats remained under the influence of multiple environmental gradients, acting at both the local and 391 regional scales (Kuglerová et al., 2015; Renofalt et al., 2005).

#### **4.3.** Traits diversity and functional composition remain stable in space and time

393 In contrast to species richness, functional dispersion values remained stable over time at the corridor and 394 gravel bar scales, with the exception of GB2. This suggests that the herbaceous species that disappeared 395 between 2017 and 2019 on the Rhone gravel bars had redundant trait values in the community, showing some 396 functional resilience of this regulated system to environmental changes (Su et al., 2020). Also, trait dispersion 397 for specific leaf area (SLA) was higher on GB1, which is located in the only bypassed reach where bedload 398 transport is still active (Vázquez-Tarrío et al., 2019). Greater co-occurrence of species with different resource 399 acquisition and conservation strategies on "active" bars may confirm that flood disturbance increases 400 functional diversity (Biswas and Mallik, 2010; Lawson et al., 2015) and suggests the use of leaf traits as 401 potential indicators of riparian habitat dynamics. Conversely, the functional dispersion value for seed mass was 402 strictly influenced by maintenance measures. Indeed, plowing by increasing microtopographic heterogeneity, 403 allows a variation in flood frequencies and velocity conditions (Pollock et al., 1998), thus the deposition of light and heavy seeds on restricted surfaces. Controlling for this spatial effect, community-wide analyses confirmed 404 405 that trait dispersion was nearly unaffected by changes in environmental conditions between the two years 406 studied. However, functional diversity peaked under the same environmental conditions as taxonomic diversity. Specifically, we showed that bars covered by a thick layer of fine sediments supported species with a 407

408 narrow range of trait values (see also, Janssen et al., 2019). Thus, even for fixed bars in a highly regulated river
409 system, as is the case of the Rhône, the co-occurrence of herbaceous species with different functional
410 strategies was greater when they consisted of coarse sediments (Janssen et al., 2020a). This is particularly true
411 at low elevations, on surfaces more intensely flooded (Lawson et al., 2015).

Functional composition did not vary between years at the corridor scale. Within bars, changes in trait 412 413 composition between 2017 and 2019 was restricted to one site. This confirms the stability of functional 414 assemblages in space and time, probably due to the low amplitude of hydrological conditions over the two 415 years studied on the Rhone River, and emphasizes that changes in species identity between gravel bars along the Rhône River do not result in changes in trait values. Regarding the influence of environmental conditions, 416 417 soil factors explained up to 60% of the variation in the functional composition of riparian herbaceous 418 communities. Specifically, we showed that accumulation of fine sediment on gravel bars resulted in an increase 419 in mean plant height (Kyle and Leishman, 2009). Indeed, silt deposits, by promoting fertile environments 420 (Asaeda and Rashid, 2012), favoured the development of taller dominant-species capable of pre-empting 421 resources, such as Phragmites australis (Minchinton and Bertness, 2003) and Phalaris arundinacea (Chen et al., 422 2017). Moreover, as these taller species are more typical of lentic aquatic ecosystems, the high cover measured 423 on the bars of the Rhone affected by fine sediment deposits may also reflect their adaptation to the fairly 424 constant minimum flow that is observed most of the year in the bypassed sections. In addition to soil texture, 425 elevation level also explained much of the functional composition, with an increase in mean seed mass from 426 low to high elevation (Kyle and Leishman, 2009; McCoy-Sulentic et al., 2017). This tends to confirm that the 427 colonization potential of light-seeded species (e.g., Artemisia annua, Juncus articulates, Veronica anagallis-428 aquatica) is higher in more frequently disturbed environments (Gurnell et al., 2008), this even in a highly 429 regulated river system. Beside bar characteristics and certainly due to the low magnitude of inundations in the 430 months preceding the field observations in 2017 and 2019, hydrological conditions did not explain much of the functional composition of communities. Overall, we showed that at the corridor scale, high species turnover 431 432 along the longitudinal-climatic gradient did not lead to a shift in the functional composition of riparian 433 communities, whereas at the gravel bar scale, trait-environment relationships varied strongly and consistently 434 with local gradients in elevation and soil texture. This underlines, at least for highly regulated river systems and 435 for the range of hydrological conditions studied, the greater relative importance of local processes in 436 controlling the functioning of riparian communities.

437

#### 438 5. CONCLUSION

439 By studying spatial and temporal variations in riparian herbaceous communities along the heavily modified 440 Rhône River, we showed (1) that different gravel bars supported different species pools, which shared almost 441 the same set of traits, and (2) that changes in richness over the two-year period were restricted to species with 442 redundant trait values in the communities, suggesting some functional resilience to environmental changes. Furthermore, by investigating the relative influence of different environmental factors on assemblage patterns, 443 444 we showed that the temporal change in taxonomic diversity was spatially restricted to communities established on low-lying bar surfaces composed of coarse-grained sediments, which were more intensely and 445 446 frequently flooded. In addition to being the most sensitive to environmental changes, the riparian communities 447 established on these surfaces were also the most diverse in species and trait values, highlighting the 448 importance of these habitats for the Rhone river biodiversity and functionality. However, in this regulated river 449 system where water and sediment exchanges are very limited, we showed that fine sediment deposition 450 threatens the functional and ecological integrity of gravel bars by inducing a strong decrease in species and 451 trait diversity. Although consistent and supported by other studies on different systems, these results are still 452 limited to a short observation period and concerns an intensively managed river system, where a lagged 453 response of the vegetation to environmental changes cannot be excluded. To better disentangle the relative 454 influence of the many environmental parameters that condition riparian community dynamics, longer, 455 replicated observational studies are needed. Finally, from an applied perspective and in order to improve the 456 biodiversity and functionality of riparian habitats in regulated rivers, our results highlight the urgent 457 importance of conserving still-functional gravel bars and promoting more morphologically complex riparian 458 habitats. This could be achieved by developing management plans, at the corridor and reach scales, aiming at 459 renaturalize the flow regime (e.g., by increasing and irregularizing baseflows, Lawson et al., 2015) and reactivate bedload supply and transport (e.g., through gravel augmentation, Staentzel et al., 2020) to maximize 460 461 scouring and prevent fine sediment deposits. Applied to restoration operations aimed at improving lateral 462 connectivity through the removals of dykes, our results suggest that it would be more beneficial for biodiversity 463 to profile banks with a gentle slope and to favor materials with a predominance of coarse-grained sediments. 464

465 ACKNOWLEDGEMENTS

466 We thank Gilles Favier, Delphine Jaymond and Sophie Labonne for help in the field as well as Laura Troudet and 467 Jérôme Le Coz from INRAE for extracting data from the hydrodynamic model. This work was cofunded by the 468 Labex DRIIHM, French program "Investissements d'Avenir" (ANR-11-LABX-0010), which is managed by the 469 French National Research Agency (ANR), the OHM Vallée du Rhône, the Agence de l'eau Rhône Méditerranée 470 Corse and INRAE. It was also conducted within the Rhône Sediment Observatory (OSR), a multi-partner 471 research program funded through the Plan Rhône by the European Regional Development Fund (contract: 472 CNRS nº144140), Agence de l'eau RMC (France, contract: 2018 1063), CNR, EDF and three regional councils 473 (Auvergne-Rhône-Alpes, PACA and Occitanie). This work was performed within the framework of the EUR H2O'Lyon (ANR-17-EURE-0018) of Université de Lyon, part of the program "Investissements d'Avenir" (ANR-11-474 475 IDEX-0007) operated by the ANR. 476 477 DATA AVAILABILITY STATEMENT 478 Data are available from the authors upon reasonable request. 479 480 REFERENCES 481 Aguiar, F.C., Segurado, P., Martins, M.J., Bejarano, M.D., Nilsson, C., Portela, M.M., Merritt, D.M., 2018. The 482 abundance and distribution of guilds of riparian woody plants change in response to land use and flow 483 regulation. J. Appl. Ecol. 55, 2227–2240. https://doi.org/10.1111/1365-2664.13110 484 Allan, J.D., Castillo, M.M., 2007. Stream ecology: structure and function of running waters, Second Edition. 485 Springer, Dordrecht, the Netherlands. 486 Anderson, M.J., Walsh, D.C., 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous 487 dispersions: What null hypothesis are you testing? Ecol. Monogr. 83, 557–574. 488 Anderson, M.J., Willis, T.J., 2003. Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. Ecology 84, 511–525. 489 490 Asaeda, T., Rashid, M.H., 2012. The impacts of sediment released from dams on downstream sediment bar vegetation. J. Hydrol. 430-431, 25-38. https://doi.org/10.1016/j.jhydrol.2012.01.040 491 492 Barton, K., 2015. MuMIn: Multi-Model Inference. R package version 1.13.4.

- 493 Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using Ime4. J. Stat.
- 494 Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

495	Bejarano, M.D., Nilsson, C., Aguiar, F.C., 2018. Riparian plant guilds become simpler and most likely fewer
496	following flow regulation. J. Appl. Ecol. 55, 365–376. https://doi.org/10.1111/1365-2664.12949
497	Biswas, S.R., Mallik, A.U., 2010. Disturbance effects on species diversity and functional diversity in riparian and
498	upland plant communities. Ecology 91, 28–35.
499	Braun-Blanquet, J., 1932. Plant sociology. The study of plant communities, McGraw-Hill Book Co. ed. NewYork,
500	NY, USA.
501	Bravard, JP., Gaydou, P., 2015. Historical development and integrated management of the Rhône River
502	floodplain, from the Alps to the Camargue delta, France, in: Hudson, P.F., Middelkoop, H. (Eds.),
503	Geomorphic Approaches to Integrated Floodplain Management of Lowland Fluvial Systems in North
504	America and Europe. Springer New York, New York, NY, pp. 289–320. https://doi.org/10.1007/978-1-
505	4939-2380-9_12
506	Catford, J.A., Downes, B.J., Gippel, C.J., Vesk, P.A., 2011. Flow regulation reduces native plant cover and
507	facilitates exotic invasion in riparian wetlands. J. Appl. Ecol. 48, 432–442.
508	https://doi.org/10.1111/j.1365-2664.2010.01945.x
509	Chen, X., Liao, Y., Xie, Y., Wu, C., Li, F., Deng, Z., Li, X., 2017. The combined effects of sediment accretion (burial)
510	and nutrient enrichment on the growth and propagation of Phalaris arundinacea. Sci. Rep. 7, 39963.
511	https://doi.org/10.1038/srep39963
512	Debay, P., Legland, T., Pache, G., 2020. Liste actualisée et hiérarchisée des espèces exotiques envahissantes,
513	bilan de la problématique végétale invasive en Rhône-Alpes. Conservatoire botanique national alpin,
514	Gap, France.
515	Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, ZI., Knowler, D.J., Lévêque, C., Naiman, R.J., Prieur-
516	Richard, AH., Soto, D., Stiassny, M.L.J., Sullivan, C.A., 2006. Freshwater biodiversity: importance,
517	threats, status and conservation challenges. Biol. Rev. 81, 163–182.
518	https://doi.org/10.1017/S1464793105006950
519	Dugué, V., Walter, C., Andries, E., Launay, M., Coz, J.L., Camenen, B., Faure, JB., 2015. Accounting for
520	hydropower schemes' operation rules in the 1D hydrodynamic modeling of the Rhône River from lake
521	Geneva to the Mediterranean sea, in: E-Proceedings of the 36th IAHR World Congress. Presented at
522	the 36th IAHR World Congress, The Hague, the Netherlands, pp. 1–9.

Gilvear, D., Francis, R., Willby, N., Gurnell, A., 2008. 26 Gravel bars: a key habitat of gravel-bed rivers for
vegetation, in: Gravel-Bed Rivers VI: From Process Understanding to River Restoration (H. Habersack,
H. Piégay, M. Rinaldi, Editors), Developments in Earth Surface Processes. Elsevier, Amsterdam, The

526 Netherlands, pp. 677–700. https://doi.org/10.1016/S0928-2025(07)11154-8

527 Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., Babu, S., Borrelli, P., Cheng, L.,

- 528 Crochetiere, H., Ehalt Macedo, H., Filgueiras, R., Goichot, M., Higgins, J., Hogan, Z., Lip, B., McClain,
- 529 M.E., Meng, J., Mulligan, M., Nilsson, C., Olden, J.D., Opperman, J.J., Petry, P., Reidy Liermann, C.,
- 530 Sáenz, L., Salinas-Rodríguez, S., Schelle, P., Schmitt, R.J.P., Snider, J., Tan, F., Tockner, K., Valdujo, P.H.,
- 531 van Soesbergen, A., Zarfl, C., 2019. Mapping the world's free-flowing rivers. Nature 569, 215–221.
- 532 https://doi.org/10.1038/s41586-019-1111-9
- Gurnell, A., Thompson, K., Goodson, J., Moggridge, H., 2008. Propagule deposition along river margins: linking
  hydrology and ecology. J. Ecol. 96, 553–565. https://doi.org/10.1111/j.1365-2745.2008.01358.x
- Janssen, J.A.M., Rodwell, J.S., García Criado, M., Gubbay, S., Haynes, T., Nieto, A., Sanders, N., Landucci, F.,
- 536 Loidi, J., Ssymank, A., Tahvanainen, T., Valderrabano, M., Acosta, A., Aronsson, M., Arts, G., Attorre, F.,
- 537 Bergmeier, E., Bijlsma, R.-J., Bioret, F., Biţă-Nicolae, C., Biurrun, I., Calix, M., Capelo, J., Čarni, A.,
- 538 Chytrý, M., Dengler, J., Dimopoulos, P., Essl, F., Gardfjell, H., Gigante, D., Giusso del Galdo, G., Hájek,
- 539 M., Jansen, F., Jansen, J., Kapfer, J., Mickolajczak, A., Molina, J.A., Molnár, Z., Paternoster, D., Piernik,
- 540 A., Poulin, B., Renaux, B., Schaminée, J.H.J., Šumberová, K., Toivonen, H., Tonteri, T., Tsiripidis, I.,
- 541 Tzonev, R., Valachovič, M., 2016. European red list of habitats. Part 2, Terrestrial and freshwater
- 542 habitats. Publications Office of the European Union, Luxembourg.
- Janssen, P., Piégay, H., Evette, A., 2020a. Fine-grained sediment deposition alters the response of plant CSR
  strategies on the gravel bars of a highly regulated river. Appl. Veg. Sci. 23, 452–463.
- 545 https://doi.org/10.1111/avsc.12494
- Janssen, P., Piégay, H., Pont, B., Evette, A., 2019. How maintenance and restoration measures mediate the
- 547 response of riparian plant functional composition to environmental gradients on channel margins:
- 548 Insights from a highly degraded large river. Sci. Total Environ. 656, 1312–1325.
- 549 https://doi.org/10.1016/j.scitotenv.2018.11.434
- Janssen, P., Stella, J.C., Piégay, H., Räpple, B., Pont, B., Faton, J.-M., Cornelissen, J.H.C., Evette, A., 2020b.
- 551 Divergence of riparian forest composition and functional traits from natural succession along a

- 552 degraded river with multiple stressor legacies. Sci. Total Environ. 721, 137730.
- 553 https://doi.org/10.1016/j.scitotenv.2020.137730
- Jansson, R., Zinko, U., Merritt, D.M., Nilsson, C., 2005. Hydrochory increases riparian plant species richness: a
   comparison between a free-flowing and a regulated river. J. Ecol. 93, 1094–1103.
- 556 https://doi.org/10.1111/j.1365-2745.2005.01057.x
- Johnson, W.C., Dixon, M.D., Scott, M.L., Rabbe, L., Larson, G., Volke, M., Werner, B., 2012. Forty years of
   vegetation change on the Missouri River floodplain. BioScience 62, 123–135.
- 559 https://doi.org/10.1525/bio.2012.62.2.6
- Julve, P., 1998. Baseflor. Index botanique, écologique et chorologique de la flore de France. Version : 2017.
   http://perso.wanadoo.fr/philippe.julve/catminat.htm.
- Junk, W.J., Bayley, P.B., Sparks, R.E., 1989. The flood pulse concept in river-floodplain systems. Can. Spec. Publ.
   Fish. Aquat. Sci. 106, 110–127.
- Kalníková, V., Chytrý, K., Chytrý, M., 2018. Early vegetation succession on gravel bars of Czech Carpathian
   streams. Folia Geobot. 53, 317–332. https://doi.org/10.1007/s12224-018-9323-6
- 566 Kattge, J., Díaz, S., Lavorel, S., Prentice, I.C., Leadley, P., Bönisch, G., Garnier, E., Westoby, M., Reich, P.B.,
- 567 Wright, I.J., Cornelissen, J.H.C., Violle, C., Harrison, S.P., Van Bodegon, P.M., Reichstein, M., Enquist,
- 568 B.J., Soudzilovskaia, N.A., Ackerly, D.D., Anand, M., Atkin, O., Bahn, M., Baker, T.R., Baldocchi, D.,
- 569 Bekker, R., Blanco, C.C., Blonder, B., Bond, W.J., Bradstock, R., Bunker, D.E., Casanoves, F., Cavender-
- 570 Bares, J., Chambers, J.Q., Chapin III, F.S., Chave, J., Coomes, D., Cornwell, W.K., Craine, J.M., Dobrin,
- 571 B.H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W.F., Fang, J., Fernández-Méndez, F.,
- 572 Fidelis, A., Finegan, B., Flores, O., Ford, H., Frank, D., Freschet, G.T., Fyllas, N.M., Gallagher, R.V.,
- 573 Green, W.A., Gutierrez, A.G., Hickler, T., Higgins, S.I., Hodgson, J.G., Jalili, A., Jansen, S., Joly, C.A.,
- 574 Kerkhoff, A.J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J.M.H., Kramer, K., Kühn, I.,
- 575 Kurokawa, H., Laughlin, D., Lee, T.D., Leishman, M., Lens, F., Lenz, T., Lewis, S.L., Lloyd, J., Llusià, J.,
- 576 Louault, F., Ma, S., Mahecha, M.D., Manning, P., Massad, T., Medlyn, B.E., Messier, J., Moles, A.T.,
- 577 Müller, S.C., Nadrowski, K., Naeem, S., Niinemets, U., Nöllert, S., Nüske, A., Ogaya, R., Oleksyn, J.,
- 578 Onipchenko, V.G., Onoda, Y., Ordoñez, J., Overbeck, G., Ozinga, W.A., Patiño, S., Paula, S., Pausas, J.G.,
- 579 Peñuelas, J., Phillips, O.L., Pillar, V., Poorter, H., Poorter, L., Poschlod, P., Prinzing, A., Proulx, R.,
- 580 Rammig, A., Reinsch, S., Reu, B., Sack, L., Salgado-Negret, B., Sardans, J., Shiodera, S., Shipley, B.,

581	Siefert, A., Sosinski, E., Soussana, JF., Swaine, E., Swenson, N., Thompson, K., Thornton, P., Waldram,
582	M., Weiher, E., White, M., White, S., Wright, S.J., Yguel, B., Zaehle, S., Zanne, A.E., Wirth, C., 2011. TRY
583	- a global database of plant traits. Glob. Change Biol. 17, 2905–2935. https://doi.org/10.1111/j.1365-
584	2486.2011.02451.x
585	Kuglerová, L., Jansson, R., Sponseller, R.A., Laudon, H., Malm-Renöfält, B., 2015. Local and regional processes
586	determine plant species richness in a river-network metacommunity. Ecology 96, 381–391.
587	https://doi.org/10.1890/14-0552.1
588	Kyle, G., Leishman, M.R., 2009. Plant functional trait variation in relation to riparian geomorphology: The
589	importance of disturbance. Austral Ecol. 34, 793–804. https://doi.org/10.1111/j.1442-
590	9993.2009.01988.x
591	Laliberté, E., Legendre, P., 2010. A distance-based framework for measuring functional diversity from multiple
592	traits. Ecology 91, 299–305.
593	Laliberté, E., Legendre, P., Shipley, B., 2014. FD: measuring functional diversity from multiple traits, and other
594	tools for functional ecology. R package version 1.0-12.
595	Launay, M., Dugué, V., Faure, JB., Coquery, M., Camenen, B., Le Coz, J., 2019. Numerical modelling of the
596	suspended particulate matter dynamics in a regulated river network. Sci. Total Environ. 665, 591–605.
597	https://doi.org/10.1016/j.scitotenv.2019.02.015
598	Lavorel, S., Grigulis, K., McIntyre, S., Williams, N.S.G., Garden, D., Dorrough, J., Berman, S., Quétier, F.,
599	Thébault, A., Bonis, A., 2008. Assessing functional diversity in the field – methodology matters! Funct.
600	Ecol. 22, 134–147. https://doi.org/10.1111/j.1365-2435.2007.01339.x
601	Lawson, J.R., Fryirs, K.A., Lenz, T., Leishman, M.R., 2015. Heterogeneous flows foster heterogeneous
602	assemblages: relationships between functional diversity and hydrological heterogeneity in riparian
603	plant communities. Freshw. Biol. 60, 2208–2225. https://doi.org/10.1111/fwb.12649
604	McCoy-Sulentic, M.E., Kolb, T.E., Merritt, D.M., Palmquist, E., Ralston, B.E., Sarr, D.A., Shafroth, P.B., 2017.
605	Changes in community-level riparian plant traits over inundation gradients, Colorado River, Grand
606	Canyon. Wetlands 37, 635–646. https://doi.org/10.1007/s13157-017-0895-3
607	Minchinton, T.E., Bertness, M.D., 2003. Disturbance-mediated competition and the spread of Phragmites
608	australis in a coastal marsh. Ecol. Appl. 13, 1400–1416. https://doi.org/10.1890/02-5136
609	Naiman, R.J., Decamps, H., 1997. The ecology of interfaces: riparian zones. Annu. Rev. Ecol. Syst. 28, 621–658.

- Nilsson, C., Reidy, C.A., Dynesius, M., Revenga, C., 2005. Fragmentation and flow regulation of the world's large
  river systems. Science 308, 405–408.
- 612 Oksanen, J., Blanchet, G.F., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P.,
- 613 Stevens, M.H.H., Wagner, H., 2013. vegan: Community Ecology Package. R package version 2.0-10.
- Palmer, M., Ruhi, A., 2019. Linkages between flow regime, biota, and ecosystem processes: Implications for
- 615 river restoration. Science 365, eaaw2087. https://doi.org/10.1126/science.aaw2087
- Poff, N.L., Olden, J.D., Merritt, D.M., Pepin, D.M., 2007. Homogenization of regional river dynamics by dams
  and global biodiversity implications. Proc. Natl. Acad. Sci. 104, 5732–5737.
- 618 Pollock, M.M., Naiman, R.J., Hanley, T.A., 1998. Plant species richness in riparian wetlands A test of
- 619 biodiversity theory. Ecology 79, 94–105. https://doi.org/10.1890/0012-
- 620 9658(1998)079[0094:PSRIRW]2.0.CO;2
- 621 R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical
- 622 Computing, Vienna, Austria.
- Raunkiaer, C., 1934. The life forms of plants and statistical plant geography, Oxford University Press. ed. Oxford
   University Press, Oxford, UK.
- Renofalt, B.M., Nilsson, C., Jansson, R., 2005. Spatial and temporal patterns of species richness in a riparian
  landscape. J. Biogeogr. 32, 2025–2037. https://doi.org/10.1111/j.1365-2699.2005.01328.x
- 627 Richardson, D.M., Holmes, P.M., Esler, K.J., Galatowitsch, S.M., Stromberg, J.C., Kirkman, S.P., Pyšek, P., Hobbs,
- R.J., 2007. Riparian vegetation: degradation, alien plant invasions, and restoration prospects. Divers.
  Distrib. 13, 126–139.
- 630 Riis, T., Kelly-Quinn, M., Aguiar, F.C., Manolaki, P., Bruno, D., Bejarano, M.D., Clerici, N., Fernandes, M.R.,
- 631 Franco, J.C., Pettit, N., Portela, A.P., Tammeorg, O., Tammeorg, P., Rodríguez-González, P.M., Dufour,
- S., 2020. Global overview of ecosystem services provided by riparian vegetation. BioScience 70, 501–
  514. https://doi.org/10.1093/biosci/biaa041
- 634 Rottenborn, J., Vítovcová, K., Prach, K., 2018. Interannual dynamics of a rare vegetation on emerged river
- 635 gravels with special attention to the critically endangered species Corrigiola litoralis L. Folia Geobot.
- 636 53, 213–225. https://doi.org/10.1007/s12224-018-9316-5
- Škornik, S., Meznarič, M., Kaligarič, M., 2017. Factors affecting composition of gravel bar vegetation in middle
   reach of a lowland river. Pol. J. Ecol. 65, 194–210. https://doi.org/10.3161/15052249PJE2017.65.2.003

- Staentzel, C., Kondolf, G.M., Schmitt, L., Combroux, I., Barillier, A., Beisel, J.-N., 2020. Restoring fluvial forms
   and processes by gravel augmentation or bank erosion below dams: A systematic review of ecological
- 641 responses. Sci. Total Environ. 706, 135743. https://doi.org/10.1016/j.scitotenv.2019.135743
- 642 Steiger, J., Tabacchi, E., Dufour, S., Corenblit, D., Peiry, J.-L., 2005. Hydrogeomorphic processes affecting
- 643 riparian habitat within alluvial channel-floodplain river systems: a review for the temperate zone.

644 River Res. Appl. 21, 719–737. https://doi.org/10.1002/rra.879

- 645 Su, X., Bejarano, M.D., Yi, X., Lin, F., Ayi, Q., Zeng, B., 2020. Unnatural flooding alters the functional diversity of
- 646 riparian vegetation of the Three Gorges Reservoir. Freshw. Biol. 65, 1585–1595.

647 https://doi.org/10.1111/fwb.13523

- Tabacchi, E., González, E., Corenblit, D., Garófano-Gómez, V., Planty-Tabacchi, A., Steiger, J., 2019. Species
- 649 composition and plant traits: Characterization of the biogeomorphological succession within

650 contrasting river corridors. River Res. Appl. 35, 1228–1240. https://doi.org/10.1002/rra.3511

Tena, A., Piégay, H., Seignemartin, G., Barra, A., Berger, J.F., Mourier, B., Winiarski, T., 2020. Cumulative effects
 of channel correction and regulation on floodplain terrestrialisation patterns and connectivity.

653 Geomorphology 354, 107034. https://doi.org/10.1016/j.geomorph.2020.107034

- 554 Tonkin, J.D., Merritt, David.M., Olden, J.D., Reynolds, L.V., Lytle, D.A., 2018. Flow regime alteration degrades
- ecological networks in riparian ecosystems. Nat. Ecol. Evol. 2, 86–93. https://doi.org/10.1038/s41559017-0379-0
- Vauclin, S., Mourier, B., Tena, A., Piégay, H., Winiarski, T., 2020. Effects of river infrastructures on the floodplain
   sedimentary environment in the Rhône River. J. Soils Sediments 20, 2697–2708.

659 https://doi.org/10.1007/s11368-019-02449-6

660 Vázquez-Tarrío, D., Tal, M., Camenen, B., Piégay, H., 2019. Effects of continuous embankments and successive

- run-of-the-river dams on bedload transport capacities along the Rhône River, France. Sci. Total
  Environ. 658, 1375–1389. https://doi.org/10.1016/j.scitotenv.2018.12.109
- 663 Ward, J.V., 1989. The four-dimensional nature of lotic ecosystems. J. North Am. Benthol. Soc. 8, 2–8.

664 https://doi.org/10.2307/1467397

665 Westoby, M., 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. Plant Soil 199, 213–227.

- 666 Wohl, E., Bledsoe, B.P., Jacobson, R.B., Poff, N.L., Rathburn, S.L., Walters, D.M., Wilcox, A.C., 2015. The natural
- 667 sediment regime in rivers: broadening the foundation for ecosystem management. BioScience 65,
- 668 358–371. https://doi.org/10.1093/biosci/biv002
- Zeng, Q., Shi, L., Wen, L., Chen, J., Duo, H., Lei, G., 2015. Gravel bars can be critical for biodiversity
- 670 conservation: A case study on scaly-sided Merganser in south China. PLOS ONE 10, e0127387.
- 671 https://doi.org/10.1371/journal.pone.0127387
- 572 Zheng, J., Arif, M., Zhang, S., Yuan, Z., Zhang, L., Li, J., Ding, D., Li, C., 2021. Dam inundation simplifies the plant
- 673 community composition. Sci. Total Environ. 801, 149827.
- 674 https://doi.org/10.1016/j.scitotenv.2021.149827

**Table 1.** Variations in environmental conditions, taxonomic and functional diversity between years and maintenance measures, as well as the interaction between both

677 factors, along the Rhône River, France (*F*-value<sub>(3,188)</sub> and *p*-value based on two-way ANOVAs, SD = standard deviation, FDis = Functional Dispersion, SLA = Specific Leaf Area)

Variables	2017 : m	ean (±SD)	2019 : me	Y	ear	Management		Interaction		
variables	Clearing	Plowing	Clearing	Plowing	F	<i>p</i> -value	F	<i>p</i> -value	F	<i>p</i> -value
Environmental conditions										
Elevation above water line (m)	1.14 (±0.78)	1.00 (±0.72)	1.20 (±0.87)	0.88 (±0.74)	0.086	0.770	4.249	0.041	0.667	0.415
Distance to water line (m)	28.43 (±21.06)	34.92 (±25.84)	28.56 (±20.60)	35.54 (±25.98)	0.012	0.912	3.941	0.049	0.005	0.942
Proportion of fine sediments (%)	43.20 (±34.73)	37.16 (±34.79)	39.06 (±37.61)	32.95 (±31.24)	0.014	0.906	0.607	0.437	0.185	0.667
Thickness of fine sediments (cm)	22.81 (±27.11)	18.14 (±27.35)	28.35 (±32.56)	18.36 (±26.86)	0.005	0.946	7.404	0.007	0.254	0.615
Inundation duration (days)	15.20 (±12.69)	11.48 (±9.30)	19.26 (±22.60)	10.20 (±11.81)	1.400	0.238	9.426	0.002	1.038	0.310
Inundation intensity (m)	2.11 (±0.89)	2.97 (±1.81)	1.93 (±0.94)	1.74 (±0.92)	16.546	0.000	3.664	0.057	9.106	0.003
Taxonomic diversity										
Total species richness	18.35 (±9.12)	31.29 (±11.61)	11.96 (±4.92)	25.00 (±10.95)	21.336	0.000	89.456	0.000	0.001	0.970
Richness of annual species	6.19 (±5.83)	14.92 (±6.09)	3.69 (±2.49)	11.71 (±6.04)	13.758	0.000	118.459	0.000	0.212	0.646
Richness of exotic species	4.21 (±3.38)	8.94 (±3.45)	2.67 (±1.74)	7.54 (±3.91)	9.956	0.002	106.425	0.000	0.025	0.876
Functional diversity										
FDis of SLA trait values	0.64 (±0.32)	0.61 (±0.25)	0.59 (±0.29)	0.59 (±0.21)	0.925	0.337	0.242	0.623	0.110	0.741
FDis of plant height trait values	0.70 (±0.26)	0.62 (±0.18)	0.64 (±0.35)	0.64 (±0.24)	0.125	0.725	1.048	0.307	1.022	0.313

Table 2. Top three among 13 top-ranking models predicting taxonomic and functional diversity of riparian
plants on gravel bars, at the community scale, along the Rhône River (France), as assessed with the Akaike
information criterion corrected for small sample size (AICc). Number of estimated parameters including the
intercept and random effect (k), AICc value, difference in AICc (ΔAICc), AICc weight (*W*) and marginal coefficient
of determination for fixed effect (R<sup>2</sup><sub>marginal</sub>) are provided (for the description of model parameters, please refer

684 to Table 1)

Dependent variable	Model (fixed-effects)	k	AICc	Δ AICc	W	R <sup>2</sup> marginal
Total richness	Manag + Year * Dura_Inund	6	1601.5	0.000	0.390	0.651
Total richness	Manag + Year + Elevation	5	1602.1	0.627	0.285	0.658
Total richness	Manag + Year * Thick _Sedim	6	1603.9	2.435	0.115	0.648
Annual richness	Manag + Year * Thick _Sedim	6	1262.3	0.000	0.887	0.710
Annual richness	Manag + Year + Thick _Sedim	5	1266.4	4.148	0.112	0.716
Annual richness	Manag + Year * Prop_Sedim	6	1275.4	13.160	0.001	0.691
Exotic richness	Manag + Year + Thick _Sedim	5	983.1	0.000	0.477	0.555
Exotic richness	Manag + Year * Thick _Sedim	6	983.7	0.621	0.350	0.551
Exotic richness	Manag + Year * Prop_Sedim	6	986.6	3.511	0.082	0.539
FDis SLA	Manag + Year * Dura_Inund	7	16.2	0.000	0.368	0.130
FDis SLA	Manag + Year + Thick _Sedim	6	17.8	1.626	0.163	0.144
FDis SLA	Manag + Year + Prop_Sedim	6	18.2	2.009	0.135	0.142
FDis height	Manag + Year + Prop_Sedim	6	30.3	0.000	0.571	0.061
FDis height	Manag + Year * Prop_Sedim	7	32.2	1.931	0.217	0.062
FDis height	Manag + Year + Thick _Sedim	6	33.1	2.871	0.136	0.047
FDis seed	Manag + Year * Thick _Sedim	7	-8.9	0.000	0.415	0.184
FDis seed	Manag + Year + Thick _Sedim	6	-8.8	0.067	0.401	0.177
FDis seed	Manag + Year + Prop_Sedim	6	-4.8	4.027	0.055	0.156

Table 3. Relative importance (Imp.), estimate coefficients (Estimate (± SE)) and confidence intervals (95% CI) for each variable used to predict variations in taxonomic and
 functional diversity of riparian plants on gravel bars along the Rhône River, France. The 95% confidence interval of coefficients in bold excluded 0. For the maintenance
 measures (Manag) and year of sampling (Year) factors, positive estimates represent plowing or 2019, while negative estimates represent brush clearing or 2018 (refer to
 Table 1 for the description of the other variables)

	Total richnoss				A					
_		lotal richn	less		Annual rich	ness	Exotic richness			
Parameters	Imp.	Estimate (±SE)	(95% CI)	Imp.	Estimate (±SE)	(95% CI)	Imp.	Estimate (±SE)	(95% CI)	
Year	1.00	-0.293 (±0.033)	(-0.357; -0.229)	1.00	-0.299 (±0.051)	(-0.400; -0.199)	1.00	-0.253 (±0.062)	(-0.373; -0.132)	
Management	1.00	0.599 (±0.105)	(0.393; 0.804)	1.00	0.944 (±0.149)	(0.652; 1.237)	1.00	0.828 (±0.111)	(0.610; 1.046)	
Elevation	0.38	-0.078 (±0.019)	(-0.115; -0.040)	0.00	-0.007 (±0.030)	(-0.065; 0.052)	0.03	-0.092 (±0.038)	(-0.166; -0.018)	
Distance	0.00	-0.070 (±0.022)	(-0.112; -0.027)	0.00	-0.033 (±0.034)	(-0.100; 0.033)	0.00	-0.062 (±0.044)	(-0.148; 0.024)	
Prop_Sedim	0.03	-0.106 (±0.028)	(-0.161; -0.050)	0.00	-0.280 (±0.045)	(-0.368; -0.193)	0.09	-0.163 (±0.055)	(-0.271; -0.055)	
Thick_Sedim	0.20	-0.111 (±0.029)	(-0.168; -0.054)	1.00	-0.307 (±0.047)	(-0.400; -0.215)	0.83	-0.162 (±0.049)	(-0.259; -0.066)	
Dura_Inund	0.39	0.132 (±0.029)	(0.076; 0.189)	0.00	0.032 (±0.045)	(-0.056; 0.121)	0.05	0.156 (±0.057)	(0.045; 0.267)	
Inte_Inund	0.00	0.061 (±0.025)	(0.012; 0.111)	0.00	0.094 (±0.037)	(0.022; 0.166)	0.00	0.056 (±0.044)	(-0.030; 0.142)	
Year * Elevation	0.10	0.003 (±0.033)	(-0.061; 0.068)	0.00	-0.023 (±0.051)	(-0.122; 0.076)	0.01	0.026 (±0.065)	(-0.101; 0.154)	
Year * Distance	0.00	0.085 (±0.030)	(0.026; 0.144)	0.00	0.111 (±0.045)	(0.022; 0.199)	0.00	0.126 (±0.057)	(0.014; 0.238)	
Year * Prop_Sedim	0.03	0.121 (±0.034)	(0.053; 0.188)	0.00	0.206 (±0.055)	(0.098; 0.315)	0.08	0.176 (±0.067)	(0.045; 0.307)	
Year * Thick_Sedim	0.12	0.054 (±0.032)	(-0.009; 0.116)	0.89	0.125 (±0.050)	(0.027; 0.224)	0.35	0.076 (±0.062)	(-0.046; 0.197)	

 Year \* Dura\_Inund
 0.39
 -0.131 (±0.034)
 (-0.198; -0.065)
 0.00
 -0.175 (±0.051)
 (-0.276; -0.074)
 0.05
 -0.200 (±0.065)
 (-0.328; -0.073)

 Year \* Inte\_Inund
 0.00
 -0.070 (±0.040)
 (-0.148; 0.008)
 0.00
 -0.236 (±0.060)
 (-0.354; -0.118)
 0.00
 -0.132 (±0.075)
 (-0.279; 0.014)

		FDis SLA	Ą	FDis Height				FDis seed			
Parameters	Imp.	Estimate (±SE)	(95% CI)	Imp.	Estimate (±SE)	(95% CI)	Imp.	Estimate (±SE)	(95% CI)		
Year	1.00	-0.030 (±0.036)	(-0.100; 0.040)	1.00	-0.014 (±0.037)	(-0.086; 0.058)	1.00	-0.021 (±0.032)	(-0.084; 0.042)		
Management	1.00	-0.013 (±0.087)	(-0.184; 0.158)	1.00	-0.048 (±0.038)	(-0.122; 0.026)	1.00	0.164 (±0.076)	(0.014; 0.314)		
Elevation	0.09	-0.090 (±0.026)	(-0.140; -0.039)	0.00	-0.003 (±0.023)	(-0.048; 0.042)	0.03	0.014 (±0.028)	(-0.040; 0.068)		
Distance	0.00	-0.042 (±0.025)	(-0.091; 0.007)	0.00	-0.004 (±0.026)	(-0.055; 0.047)	0.01	0.012 (±0.019)	(-0.025; 0.050)		
Prop_Sedim	0.20	-0.104 (±0.025)	(-0.152; -0.056)	0.79	-0.064 (±0.021)	(-0.105; -0.022)	0.08	-0.039 (±0.025)	(-0.087; 0.009)		
Thick_Sedim	0.22	-0.106 (±0.025)	(-0.155; -0.056)	0.19	-0.056 (±0.022)	(-0.099; -0.013)	0.82	-0.049 (±0.028)	(-0.105; 0.006)		
Dura_Inund	0.49	0.128 (±0.035)	(0.059; 0.197)	0.01	0.024 (±0.024)	(-0.023; 0.071)	0.01	0.000 (±0.026)	(-0.051; 0.052)		
Inte_Inund	0.00	0.069 (±0.025)	(0.020; 0.118)	0.00	0.001 (±0.021)	(-0.040; 0.042)	0.03	-0.030 (±0.026)	(-0.080; 0.021)		
Year * Elevation	0.04	0.046 (±0.035)	(-0.021; 0.114)	0.00	0.024 (±0.038)	(-0.051; 0.098)	0.02	-0.064 (±0.033)	(-0.128; 0.000)		
Year * Distance	0.00	0.046 (±0.036)	(-0.024; 0.117)	0.00	0.049 (±0.038)	(-0.025; 0.123)	0.00	0.002 (±0.033)	(-0.062; 0.066)		
Year * Prop_Sedim	0.06	0.026 (±0.035)	(-0.043; 0.096)	0.22	0.017 (±0.037)	(-0.055; 0.090)	0.03	-0.031 (±0.033)	(-0.096; 0.034)		
Year * Thick_Sedim	0.06	0.013 (±0.035)	(-0.055; 0.082)	0.05	0.016 (±0.037)	(-0.057; 0.089)	0.41	-0.048 (±0.032)	(-0.111; 0.015)		
Year * Dura_Inund	0.37	-0.075 (±0.036)	(-0.145; -0.005)	0.00	-0.021 (±0.039)	(-0.098; 0.055)	0.00	0.037 (±0.034)	(-0.030; 0.105)		

- 690 Figure 1. Location of the study area within the Rhône River watershed (A), distribution of sampled gravel bars
- along the Rhône River corridor (B), typical arrangement of a diversion scheme on the Rhône (C) and form of the
- 692 sampling design used to survey riparian plant communities (D)



Figure 2. Variation in topographic, pedological and hydrological variables, for each gravel bar, between 2017
and 2019 along the Rhône River, France (\* = adjusted *p*-value < 0.05 based on paired samples *t*-test, see
Appendix S7 for details and refer to Table 1 for the description of the variables). Gravel bars are aligned in a
north-south direction of the river corridor (Figure 1), with GB1 to GB4 managed by brush clearing and GB5 to
GB8 managed by plowing



Figure 3. Variation in taxonomic and functional diversity of riparian plants, for each gravel bar, between 2017
and 2019, along the Rhône River, France (\* = adjusted *p*-value < 0.05 based on paired samples *t*-test, see
Appendix S9 for details and refer to Table 1 for the description of the variables). Gravel bars are aligned in a
north-south direction of the river corridor (Figure 1), with GB1 to GB4 managed by brush clearing and GB5 to
GB8 managed by plowing (FDis = functional dispersion)



Figure 4. Constrained canonical analysis of principal coordinates of A) taxonomic and B) functional composition
of riparian communities, for each gravel bars, between 2017 and 2019 along the Rhône River, France. Small
squares and circles represent the position of riparian communities at the quadrat scale, respectively in 2017
and 2019, colored lines connect each quadrat to the centroids in the principal coordinate space of each
individual bar per year, represented by a rectangle with the site name



Figure 5. Constrained canonical analysis of principal coordinates of A) taxonomic and B) functional composition
of riparian communities in relation to topographic, pedological and hydrological variables along the Rhône
River, France (refer to Table 1 for the description of the variables). To facilitate graphical interpretation, the
centroids in principle coordinate space of each individual bar by year, the 10 most abundant species and the
three functional traits (i.e., CWM) are provided

