



HAL
open science

Tn-phyto: essential genomes of nine bacterial phytopathogens

Julie Baltenneck, Hajar Ben Moussa, Loïc Couderc, Areski Flissi, Jacques Pédrón, Cindy E. Morris, Marie-Anne Barny, Odile Berge, Hélène Touzet, Erwan Gueguen, et al.

► To cite this version:

Julie Baltenneck, Hajar Ben Moussa, Loïc Couderc, Areski Flissi, Jacques Pédrón, et al.. Tn-phyto: essential genomes of nine bacterial phytopathogens. 4. International Ewinia workshop (IEW), Jul 2022, Assise, Italy. hal-03721522

HAL Id: hal-03721522

<https://hal.science/hal-03721522v1>

Submitted on 19 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TN-PHYTO: ESSENTIAL GENOMES OF NINE BACTERIAL PHYTOPATHOGENS

Baltenneck J.¹, Ben Moussa H.², Couderc L.³, Flissi A.⁵, Pédrón J.², Morris C.E.⁴, Barny MA.², Berge O.⁴, Touzet H.⁵, Gueguen E.^{1*}, **Condemine G.^{1*}**

¹ University of Lyon, Université Claude Bernard Lyon 1, INSA of Lyon, MAP CNRS UMR 5240, 69622 Villeurbanne, France

² Sorbonne Université, INRAE, Institute of Ecology and Environmental Sciences-Paris, 4 place 7 Jussieu, F-75 252 Paris, France.


³ University of Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, US 41 - UMS 2014 - PLBS, Lille, France.

⁴ INRAE, Pathologie Végétale, 84140, Monfavet, France


⁵ University of Lille, CNRS, Centrale Lille, UMR 9189 - CRISTAL - Centre de Recherche en Informatique Signal et Automatique de Lille, Lille, France.

The aim of our project is to identify virulence factors of phytopathogenic bacteria isolated from the environment as well as those allowing their survival in water. For this purpose, we use Tn-seq (2), an innovative mutagenesis technique coupled with high throughput sequencing chosen to rapidly identify, on a large panel of strains, genes necessary for growth in a given condition, the plant or water. This study is carried out on strains of *Dickeya*, *Pectobacterium* and on the *Pseudomonas syringae* complex, all of which are capable of growing on a wide range of hosts as well as in water, thus allowing the most complete identification of genes of interest. Another axis of this ANR is the development of a complete bioinformatics pipeline, allowing to process data obtained under multiple conditions and for strains of different species. The results obtained will help us to predict the virulence of strains and/or to develop antibacterial strategies. Overall, this project will pave the way for studies of other environmental pathogens by proposing an original conceptual and technological framework that is extremely promising.


Materials and methods




Mutant libraries are constructed by conjugation between an *Escherichia coli* MDFpir strain and each of the strains of interest. During this conjugation, the plasmid pSAM-Ec, which contains the mariner Himar9 transposon that confers kanamycin resistance (3), a transposase and an ampicillin resistance gene, is transferred to the strain of interest. It is a suicide plasmid that cannot replicate in the recipient strain. Thus, the transposon will integrate uniquely and randomly into the genome of the strain of interest. The mutants obtained can then be selected on medium containing kanamycin.



Genomic DNA from the mutant libraries is extracted and digested with the enzyme MmeI. Indeed, the transposon has MmeI restriction sites in each of its extremities. The particularity is that the enzyme will cut at 16 bp of its site. This will generate fragments containing the transposon and the adjacent 16 or 17 bp which corresponds to gDNA flanking the transposon. The P5 adaptors containing a barcode are ligated to the digested DNA fragments using a T4 DNA ligase. Then, PCRs are used to amplify the fragment which will be sequenced to determine the insertion gene of the transposon.




Results



The quality of the libraries created for each strain was analyzed by Novaseq sequencing (MGX, Montpellier). It allowed to determine the percentage of transposon insertion in the genome of each strain. A good quality library is considered to have an insertion density of more than 30%. The correlation coefficient shows a high reproducibility of the transposition. These two elements attest to the good quality of the mutant libraries constructed.

Conclusions and perspectives



(A) Now that the quality of the library is checked, the objective is to apply different conditions to the mutant libraries in order to determine the essential genes for bean infection or bacterial survival in a specific medium (TSB 50%, sterilized river water).

(B) Figure from Royet et al. (2019) (1). The results that will be obtained will provide a read number of the sequenced fragment. The presence of 16 bp adjacent to the transposon will allow to align these reads to the bacterial genome. The number of reads will reflect the importance of the mutated gene in under a certain condition. If the read count is low for a given gene, the mutant is not represented and the gene has an important role in that condition.

References:

- (1) Royet et al. 2019. *Molecular Plant Pathology* : 287-306.
- (2) van Opijnen. 2009. *Nat Methods* : 767-772.
- (3) Wiles TJ. 2013. *PLoS Genet* 9(8): e1003716.

Funding: ANR- 19-CE35-0016