

Singular behavior of the solution of the Helmholtz equation in weighted L^p -Sobolev spaces

Colette De Coster and Serge Nicaise

Université de Valenciennes et du Hainaut Cambrésis
LAMAV, FR CNRS 2956,
Institut des Sciences et Techniques de Valenciennes
F-59313 Valenciennes Cedex 9, France
Colette.DeCoster,Serge.Nicaise@univ-valenciennes.fr

September 2, 2010

Abstract

We study the Helmholtz equation

$$(1) \quad -\Delta u + zu = g \text{ in } \Omega,$$

with Dirichlet boundary conditions in a polygonal domain Ω , where z is a complex number. Here g belongs to $L^p_\mu(\Omega) = \{v \in L^p_{loc}(\Omega) : r^\mu v \in L^p(\Omega)\}$, with a real parameter μ and $r(x)$ the distance from x to the set of corners of Ω . We give sufficient conditions on μ, p and Ω that guarantee that problem (1) has a unique solution $u \in H^1_0(\Omega)$ that admits a decomposition into a regular part in weighted L^p -Sobolev spaces and an explicit singular part. We further obtain some estimates where the explicit dependence on $|z|$ is given.

Keywords: Helmholtz equation, singular behavior, nonsmooth domains.

AMS Subject Classification: 35J05, 35B65.

1 Introduction

This paper is the first one of a large program of research devoted to the study of (nonlinear) heat equation in nonsmooth domains in weighted L^p -Sobolev spaces. Our final goal requires precise informations about the solution of the linear heat equation, in particular its decomposition into a regular part and an explicit singular part. Although this theory is well developed in weighted L^2 -Sobolev spaces [8, 11, 10, 3] or in L^p -Sobolev spaces [9], to our best knowledge such a result does not exist in the framework of weighted L^p -Sobolev spaces. For maximal regularity type results in weighted L^p -Sobolev spaces, we refer to [5, 15, 20, 16, 19].

According to the approach of [9], the study of the linear heat equation in non-hilbertian Sobolev spaces can be performed with the help of the theory of sums of operators. This theory requires in a first step to obtain uniform estimates of the solution of the Helmholtz equation. Hence the goal of this paper is to make this analysis in $L^p_\mu(\Omega)$ for a large range of values of μ and p . Our results extend the ones from [8] to the $L^p_\mu(\Omega)$ setting.

For the sake of simplicity we have restricted ourselves to the two-dimensional situation. The results of this paper can be easily extended to the case of domains with conical points.

The paper is organized as follows: In section 2 after some recalls, we prove some embeddings and some a priori estimates in $H^1 - 0$ and in L^p_μ norms. Section 3 is devoted to the proof of the decomposition of the solution of (1) with uniform estimates with respect to the parameter z .

2 Some preliminary results

2.1 Some notations and definitions

In this paper we consider polygonal domains of \mathbb{R}^2 in the following sense.

Definition 2.1. *Let Ω be a bounded domain of \mathbb{R}^2 . We say that Ω is a polygonal domain if its boundary is the union of a finite number of line segments $\bar{\Gamma}_j$, $j \in \{1, \dots, J\}$ (Γ_j being supposed to be open). Hence we do not assume that Ω is a Lipschitz domain, that is we include the presence of cracks.*

Denote by $S_j, j = 1, \dots, J$ the vertices of $\partial\Omega$ enumerated clockwise. Without loss of generality we may assume that $B(S_j, 1) \cap \Omega$ does not contain

any other vertex of Ω . For $j \in \{1, 2, \dots, J\}$, let ψ_j be the interior angle of Ω at the vertex S_j , $\lambda_j = \frac{\pi}{\psi_j}$ and (r_j, θ_j) the polar coordinates centered at S_j such that

$$B(S_j, 1) \cap \Omega = \{(r_j \cos \theta_j, r_j \sin \theta_j) \mid 0 < r_j < 1, 0 < \theta_j < \psi_j\} =: D_j.$$

For $\vec{\mu} = (\mu_j)_{j=1}^J$, we define the spaces $L_{\vec{\mu}}^p(\Omega) = \{f \in L_{loc}^p(\Omega) \mid wf \in L^p(\Omega)\}$ with

$$w = 1 + \sum_{j=1}^J \eta_j (r_j^{\mu_j} - 1), \quad (2.1)$$

where $r_j(x)$ is the distance from x to the vertex S_j and $\eta_j \in \mathcal{D}(\mathbb{R}^2)$ are such that

$$\eta_j \equiv 1 \text{ in } D_j(1/2), \quad \eta_j \equiv 0 \text{ on } \Omega \setminus D_j(1),$$

where $D_j(r)$ is the truncated cone $D_j(r) = \Omega \cap B(S_j, r)$. Note that the weight w satisfies

$$w = r_j^{\mu_j} \text{ on } D_j(1/2) \quad \text{and} \quad w = 1 \text{ on } \Omega \setminus \cup_{j=1}^J D_j(1).$$

The space $L_{\vec{\mu}}^p(\Omega)$ is a Banach space for the norm

$$\|f\|_{L_{\vec{\mu}}^p(\Omega)} = \left(\int_{\Omega} |f(x)|^p w^p(x) dx \right)^{1/p}.$$

$V_{\vec{\mu}}^{k,p}(\Omega)$ is now defined as the closure of

$$\mathcal{C}_S^\infty(\Omega) = \{v \in \mathcal{C}^\infty(\bar{\Omega}) \mid S_j \notin \text{supp } v\}$$

with respect to the norm

$$\|u\|_{V_{\vec{\mu}}^{k,p}(\Omega)} = \left(\sum_{|\gamma| \leq k} \int_{\Omega} |D^\gamma u(x)|^p w^p(x) r^{(|\gamma|-k)p}(x) dx \right)^{1/p}.$$

We denote

$$|u|_{V_{\vec{\mu}}^{k,p}(\Omega)} = \left(\sum_{|\gamma|=k} \int_{\Omega} |D^\gamma u(x)|^p w^p(x) r^{(|\gamma|-k)p}(x) dx \right)^{1/p}.$$

In $H_0^1(\Omega)$ we will denote the norms in the following way

$$|u|_{H_0^1}^2 = \int_{\Omega} |\nabla u|^2 \quad \text{and} \quad \|u\|_{H_0^1}^2 = \int_{\Omega} (|\nabla u|^2 + |u|^2).$$

For $\vec{\mu}$ and $\vec{\gamma}$, we write $\vec{\mu} > \vec{\gamma}$ in case, for all $j \in \{1, \dots, J\}$, $\mu_j > \gamma_j$.

2.2 Embeddings and consequences

Lemma 2.1. *Let Ω be a polygonal domain of \mathbb{R}^2 , $p \geq 2$ and $\vec{\mu}$ satisfies,*

$$\forall j = 1, \dots, J, \quad \begin{aligned} \mu_j &< \frac{2p-2}{p}, & \text{if } p > 2, \\ \mu_j &\leq 1, & \text{if } p = 2. \end{aligned} \quad (2.2)$$

Then the next continuous embeddings hold:

1. $L_{\vec{\mu}}^p(\Omega) \hookrightarrow L_{\vec{1}}^2(\Omega)$,
2. $L_{-\vec{1}}^2(\Omega) \hookrightarrow (L_{\vec{\mu}}^p(\Omega))' = L_{-\vec{\mu}}^q(\Omega)$, with $q = \frac{p}{p-1}$,
3. $H_0^1(\Omega) \hookrightarrow L_{-\vec{\mu}}^q(\Omega)$.

Proof. The first assertion follows from the identity

$$r_j f = r_j^{\mu_j} f r_j^{1-\mu_j},$$

Hölder's inequality and the fact that $r_j^{1-\mu_j}$ belongs to $L^q(\Omega)$ if $1 - \mu_j > -\frac{2}{q}$.

The second assertion is a consequence of the first one by using duality.

For the last assertion, in case $p = 2$, it is known (see [4, Thm 14.5.5] or Lemma 2.11 below) that $H_0^1(\Omega)$ is continuously embedded in $L_{-\vec{1}}^2(\Omega)$. We then conclude observing that, for $\mu_j \leq 1$, we have $r_j^{2(-\mu_j+1)} \in L^\infty(\Omega)$.

In the case $p > 2$, we use the embedding $H_0^1(\Omega) \hookrightarrow L_{-\vec{1}}^2(\Omega)$ and the second assertion. \square

Lemma 2.2. *Let Ω be a polygonal domain of \mathbb{R}^2 , $p \geq 2$ and $\vec{\mu}$ satisfies, for all $j = 1, \dots, J$,*

$$\begin{aligned} \mu_j &> \frac{-2}{p}, & \text{if } p > 2, \\ \mu_j &\geq -1, & \text{if } p = 2. \end{aligned}$$

Then $H_0^1(\Omega)$ is continuously embedded in $L_{\vec{\mu}}^p(\Omega)$.

Proof. We have the continuous embedding of $H_0^1(\Omega)$ into $L^p(\Omega)$ for all $p > 1$ and $r_j^{\mu p} \in L^s(\Omega)$ for all $s > 1$ satisfying $\mu p s + 2 > 0$. The result follows by Hölder inequality as we have by assumption $s > 1$ such that $\mu p s + 2 > 0$ and, for $f \in H_0^1(\Omega)$,

$$\int_{\Omega} r_j^{\mu p} |f|^p \leq \| |f|^p \|_{L^{s'}} \| r_j^{\mu p} \|_{L^s},$$

with $\frac{1}{s} + \frac{1}{s'} = 1$. \square

Lemma 2.3. *Let Ω be a polygonal domain of \mathbb{R}^2 , $p \geq 2$, $\vec{\mu}$ satisfy (2.2) and $\theta_A \in]0, \pi[$. Then for all $g \in L^p_{\vec{\mu}}(\Omega)$ and all $z \in \mathbb{C}$ with $|\arg z| \leq \theta_A$, the problem*

$$\forall \varphi \in H_0^1(\Omega), \quad \int_{\Omega} \nabla u \cdot \nabla \bar{\varphi} + z \int_{\Omega} u \bar{\varphi} = \int_{\Omega} g \bar{\varphi}, \quad (2.3)$$

has a unique solution $u \in H_0^1(\Omega)$.

Proof. For all $g \in L^p_{\vec{\mu}}(\Omega)$, we know by Lemma 2.1 that the mapping

$$H : H_0^1(\Omega) \rightarrow \mathbb{C} : u \mapsto \int_{\Omega} g \bar{u}$$

is linear and continuous. Observe that, for all $z \in \mathbb{C}$ with $|\arg z| \leq \theta_A$, there exists $\theta \in [0, 2\pi]$ such that $\cos \theta > 0$ and $\Re(ze^{i\theta}) \geq 0$. Hence the conclusion follows from the Lax-Milgram Lemma as, for all $u \in H_0^1(\Omega)$,

$$\Re[e^{i\theta} \int_{\Omega} (|\nabla u|^2 + z|u|^2)] \geq \cos \theta \int_{\Omega} |\nabla u|^2 \geq \alpha \|u\|_{H_0^1}^2,$$

for some $\alpha > 0$, due to Poincaré's inequality. \square

Remark 2.1 The unique solution $u \in H_0^1(\Omega)$ of (2.3) is called a weak solution of the problem

$$\begin{cases} -\Delta u + zu = g, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

since $-\Delta u + zu = g$ in the distributional sense.

2.3 Some inequalities

For $R > 0$ and $\theta_A \in]\frac{\pi}{2}, \pi[$ fixed, we define

$$\pi^+ = \{z \in \mathbb{C} \mid \Re(z) \geq 0\}, \quad S_A = \{z \in \mathbb{C} \mid |z| \geq R \text{ and } |\arg z| \leq \theta_A\}. \quad (2.4)$$

For a solution $u \in H_0^1(\Omega)$ of (2.3) we will use the following notation: $\|u\| \lesssim \|g\|$ means the existence of a positive constant C , which is independent of the quantities $\|u\|$, $\|g\|$ and z under consideration such that $\|u\| \leq C\|g\|$ and $\|u\| \sim \|g\|$ means $\|u\| \lesssim \|g\|$ and $\|g\| \lesssim \|u\|$.

Lemma 2.4. *Let $R > 0$ and $\theta_A \in]\frac{\pi}{2}, \pi[$ be fixed. Let Ω be a polygonal domain of \mathbb{R}^2 , $p \geq 2$, $\vec{\mu}$ satisfy (2.2), $z \in \pi^+ \cup S_A$, and $u \in H_0^1(\Omega)$ be the solution of (2.3). Then u satisfies the inequality*

$$|u|_{H_0^1(\Omega)} \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}. \quad (2.5)$$

Proof. Applying (2.3) with $w = u$ we have

$$|u|_{H_0^1(\Omega)}^2 + z \int_{\Omega} |u|^2 = \int_{\Omega} g\bar{u}. \quad (2.6)$$

By Lemma 2.1, taking the real and the imaginary parts of (2.6), we obtain

$$|u|_{H_0^1(\Omega)}^2 + \Re(z) \int_{\Omega} |u|^2 \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)}. \quad (2.7)$$

and

$$|\Im(z)| \int_{\Omega} |u|^2 \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)}. \quad (2.8)$$

Case 1: $\Re(z) \geq 0$. In that case the result can be directly deduced from (2.7).

Case 2: $\Re(z) < 0$. As $z \in S_A \cup \pi^+$, we have $\Re(z) = \rho \cos \theta$, $\Im(z) = \rho \sin \theta$ with $\rho > R$ and $|\Im(z)| > \rho \sin \theta_A$. Hence we deduce from (2.8) that

$$\int_{\Omega} |u|^2 \lesssim \frac{1}{\rho} \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)}. \quad (2.9)$$

Then (2.7) together with $\Re(z) < 0$ gives

$$\begin{aligned} |u|_{H_0^1(\Omega)}^2 &\lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)} - \Re(z) \int_{\Omega} |u|^2 \\ &\lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)} - \Re(z) \frac{1}{\rho} \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)}. \end{aligned}$$

We conclude using the inequality $-\Re(z) \leq \rho$. \square

Corollary 2.5. *Let $R > 0$ and $\theta_A \in]\frac{\pi}{2}, \pi[$ be fixed. Let Ω be a polygonal domain of \mathbb{R}^2 , $p \geq 2$, $\vec{\mu}$ satisfy (2.2), $g \in L_{\vec{\mu}}^p(\Omega)$, $z \in \pi^+ \cup S_A$, and $u \in H_0^1(\Omega)$ be the solution of (2.3). Then we have the inequalities*

$$|u|_{L^2(\Omega)}^2 \lesssim \frac{1}{|z|} \|g\|_{L_{\vec{\mu}}^p(\Omega)} |u|_{H_0^1(\Omega)} \lesssim \frac{1}{|z|} \|g\|_{L_{\vec{\mu}}^p(\Omega)}^2 \quad (2.10)$$

and

$$(1 + |z|^{1/2}) |u|_{L^2(\Omega)} \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}. \quad (2.11)$$

Proof. By (2.6), we have

$$z \int_{\Omega} |u|^2 = \int_{\Omega} g\bar{u} - |u|_{H_0^1(\Omega)}^2,$$

from which we deduce, using Lemmas 2.1 and 2.4,

$$|z| \int_{\Omega} |u|^2 \lesssim \|g\|_{L_{\mu}^p(\Omega)} |u|_{H_0^1(\Omega)} + |u|_{H_0^1(\Omega)}^2 \lesssim \|g\|_{L_{\mu}^p(\Omega)} |u|_{H_0^1(\Omega)}.$$

We then prove (2.10) by a second application of Lemma 2.4 and (2.11) is obtained by (2.10), Lemma 2.4 and Poincaré inequality. \square

Corollary 2.6. *Let $R > 0$ and $\theta_A \in]\frac{\pi}{2}, \pi[$ be fixed. Let Ω be a bounded domain of \mathbb{R}^2 , $g \in L^2(\Omega)$, $z \in \pi^+ \cup S_A$, and $u \in H_0^1(\Omega)$ be the solution of (2.3). Then we have the inequality*

$$(1 + |z|) \|u\|_{L^2(\Omega)} \lesssim \|g\|_{L^2(\Omega)}.$$

Proof. As in Lemma 2.4, we obtain (2.6).

Case 1: $\Re(z) \geq 0$. In that case, we also have

$$|u|_{H_0^1(\Omega)}^2 \leq \Re\left(\int_{\Omega} g\bar{u}\right) \leq \|g\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)}, \quad (2.12)$$

and

$$z \int_{\Omega} |u|^2 = \int_{\Omega} g\bar{u} - \int_{\Omega} |\nabla u|^2.$$

Using (2.12), this implies

$$|z| \int_{\Omega} |u|^2 \leq \left| \int_{\Omega} g\bar{u} \right| + |u|_{H_0^1(\Omega)}^2 \lesssim \|g\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)},$$

from which we deduce

$$|z| \|u\|_{L^2(\Omega)} \lesssim \|g\|_{L^2(\Omega)}.$$

By Poincaré inequality and (2.12), we have,

$$\|u\|_{L^2(\Omega)}^2 \lesssim |u|_{H_0^1(\Omega)}^2 \leq \|g\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)},$$

and the result follows.

Case 2: $\Re(z) < 0$. In that case $z \in S_A$. As in the proof of Lemma 2.4, we have

$$\rho \|u\|_{L^2(\Omega)} \lesssim \|g\|_{L^2(\Omega)},$$

where $\rho = |z| > R$. Hence we deduce

$$(1 + |z|) \|u\|_{L^2(\Omega)} \lesssim |z| \|u\|_{L^2(\Omega)} \lesssim \|g\|_{L^2(\Omega)},$$

which concludes the proof. \square

Lemma 2.7. *Let Ω be a polygonal domain of \mathbb{R}^2 , $p \geq 2$, $\vec{\mu}$ satisfy (2.2), $f \in L^p_{\vec{\mu}}(\Omega)$ and $u \in H^1_0(\Omega)$ be the solution of*

$$\forall \varphi \in H^1_0(\Omega), \quad \int_{\Omega} \nabla u \cdot \nabla \bar{\varphi} = \int_{\Omega} f \bar{\varphi}. \quad (2.13)$$

Then, for all $\vec{\gamma} \geq \vec{\mu}$ with $\vec{\gamma} > 2 - \frac{2}{p} - \vec{\lambda}$ and $\vec{\lambda} = (\lambda_j)_{j=1}^J = (\frac{\pi}{\lambda_j})_{j=1}^J$,

$$u \in V_{\vec{\gamma}}^{2,p}(\Omega),$$

and, in particular,

$$r_j^{\gamma_j - 2} u \in L^p(D_j), \quad r_j^{\gamma_j - 1} \nabla u \in (L^p(D_j))^2, \quad r_j^{\gamma_j} \Delta u \in L^p(D_j).$$

If moreover $\vec{\mu} > -\vec{\lambda}$ then we have in particular

$$u \in V_{\vec{\mu} + 2 - \frac{2}{p}}^{2,p}(\Omega),$$

and hence,

$$r_j^{\mu_j - \frac{2}{p}} u \in L^p(D_j), \quad r_j^{\mu_j + 1 - \frac{2}{p}} \nabla u \in (L^p(D_j))^2, \quad r_j^{\mu_j + 2 - \frac{2}{p}} \Delta u \in L^p(D_j).$$

Proof. Using regularity results far from the corners of Ω for the Laplace equation with Dirichlet boundary conditions (see [6, 7]), $u \in W^{2,p}(\tilde{\Omega})$, where, for some $\delta > 0$, $\tilde{\Omega}$ is a subdomain of $\bar{\Omega}$ with a smooth boundary, its boundary being the same as Ω except in $\bigcup_{j=1}^J B(S_j, \delta)$, with the estimate

$$\|u\|_{W^{2,p}(\tilde{\Omega})} \lesssim \|f\|_{L^p_{\vec{\mu}}(\Omega)}.$$

September 2, 2010

It then remains to look at the behaviour of u near the corners. For any $j = 1, \dots, J$, consider the cut-off function $\eta_j \in \mathcal{D}(\mathbb{R}^2)$ used in (2.1). For shortness we write $D = D_j(1)$ and drop the index j . Let us set

$$\tilde{u} = \eta u.$$

This function satisfies

$$\begin{cases} -\Delta \tilde{u} = \tilde{h}, & \text{in } D, \\ \tilde{u} = 0, & \text{on } \partial D, \end{cases} \quad (2.14)$$

where $\tilde{h} = \eta h - 2\nabla \eta \cdot \nabla u - u \Delta \eta$. Moreover due to the above results (regularity far from the corners), \tilde{h} belongs to $L^p_\mu(D)$ and

$$\|\tilde{h}\|_{L^p_\mu(D)} \lesssim \|h\|_{L^p_\mu(\Omega)}.$$

For $\gamma \geq \mu$ we have $L^p_\mu(D) \subset L^p_\gamma(D)$. As, by [12, Remark 9.11], $L^p_\gamma(D) = V^{0,p}_\gamma(D)$, applying [13, Lemma 11.2 (ii)] (as in [13, Example 11.3]) we prove that $u \in V^{2,2}_1(D)$. Let γ_1 be such that $\mu \leq \gamma_1 \leq \gamma$, $\gamma_1 > 2 - \frac{2}{p} - \lambda$ and $\gamma_1 \leq 2 - \frac{2}{p}$. As $f \in L^p_{\gamma_1}(D)$, by [13, Thm 9.3] we have $u \in V^{2,p}_{\gamma_1}(D)$ if

$$0 \leq 2 - \frac{2}{p} - \gamma_1 < \lambda,$$

which is equivalent to

$$2 - \frac{2}{p} - \lambda < \gamma_1 \leq 2 - \frac{2}{p}.$$

This gives the first part of the result observing that $V^{2,p}_{\gamma_1}(D) \subset V^{2,p}_\gamma(D)$ if $\gamma_1 \leq \gamma$.

To prove the second part of the result, we just have to observe that, if $\mu > -\lambda$ then $\mu - \frac{2}{p} + 2 \geq \mu$ and $\mu + 2 - \frac{2}{p} > -\lambda + 2 - \frac{2}{p}$. \square

Corollary 2.8. *Let Ω be a polygonal domain, $p \geq 2$, $\vec{\mu} > -\vec{\lambda}$ satisfy (2.2), $f \in L^p_{\vec{\mu}}(\Omega)$, $u \in H^1_0(\Omega)$ be the solution of (2.13) and define $v = w u$ with w defined by (2.1). Then we have*

$$v \in V^{2,p}_{2-\frac{2}{p}}(\Omega), \quad r^{-\frac{2}{p}}v \in L^p(\Omega), \quad r^{1-\frac{2}{p}}\nabla v \in (L^p(\Omega))^2, \quad r^{2-\frac{2}{p}}\Delta v \in L^p(\Omega).$$

Proof. This can be easily deduced from Lemma 2.7. \square

Definition 2.2. For Ω a polygonal domain, $p \geq 2$ and $\vec{\mu} \in \mathbb{R}^J$, we define

$$D(\Delta_{p,\vec{\mu}}) = \{u \in H_0^1(\Omega) \mid \Delta u \in L_{\vec{\mu}}^p(\Omega)\}.$$

Before going on let us show that Lemma 2.7 furnishes an existence result in $D(\Delta_{p,\vec{\mu}})$ for problem (2.3).

Lemma 2.9. Let $\theta_A \in]0, 2\pi[$, $p \geq 2$, $\vec{\mu}$ satisfies (2.2) as well as $\vec{\mu} > -\vec{\lambda} - \frac{2}{p}$, $z \in \mathbb{C}$ with $|\arg z| \leq \theta_A$, and $u \in H_0^1(\Omega)$ be the solution of (2.3) with $g \in L_{\vec{\mu}}^p(\Omega)$. Then $u \in D(\Delta_{p,\vec{\mu}})$.

Proof. We only need to show that

$$-\Delta u \in L_{\vec{\mu}}^p(\Omega). \quad (2.15)$$

Using regularity results far from the corners of Ω for the Laplace equation with Dirichlet boundary conditions (see [6, 7]), $u \in W^{2,p}(\tilde{\Omega})$, where $\tilde{\Omega}$ was introduced in the proof of Lemma 2.7. We directly deduce that $u \in W^{2,p}(\tilde{\Omega})$. Hence it remains to show (2.15) near each corner S_j . For a fixed $j \in \{1, 2, \dots, J\}$, we then set $u_j = \eta_j u \in H_0^1(D_j)$ with η_j the cut-off function used in (2.1). The function u_j is a weak solution of

$$\begin{cases} -\Delta u_j + z u_j = g_j, & \text{in } D_j, \\ u_j = 0, & \text{on } \partial D_j, \end{cases} \quad (2.16)$$

where $g_j = \eta_j g - 2\nabla \eta_j \cdot \nabla u - u \Delta \eta_j$. By the regularity $u \in W^{2,p}(\tilde{\Omega})$ mentioned before, we have

$$\nabla u \in W^{1,p}(\tilde{\Omega})^2, \quad u \in L^p(\tilde{\Omega}),$$

and consequently $g_j \in L_{\mu_j}^p(D_j)$ with

$$\|g_j\|_{L_{\mu_j}^p(D_j)} \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}. \quad (2.17)$$

Hence we can concentrate on (2.16) and prove that

$$-\Delta u_j \in L_{\mu_j}^p(D_j). \quad (2.18)$$

For the rest of the proof we drop the index j and write D for D_j , μ for μ_j , ...

As Lemma 2.2 guarantees that $H_0^1(D) \hookrightarrow L_{\mu'}^p(D)$ for all $\mu' > -\frac{2}{p}$, we distinguish different cases:

Case 1. $\mu > -\frac{2}{p}$. In that case $-\Delta u = g - zu \in L_{\mu}^p(D)$.

Case 2. $-2 - \frac{2}{p} < \mu \leq -\frac{2}{p}$. In that case, we take $\mu_1 = \mu + 2$ if $\mu < -\frac{2}{p}$ and $\mu_1 = 2 - \frac{2}{p} - \epsilon$, with $\epsilon \in]0, 2[$ if $\mu = -\frac{2}{p}$. Since $\mu_1 > -\frac{2}{p}$, $u \in H_0^1(\Omega)$ is solution of

$$-\Delta u = g - zu \in L_{\mu_1}^p(D).$$

This implies by Lemma 2.7 that

$$u \in V_{\mu_1}^{2,p}(D).$$

Accordingly

$$r^{\mu_1-2}u \in L^p(D),$$

which implies that

$$u \in L_{\mu}^p(D), \tag{2.19}$$

due to $\mu_1 - 2 \leq \mu$. This again guarantees (2.18) because $-\Delta u = g - zu$.

Case 3. $-4 - \frac{2}{p} < \mu \leq -2 - \frac{2}{p}$. then $\mu_2 = \mu + 2$ enters in the framework of Case 2, and therefore by Case 2, we obtain by (2.19) that

$$u \in L_{\mu_2}^p(D).$$

Now we use the same argument as before: look at $u \in H_0^1(D)$ as a solution of

$$-\Delta u = g - zu \in L_{\mu_2}^p(\Omega).$$

This implies by Lemma 2.7 that

$$u \in V_{\mu_2}^{2,p}(\Omega).$$

Accordingly

$$r^{\mu_2-2}u \in L^p(\Omega),$$

and we conclude that (2.19) holds.

The general case follows by induction. \square

Corollary 2.10. *Let $\theta_A \in]0, 2\pi[$, $D = \{(r \cos \theta, r \sin \theta) \mid 0 < r < 1, 0 < \theta < \psi\}$, $\lambda = \frac{\pi}{\psi}$, $p \geq 2$, $\mu \in \mathbb{R}$, $f \in L_{\mu}^p(D)$, $z \in \mathbb{C}$ with $|\arg z| \leq \theta_A$ and $u \in D(\Delta_{p,\mu})$ be the solution of*

$$\begin{cases} -\Delta u + zu = f, & \text{in } D, \\ u = 0, & \text{on } \partial D. \end{cases} \tag{2.20}$$

September 2, 2010

Assume $v = r^\mu u \in V_{2-\frac{2}{p}}^{2,p}(D)$. Then v satisfies

$$(a) \Re \left(\int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} \right) = 0;$$

$$(b) \frac{p}{2} \int_D |\nabla v|^2 |v|^{p-2} + \frac{p-2}{2} \int_D |v|^{p-4} \bar{v}^2 (\nabla v)^2 - \mu^2 \int_D r^{-2} |v|^p \\ + 2\mu \int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} + z \int_D |v|^p = \int_D r^\mu f |v|^{p-2} \bar{v}.$$

Remark 2.2 Recall that, by Corollary 2.8, if $\mu > -\lambda$ satisfies (2.2) then $v \in V_{2-\frac{2}{p}}^{2,p}(D)$ as $u \in D(\Delta_{p,\mu})$ is a solution of (2.20).

Proof. Recall that $V_{2-\frac{2}{p}}^{2,p}(D) = \overline{\mathcal{C}_S^\infty(D)}$ with $\mathcal{C}_S^\infty(D) = \{v \in \mathcal{C}^\infty(\bar{D}) \mid S \notin \text{supp } v\}$.

Step 1: Proof of (a). For $v \in \mathcal{C}_S^\infty(D)$, we have

$$\int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} r dr d\theta = - \int_D v \frac{\partial}{\partial r} (|v|^{p-2} \bar{v}) dr d\theta + \int_0^\psi |v(1, \theta)|^p d\theta$$

and hence

$$\begin{aligned} \int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} r dr d\theta &= \int_0^\psi |v(1, \theta)|^p d\theta \\ &\quad - \int_D v \left[\frac{\partial}{\partial r} ((v \bar{v})^{\frac{p-2}{2}}) \bar{v} + |v|^{p-2} \frac{\partial \bar{v}}{\partial r} \right] dr d\theta \\ &= \int_0^\psi |v(1, \theta)|^p d\theta \\ &\quad - \int_D v \left(\frac{p-2}{2} |v|^{p-4} (\bar{v})^2 \frac{\partial v}{\partial r} \right. \\ &\quad \left. + \frac{p-2}{2} |v|^{p-2} \frac{\partial \bar{v}}{\partial r} + |v|^{p-2} \frac{\partial \bar{v}}{\partial r} \right) dr d\theta \\ &= \int_0^\psi |v(1, \theta)|^p d\theta \\ &\quad - \int_D \frac{p-2}{2} |v|^{p-2} \bar{v} \frac{\partial v}{\partial r} dr d\theta - \int_D \frac{p}{2} |v|^{p-2} v \frac{\partial \bar{v}}{\partial r} dr d\theta. \end{aligned}$$

We obtain then

$$\frac{p}{2} \left(\int_D r^{-1} |v|^{p-2} \bar{v} \frac{\partial v}{\partial r} r dr d\theta + \int_D r^{-1} |v|^{p-2} v \frac{\partial \bar{v}}{\partial r} r dr d\theta \right) = \int_0^\psi |v(1, \theta)|^p d\theta$$

By density and [17, Thm 1.31, p.27 and Def 1.9 p.15], we can pass to the limit and we see that the previous equality is also valid for $v \in V_{2-2/p}^{2,p}(D)$. As $v = r^\mu u$ with $u \in H_0^1(D)$ we have $\gamma_0 v = 0$ on ∂D which gives the equality

$$\frac{p}{2} \left(\int_D r^{-1} |v|^{p-2} \bar{v} \frac{\partial v}{\partial r} + \int_D r^{-1} |v|^{p-2} v \frac{\partial \bar{v}}{\partial r} \right) = 0. \quad (2.21)$$

Then (a) follows by taking the real part of (2.21).

Step 2: Proof of (b). Observe that v satisfies

$$\begin{cases} -\Delta v - \mu^2 r^{-2} v + 2\mu r^{-1} \frac{\partial v}{\partial r} + z v = g, & \text{in } D, \\ v = 0, & \text{on } \partial D, \end{cases} \quad (2.22)$$

with $g = r^\mu f \in L^p(D)$. As $v \in V_{2-2/p}^{2,p}(D)$, it is meaningful to multiply (2.22) by $|v|^{p-2} \bar{v}$ and integrate. We then obtain

$$\begin{aligned} - \int_D \Delta v |v|^{p-2} \bar{v} - \mu^2 \int_D r^{-2} |v|^p + 2\mu \int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} + z \int_D |v|^p \\ = \int_D g |v|^{p-2} \bar{v}. \end{aligned} \quad (2.23)$$

For $v \in \mathcal{C}_S^\infty(D)$ we have

$$\begin{aligned} - \int_D \Delta v |v|^{p-2} \bar{v} &= \int_D \nabla v \cdot \nabla (|v|^{p-2} \bar{v}) + \int_{\partial D} \gamma_0 (|v|^{p-2} \bar{v}) \gamma_0 (\nabla v) \cdot \nu \\ &= \int_D |\nabla v|^2 |v|^{p-2} + \frac{p-2}{2} \left(\int_D |\nabla v|^2 |v|^{p-2} + \int_D |v|^{p-4} \bar{v}^2 (\nabla v)^2 \right) \\ &\quad + \int_{\partial D} \gamma_0 (|v|^{p-2} \bar{v}) \gamma_0 (\nabla v) \cdot \nu. \end{aligned}$$

As in Step 1, we show that we can pass to the limit and the previous equality is also valid for $v \in V_{2-2/p}^{2,p}(D)$. Again the boundary term is equal to zero and we obtain the equality

$$- \int_D \Delta v |v|^{p-2} \bar{v} = \frac{p}{2} \int_D |\nabla v|^2 |v|^{p-2} + \frac{p-2}{2} \int_D |v|^{p-4} \bar{v} (\nabla v)^2. \quad (2.24)$$

The result follows then from (2.23) and (2.24). \square

Lemma 2.11. *Let $D = \{(r \cos \theta, r \sin \theta) \mid 0 < r < 1, 0 < \theta < \psi\}$. For all $w \in H_0^1(D)$, we have*

$$\int_D |\nabla w|^2 \geq \frac{\pi^2}{\psi^2} \int_D \frac{1}{r^2} w^2.$$

Proof. First observe that

$$\int_D |\nabla w|^2 = \int_D \left(\left(\frac{\partial w}{\partial r} \right)^2 + \frac{1}{r^2} \left(\frac{\partial w}{\partial \theta} \right)^2 \right).$$

Moreover, for all $r \in]0, 1[$ we have $w(r, \cdot) \in H_0^1(0, \psi)$ and hence, by applying Poincaré inequality in $(0, \psi)$,

$$\int_0^\psi \left(\frac{\partial w}{\partial \theta} \right)^2(r, \theta) d\theta \geq \frac{\pi^2}{\psi^2} \int_0^\psi w^2(r, \theta) d\theta.$$

We deduce from this inequality that

$$\begin{aligned} \int_0^1 \int_0^\psi \left(\left(\frac{\partial w}{\partial r} \right)^2 + \frac{1}{r^2} \left(\frac{\partial w}{\partial \theta} \right)^2 \right) r d\theta dr &\geq \int_0^1 \int_0^\psi \left(\frac{\partial w}{\partial \theta} \right)^2 d\theta \frac{1}{r} dr \\ &\geq \frac{\pi^2}{\psi^2} \int_0^1 \int_0^\psi w^2(r, \theta) d\theta \frac{1}{r} dr \\ &= \frac{\pi^2}{\psi^2} \int_D w^2 \frac{1}{r^2}. \quad \square \end{aligned}$$

Remark 2.3 Observe that for $w(r, \theta) = r^\beta (1-r)^\alpha \sin(\lambda\theta)$ with $\beta > 0$ and $\alpha > 1/2$ we have $w \in H_0^1(D)$ and

$$\int_D |\nabla w|^2 = \left(\frac{\pi^2}{\psi^2} + \frac{\beta(\alpha + \beta)}{2\alpha - 1} \right) \int_D \frac{1}{r^2} w^2,$$

which proves the optimality of the previous inequality.

Corollary 2.12. *Let $D = \{(r \cos \theta, r \sin \theta) \mid 0 < r < 1, 0 < \theta < \psi\}$, $\lambda = \frac{\pi}{\psi}$, $p \geq 2$, $\mu \in \mathbb{R}$ satisfy*

$$4(p-1)\lambda^2 - \mu^2 p^2 > 0. \quad (2.25)$$

Let $f \in L_\mu^p(D)$, $z \in \mathbb{C}$ with $\Re(z) \geq 0$ and $u \in D(\Delta_{p,\mu})$ be the solution of

$$\begin{cases} -\Delta u + zu = f, & \text{in } D, \\ u = 0, & \text{on } \partial D. \end{cases} \quad (2.26)$$

Assume $v = r^\mu u \in V_{2-\frac{2}{p}}^{2,p}(D)$. Then we have the inequalities

$$\Re(z) \|u\|_{L_\mu^p(D)} \leq \|f\|_{L_\mu^p(D)} \quad \text{and} \quad |\Im(z)| \|u\|_{L_\mu^p(D)} \lesssim \|f\|_{L_\mu^p(D)}.$$

Proof. By Corollary 2.10, $v = r^\mu u$ satisfies

$$\begin{aligned} \frac{p}{2} \int_D |\nabla v|^2 |v|^{p-2} + \frac{p-2}{2} \int_D |v|^{p-4} \bar{v}^2 (\nabla v)^2 - \mu^2 \int_D r^{-2} |v|^p \\ + 2\mu \int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} + z \int_D |v|^p = \int_D g |v|^{p-2} \bar{v}, \end{aligned} \quad (2.27)$$

with $g = r^\mu f$ and

$$\Re \left(\int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} \right) = 0. \quad (2.28)$$

Writing $v = v_1 + i v_2$ with $v_i : D \rightarrow \mathbb{R}$, (2.27) becomes

$$\begin{aligned} \frac{p}{2} \int_D \left[\left(\frac{\partial v_1}{\partial r} \right)^2 + \left(\frac{\partial v_2}{\partial r} \right)^2 + \frac{1}{r^2} \left(\frac{\partial v_1}{\partial \theta} \right)^2 + \frac{1}{r^2} \left(\frac{\partial v_2}{\partial \theta} \right)^2 \right] (v_1^2 + v_2^2) |v|^{p-4} \\ + \frac{p-2}{2} \int_D |v|^{p-4} (v_1^2 - v_2^2 - 2i v_1 v_2) \left[\left(\frac{\partial v_1}{\partial r} \right)^2 - \left(\frac{\partial v_2}{\partial r} \right)^2 + 2i \frac{\partial v_1}{\partial r} \frac{\partial v_2}{\partial r} \right. \\ \left. + \frac{1}{r^2} \left(\frac{\partial v_1}{\partial \theta} \right)^2 - \frac{1}{r^2} \left(\frac{\partial v_2}{\partial \theta} \right)^2 + \frac{2i}{r^2} \frac{\partial v_1}{\partial \theta} \frac{\partial v_2}{\partial \theta} \right] \\ - \mu^2 \int_D r^{-2} |v|^p + 2\mu \int_D r^{-1} \frac{\partial v}{\partial r} |v|^{p-2} \bar{v} + z \int_D |v|^p = \int_D g |v|^{p-2} \bar{v}. \end{aligned} \quad (2.29)$$

By taking the real part of (2.29) and using (2.28) we obtain

$$\begin{aligned} \int_D |v|^{p-4} \{ (p-1) \left[\left(v_1 \frac{\partial v_1}{\partial r} + v_2 \frac{\partial v_2}{\partial r} \right)^2 + \left(v_1 \frac{1}{r} \frac{\partial v_1}{\partial \theta} + v_2 \frac{1}{r} \frac{\partial v_2}{\partial \theta} \right)^2 \right] \right. \\ \left. + \left[\left(v_2 \frac{\partial v_1}{\partial r} - v_1 \frac{\partial v_2}{\partial r} \right)^2 + \left(v_2 \frac{1}{r} \frac{\partial v_1}{\partial \theta} - v_1 \frac{1}{r} \frac{\partial v_2}{\partial \theta} \right)^2 \right] \right\} \\ - \mu^2 \int_D r^{-2} |v|^p + \Re(z) \int_D |v|^p = \Re \left(\int_D g |v|^{p-2} \bar{v} \right). \end{aligned} \quad (2.30)$$

Denoting $w = (v_1^2 + v_2^2)^{p/4} = |v|^{p/2}$, then (2.30) gives

$$\begin{aligned} \frac{4(p-1)}{p^2} \int_D \left[\left(\frac{\partial w}{\partial r} \right)^2 + \left(\frac{1}{r} \frac{\partial w}{\partial \theta} \right)^2 \right] - \mu^2 \int_D r^{-2} w^2 \\ + \Re(z) \int_D w^2 \leq \Re \left(\int_D g |v|^{p-2} \bar{v} \right). \end{aligned} \quad (2.31)$$

By Lemma 2.11, we have

$$\frac{4(p-1)}{p^2} \int_D |\nabla w|^2 - \mu^2 \int_D r^{-2} w^2 \geq \left(\frac{4(p-1)\lambda^2}{p^2} - \mu^2 \right) \int_D r^{-2} w^2.$$

Hence, we deduce from (2.31) that

$$\Re(z) \|v\|_{L^p}^p \leq \|g\|_{L^p} \|v\|_{L^p}^{p-1},$$

in case

$$\frac{4(p-1)\lambda^2}{p^2} - \mu^2 \geq 0.$$

This gives the first inequality.

As $\frac{4(p-1)\lambda^2}{p^2} - \mu^2 > 0$ and $\Re(z) \geq 0$, we also have

$$\int_D w^2 \frac{1}{r^2} = \int_D |v|^p \frac{1}{r^2} \lesssim \|g\|_{L^p} \|v\|_{L^p}^{p-1}, \quad (2.32)$$

$$\begin{aligned} \int_D |\nabla w|^2 &= \frac{p^2}{4} \int_D |v|^{p-4} \left[\left(v_1 \frac{\partial v_1}{\partial r} + v_2 \frac{\partial v_2}{\partial r} \right)^2 + \left(v_1 \frac{1}{r} \frac{\partial v_1}{\partial \theta} + v_2 \frac{1}{r} \frac{\partial v_2}{\partial \theta} \right)^2 \right] \\ &\lesssim \|g\|_{L^p} \|v\|_{L^p}^{p-1} \end{aligned} \quad (2.33)$$

and

$$\int_D |v|^{p-4} \left[\left(v_2 \frac{\partial v_1}{\partial r} - v_1 \frac{\partial v_2}{\partial r} \right)^2 + \left(v_2 \frac{1}{r} \frac{\partial v_1}{\partial \theta} - v_1 \frac{1}{r} \frac{\partial v_2}{\partial \theta} \right)^2 \right] \lesssim \|g\|_{L^p} \|v\|_{L^p}^{p-1}. \quad (2.34)$$

By taking the imaginary part of (2.29) we obtain

$$\begin{aligned} (p-2) \int_D |v|^{p-4} &\left[v_1^2 \frac{\partial v_1}{\partial r} \frac{\partial v_2}{\partial r} - v_2^2 \frac{\partial v_1}{\partial r} \frac{\partial v_2}{\partial r} - v_1 v_2 \left(\frac{\partial v_1}{\partial r} \right)^2 + v_1 v_2 \left(\frac{\partial v_2}{\partial r} \right)^2 \right. \\ &\quad \left. + v_1^2 \frac{1}{r} \frac{\partial v_1}{\partial \theta} \frac{1}{r} \frac{\partial v_2}{\partial \theta} - v_2^2 \frac{1}{r} \frac{\partial v_1}{\partial \theta} \frac{1}{r} \frac{\partial v_2}{\partial \theta} - v_1 v_2 \left(\frac{1}{r} \frac{\partial v_1}{\partial \theta} \right)^2 + v_1 v_2 \left(\frac{1}{r} \frac{\partial v_2}{\partial \theta} \right)^2 \right] \\ &+ 2\mu \int_D r^{-1} |v|^{p-2} \left(v_1 \frac{\partial v_2}{\partial r} - v_2 \frac{\partial v_1}{\partial r} \right) + \Im(z) \int_D |v|^p = \Im \left(\int_D g |v|^{p-2} \bar{v} \right). \end{aligned}$$

Hence, we deduce from (2.32)-(2.33)-(2.34) that

$$|\Im(z)| \int_D |v|^p \lesssim \|g\|_{L^p} \|v\|_{L^p}^{p-1}$$

and hence

$$|\Im(z)| \|v\|_{L^p} \lesssim \|g\|_{L^p}.$$

The conclusion follows observing that $\|v\|_{L^p} = \|u\|_{L_\mu^p}$ and $\|g\|_{L^p} = \|f\|_{L_\mu^p}$. \square

Before going on let us show that this result is mainly optimal.

Lemma 2.13. *Let D and $\lambda = \frac{\pi}{\psi}$ as in Corollary 2.12, and let $p \geq 2$ and $\mu \in \mathbb{R}$. If*

$$\lambda^2 < 2\left(1 - \frac{1}{p}\right)^2, \quad (2.35)$$

then, for all $\mu \in]\sqrt{2(1 - \frac{1}{p})^2 + \lambda^2}, 2 - \frac{2}{p}[$, there exist $f \in L_\mu^p(D)$, $z \in \mathbb{R}_+$ and $u \in D(\Delta_{p,\mu})$ solution of (2.26) such that $r^\mu u \in V_{2-\frac{2}{p}}^{2,p}(D)$ and

$$z \|u\|_{L_\mu^p(D)} > \|f\|_{L_\mu^p(D)}. \quad (2.36)$$

Proof. Take $p \geq 2$, $z \in \mathbb{R}_+$, and $u \in D(\Delta_{p,\mu})$ such that $r^\mu u \in V_{2-\frac{2}{p}}^{2,p}(D)$, $u > 0$ on D and $\|u\|_{L_\mu^p(D)} = 1$. Let us consider the function $q : \mathbb{R}_0^+ \rightarrow \mathbb{R}^+$ defined by

$$q(X) = \frac{\| -X\Delta u + u \|_{L_\mu^p(D)}^p}{\|u\|_{L_\mu^p(D)}^p},$$

in order that

$$\frac{\| -\Delta u + zu \|_{L_\mu^p(D)}^p}{z^p \|u\|_{L_\mu^p(D)}^p} = q(1/z).$$

Clearly we have $q(0) = 1$ and therefore if q satisfies

$$q'(0) < 0,$$

then there exists a positive real number X close enough to 0 (and then a positive real number z large enough) such that $q(X) < 1$ which will show (2.36).

In view of the definition of q , we directly see that

$$q'(0) = p \int_D u^{p-1} (-\Delta u) r^{\mu p} dx.$$

Hence we are reduced to find a non zero positive function $u \in D(\Delta_{p,\mu})$ such that

$$\int_D u^{p-1} (-\Delta u) r^{\mu p} dx < 0. \quad (2.37)$$

For that purpose, according to the proof of the previous Corollary and to Remark 2.3 we take

$$u(r, \theta) = \varphi(r) \sin(\lambda\theta),$$

September 2, 2010

with $\varphi(r) = r^{-\mu+\beta}(1-r)^\alpha$. Then we see that $u \in H_0^1(D)$ if

$$\mu < \beta \quad \text{and} \quad \alpha > 1/2. \quad (2.38)$$

On the other hand by direct calculation we check that $\Delta u \in L_\mu^p(D)$ if

$$\beta > 2 - \frac{2}{p} \quad \text{and} \quad \alpha > 2 - \frac{1}{p}, \quad (2.39)$$

and $r^\mu u \in V_{2-\frac{2}{p}}^{2,p}(D)$ if

$$\beta > 0 \quad \text{and} \quad \alpha > 2 - \frac{1}{p}.$$

Now we come back to (2.37). By direct calculation, (2.37) reduces to

$$\begin{aligned} I &:= -[(\beta - \mu)^2 - \lambda^2] \int_0^1 r^{\beta p - 2} (1 - r)^{\alpha p} r \, dr \\ &\quad + [2(\beta - \mu) + 1] \alpha \int_0^1 r^{\beta p - 1} (1 - r)^{\alpha p - 1} r \, dr \\ &\quad - [\alpha(\alpha - 1)] \int_0^1 r^{\beta p} (1 - r)^{\alpha p - 2} r \, dr \\ &< 0 \end{aligned} \quad (2.40)$$

By using the definition of the Beta function, we get that

$$I = -\text{B}(p\beta, p\alpha + 1)[(\beta - \mu)^2 - \lambda^2] + \text{B}(p\beta + 1, p\alpha)[2(\beta - \mu) + 1]\alpha - \text{B}(p\beta + 2, p\alpha - 1)[\alpha(\alpha - 1)].$$

Finally the relation $\text{B}(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$ and the recurrence relation $\Gamma(z + 1) = z\Gamma(z)$ lead to

$$\begin{aligned} I &= \alpha p \frac{\Gamma(p\beta)\Gamma(p\alpha - 1)}{\Gamma(p\alpha + p\beta + 1)} \\ &\quad \left(\{ -[(\beta - \mu)^2 - \lambda^2] + [2(\beta - \mu) + 1]\beta \} (p\alpha - 1) - (\alpha - 1)(p\beta + 1)\beta \right). \end{aligned}$$

In conclusion we are looking for parameter α , β and μ satisfying

$$\beta > \max\left(\mu, 2 - \frac{2}{p}\right), \quad \alpha > 2 - \frac{1}{p}$$

and

$$[(p-1)\beta + p(\lambda^2 - \mu^2)]\alpha + (p-1)\beta^2 - (\lambda^2 - \mu^2) < 0.$$

Hence, if $2(p-1)^2 + p^2(\lambda^2 - \mu^2) < 0$, then for $\beta > 2 - \frac{2}{p}$ close enough to $2 - \frac{2}{p}$, $\mu \leq 2 - \frac{2}{p}$ and α large, the above conditions are satisfied. \square

Remark 2.4 Observe that, in the notations of the previous proof, if $2(p-1)^2 + p^2(\lambda^2 - \mu^2) \geq 0$, $\beta > 2 - \frac{2}{p}$ and $\alpha > 2 - \frac{1}{p}$ then we have

$$[(p-1)\beta + p(\lambda^2 - \mu^2)]\alpha + (p-1)\beta^2 - (\lambda^2 - \mu^2) \geq 0.$$

In order to go further, we need estimates like in Corollary 2.12 for z in a larger part of the complex plane.

Corollary 2.14. *Let $D = \{(r \cos \theta, r \sin \theta) \mid 0 < r < 1, 0 < \theta < \psi\}$, $\lambda = \frac{\pi}{\psi}$, $p \geq 2$, $\mu > -\lambda$ satisfy (2.2) and (2.25). Then there exists $\theta_A \in]\frac{\pi}{2}, \pi[$ such that, for all $f \in L_\mu^p(D)$, all $z \in \mathbb{C}$ with $|\arg z| \leq \theta_A$ and $u \in D(\Delta_{p,\mu})$ solution of (2.26), we have the estimate*

$$|z| \|u\|_{L_\mu^p(D)} \lesssim \|f\|_{L_\mu^p(D)}.$$

Proof. Observe that by Lemma 2.7, we have that $D(\Delta_{p,\mu}) \subset L_\mu^p(D)$. Hence we can define the operator

$$A : D(\Delta_{p,\mu}) \subset L_\mu^p(D) \rightarrow L_\mu^p(D) : u \mapsto -\Delta u.$$

First Lemma 2.9 guarantees that, for all $z > 0$, the range of $A + zI$ is $L_\mu^p(D)$. Second by Corollary 2.12 we have, for all $z > 0$,

$$z \|(A + zI)^{-1}\| \leq 1.$$

By [18, Thm I-4.2], this implies that $-A$ is dissipative. As $L_\mu^p(D)$ is reflexive, we have by [18, Thm I-4.6] that $D(\Delta_{p,\mu})$ is dense in $L_\mu^p(D)$. Hence by Hille-Yosida Theorem [18, Thm I-3.1], $-A$ is the infinitesimal generator of a C_0 semigroup of contractions $T(t)$ for $t \geq 0$.

By Corollary 2.12, we have also a constant $C > 0$ such that, for all $\sigma \geq 0$ and all $\tau \in \mathbb{R}$

$$|\tau| \|(A + (\sigma + i\tau)I)^{-1}\| \leq C,$$

hence, by [18, Thm II-5.2], there exists $\delta \in]0, \frac{\pi}{2}[$ and $M > 0$ such that

$$\rho(-A) \supset \Sigma := \{z \in \mathbb{C} \mid |\arg z| < \frac{\pi}{2} + \delta\} \cup \{0\},$$

September 2, 2010

and, for all $z \in \Sigma$,

$$|z| \|(A + zI)^{-1}\| \leq M.$$

This proves the required inequality. \square

Lemma 2.15. *Let $\tilde{\Omega}$ be a bounded domain with a smooth boundary and $p \geq 2$. Then there exists $\tilde{\theta}_A \in]\frac{\pi}{2}, \pi[$ such that, for all $h \in L^p(\tilde{\Omega})$, all $z \in \mathbb{C}$ with $|\arg z| \leq \tilde{\theta}_A$, and $v \in H_0^1(\tilde{\Omega})$ weak solution of*

$$\begin{cases} -\Delta v + z v = h, & \text{in } \tilde{\Omega}, \\ v = 0, & \text{on } \partial\tilde{\Omega}, \end{cases}$$

we have

$$(1 + |z|) \|v\|_{L^p(\tilde{\Omega})} \lesssim \|h\|_{L^p(\tilde{\Omega})}.$$

Proof. Recall that v satisfies

$$\forall \varphi \in H_0^1(\tilde{\Omega}), \quad \int_{\tilde{\Omega}} \nabla v \cdot \nabla \bar{\varphi} + z \int_{\tilde{\Omega}} v \bar{\varphi} = \int_{\tilde{\Omega}} h \bar{\varphi}.$$

Step 1: $\Re z \geq 0$. By [1, Theorem 1.6], we have

$$|z| \|v\|_{L^p(\tilde{\Omega})} \lesssim \|h\|_{L^p(\tilde{\Omega})}.$$

As moreover

$$\|v\|_{L^p(\tilde{\Omega})} \lesssim \|v\|_{H_0^1(\tilde{\Omega})} \lesssim \|h\|_{L^p(\tilde{\Omega})},$$

we obtain the result.

Step 2. Extension. The extension to $\{z \in \mathbb{C} \mid |\arg z| \leq \tilde{\theta}_A\}$ can be made as in Corollary 2.14. \square

Corollary 2.16. *Under the assumptions of Corollary 2.14, for all $f \in L_{-\bar{\mu}}^q(\Omega)$, there exists a unique $\phi \in L_{-\bar{\mu}}^q(\Omega)$ solution of*

$$\forall v \in D(\Delta_{p, \bar{\mu}}), \quad \int_{\Omega} \phi ((-\Delta + \bar{z})v) = \int_{\Omega} f \bar{v}.$$

Moreover ϕ satisfies

$$|z| \|\phi\|_{L_{-\bar{\mu}}^q(\Omega)} \lesssim \|f\|_{L_{-\bar{\mu}}^q(\Omega)}.$$

Proof. By Corollary 2.14 and Lemma 2.15, the linear operator

$$A : D(\Delta_{p,\vec{\mu}}) \subset L_{\vec{\mu}}^p(\Omega) \rightarrow L_{\vec{\mu}}^p(\Omega) : u \mapsto -\Delta u$$

satisfies, for $z \in \mathbb{C}$ with $|\arg(z)| \leq \theta_A$,

$$|z| \|(A + zI)^{-1}f\|_{L_{\vec{\mu}}^p(\Omega)} \lesssim \|f\|_{L_{\vec{\mu}}^p(\Omega)}.$$

Moreover $D(\Delta_{p,\vec{\mu}})$ is dense in $L_{\vec{\mu}}^p(\Omega)$ (because $\mathcal{D}(\Omega) \subset D(\Delta_{p,\vec{\mu}})$). Hence we can define the adjoint A^* of the operator A

$$A^* : D(A^*) \subset (L_{\vec{\mu}}^p(\Omega))^* \rightarrow (L_{\vec{\mu}}^p(\Omega))^*$$

where

$$D(A^*) = \{x^* \in (L_{\vec{\mu}}^p(\Omega))^* \mid \exists y^* \in (L_{\vec{\mu}}^p(\Omega))^*, \forall x \in D(A), \int_{\Omega} x^* \overline{Ax} = \int_{\Omega} y^* \overline{x}\}$$

$$A^*x^* = y^*, \text{ for } x^* \in D(A^*).$$

For $z \in \mathbb{C}$ with $|\arg(z)| \leq \theta_A$, we have $z \in \rho(A)$ and by [18, Lemma I-10.2], $\bar{z} \in \rho(A^*)$ and

$$(\bar{z}I + A^*)^{-1} = ((zI + A)^{-1})^*.$$

The result follows from [18, Lemma I-10.1] observing that A^* is such that $u \in D(A^*) \subset L_{-\vec{\mu}}^q(\Omega)$ if and only if there exists $g \in L_{-\vec{\mu}}^q(\Omega)$ such that

$$\forall v \in D(\Delta_{p,\vec{\mu}}), \quad \int_{\Omega} g \bar{v} = - \int_{\Omega} u \overline{\Delta v}.$$

and, for $u \in D(A^*)$, in the above notations, $A^*(u) = g$. □

3 Uniform decomposition

Theorem 3.1. *Let $R > 0$ be fixed. Let $p \geq 2$, $\epsilon > 0$ fixed and Ω be a bounded polygonal domain of \mathbb{R}^2 . Denote by $S_j, j = 1, \dots, J$, the vertices of $\partial\Omega$. Set*

$$\Omega_{\epsilon} = \{x \in \Omega \mid \forall j = 1, \dots, J, \quad \text{dist}(x, S_j) > \epsilon\},$$

and let $\vec{\mu} \in \mathbb{R}^J$.

Then, there exists $\theta_A \in]\frac{\pi}{2}, \pi[$ such that, for all $g \in L_{\vec{\mu}}^p(\Omega)$, all $z \in \pi^+ \cup S_A$, the solution $u \in H_0^1(\Omega)$ of (2.3) satisfies $u \in W^{2,p}(\Omega_{\epsilon})$ and

$$|u|_{W^{2,p}(\Omega_{\epsilon})} + (1 + |z|^{1/2}) |u|_{W^{1,p}(\Omega_{\epsilon})} + (1 + |z|) |u|_{L^p(\Omega_{\epsilon})} \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}. \quad (3.1)$$

Proof. Let θ_A be given by Corollary 2.15.

September 2, 2010

Step 1: Regularity H^2 . Let us fix η a cut-off function such that $\eta = 0$ on $\bigcup_{j=1, \dots, J} B(S_j, \epsilon/2)$ and $\eta = 1$ on Ω_ϵ . Let $\tilde{\Omega}$ a regular domain such that $\Omega_\epsilon \subset \tilde{\Omega} \subset \bar{\Omega}$, the boundary of $\tilde{\Omega}$ satisfying $\partial\tilde{\Omega} = \partial\Omega$ except near the corner, and $v = \eta u \in H_0^1(\tilde{\Omega})$. In that case v is a weak solution of

$$\begin{cases} -\Delta v + z v = \eta g - 2\nabla\eta \cdot \nabla u - \Delta\eta u =: h, & \text{in } \tilde{\Omega}, \\ v = 0, & \text{on } \partial\tilde{\Omega}. \end{cases} \quad (3.2)$$

By Lemma 2.4 and Poincaré inequality, we have $h \in L^2(\tilde{\Omega})$ and satisfies

$$\|h\|_{L^2(\tilde{\Omega})} \lesssim \|g\|_{L^p_\mu(\Omega)} + \|\nabla u\|_{L^2(\tilde{\Omega})} + \|u\|_{L^2(\tilde{\Omega})} \lesssim \|g\|_{L^p_\mu(\Omega)}. \quad (3.3)$$

By Lemma 2.4 and Corollary 2.6 we obtain

$$|v|_{H_0^1(\tilde{\Omega})} + (1 + |z|) \|v\|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})}. \quad (3.4)$$

We can also consider $v \in H_0^1(\tilde{\Omega})$ as a weak solution of

$$\begin{cases} -\Delta v + v = h + (1 - z) v =: h_1, & \text{in } \tilde{\Omega}, \\ v = 0, & \text{on } \partial\tilde{\Omega}, \end{cases}$$

where $h_1 \in L^2(\tilde{\Omega})$ and, by (3.4),

$$\|h_1\|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})} + (1 + |z|) \|v\|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})} \lesssim \|g\|_{L^p_\mu(\Omega)}.$$

By the strong ellipticity of $-\Delta$ and the fact that the boundary conditions cover $-\Delta$ we have (see [4]), $v \in H^2(\tilde{\Omega})$ and

$$\|v\|_{H^2(\tilde{\Omega})} \lesssim \|h_1\|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})}. \quad (3.5)$$

As $v \in H^2(\tilde{\Omega})$ we have $\Delta v \in L^2(\tilde{\Omega})$. Multiplying (3.2) by $-\Delta\bar{v}$ and integrating, we obtain

$$z \int_{\tilde{\Omega}} |\nabla v|^2 = -z \int_{\tilde{\Omega}} v \Delta\bar{v} = - \int_{\tilde{\Omega}} h \Delta\bar{v} - \int_{\tilde{\Omega}} |\Delta v|^2.$$

This implies, using (3.5),

$$|z| \int_{\tilde{\Omega}} |\nabla v|^2 \leq \|h\|_{L^2(\tilde{\Omega})} \|\Delta\bar{v}\|_{L^2(\tilde{\Omega})} + \|\Delta\bar{v}\|_{L^2(\tilde{\Omega})}^2 \lesssim \|h\|_{L^2(\tilde{\Omega})}^2.$$

Combining (3.3), (3.4) and (3.5) we prove the inequality

$$|v|_{H^2(\tilde{\Omega})} + (1 + \sqrt{|z|}) |v|_{H^1(\tilde{\Omega})} + (1 + |z|) |v|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})} \lesssim \|g\|_{L^p_{\mu}(\Omega)}.$$

As $v = u$ on Ω_{ϵ} , this implies $u \in H^2(\Omega_{\epsilon})$ and

$$|u|_{H^2(\Omega_{\epsilon})} + (1 + \sqrt{|z|}) |u|_{H^1(\Omega_{\epsilon})} + (1 + |z|) |u|_{L^2(\Omega_{\epsilon})} \lesssim \|g\|_{L^p_{\mu}(\Omega)}.$$

Step 2: Regularity $W^{2,p}(\Omega_{\epsilon})$. As in Step 1, let η_1 be a cut-off function such that $\eta_1 = 0$ near the corners and $\eta_1 = 1$ on $\tilde{\Omega}$. Let Ω_1 a regular domain such that $\tilde{\Omega} \subset \Omega_1 \subset \bar{\Omega}$, the boundary of Ω_1 satisfying $\partial\Omega_1 = \partial\Omega$ except near the corner, and $w = \eta_1 u \in H_0^1(\Omega_1)$. In that case w is a weak solution of

$$\begin{cases} -\Delta w + z w = \eta_1 g - 2\nabla\eta_1 \cdot \nabla u - \Delta\eta_1 u =: \tilde{h}_1, & \text{in } \Omega_1, \\ w = 0, & \text{on } \partial\Omega_1. \end{cases}$$

By Step 1, we have that $w \in H^2(\Omega_1)$ and

$$|w|_{H^2(\Omega_1)} + (1 + \sqrt{|z|}) |w|_{H^1(\Omega_1)} + (1 + |z|) |w|_{L^2(\Omega_1)} \lesssim \|g\|_{L^p_{\mu}(\Omega)}.$$

As $w = u$ on $\tilde{\Omega}$ this shows $u \in H^2(\tilde{\Omega})$ and

$$|u|_{H^2(\tilde{\Omega})} + (1 + \sqrt{|z|}) |u|_{H^1(\tilde{\Omega})} + (1 + |z|) |u|_{L^2(\tilde{\Omega})} \lesssim \|g\|_{L^p_{\mu}(\Omega)}. \quad (3.6)$$

This implies that

$$h := \eta g - 2\nabla\eta \cdot \nabla u - \Delta\eta u \in L^p(\tilde{\Omega})$$

with

$$\|h\|_{L^p(\tilde{\Omega})} \lesssim \|g\|_{L^p_{\mu}(\Omega)} + |u|_{W^{1,p}(\tilde{\Omega})} + |u|_{L^p(\tilde{\Omega})} \lesssim \|g\|_{L^p_{\mu}(\Omega)}, \quad (3.7)$$

where we have used Sobolev inequality and (3.6).

We can now proceed in the same way as before but with a given function $h \in L^p(\tilde{\Omega})$. Hence $v \in H_0^1(\tilde{\Omega})$ is a weak solution of

$$\begin{cases} -\Delta v + v = h + (1 - z) v =: h_1, & \text{in } \tilde{\Omega}, \\ v = 0, & \text{on } \partial\tilde{\Omega}. \end{cases}$$

By the elliptic regularity, for all $\tilde{p} \in [2, p]$ we have $v \in W^{2,\tilde{p}}(\tilde{\Omega})$ and

$$\|v\|_{W^{2,\tilde{p}}(\tilde{\Omega})} \lesssim \|h_1\|_{L^{\tilde{p}}(\tilde{\Omega})}.$$

September 2, 2010

Using Corollary 2.6, we obtain

$$(1 + |z|) \|v\|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})},$$

which allows to prove in particular

$$\|v\|_{H^2(\tilde{\Omega})} \lesssim \|h_1\|_{L^2(\tilde{\Omega})} \leq \|h\|_{L^2(\tilde{\Omega})} + (1 + |z|) \|v\|_{L^2(\tilde{\Omega})} \lesssim \|h\|_{L^2(\tilde{\Omega})} \leq \|h\|_{L^p(\tilde{\Omega})}.$$

By Corollary 2.15, we deduce that

$$(1 + |z|) \|v\|_{L^p(\tilde{\Omega})} \lesssim \|h\|_{L^p(\tilde{\Omega})}$$

and hence

$$\|h_1\|_{L^p(\tilde{\Omega})} \lesssim \|g\|_{L^p_\mu(\Omega)}.$$

By interpolation (see [7, Thm 1.4.3.3]) we obtain

$$\|v\|_{W^{1,p}(\tilde{\Omega})} \leq \epsilon \|v\|_{W^{2,p}(\tilde{\Omega})} + K \epsilon^{-1} \|v\|_{L^p(\tilde{\Omega})}.$$

Applying this inequality with $\epsilon = \frac{1}{1 + \sqrt{|z|}}$, this gives

$$|v|_{W^{2,p}(\tilde{\Omega})} + (1 + \sqrt{|z|}) |v|_{W^{1,p}(\tilde{\Omega})} + (1 + |z|) |v|_{L^p(\tilde{\Omega})} \lesssim \|g\|_{L^p_\mu(\Omega)}.$$

The result follows as $v = u$ on Ω_ϵ . \square

Lemma 3.2. *Let $R > 0$ and $\theta_A \in]\frac{\pi}{2}, \pi[$ be fixed. Let C be the cone with interior angle ψ and $\lambda = \frac{\pi}{\psi}$. For $z \in \pi^+ \cup S_A$ and $u \in V_\mu^{2,p}(C) \cap H_0^1(C)$ with $u = 0$ for $r > r_0$, if μ satisfies (2.2) and, for all $k \in \mathbb{Z}^*$, $2 - \frac{2}{p} - \mu \neq k\lambda$, then we have*

$$\begin{aligned} |u|_{V_\mu^{2,p}(C)} + |u|_{V_{\mu-1}^{1,p}(C)} + \|u\|_{L_{\mu-2}^p(C)} + |z|^{1/2} |u|_{V_\mu^{1,p}(C)} + |z| \|u\|_{L_\mu^p(C)} \\ \lesssim \|(-\Delta + z)u\|_{L_\mu^p(C)} + |z| \|u\|_{L_\mu^p(C)}. \end{aligned}$$

Proof. By [14, Thm 6.2], we have, for $\theta \in [-\theta_A, \theta_A]$ and $u \in E_\mu^{2,p}(C)$,

$$\|u\|_{E_\mu^{2,p}(C)} \lesssim \|(-\Delta + e^{i\theta})u\|_{L_\mu^p(C)} + \|u\|_{L^p(S)}, \quad (3.8)$$

where $S = \{x \in \mathbb{R}^2 \mid \delta_1 < |x| < \delta_2\} \cap C$ with $0 < \delta_1 < \delta_2$, $E_\mu^{2,p}(C) = \text{adh}_{\|\cdot\|_{E_\mu^{2,p}}}(\mathcal{C}_0^\infty(\overline{C} \setminus \{0\}))$ where

$$\|u\|_{E_\mu^{2,p}} = \left(\int_C r^{p\mu} \sum_{|\alpha|=0}^2 (r^{p(|\alpha|-2)} + 1) |D_y^\alpha u(y)|^p \right)^{1/p}.$$

As $\|u\|_{L^p(S)} \lesssim \|u\|_{L_\mu^p(C)}$, the result can be deduced from a change of variable. \square

Theorem 3.3. *Let $R > 0$, $p \geq 2$, Ω be a bounded polygonal domain of \mathbb{R}^2 . Denote by $S_j, j = 1, \dots, J$, the vertices of $\partial\Omega$ enumerated clockwise and, for $j \in \{1, 2, \dots, J\}$, let ψ_j be the interior angle of Ω at the vertex S_j , $\lambda_j = \frac{\pi}{\psi_j}$ and $\vec{\lambda} = (\lambda_j)_{1 \leq j \leq J}$.*

Let $\vec{\mu} > -\vec{\lambda}$ satisfies (2.2), (2.25) and, for all $k \in \mathbb{Z}^$ and all $j \in \{1, 2, \dots, J\}$, $2 - \frac{2}{p} - \mu_j \neq k\lambda_j$.*

Then, there exists $\theta_A \in]\frac{\pi}{2}, \pi[$ such that, for all $g \in L_{\vec{\mu}}^p(\Omega)$, all $z \in \pi^+ \cup S_A$, the unique solution $u \in D(\Delta_{p, \vec{\mu}})$ of

$$\begin{cases} -\Delta u + zu = g, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

admits the decomposition

$$u = u_R + \sum_{j=1}^J \eta_j(r) \sum_{\substack{0 < \lambda'_j < 2 - \frac{2}{p} - \mu_j \\ \exists k \in \mathbb{N}, \lambda'_j = k\lambda_j}} c_{\lambda'_j}(z) P_{j, \lambda'_j}(r\sqrt{z}) e^{-r\sqrt{z}} r^{\lambda'_j} \sin(\lambda'_j \theta), \quad (3.9)$$

with $u_R \in V_{\vec{\mu}}^{2,p}(\Omega)$, $c_{\lambda'_j}(z) \in \mathbb{C}$, $\eta_j \in \mathcal{D}(\mathbb{R}^2)$ is a cut-off function such that

$$\eta_j \equiv 1 \text{ in } D_j(1/2), \quad \eta_j \equiv 0 \text{ on } \Omega \setminus D_j(1),$$

and $P_{j, \lambda'_j}(s) = \sum_{i=0}^{l_{j, \lambda'_j}-1} \frac{s^i}{i!}$ with $l_{j, \lambda'_j} > 2 - \mu_j - \frac{2}{p} - \lambda'_j$.

Moreover, the following inequalities are satisfied

- (a) $|u_R|_{V_{\vec{\mu}}^{2,p}(\Omega)} + |u_R|_{V_{\vec{\mu}-1}^{1,p}(\Omega)} + |u_R|_{L_{\vec{\mu}-2}^p(\Omega)} \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}$;
- (b) $|u_R|_{V_{\vec{\mu}}^{2,p}(\Omega)} + |z|^{1/2} |u_R|_{V_{\vec{\mu}}^{1,p}(\Omega)} + |z| |u_R|_{L_{\vec{\mu}}^p(\Omega)} \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}$;
- (c) $\sum_{j=1}^J \sum_{\substack{0 < \lambda'_j < 2 - \frac{2}{p} - \mu_j \\ \exists k \in \mathbb{N}, \lambda'_j = k\lambda_j}} |c_{\lambda'_j}(z)| (1 + |z|^{1 - \frac{1}{p} - \frac{\mu_j + \lambda'_j}{2}}) \lesssim \|g\|_{L_{\vec{\mu}}^p(\Omega)}$.

Proof. Let θ_A be the minimum of the ones obtained by Theorem 3.1 and by Corollary 2.14. By Theorem 3.1, we have the required result in the interior of Ω . It remains to prove the regularity near the corners.

Let $j \in \{1, 2, \dots, J\}$ be fixed and observe that $v = \eta_j u$ is the solution of

$$\begin{cases} -\Delta v + zv = g_1, & \text{in } D_j, \\ v = 0, & \text{on } \partial D_j, \end{cases} \quad (3.10)$$

September 2, 2010

with $D_j = \{(r_j \cos \theta_j, r_j \sin \theta_j) \mid 0 < r_j < 1, 0 < \theta_j < \psi_j\}$ and $g_1 = \eta_j g - 2\nabla\eta_j \cdot \nabla u - \Delta\eta_j u$. By Theorem 3.1, we have

$$\nabla u \in W^{1,p}(\Omega_\epsilon), \quad u \in L^p(\Omega_\epsilon),$$

and $g_1 \in L^p_{\mu_j}(D_j)$ with

$$\|g_1\|_{L^p_{\mu_j}(D_j)} \lesssim \|g\|_{L^p_\mu(\Omega)}. \quad (3.11)$$

Hence we can concentrate on (3.10). For the rest of the proof we drop the index j and write D for D_j , μ for μ_j , ...

Step 1: $v \in V_1^{2,2}(D)$ and

$$|v|_{V_1^{2,2}(D)} + |v|_{V_0^{1,2}(D)} + |v|_{V_{-1}^{0,2}(D)} \lesssim \|g_1\|_{L_1^2(D)} \lesssim \|g\|_{L^p_\mu(\Omega)}. \quad (3.12)$$

Using Lemma 2.4 and Corollary 2.5, we have

$$\begin{aligned} \|g_1\|_{L_1^2(D)} &\lesssim \|rg\|_{L^2(D)} + \|u\|_{H^1(\Omega)} \\ &\lesssim \left(\int_D |r|^{\frac{2(1-\mu)p}{p-2}} \right)^{\frac{p-2}{2p}} \|r^\mu g\|_{L^p(D)} + \|u\|_{H^1(\Omega)} \lesssim \|g\|_{L^p_\mu(\Omega)}. \end{aligned}$$

This proves

$$\|g_1\|_{L_1^2(D)} \lesssim \|g\|_{L^p_\mu(\Omega)}. \quad (3.13)$$

Moreover v can be considered as a solution of

$$\begin{cases} -\Delta v = g_1 - z v =: g_2, & \text{in } D, \\ v = 0, & \text{on } \partial D. \end{cases} \quad (3.14)$$

By Lemma 2.7, $v \in L^p_\mu(D)$ and by Corollary 2.14, we obtain

$$\|g_2\|_{L^p_\mu(D)} \lesssim \|g_1\|_{L^p_\mu(D)} + |z| \|v\|_{L^p_\mu(D)} \lesssim \|g_1\|_{L^p_\mu(D)} \lesssim \|g\|_{L^p_\mu(\Omega)}. \quad (3.15)$$

Applying [13, Lemma 11.2(ii)] (as in [13, Example 11.3]), this allows to conclude $v \in V_1^{2,2}(D)$ and

$$\|v\|_{V_1^{2,2}(D)} = |v|_{V_1^{2,2}(D)} + |v|_{V_0^{1,2}(D)} + |v|_{V_{-1}^{0,2}(D)} \lesssim \|g_2\|_{L_1^2(D)} \lesssim \|g\|_{L^p_\mu(\Omega)}.$$

The case $p = 2$ and $\mu = 1$ is proved.

Step 2: L^p -decomposition in case $p > 2$ or in case $p = 2$ and $\mu < 1$. As $\mu < \frac{2p-2}{p}$ and, by Step 1, $v \in V_1^{2,2}(D)$, applying [13, Cor (iv) of Thm 10.2 and Thm 10.3], we have

$$v = v_R^{(1)} + \sum_{\substack{0 < \lambda' < 2 - \frac{2}{p} - \mu \\ \exists k \in \mathbb{N}, \lambda' = k\lambda}} c_{\lambda'}(z) \eta r^{\lambda'} \sin(\lambda' \theta) =: v_R^{(1)} + v_S \quad (3.16)$$

with $v_R^{(1)} \in V_\mu^{2,p}(D)$ satisfying, by (3.15),

$$\|v_R^{(1)}\|_{V_\mu^{2,p}(D)} \lesssim \|g_2\|_{L_\mu^p(D)} \lesssim \|g_1\|_{L_\mu^p(D)} \lesssim \|g\|_{L_\mu^p(\Omega)}. \quad (3.17)$$

As $\mu < \frac{2p-2}{p}$, by [13, section 0.10], we have $V_\mu^{2,p}(D) \hookrightarrow V_1^{2,2}(D)$ and $L_\mu^p(D) \hookrightarrow L_1^2(D)$. Hence, using (3.12), (3.15) and (3.17), we obtain

$$\|v_S\|_{V_1^{2,2}(D)} \lesssim \|v\|_{V_1^{2,2}(D)} + \|v_R^{(1)}\|_{V_\mu^{2,p}(D)} \lesssim \|g_1\|_{L_\mu^p(D)} \lesssim \|g\|_{L_\mu^p(\Omega)}.$$

The space $V_S = \text{span}\{r^{\lambda'} \sin(\lambda' \theta) \mid 0 < \lambda' < 2 - \frac{2}{p} - \mu \text{ and } \exists k \in \mathbb{N}, \lambda' = k\lambda\}$ being of finite dimension, we have

$$\sum_{\substack{0 < \lambda' < 2 - \frac{2}{p} - \mu \\ \exists k \in \mathbb{N}, \lambda' = k\lambda}} |c_{\lambda'}(z)| \lesssim \|v_S\|_{V_1^{2,2}(D)} \lesssim \|g_1\|_{L_\mu^p(D)}. \quad (3.18)$$

Now in order to have uniform estimates with respect to $|z|$, we choose another decomposition. We rewrite (3.16) in the following way

$$v = v_R + \sum_{\substack{0 < \lambda' < 2 - \frac{2}{p} - \mu \\ \exists k \in \mathbb{N}, \lambda' = k\lambda}} c_{\lambda'}(z) \eta(r) \psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda' \theta), \quad (3.19)$$

where $v_R = v_R^{(1)} + \sum_{\substack{0 < \lambda' < 2 - \frac{2}{p} - \mu \\ \exists k \in \mathbb{N}, \lambda' = k\lambda}} c_{\lambda'}(z) \eta(r) (1 - \psi_{\lambda'}(r\sqrt{z})) r^{\lambda'} \sin(\lambda' \theta)$ and $\psi_{\lambda'}(s) = P_{\lambda'}(s)e^{-s}$ with $P_{\lambda'}(s) = \sum_{i=0}^{l_{\lambda'}-1} \frac{s^i}{i!}$ and $l_{\lambda'} > 2 - \mu - \frac{2}{p} - \lambda'$ is chosen in such a way that $(1 - \psi_{\lambda'}(r\sqrt{z})) r^{\lambda'} \sin(\lambda' \theta) \in V_\mu^{2,p}(D)$.

Step 3: Estimates (c) on $|c_{\lambda'}(z)|$. By (3.18) we know that, for all $z \in \pi^+ \cup S_A$, the mapping

$$L_{\mu}^p(D) \rightarrow \mathbb{C} : g_1 \mapsto c_{\lambda'}(z)$$

is linear and continuous. Hence by Riesz theorem, there exists a unique $w_z^{\lambda'} \in L_{-\mu}^q(D) = (L_{\mu}^p(D))'$ such that

$$c_{\lambda'}(z) = \int_D g_1 \bar{w}_z^{\lambda'} dx.$$

Claim: The function $w_z^{\lambda'}$ is characterized by

(i) for all $v \in V_{\mu}^{2,p}(D) \cap D(\Delta_{p,\mu})$, $\int_D (-\Delta v + z v) \bar{w}_z^{\lambda'} = 0$;

(ii) for all $\xi \in]0, 2 - \frac{2}{p} - \mu[$ such that $\exists k \in \mathbb{N}$, $\lambda' = k\lambda$, we have

$$\int_D (-\Delta + z)(\eta(r) r^{\xi} \sin(\xi\theta)) \bar{w}_z^{\lambda'} = \delta_{\xi\lambda'}.$$

This can be easily deduced from (3.16) as this implies

$$\begin{aligned} v \in V_{\mu}^{2,p}(D) \cap D(\Delta_{p,\mu}) \\ \iff \forall \lambda' \in]0, 2 - \frac{2}{p} - \mu[\text{ with } \exists k \in \mathbb{N}, \lambda' = k\lambda, c_{\lambda'}(z) = 0. \end{aligned}$$

and

$$\int_D (-\Delta + z)(\eta(r) r^{\xi} \sin(\xi\theta)) \bar{w}_z^{\lambda'} = \delta_{\xi\lambda'}.$$

The reversed implications can be deduced by uniqueness of $w_z^{\lambda'}$.

To obtain the good estimate on $c_{\lambda'}(z)$, we will decompose $w_z^{\lambda'}$ in an appropriate way. To this aim define the function

$$\psi^{\lambda'} = A_{\lambda'} e^{-r\sqrt{\bar{z}}} Q(\sqrt{\bar{z}}r) r^{-\lambda'} \sin(\lambda'\theta), \quad (3.20)$$

with $A_{\lambda'} = \frac{1}{\lambda'\psi}$, $Q(s) = \sum_{i=0}^m \alpha_i s^i$, with m fixed below and α_i such that

$$\begin{aligned} \alpha_0 = \alpha_1 = 1, \\ \forall i \in \{1, \dots, m-1\}, \quad (2\lambda' - 1 - 2i) \alpha_i = (i+1)(2\lambda' - i - 1) \alpha_{i+1}. \end{aligned}$$

The positive integer m is fixed as follows: if $\lambda' - 1 \geq 1$ then we take $m > \lambda' - 1 \geq m - 1$, hence we have $m \geq 1$ and for $i \in \{1, \dots, m-1\}$ we have $i+1 \leq m \leq \lambda'$ and hence $i+1 \neq 2\lambda'$. If $\lambda' - 1 < 1$ then we take $m = 1$.

The above choice leads to

$$(-\Delta + \bar{z})(\eta\psi^{\lambda'}) = -\eta A_{\lambda'} (2\lambda' - 1 - 2m)\alpha_m r^{m-\lambda'-1} \bar{z}^{\frac{m+1}{2}} e^{-r\sqrt{\bar{z}}} \sin(\lambda'\theta) \\ + (-2\nabla\psi^{\lambda'} \cdot \nabla\eta - \psi^{\lambda'}\Delta\eta).$$

Let $t > 1$ be such that $(m - \lambda' - 1)t + 2 > 0$, by the above computation, we see that $(-\Delta + \bar{z})(\eta\psi^{\lambda'}) \in L^t(D)$. Hence we can define $\phi^{\lambda'} \in H_0^1(D)$ such that

$$\forall v \in H_0^1(D), \quad \int_D (\nabla\phi^{\lambda'} \cdot \nabla\bar{v} + \bar{z}\phi^{\lambda'}\bar{v}) = \int_D (-\Delta + \bar{z})(\eta\psi^{\lambda'})\bar{v}. \quad (3.21)$$

Let us show that

$$w_z^{\lambda'} = \eta\psi^{\lambda'} - \phi^{\lambda'}. \quad (3.22)$$

To this aim, let us prove that $\eta\psi^{\lambda'} - \phi^{\lambda'}$ satisfies the conditions (i) and (ii) above.

The condition (i) can be deduced from the Green formula, see Lemma 3.5 below. Consider then the condition (ii). By the Green formula (apply [17, Corollary 1.42] and remind that $H_0^1(D) \hookrightarrow L_{-1}^2(D)$ (see Lemma 2.11)) we deduce

$$\int_D (-\Delta + z)(\eta(r)r^\xi \sin(\xi\theta))\bar{\phi}^{\lambda'} \\ = \int_D \nabla(\eta(r)r^\xi \sin(\xi\theta)) \cdot \nabla\bar{\phi}^{\lambda'} + z \int_D \eta(r)r^\xi \sin(\xi\theta)\bar{\phi}^{\lambda'}$$

By definition of $\phi^{\lambda'}$ we obtain

$$\int_D \nabla(\eta(r)r^\xi \sin(\xi\theta)) \cdot \nabla\bar{\phi}^{\lambda'} + z \int_D \eta(r)r^\xi \sin(\xi\theta)\bar{\phi}^{\lambda'} \\ = \int_D \overline{(-\Delta + \bar{z})(\eta\psi^{\lambda'})} \eta(r)r^\xi \sin(\xi\theta).$$

Hence we have

$$\int_D (-\Delta + z)(\eta(r)r^\xi \sin(\xi\theta)) (\eta\bar{\psi}^{\lambda'} - \bar{\phi}^{\lambda'}) \\ = \int_D (-\Delta + z)(\eta(r)r^\xi \sin(\xi\theta)) \eta\bar{\psi}^{\lambda'} \\ - \int_D \overline{(-\Delta + \bar{z})(\eta\psi^{\lambda'})} \eta(r)r^\xi \sin(\xi\theta) \\ = - \int_D \Delta(\eta(r)r^\xi \sin(\xi\theta)) \eta\bar{\psi}^{\lambda'} + \int_D \Delta(\eta\bar{\psi}^{\lambda'}) \eta(r)r^\xi \sin(\xi\theta).$$

Let us denote by $D_\epsilon = D \setminus B(0, \epsilon)$, then we can write by the Green formula

$$\begin{aligned}
& - \int_D \Delta(\eta(r)r^\xi \sin(\xi\theta)) \eta \bar{\psi}^{\lambda'} + \int_D \Delta(\eta \bar{\psi}^{\lambda'}) \eta(r)r^\xi \sin(\xi\theta) \\
&= \lim_{\epsilon \rightarrow 0} \left[- \int_{D_\epsilon} \Delta(\eta(r)r^\xi \sin(\xi\theta)) \eta \bar{\psi}^{\lambda'} + \int_{D_\epsilon} \Delta(\eta \bar{\psi}^{\lambda'}) \eta(r)r^\xi \sin(\xi\theta) \right] \\
&= \lim_{\epsilon \rightarrow 0} \left[- \int_{\partial D_\epsilon} \frac{\partial}{\partial \nu} (\eta(r)r^\xi \sin(\xi\theta)) \eta \bar{\psi}^{\lambda'} + \int_{\partial D_\epsilon} \frac{\partial}{\partial \nu} (\eta \bar{\psi}^{\lambda'}) \eta(r)r^\xi \sin(\xi\theta) \right] \\
&= \lim_{\epsilon \rightarrow 0} \epsilon \int_0^\psi \left[\frac{\partial}{\partial r} (\eta(r)r^\xi \sin(\xi\theta)) \eta \bar{\psi}^{\lambda'} - \frac{\partial}{\partial r} (\eta \bar{\psi}^{\lambda'}) \eta(r)r^\xi \sin(\xi\theta) \right] \Big|_{r=\epsilon} d\theta.
\end{aligned}$$

By definition of $\psi^{\lambda'}$ we deduce that, for $\xi \neq \lambda'$,

$$\int_D (-\Delta + z)(\eta(r)r^\xi \sin(\xi\theta)) \bar{w}_z^{\lambda'} = 0,$$

as $\int_0^\psi \sin(\xi\theta) \sin(\lambda'\theta) d\theta = 0$ if $\xi \neq \lambda'$.

In case $\xi = \lambda'$ we have

$$\begin{aligned}
& \int_D (-\Delta + z)(\eta(r)r^{\lambda'} \sin(\lambda'\theta)) (\eta \bar{\psi}^{\lambda'} - \bar{\phi}^{\lambda'}) \\
&= A_{\lambda'} \frac{\psi}{2} \lim_{\epsilon \rightarrow 0} \left[2\lambda' e^{-\epsilon\sqrt{z}} Q(\sqrt{z}\epsilon) + \sqrt{z} e^{-\epsilon\sqrt{z}} \epsilon (Q(\sqrt{z}\epsilon) - Q'(\sqrt{z}\epsilon)) \right] \\
&= A_{\lambda'} \psi \lambda' = 1.
\end{aligned}$$

This proves that $w_z^{\lambda'} = \eta \psi^{\lambda'} - \phi^{\lambda'}$.

Estimate on $\|w_z^{\lambda'}\|_{L^q_{-\mu}(D)}$. Observe that, as $z \in \pi^+ \cup S_A$, we can write $z = |z|e^{i\theta}$ with $\theta \in [-\theta_A, \theta_A]$ and we have $|e^{-qr\sqrt{z}}| = e^{-qr|z|^{1/2} \cos(\theta/2)} \lesssim e^{-\gamma qr|z|^{1/2}}$ with $\gamma = \cos(\theta_A/2) > 0$. Hence we obtain

$$\|\psi^{\lambda'}\|_{L^q_{-\mu}(D)}^q \lesssim \int_0^1 e^{-\gamma qr|z|^{1/2}} |Q(\sqrt{z}r)|^q r^{(-\lambda' - \mu)q+1} dr.$$

Making the change of variable $s = r|z|^{1/2}$ we obtain

$$\|\psi^{\lambda'}\|_{L^q_{-\mu}(D)}^q \lesssim |z|^{-\frac{1}{2}((-\mu - \lambda')q+2)} \int_0^{+\infty} e^{-\gamma qs} s^{(-\lambda' - \mu)q+1} |Q(s)|^q ds.$$

where the integral $\int_0^{+\infty} e^{-\gamma qs} s^{(-\lambda' - \mu)q+1} |Q(s)|^q ds < +\infty$ as $(-\lambda' - \mu)q + 2 > 0$. Hence we have

$$\|\psi^{\lambda'}\|_{L^q_{-\mu}(D)} \lesssim |z|^{-\frac{1}{q} + \frac{\mu + \lambda'}{2}}. \quad (3.23)$$

Recall that

$$\begin{aligned} (-\Delta + \bar{z})(\eta\psi^{\lambda'}) &= -\eta A_{\lambda'} (2\lambda' - 1 - 2m)\alpha_m r^{m-\lambda'-1} \bar{z}^{\frac{m+1}{2}} e^{-r\sqrt{\bar{z}}} \sin(\lambda'\theta) \\ &\quad + (-2\nabla\psi^{\lambda'} \cdot \nabla\eta - \psi^{\lambda'}\Delta\eta). \end{aligned}$$

Hence, we see that $(-\Delta + \bar{z})(\eta\psi^{\lambda'}) \in L^q_{-\mu}(D)$ as $\lambda' < -\mu + 2 - \frac{2}{p}$ and $m \geq 1$. Making the change of variables $s = r|z|^{1/2}$ we obtain

$$\begin{aligned} \|e^{-r\sqrt{\bar{z}}} r^{m-\lambda'-1} \sin(\lambda'\theta)\|_{L^q_{-\mu}(D)} &\lesssim \int_0^1 |e^{-qr\sqrt{\bar{z}}}| r^{(m-\mu-\lambda'-1)q+1} dr \\ &\lesssim \left(\int_0^{+\infty} e^{-\gamma qs} s^{(m-\mu-\lambda'-1)q+1} ds \right) |z|^{-\frac{q}{2}(m-\mu-\lambda'-1)-1}, \end{aligned}$$

where the integral $\int_0^{+\infty} e^{-\gamma qs} s^{(m-\mu-\lambda'-1)q+1} ds < +\infty$. Hence we obtain

$$\|e^{-r\sqrt{\bar{z}}} r^{m-\lambda'-1} \bar{z}^{\frac{m+1}{2}} \sin(\lambda'\theta)\|_{L^q_{-\mu}(D)} \lesssim |z|^{-\frac{1}{2}((-\mu-\lambda'-2)+\frac{2}{q})}. \quad (3.24)$$

Let us denote by $r_{\lambda'} := (-2\nabla\psi^{\lambda'} \cdot \nabla\eta - \psi^{\lambda'}\Delta\eta)$. For $|z| \geq 1$, we have

$$\begin{aligned} \|r_{\lambda'}\|_{L^q_{-\mu}(D)}^q &\lesssim \int_{r_0}^1 |e^{-rq\sqrt{\bar{z}}}| r^{(-\lambda'-\mu)q+1} |Q(r\sqrt{\bar{z}})|^q dr \\ &\quad + \int_{r_0}^1 |e^{-rq\sqrt{\bar{z}}}| r^{(-\lambda'-\mu-1)q+1} |Q(r\sqrt{\bar{z}})|^q dr \\ &\quad + \int_{r_0}^1 |e^{-rq\sqrt{\bar{z}}}| r^{(-\lambda'-\mu)q+1} |Q(r\sqrt{\bar{z}})|^q |z|^{\frac{q}{2}} dr \\ &\quad + \int_{r_0}^1 |e^{-rq\sqrt{\bar{z}}}| r^{(-\lambda'-\mu-2)q+1} |Q(r\sqrt{\bar{z}})|^q dr \\ &\quad + \int_{r_0}^1 |e^{-rq\sqrt{\bar{z}}}| r^{(-\lambda'-\mu)q+1} |Q'(r\sqrt{\bar{z}})|^q |z|^{\frac{q}{2}} dr \\ &\lesssim \int_{r_0}^1 e^{-\gamma rq|z|^{\frac{1}{2}}} |z|^{\frac{q}{2}} (|Q(r\sqrt{\bar{z}})|^q + |Q'(r\sqrt{\bar{z}})|^q) dr \\ &\quad + \int_{r_0}^1 e^{-\gamma rq|z|^{\frac{1}{2}}} |Q(r\sqrt{\bar{z}})|^q dr \\ &\lesssim e^{-\gamma r_0 q|z|^{\frac{1}{2}}} (|z|^{\frac{(l+1)q}{2}} + 1). \end{aligned}$$

September 2, 2010

Hence we obtain

$$\|r_{\lambda'}\|_{L^q_{-\mu}(D)} \lesssim e^{-|z|^{1/2}\tilde{\gamma}}, \quad (3.25)$$

for some $\tilde{\gamma} > 0$.

We then deduce from (3.24) and (3.25) that

$$\|(-\Delta + \bar{z})(\eta\psi^{\lambda'})\|_{L^q_{-\mu}(D)} \lesssim |z|^{1-\frac{1}{q}+\frac{\mu+\lambda'}{2}}.$$

As $(-\Delta + \bar{z})(\eta\psi^{\lambda'}) \in L^q_{-\mu}(D)$, by Lemma 3.4 below, $\phi^{\lambda'} \in H_0^1(D) \hookrightarrow L^q_{-\mu}(D)$ satisfies the assumptions of Corollary 2.16, hence it satisfies

$$|z| \|\phi^{\lambda'}\|_{L^q_{-\mu}(D)} \lesssim \|(-\Delta + \bar{z})(\eta\psi^{\lambda'})\|_{L^q_{-\mu}(D)} \lesssim |z|^{1-\frac{1}{q}+\frac{\mu+\lambda'}{2}}$$

i.e.

$$\|\phi^{\lambda'}\|_{L^q_{-\mu}(D)} \lesssim |z|^{-\frac{1}{q}+\frac{\mu+\lambda'}{2}}. \quad (3.26)$$

By (3.23) and (3.26) we deduce

$$\|w^{\lambda'}\|_{L^q_{-\mu}(D)} \lesssim |z|^{-\frac{1}{q}+\frac{\mu+\lambda'}{2}}$$

and by Hölder inequality

$$|c_{\lambda'}| = \left| \int_D g_1 \bar{w}^{\lambda'} \right| \leq \|g_1\|_{L^p_{\mu}(D)} \|w^{\lambda'}\|_{L^q_{-\mu}(D)} \lesssim |z|^{-\frac{1}{q}+\frac{\mu+\lambda'}{2}} \|g_1\|_{L^p_{\mu}(D)}.$$

Hence, using (3.11), we conclude that, for $|z| \geq 1$,

$$|c_{\lambda'}| \lesssim |z|^{-\frac{1}{q}+\frac{\mu+\lambda'}{2}} \|g\|_{L^p_{\mu}(\Omega)}. \quad (3.27)$$

which proves (c).

Step 4: Estimates (a) and (b) on v_R . By (3.19) we know that $v_R \in V^{2,p}(C) \cap H_0^1(C)$ with C the cone with interior angle ψ . By Lemma 3.2, to obtain the estimates (a) and (b), it remains to estimate $\|(-\Delta + z)v_R\|_{L^p_{\mu}(C)} + |z| \|v_R\|_{L^p_{\mu}(C)}$. We have

$$\begin{aligned} & \|(-\Delta + z)v_R\|_{L^p_{\mu}(C)} + |z| \|v_R\|_{L^p_{\mu}(C)} \\ & \lesssim \|(-\Delta + z)v\|_{L^p_{\mu}(D)} + |z| \|v\|_{L^p_{\mu}(D)} \\ & \quad + \sum_{\substack{0 < \lambda' < 2 - \frac{2}{p} - \mu \\ \exists k \in \mathbb{N}, \lambda' = k\lambda}} |c_{\lambda'}(z)| \|(-\Delta + z)(\eta(r)\psi_{\lambda'}(r\sqrt{z})r^{\lambda'} \sin(\lambda'\theta))\|_{L^p_{\mu}(D)} \\ & \quad + |z| \sum_{\substack{0 < \lambda' < 2 - \frac{2}{p} - \mu \\ \exists k \in \mathbb{N}, \lambda' = k\lambda}} |c_{\lambda'}(z)| \|\eta(r)\psi_{\lambda'}(r\sqrt{z})r^{\lambda'} \sin(\lambda'\theta)\|_{L^p_{\mu}(D)}. \end{aligned}$$

September 2, 2010

Recall that v is a solution of (3.10) and hence, by Corollary 2.14 and (3.11),

$$\|(-\Delta + z)v\|_{L_\mu^p(D)} + |z| \|v\|_{L_\mu^p(D)} \lesssim \|g_1\|_{L_\mu^p(D)} \lesssim \|g\|_{L_\mu^p(\Omega)}.$$

It remains to estimate the two last terms.

Since $\lambda'p + \mu p + 2 > 0$, we have

$$\begin{aligned} & \|\eta(r) \psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda'\theta)\|_{L_\mu^p(D)}^p \\ & \lesssim \int_0^1 |P_{\lambda'}(r\sqrt{z}) e^{-\gamma r|z|^{\frac{1}{2}}} r^{\lambda'} \sin(\lambda'\theta)|^p r^{\mu p+1} dr \\ & \lesssim \int_0^\infty |P_{\lambda'}(s)|^p e^{-\gamma s p} s^{\lambda'p+\mu p+1} ds |z|^{-\frac{1}{2}(\lambda'p+\mu p+2)} \\ & \lesssim |z|^{-\frac{1}{2}(\lambda'p+\mu p+2)}. \end{aligned}$$

We then deduce from (3.27) that

$$\begin{aligned} & |z| \sum_{0 < \lambda' < 2 - \frac{2}{p} - \mu} |c_{\lambda'}(z)| \|\eta(r) \psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda'\theta)\|_{L_\mu^p(D)} \\ & \lesssim \sum_{0 < \lambda' < 2 - \frac{2}{p} - \mu} |z|^{1 - \frac{1}{q} + \frac{\lambda'+\mu}{2}} \|g\|_{L_\mu^p(\Omega)} |z|^{-\frac{1}{2}(\lambda'+\mu) - \frac{1}{p}} \\ & \lesssim \|g\|_{L_\mu^p(\Omega)}. \end{aligned}$$

Moreover if $l_{\lambda'} \geq 2$, we have

$$\|(-\Delta + z)(\psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda'\theta))\|_{L_\mu^p(D)}^p = \|z e^{-r\sqrt{z}} r^{\lambda'} \sin(\lambda'\theta) \tilde{P}(r\sqrt{z})\|_{L_\mu^p(D)}^p$$

with \tilde{P} a polynomial function of degree $l_{\lambda'} - 2$. As before we obtain

$$\|(-\Delta + z)(\psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda'\theta))\|_{L_\mu^p(D)}^p \lesssim |z|^{-\frac{1}{2}(\lambda'p+\mu p+2-2p)}$$

since $\lambda'p + \mu p + 2 > 0$. Hence we have

$$\|(-\Delta + z)(\psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda'\theta))\|_{L_\mu^p(D)} \lesssim |z|^{1 - \frac{1}{p} - \frac{\lambda'+\mu}{2}}. \quad (3.28)$$

If $l_{\lambda'} = 1$, then we have

$$(-\Delta + z)(\psi_{\lambda'}(r\sqrt{z}) r^{\lambda'} \sin(\lambda'\theta)) = (2\lambda' + 1)\sqrt{z} e^{-r\sqrt{z}} r^{\lambda'-1} \sin(\lambda'\theta),$$

and the same arguments than before yield (3.28) since here $\lambda' - 1 + \mu + \frac{2}{p} > 0$ (reminding that if $l_{\lambda'} = 1$, then $2 - \mu - \frac{2}{p} - \lambda' < 1$).

As the function η does not interfere the result follows from (3.27). \square

Lemma 3.4. *Let $p \geq 2$, $D = \{(r \cos \theta, r \sin \theta) \mid 0 < r < 1, 0 < \theta < \psi\}$ and μ satisfy (2.2). Let $\phi \in H_0^1(D)$ and $v \in D(\Delta_{p,\mu})$ then we have*

$$\int_D (\Delta v \bar{\phi} + \nabla v \cdot \nabla \bar{\phi}) = 0. \quad (3.29)$$

Proof. For all $\epsilon \in (0, 1)$, we set $D_\epsilon = \{x \in D \mid r(x) > \epsilon\}$. Since v is regular far from the origin, we can apply Green's formula on D_ϵ and find

$$\int_{D_\epsilon} (\Delta v \bar{\phi} + \nabla v \cdot \nabla \bar{\phi}) = -\epsilon \int_{r=\epsilon} \partial_r v \bar{\phi} d\theta. \quad (3.30)$$

Since Hölder's inequality and Lemma 2.1 guarantee that the integrand of left-hand side of (3.30) is integrable on the whole domain D , Lebesgue's convergence theorem allows to conclude that the left-hand side of (3.30) tends to the left-hand side of (3.29) as $\epsilon \rightarrow 0$. Hence it remains to show that the right-hand side of (3.30) tends to 0.

Now denote by \hat{A} the annulus defined as follows

$$\hat{A} = \{(r \cos \theta, r \sin \theta) \mid 1/2 < r < 1, 0 < \theta < \psi\}.$$

For any $\hat{v} \in H^1(\hat{A})$ such that $\Delta \hat{v} \in L^p(\hat{A})$ and any $\hat{\phi} \in H^1(\hat{A})$ such that $\hat{\phi} = 0$ on $\theta = 0$ and $\theta = \psi$, let us show that

$$\left| \int_{\hat{r}=1} \partial_r \hat{v} \bar{\hat{\phi}} d\hat{\theta} \right| \lesssim |\hat{\phi}|_{H^1(\hat{A})} |\hat{v}|_{H^1(\hat{A})} + \|\hat{\phi}\|_{L_{-\mu}^q(\hat{A})} \|\Delta \hat{v}\|_{L_\mu^p(\hat{A})}. \quad (3.31)$$

Indeed by taking an arbitrary cut-off function $\eta \in \mathcal{D}(\mathbb{R})$ such that $\eta(1) = 1$ and $\eta(1/2) = 0$, Green's formula yields

$$\int_{\hat{r}=1} \partial_r \hat{v} \bar{\hat{\phi}} d\hat{\theta} = \int_{\hat{A}} (\Delta \hat{v} (\eta \bar{\hat{\phi}}) + \nabla \hat{v} \cdot \nabla (\eta \bar{\hat{\phi}})).$$

By Hölder's inequality we obtain

$$\left| \int_{\hat{r}=1} \partial_r \hat{v} \bar{\hat{\phi}} d\hat{\theta} \right| \lesssim \|\hat{\phi}\|_{H^1(\hat{A})} |\hat{v}|_{H^1(\hat{A})} + \|\hat{\phi}\|_{L^q(\hat{A})} \|\Delta \hat{v}\|_{L^p(\hat{A})}.$$

Since Poincaré's inequality guarantees that

$$\|\hat{\phi}\|_{H^1(\hat{A})} \lesssim |\hat{\phi}|_{H^1(\hat{A})},$$

September 2, 2010

and since \hat{r} is equivalent to 1 on \hat{A} , the previous estimate leads to (3.31).

Now for all $j \in \mathbb{N}$, we set

$$A_j = \{(r \cos \theta, r \sin \theta) \mid \epsilon 2^{-(j+1)} < r < \epsilon 2^{-j}, 0 < \theta < \psi\}.$$

We then see that the linear mapping

$$\hat{x} \rightarrow x = \epsilon 2^{-j} \hat{x},$$

is a bijection from \hat{A} into A_j .

Now we perform the change of variables $x = \epsilon 2^{-j} \hat{x}$ and using (3.31), we obtain

$$\left| \int_{r=\epsilon 2^{-j}} \partial_r v \bar{\phi} r d\theta \right| \lesssim |\phi|_{H^1(\hat{A}_j)} |v|_{H^1(\hat{A}_j)} + \|\phi\|_{L^q_{-\mu}(A_j)} \|\Delta v\|_{L^p_{\mu}(A_j)}. \quad (3.32)$$

Since the measure of A_j tends to zero as j goes to infinity, we deduce from this estimate that

$$I_j(\epsilon) := \int_{r=\epsilon 2^{-j}} \partial_r v \bar{\phi} r d\theta \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Moreover Green's formula on $B_j = \{(r \cos \theta, r \sin \theta) \mid \epsilon 2^{-j} < r < \epsilon, 0 < \theta < \psi\}$, yields

$$I_0(\epsilon) = \int_{B_j} (\Delta v \bar{\phi} + \nabla v \cdot \nabla \bar{\phi}) + I_j(\epsilon).$$

Accordingly passing to the limit in j , we obtain that

$$I_0(\epsilon) = \int_{r=\epsilon} \partial_r v \bar{\phi} r d\theta = \int_{D(\epsilon)} (\Delta v \bar{\phi} + \nabla v \cdot \nabla \bar{\phi}),$$

where we recall that $D(\epsilon) = \{(r \cos \theta, r \sin \theta) \mid 0 < r < \epsilon, 0 < \theta < \psi\}$. Passing to the limit as ϵ goes to zero, by Lebesgue's bounded convergence theorem, we find that

$$I_0(\epsilon) \rightarrow 0 \text{ as } \epsilon \rightarrow 0.$$

This proves (3.29) thanks to (3.30). \square

Lemma 3.5. *With the notations of the proof of Theorem 3.3, the function $\eta \psi^{\lambda'} - \phi^{\lambda'}$ (with $\psi^{\lambda'}$ and $\phi^{\lambda'}$ defined by (3.20) and (3.21)) satisfies, for all $v \in V_{\mu}^{2,p}(D) \cap D(\Delta_{p,\mu})$,*

$$\int_D (-\Delta v + z v) (\eta \bar{\psi}^{\lambda'} - \bar{\phi}^{\lambda'}) = 0. \quad (3.33)$$

September 2, 2010

Proof. For shortness we now skip the indices z and λ' and we use the notations of the Proof of Lemma 3.4. Thanks to (3.29), we may write

$$\int_D (-\Delta v + z v) \bar{\phi} = \int_D (\nabla v \cdot \nabla \bar{\phi} + z v \bar{\phi}),$$

and using (3.21), we obtain

$$\int_D (-\Delta v + z v) \bar{\phi} = \int_D v(-\Delta + z)(\eta \bar{\psi}).$$

Consequently if we prove that

$$\int_D (-\Delta v + z v) \eta \bar{\psi} = \int_D v(-\Delta + z)(\eta \bar{\psi}), \quad (3.34)$$

then the difference between these two last identity yields (3.33).

It remains to prove (3.34). For that purpose, we again apply Green's formula in $D_\epsilon = \{x \in D \mid r(x) > \epsilon\}$, and find

$$\int_{D_\epsilon} (-\Delta v + z v) \eta \bar{\psi} = \int_{D_\epsilon} v(-\Delta + z)(\eta \bar{\psi}) + J_\epsilon,$$

where

$$J_\epsilon = \epsilon \int_{r=\epsilon} (v \partial_r \bar{\psi} - \partial_r v \bar{\psi}) d\theta.$$

Hence as before it suffices to show that

$$J_\epsilon \rightarrow 0 \text{ as } \epsilon \rightarrow 0. \quad (3.35)$$

For that last purpose we remark that

$$\partial_r \psi = A[\sqrt{\bar{z}} r \frac{d}{ds} Q(\sqrt{\bar{z}} r) - \sqrt{\bar{z}} r Q(\sqrt{\bar{z}} r) - \lambda' Q(\sqrt{\bar{z}} r)] e^{-\sqrt{\bar{z}} r} r^{-\lambda'-1} \sin(\lambda' \theta),$$

consequently

$$|J_\epsilon| \lesssim \epsilon^{-\lambda'} \int_{r=\epsilon} |v(\epsilon, \theta)| d\theta + \epsilon^{-\lambda'+1} \int_{r=\epsilon} |\partial_r v(\epsilon, \theta)| d\theta. \quad (3.36)$$

Now we estimate each term of this right-hand side separately. For the first one, we perform the change of variables $x = \epsilon \hat{x}$ that maps \hat{A} into A_0 , and by setting $\hat{v}(\hat{x}) = v(x)$, we obtain

$$\int_{r=\epsilon} |v(\epsilon, \theta)| d\theta = \int_{\hat{r}=1} |\hat{v}(1, \hat{\theta})| d\hat{\theta}.$$

On \hat{A} , we notice that \hat{v} belongs to $W^{1,p}(\hat{A})$ and by a standard trace theorem, we deduce that

$$\int_{\hat{r}=1} |\hat{v}(1, \hat{\theta})| d\hat{\theta} \lesssim \left(\int_{\hat{r}=1} |\hat{v}(1, \hat{\theta})|^p d\hat{\theta} \right)^{\frac{1}{p}} \lesssim \|\hat{v}\|_{W^{1,p}(\hat{A})}.$$

Since $\hat{v} = 0$ on a part of the boundary of \hat{A} , $\|\hat{v}\|_{W^{1,p}(\hat{A})} \lesssim |\hat{v}|_{W^{1,p}(\hat{A})}$, and therefore we obtain

$$\int_{\hat{r}=1} |\hat{v}(1, \hat{\theta})| d\hat{\theta} \lesssim |\hat{v}|_{W^{1,p}(\hat{A})}.$$

Finally as \hat{r} is equivalent to 1 on \hat{A} , we arrive at

$$\int_{\hat{r}=1} |\hat{v}(1, \hat{\theta})| d\hat{\theta} \lesssim \left(\int_{\hat{A}} \hat{r}^{(\mu-1)p} |\nabla \hat{v}|^p \right)^{\frac{1}{p}}.$$

Going back to A_0 , we have proved that

$$\int_{r=\epsilon} |v(\epsilon, \theta)| d\theta \lesssim \epsilon^{2-\frac{2}{p}-\mu} \left(\int_{A_0} r^{(\mu-1)p} |\nabla v|^p \right)^{\frac{1}{p}} \lesssim \epsilon^{2-\frac{2}{p}-\mu} \|v\|_{V_\mu^{2,p}(D)}.$$

In a similar way we show that

$$\int_{r=\epsilon} |\partial_r v(\epsilon, \theta)| d\theta \lesssim \epsilon^{1-\frac{2}{p}-\mu} \left(\int_{A_0} r^{\mu p} |\nabla \partial_r v|^p \right)^{\frac{1}{p}} \lesssim \epsilon^{1-\frac{2}{p}-\mu} \|v\|_{V_\mu^{2,p}(D)}.$$

These two estimates in (3.36) lead to

$$|J_\epsilon| \lesssim \epsilon^{2-\frac{2}{p}-\mu-\lambda'} \|v\|_{V_\mu^{2,p}(D)}.$$

Since $2 - \frac{2}{p} - \mu - \lambda'$ is positive, we obtain (3.35). \square

References

- [1] **J.O. Adeyeye**, *Generation of analytic semi-group in $L^p(\Omega)$ by the Laplace operator*, Boll. U.M.I. Analisi Funzionale e Applicazioni, Serie VI, Vol. IV - C, n. 1 (1985), 113-128.
- [2] **A. Aibeche, W. Chikouche and S. Nicaise**, *L^p regularity of transmission problems in dihedral domains*, Boll. U.M.I. 10B (2007), 633-660.

- [3] **N.T. Anh and N.M. Hung**, *Asymptotic formulas for solutions of parameter-dependent elliptic boundary-value problems in domains with conical points*, Electron. J. Differential Equations, 2009 (2009), No. 125, 1-21.
- [4] **M. Costabel, M. Dauge and S. Nicaise**, *Corner Singularities and Analytic Regularity for Linear Elliptic Systems.*, book in preparation.
- [5] **G. Da Prato and P. Grisvard**, *Sommes d'opérateurs linéaires et équations différentielles opérationnelles*, J. Math. Pure Appl. 54 (1975), 305-387.
- [6] **D. Gilbarg and N.S. Trudinger**, *Elliptic partial differential equations of second order, Second Edition*, Springer-Verlag, Berlin 1983.
- [7] **P. Grisvard**, *Elliptic problems in nonsmooth domains*, Monographs and Studies in Mathematics 24, Pitman, Boston-London-Melbourne, 1985.
- [8] **P. Grisvard**, *Edge behavior of the solution of an elliptic problem*, Math. Nachr. 132 (1987), 281-299.
- [9] **P. Grisvard**, *Singular behavior of elliptic problems in non hilbertian Sobolev spaces*, J. Math. Pures Appl. 74 (1995), 3-33
- [10] **V.A. Kozlov**, *Coefficients in the asymptotic solutions of the Cauchy boundary-value parabolic problems in domains with a conical point*, Sibirskii Mat. Zhurnal 29 (1988), 75-89.
- [11] **V.A. Kozlov and V.G. Maz'ya**, *Singularities of solutions of the first boundary value problem for the heat equation in domains with conical points. II*, Soviet Math. (Iz. VUZ) 31 (1987), 49-57.
- [12] **A. Kufner**, *Weighted Sobolev spaces*, John Wiley and Sons, Chichester, 1980.
- [13] **A. Kufner and A.-M. Sändig**, *Some applications of weighted Sobolev spaces*, Teubner texte zur mathematik 100 (1987).
- [14] **V.G. Maz'ya and B.A. Plamenevskii**, *L_p -estimates of solutions of elliptic boundary value problems in domains with edged*, Trudy Moskov Mat. Obshch. 37 (1978), 49-93 = Trans. Moscow Math. Soc. 37 (1980), 49-97.

- [15] **A.I. Nazarov**, *L_p -estimates for the solution to the Dirichlet problem and to the Neumann problem for the heat equation in a wedge with edge of arbitrary codimension*, J. Math. Sciences 106 (2001), 2989-3014.
- [16] **A.I. Nazarov**, *Dirichlet problem for quasilinear parabolic equations in domains with smooth closed edges*, Amer. Math. Soc. Transl. 209 (2003), 115-141.
- [17] **S. Nicaise**, *Polygonal interface problems*, Methoden und Verfahren der mathematischen Physik, Peter Lang (1993).
- [18] **A. Pazy**, *Semigroups of linear operators and applications to partial differential equations*, Springer-Verlag (1983).
- [19] **J. Prüss and G. Simonett**, *H^∞ -Calculus for the sum of non-commuting operators*, Trans. A.M.S. 359 (2007), 3549-3565.
- [20] **V.A. Solonnikov**, *L_p -estimates for solutions of the heat equation in a dihedral angle*, Rend. Matematica, Serie VII, vol. 21, Roma (2001), 1-15.