



HAL
open science

On Cutting Planes for Signomial Programming

Liding Xu, Claudia d'Ambrosio, Leo Liberti, Sonia Haddad Vanier

► **To cite this version:**

Liding Xu, Claudia d'Ambrosio, Leo Liberti, Sonia Haddad Vanier. On Cutting Planes for Signomial Programming. 2023. hal-03723024v2

HAL Id: hal-03723024

<https://hal.science/hal-03723024v2>

Preprint submitted on 24 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cutting planes for signomial programming

Liding Xu* Claudia D’Ambrosio* Leo Liberti* Sonia Haddad-Vanier*[†]

September 12, 2023

Abstract

Cutting planes are of crucial importance when solving nonconvex nonlinear programs to global optimality, for example using the spatial branch-and-bound algorithms. In this paper, we discuss the generation of cutting planes for signomial programming. Many global optimization algorithms lift signomial programs into an extended formulation such that these algorithms can construct relaxations of the signomial program by outer approximations of the lifted set encoding nonconvex signomial term sets, i.e., hypographs, or epigraphs of signomial terms. We show that any signomial term set can be transformed into the subset of the difference of two concave power functions, from which we derive two kinds of valid linear inequalities. Intersection cuts are constructed using signomial term-free sets which do not contain any point of the signomial term set in their interior. We show that these signomial term-free sets are maximal in the nonnegative orthant, and use them to derive intersection sets. We then convexify a concave power function in the reformulation of the signomial term set, resulting in a convex set containing the signomial term set. This convex outer approximation is constructed in an extended space, and we separate a class of valid linear inequalities by projection from this approximation. We implement the valid inequalities in a global optimization solver and test them on MINLPLib instances. Our results show that both types of valid inequalities provide comparable reductions in running time, number of search nodes, and duality gap.

Key words: global optimization, signomial programming, extended formulation, cutting plane, intersection cut, convex relaxation

1 Introduction

General nonconvex nonlinear programming (NLP) problems typically admit the following formulation:

$$\min_{x \in \mathbb{R}^n} c \cdot x \quad \text{s. t.} \quad Ax + Bg(x) \leq d, \quad (1)$$

where $c \in \mathbb{R}^n$, $A \in \mathbb{R}^{m \times n}$, $B \in \mathbb{R}^{m \times k}$, $g : \mathbb{R}^n \rightarrow \mathbb{R}^k$, $d \in \mathbb{R}^m$.

The mapping $g(x)$ represents a vector $(g_1(x), \dots, g_k(x))$ of nonconvex functions on x , and we denote g_i as their *terms*. Note that the objective function is supposed to be linear, w.l.o.g., since we can always reformulate a problem with a nonlinear objective function as the problem (1) above (epigraphic reformulation).

General-purpose global optimization solvers, such as BARON [65], Couenne [10], and SCIP [11], are capable of solving the problem (1) within an ϵ -global optimality. They achieve this by

*LIX CNRS, École Polytechnique, Institut Polytechnique de Paris, Palaiseau, 91128, France. E-mail: lidingxu.ac@gmail.com, dambrosio@lix.polytechnique.fr, liberti@lix.polytechnique.fr, sonia.vanier@lix.polytechnique.fr

[†]Université Paris 1 Panthéon-Sorbonne, Paris, 75005, France.

employing the spatial branch-and-bound (sBB) algorithm, which explores the feasible region of (1) implicitly, but systematically. The sBB algorithm effectively prunes out unpromising search regions by comparing the cost of the best feasible solution found with the cost bounds associated with those regions. These cost bounds can be computed by solving convex relaxations of (1).

The backend convex relaxation algorithms implemented in many general-purpose solvers, including **BARON**, **Couenne**, and **SCIP**, are linear programming relaxations. These solvers take advantage of the separability introduced in the rows of $Ax + Bg(x)$, allowing them to relax and linearize nonlinear terms g_i individually. In the solvers' data structures, the problem (1) is transformed into an extended formulation:

$$\min_{(x,y) \in \mathbb{R}^{n+k}} c \cdot x \quad \text{s. t.} \quad Ax + By \leq d \wedge y = g(x). \quad (2)$$

All the nonlinear terms are grouped within the nonconvex constraints $y = g(x)$. These constraints give rise to a nonconvex *lifted set* defined as:

$$\mathcal{S}_{\text{lift}} := \{(x, y) \in \mathbb{R}^{n+k} : y = g(x)\}. \quad (3)$$

The relaxation algorithms used by these solvers are based on factorable programming [35, 47]: This approach treats the multivariate nonlinear terms g_i as composite functions. These algorithms typically factorize each g_i into sums and products of a collection of univariate functions. If convex and concave relaxations of those univariate functions are available, these algorithms can linearize these relaxations, and yield a linear relaxation for Eq. (1). Common lists of such univariate functions, that are usually available to all sBB solvers, include t^a (for $a \in \mathbb{N}$), $\frac{1}{t}$, $\log t$, $\exp t$. Some solvers also offer a choice of trigonometric functions, e.g. **Couenne**.

Most sBB solvers can handle *signomial term* $\psi_\alpha(x) := x^\alpha = \prod_{j \in [n]} x_j^{\alpha_j}$, where the exponent vector α is in \mathbb{R}^n , but in a way that yields poor relaxations (more about this below). In this paper, we provide a deeper treatment of the signomial term w.r.t. convexification and linearization within an sBB algorithm. The **SCIP** solver can handle signomial terms, but, as of version 8.0, the treatment was limited to recognizing convexity and concavity. The technical report [11] explicitly states that the techniques proposed in this paper are going to be integrated into **SCIP**.

When all the terms in g are signomial terms, the problem (1) falls under the category of signomial programming (SP). In this scenario, we refer to (1) as the *natural formulation* of SP. The left-hand sides of the constraints in this formulation are referred to as *signomial functions*. The lifted set $\mathcal{S}_{\text{lift}}$ in the extended formulation (2) is called a *signomial lift*.

Since negative entries may present in the exponent vector α , in general, variables of SP are assumed to be positive. The point of restriction on SP over positive variables is simply to make the theoretical treatment more readable and streamlined. We remark that the techniques in this paper can also treat signomial terms in general mixed-integer NLP problems.

In the case of SP, LP relaxations can be derived from polyhedral outer approximations of the signomial lift in its extended formulation. A typical relaxation algorithm for SP involves factorizing the signomial term $\psi_\alpha(x)$ into the product of n univariate signomial terms $x_i^{\alpha_i}$. After the factorization, the algorithm proceeds to convexify and linearize the intermediate multilinear term and univariate functions. However, this factorial programming approach can lead to weak LP relaxation and introduce additional auxiliary variables that represent intermediate functions. These problems have already been discussed in the context of pure multilinear terms [14, 21, 63].

We propose two cutting plane-based relaxation algorithms for SP. In contrast to the conventional factorable programming approach, our method uses a novel reformulation of the signomial lift. We transform each nonlinear equality constraint $y_i = g_i(x)$ in (3) to an equivalent constraint $\psi_\beta(u) - \psi_\gamma(v) = 0$, where $\beta > 0, \gamma > 0$, $\max(\|\beta\|_1, \|\gamma\|_1) = 1$, u, v are sub-vectors partitioned from (x, y) , and ψ_β, ψ_γ are concave functions. We consider approximating the following set

$$\mathcal{S}_{\text{st}} := \{(u, v) \in \mathbb{R}_+^{h+\ell} : \psi_\beta(u) - \psi_\gamma(v) \leq 0\}. \quad (4)$$

Our first cutting plane algorithm is based on the intersection cut paradigm [19]. As shown in Sec. 2, one can approximate a nonconvex set \mathcal{S} using its polyhedral outer approximation. This requires the construction of \mathcal{S} -free sets, i.e., closed convex sets containing none of the interiors of \mathcal{S} . The main insight about \mathcal{S} -free sets for a nonconvex set \mathcal{S} is that they provide an explicit and useful description of the convex parts of the complement of \mathcal{S} . In Sec. 3 we extend several general results from the literature on maximal \mathcal{S} -free sets. In Sec. 4 we give the transformation procedure leading to \mathcal{S}_{st} and construct \mathcal{S}_{st} -free sets from the transformation. We show that these sets are also signomial-lift-free and maximal in the nonnegative orthant. We also discuss the separation of intersection cuts.

To ensure convergence of the sBB algorithm, a common assumption for SP is that all variables are bounded. Our second cutting plane algorithm aims to approximate \mathcal{S}_{st} within a hypercube. In Sec. 5, we provide an extended formulation for the convex envelope of the concave function ψ_β over the hypercube. This formulation yields a convex set including \mathcal{S}_{st} (which is a convex outer approximation of \mathcal{S}_{st}), so that we can generate outer approximation cuts by projection. We prove that ψ_β is a supermodular function. For $h = 2$ we provide a closed expression for its convex envelope by exploiting supermodularity, which allows us to remove the projection step.

For the computational part of this study, we note that signomials are one of the four main types of nonlinearities found in the mixed-integer NLP library (MINLPLib) [9, 13]. Our relaxation approach does not require factorization or the introduction of intermediate functions, so implementing the proposed cutting planes in the general-purpose solver SCIP is straightforward. In Sec. 6, we perform computational tests with instances from MINLPLib and observe improvements to SCIP default settings due to the proposed valid inequalities.

1.1 Related works

The majority of relaxations for SP are derived from its generalized geometric programming (GPP) formulation, which is an exponential transformation [25] of its natural formulation. The exponential transformation replaces positive variables x by exponentials $\exp(z)$, where z are real variables. The authors of [46] show that signomial functions in GGP are difference-of-convex (DC) functions. For the signomial function in each constraint of GGP, they construct linear underestimators of its concave part; the author of [61] constructs linear underestimators of the whole function via the mean value theorem. The author of [68] proposes inner approximations of GGP via the inequality of arithmetic and geometric means (AM-GM inequality). The authors of [15, 24, 54] construct non-negativity certificates for signomial functions via the AM-GM inequality, and propose a hierarchy of convex relaxations for GGP. Exponential transformations can be combined with other variable transformations, such as power transformations, and the inverse transformations can be approximated by piece-wise linear functions, see [38, 43, 44].

The solvers SCIP [11], BARON [65], ANTIGONE [50], and MISO [51] are able to solve the natural formulation of SP or its extended formulation within a global ϵ -optimality using the sBB algorithm. More precisely, MISO is a specialized solver for SP, which uses exponential transformations of some signomial terms only when necessary. For the following reasons, exponential transformations can complicate general-purpose solvers. First, in certain NLP problems, signomial terms may appear only as a subset of the nonlinear terms of $g(x)$. In such cases, solvers may need to force the inverse transformation $x_j = \ln(z_j)$, which requires additional processing for convexification algorithms. Second, when dealing with mixed-integer SP and some variables of x are integer, exponential transformations cause certain components of z to become discrete but not necessarily integer. As a result, the sBB algorithm must adjust its branching rules.

While much attention has been paid to the construction of relaxations for GGP, the literature on relaxations for the extended natural formulation of SP is relatively limited. The convex relaxations used in the aforementioned solvers rely mainly on factorial programming [36, 47].

Since exponential transformations are nonlinear variable transformations, it is impossible to apply the relaxations developed for the GGP formulation directly to the natural formulation.

Numerous research efforts have been devoted to improving relaxation techniques for multilinear terms and univariate/bivariate functions commonly used in factorizable programming [6]. Multilinear terms over the unit hypercube are vertex polyhedral and their envelopes over the unit hypercube admits simple extended formulations [58]. In particular, there are closed forms for the convex envelopes of bilinear functions [1, 47] and trilinear functions [48, 49] over hypercubes. In [62], the author presents convex envelopes for multilinear functions (sum of multilinear terms) over the unit hypercube and specific discrete sets. For a comprehensive analysis of multilinear term factorization via bilinear terms, we refer to [42, 63]. Additionally, [14] offers an in-depth examination of quadrilinear function factorization through bilinear and trilinear terms, while [21] presents a computational study on extended formulations.

Convexifying univariate/bivariate functions plays an important role in the field of global optimization. In [37], convex envelopes for monomials with odd degrees are derived. An approach presented in [41] enables the evaluation of the convex envelope of a bivariate function over a polytope and separating its supporting hyperplane by solving low-dimensional convex optimization problems. The convex optimization problems are further reduced by solving a Karush-Kuhn-Tucker system [40]. In [39], convex envelopes for bilinear, fractional, and other bivariate functions over a polytope are constructed using a polyhedral subdivision technique. Additionally, [55, 64] employ polyhedral subdivision and lift-project methods to derive explicit forms of convex envelopes for various nonconvex functions, including a specific subclass of bivariate signomial terms.

Convexifying high-order multivariate functions is a major challenge, and the available literature on convex underestimators for trivariate functions is relatively few. In [31, 32], the authors propose a novel framework for relaxing composite functions in nonlinear programs. Another approach is to use the intersection cut paradigm [19] to approximate nonconvex functions. This paradigm can generate cutting planes to strengthen LP relaxations of NLP problems. Constructing intersection cuts involves finding an \mathcal{S} -free set, where \mathcal{S} represents a nonconvex set defined by nonconvex functions. The study of intersection cuts originated in the context of NLP [67]. Gomory later introduced the concept of corner polyhedron [30], and intersection cuts were explored in the field of integer programming [5]. The modern definition of intersection cuts for arbitrary sets \mathcal{S} is from [23, 29]. For more comprehensive details, we refer to [2, 7, 20, 22, 23, 57]. Recent research has revealed \mathcal{S} -free sets for various nonconvex sets encountered in structured NLP problems. Examples include outer product sets [12], sublevel sets of DC functions [59], quadratic sets [53], and graphs of bilinear terms [28]. Intersection cuts have also been developed for convex mixed-integer NLP problems [3, 8, 34, 52] and for bilevel programming [27].

1.2 Notation

We follow standard notation in most cases. Let $[n_1 : n_2]$ stand for $\{n_1, \dots, n_2\}$, and let $[n]$ stand for $[1 : n]$. For a vector $x \in \mathbb{R}^n$, x_j denotes the j -th entry of x ; given $J \subseteq [n]$, $x_J = (x_j)_{j \in J}$ denotes the sub-vector formed by entries indexed by J . $\|\cdot\|_p$ denotes the L_p -norm ($1 \leq p \leq +\infty$). For a set $X \subseteq \mathbb{R}^n$, $\text{conv}(X)$, $\text{cl}(X)$, $\text{int}(X)$, $\text{bd}(X)$, $|X|$, X^c denote the convex hull, closure, interior, boundary, cardinality, and complement of X , respectively. For a function f , $\text{dom}(f)$ and $\text{range}(f)$ denote the domain and range of f , respectively; $\text{graph}(f)$ denotes its graph $\{(x, t) \in \mathbb{R}^{n+1} : f(x) = t\}$, $\text{epi}(f)$ denotes its epigraph $\{(x, t) \in \mathbb{R}^{n+1} : f(x) \leq t\}$, and $\text{hypo}(f)$ denotes its hypograph $\{(x, t) \in \mathbb{R}^{n+1} : f(x) \geq t\}$; if f is differentiable, for a $\tilde{x} \in \text{dom}(f)$, $\nabla f(\tilde{x})$ denotes the gradient of f at \tilde{x} and

$$\Xi_{\tilde{x}}^f(x) := f(\tilde{x}) + \nabla f(\tilde{x}) \cdot (x - \tilde{x}). \quad (5)$$

The word *linearization* involves the replacement of a nonlinear function by its affine underestimators or overestimators. For example, the affine underestimators of convex functions f are given as $\Xi_{\tilde{x}}^f(x)$ for some \tilde{x} .

2 Preliminaries

In this section we present an overview of \mathcal{S} -free sets and intersection cut theory. The process of constructing intersection cuts involves two fundamental steps [18]: constructing \mathcal{S} -free sets and deriving cutting planes from these sets. Since maximal \mathcal{S} -free sets yield tightest cutting planes, one can include an optional step to check the maximality of \mathcal{S} -free sets.

Definition 1. *Given a set $\mathcal{S} \subsetneq \mathbb{R}^p$, a closed set \mathcal{C} is (convex) \mathcal{S} -free if \mathcal{C} is convex and $\text{int}(\mathcal{C}) \cap \mathcal{S} = \emptyset$.*

Fig. 1 shows an example of an \mathcal{S} -free set, where we find that \mathcal{C} is a convex inner approximation of $\text{cl}(\mathcal{S}^c)$. Thereby, we show how to construct \mathcal{S} -free sets from a “reverse” representation of some nonconvex sets. We look at sets involving a particular type of nonconvex function.

Definition 2. *A function f is said to be difference-of-concave (DCC), if there exist two concave functions f_1, f_2 such that $f = f_1 - f_2$.*

It is easy to show that the negative of a DCC function is also a DCC function, and any DC function can be converted into a DCC function. We call a nonconvex set a *DCC set*, if it admits a *DCC formulation*, meaning that it can be represented as the sublevel set of a DCC function. The superlevel set of a DCC function is a sublevel set of another DCC function (the negative of that function), so the superlevel set is also a DCC set. By using the *reverse-linearization technique*, the following lemma provides a collection of \mathcal{S} -free sets for DCC sets.

Lemma 1. [59, Prop. 6] *Let $\mathcal{S} := \{z \in \mathbb{R}^p : f_1(z) - f_2(z) \leq 0\}$, where f_1, f_2 are concave functions over \mathbb{R}^p . Then, for any $\tilde{z} \in \mathbb{R}^p$, $\mathcal{C} := \{z \in \mathbb{R}^p : f_1(z) - \Xi_{\tilde{z}}^{f_2}(z) \geq 0\}$ is \mathcal{S} -free. Moreover, if $\tilde{z} \in \mathbb{R}^p \setminus \mathcal{S}$, $\tilde{z} \in \text{int}(\mathcal{C})$.*

The reverse-linearization technique involves reversing the inequality that defines \mathcal{S} and linearizing its convex component $-f_2$. The point \tilde{z} is referred to as the *linearization point*. It is important to note that, when the shared domain \mathcal{G} of f_1 and f_2 is not the entire space \mathbb{R}^p , the set \mathcal{S} needs to be constrained to the *ground set* \mathcal{G} . This restriction ensures the applicability of the lemma.

To construct an intersection cut, an essential requirement is the availability of a translated simplicial cone \mathcal{R} that satisfies two conditions: (i) \mathcal{R} is generated by linearly independent vectors, (ii) \mathcal{R} contains \mathcal{S} , and (iii) the vertex \tilde{z} of \mathcal{R} does not belong to \mathcal{S} . We assume that \mathcal{R} admits a hyper-plane representation $\{z \in \mathbb{R}^p : B(z - \tilde{z}) \leq 0\}$, where $B \in \mathbb{R}^{p \times p}$ is an invertible matrix. For every $j \in [p]$, let r^j denote the j -th column of $-B^{-1}$, then r^j turns out to be an extreme ray of \mathcal{R} . Thereby, \mathcal{R} also admits a ray representation $\{z \in \mathbb{R}^p : \exists \eta \in \mathbb{R}_+^p \ z = \tilde{z} + \sum_{j=1}^p \eta_j r^j\}$. For every $j \in [p]$, we define the *step length* from \tilde{z} along ray r_j to the boundary $\text{bd}(\mathcal{C})$ as

$$\eta_j^* := \sup_{\eta_j \in [0, +\infty]} \{\eta_j : \tilde{z} + \eta_j r^j \in \mathcal{C}\}. \quad (6)$$

Then, an intersection cut admits the form

$$\sum_{j=1}^p B_j(z - \tilde{z})/\eta_j^* \leq -1, \quad (7)$$




Figure 1: An S -free set C , simplicial cone \mathcal{R} , and intersection cut.

where B_j is the j -th row of B . When all step lengths are positive, the above linear inequality cuts off \tilde{z} from S . See for an example of an intersection cut in Fig. 1.

We can obtain the sets C , \mathcal{R} and the vertex \tilde{z} by the following procedure. Suppose that we have an LP relaxation $\min_{z \in \mathcal{P}} c \cdot z$ of an SP problem, where \mathcal{P} is a polyhedral outer approximation of the feasible set of the SP problem. If the solution to the LP problem turns out to be infeasible for the SP problem, it means that the solution does not belong to the signomial lift. In such cases, we can fix \tilde{z} as the solution obtained from LP and let C be the signomial-lift-free (S_{lift} -free) set. Moreover, we can extract the cone \mathcal{R} from the optimal LP basis defining \tilde{z} , see [18].

One focus of our study is the construction of (maximal) S -free sets. The importance of finding *maximal* sets follows from the fact that if we have two S -free sets called C and C^* , where C is a subset of C^* , then the intersection cut derived from C^* dominates the cut derived from C (see [19, Remark 3.2]). To give a precise characterization, we present a formal definition of maximal S -free sets.

Definition 3. *Given a closed convex set $\mathcal{G} \subseteq \mathbb{R}^p$ such that $S \subsetneq \mathcal{G}$, an S -free set C is (inclusion-wise) maximal in \mathcal{G} , if there is no other S -free set C' such that $C \cap \mathcal{G} \subsetneq C' \cap \mathcal{G}$.*

The above definition provides a generalization of the conventional concept of maximal S -free sets, which is a special case when $\mathcal{G} = \mathbb{R}^p$. Studying maximality for S -free sets in \mathbb{R}^p can be challenging in certain scenarios. However, Defn. 3 allows us to examine the intersections of S -free sets within the ground set \mathcal{G} . This constraint is essential for our analysis, especially considering that all variables in SP are non-negative.

3 General results on maximality

In this section, we present two results on the maximality of S -free sets arising in general nonconvex NLP problems. The results are used to construct maximal signomial-lift-free sets in non-negative orthants.

3.1 Lifted sets

We consider the extended formulation (2) of a general NLP problem and focus on the associated lifted set S_{lift} in (3). We show a lifting result on constructing maximal S_{lift} -free sets.

Let $z := (x, y)$ denote the vector variable in the extended formulation (2), with its index set being $[n + k]$. Consequently, we have $z_{[n]} = x$ and $z_{[n+1:n+k]} = y$. Consider a closed subset \mathcal{X} of the domain $\bigcap_{i \in [k]} \text{dom}(g_i)$ for x , and let \mathcal{Y} be a closed subset of the domain $\bigtimes_{i \in [k]} \text{range}(g_i)$ for y . The ground set \mathcal{G} can, thus, be set as $\mathcal{X} \times \mathcal{Y}$. Consequently, the lifted set $\mathcal{S}_{\text{lift}}$ in (3) admits the form $\{(x, y) \in \mathcal{G} : y = g(x)\}$.

Given that each $g_i(x)$ (for $i \in [k]$) may only depend on a subset of variables indexed by $J_i \subseteq [n]$, we can express $g_i(x)$ as a lower order function $g'_i(x_{J_i})$ defined over \mathbb{R}^{J_i} . Let $I_i := J_i \cup \{i + n\}$. As above, we consider a closed subset \mathcal{X}^i of $\text{dom}(g'_i)$ and \mathcal{Y}^i of $\text{range}(g'_i)$. Consequently, the graph, epigraph, and hypograph of g'_i reside within sets $\mathcal{G}^i := \mathcal{X}^i \times \mathcal{Y}^i$, e.g., $\text{epi}(g'_i) = \{(x_{J_i}, y_i) \in \mathcal{G}^i : g'_i(x_{J_i}) \leq y_i\}$.

We refer to $\mathcal{X}, \mathcal{Y}, \{\mathcal{X}^i, \mathcal{Y}^i\}_{i \in [k]}$ as the *underlying sets* of the lifted set $\mathcal{S}_{\text{lift}}$. The sets are said to be *1d-convex decomposable* by a collection $\{\mathcal{D}_j\}_{j \in [n+k]}$ of closed convex sets in \mathbb{R} , if $\mathcal{X} = \bigtimes_{j \in [n]} \mathcal{D}_j, \mathcal{Y} = \bigtimes_{j \in [n+1:n+k]} \mathcal{D}_j$, and, for all $i \in [k]$, $\mathcal{X}^i = \bigtimes_{j \in J_i} \mathcal{D}_j, \mathcal{Y}^i = \mathcal{D}_{n+i}$. This decomposability condition restricts the domains to Cartesian products of real lines, intervals, or half rays, thereby excluding complicated domain structures.

The decomposability condition allows the analysis of sets with fewer variables. The construction of $\text{epi}(g'_i)$ -free sets and $\text{hypo}(g'_i)$ -free sets is in general simpler than the construction of $\mathcal{S}_{\text{lift}}$ -free sets. We show that any maximal $\text{epi}(g'_i)$ -free or $\text{hypo}(g'_i)$ -free set can be transformed into a maximal $\mathcal{S}_{\text{lift}}$ -free set.

Theorem 1. *Suppose the underlying sets of $\mathcal{S}_{\text{lift}}$ are 1d-convex decomposable and g is continuous. For some $i \in [k]$, let \mathcal{C} be a maximal $\text{epi}(g'_i)$ -free set or a maximal $\text{hypo}(g'_i)$ -free set in \mathcal{G}^i . Then, $\tilde{\mathcal{C}} := \mathcal{C} \times \mathbb{R}^{I_i^c}$ is a maximal $\mathcal{S}_{\text{lift}}$ -free set in \mathcal{G} .*

See the proof in the appendix. For any $i \in [k]$, we call the operation $\mathcal{C} \times \mathbb{R}^{I_i^c}$ the *orthogonal lifting* of \mathcal{C} with respect to g_i . A similar lifting result for integer programming is given by [19, Lemma 4.1]: given $\mathcal{S} := \mathbb{Z}^n \times \mathbb{R}^h$, any maximal lattice-free set (i.e., \mathbb{Z}^n -free set) can be transformed into a maximal \mathcal{S} -free set by orthogonal lifting. Therefore, Thm. 1 serves as the NLP counterpart to this lemma (whose proof is also similar). This theorem allows us to focus on low-dimensional projections of the lifted set. We will show in Cor. 1 that the signomial lift satisfies the prerequisites of Thm. 1. The following example illustrates the application of Thm. 1.

Example 1. *Consider a lifted set $\mathcal{S}_{\text{lift}}$ defined as*

$$\{(x_1, x_2, x_3, x_4, y_1, y_2, y_3) : y_1 = \exp(x_1 - x_2/x_3) \wedge y_2 = \log(x_1) \wedge y_3 = \sin(x_1/x_4)\}.$$

One can verify that the 1d-convex decomposable condition holds for $\mathcal{D}_1 = \mathbb{R}_+, \mathcal{D}_j = \mathbb{R}$ (for $j \in [2 : 7]$). Then $\mathcal{G} := \mathbb{R}_+^1 \times \mathbb{R}^6$. We use $\log(x_1)$ to construct a $\mathcal{S}_{\text{lift}}$ -free set. A maximal $\mathcal{S}_{\text{lift}}$ -free set can be $\{(x_1, x_2, x_3, x_4, y_1, y_2, y_3) \in \mathcal{G} : y_2 \leq \log(x_1)\}$. Since $\log(x_1)$ is defined over positive reals, this example gives a reason to restrict maximality over \mathcal{G} .

3.2 Sufficient conditions on maximality

We provide sufficient conditions for the maximality of \mathcal{S} -free sets for two general classes of nonconvex sets \mathcal{S} . At the beginning, we give an overview of some basic results of convex analysis. Our subsequent exposition relies on the use of support functions of convex sets. The properties of support functions can be summarized as follows.

Lemma 2. [33, Chap. C] *For a full-dimensional closed convex set $\mathcal{C} \subsetneq \mathbb{R}^p$, let $\sigma_{\mathcal{C}} : \mathbb{R}^p \rightarrow \mathbb{R}, \lambda \mapsto \sup_{z \in \mathcal{C}} \lambda \cdot z$ be the support function of \mathcal{C} . Then: (i) $\mathcal{C} = \{z \in \mathbb{R}^p : \forall \lambda \in \text{dom}(\sigma_{\mathcal{C}}) \lambda \cdot z \leq \sigma_{\mathcal{C}}(\lambda)\}$, (ii) $\text{int}(\mathcal{C}) = \{z \in \mathbb{R}^p : \forall \lambda \in \text{dom}(\sigma_{\mathcal{C}}) \setminus \{0\} \lambda \cdot z < \sigma_{\mathcal{C}}(\lambda)\}$, (iii) $\sigma_{\mathcal{C}}(\rho\lambda) = \rho\sigma_{\mathcal{C}}(\lambda)$ for any $\rho > 0$. Moreover, for any closed convex set \mathcal{C}' including \mathcal{C} , $\sigma_{\mathcal{C}} \leq \sigma_{\mathcal{C}'}$.*

A valid inequality $a \cdot z \leq b$ of \mathcal{C} is called a *supported valid inequality*, if there exists a *supporting point* $z' \in \text{bd}(\mathcal{C})$ such that $a \cdot z' = b$. Geometrically, a closed convex set is the intersection of half-spaces associated with supported valid inequalities.

Observation 1. *It follows from Lemma 2 that every supported valid inequality of \mathcal{C} must admit the form $\lambda \cdot z \leq \sigma_{\mathcal{C}}(\lambda)$ for some $\lambda \in \text{dom}(\sigma_{\mathcal{C}})$, where the supremum $\sigma_{\mathcal{C}}(\lambda)$ is attained at its supporting points.*

An inequality of the form $\lambda \cdot z \leq \sigma_{\mathcal{C}}(\lambda)$, for $\lambda \in \text{dom}(\sigma_{\mathcal{C}})$, is referred to as an *exposed valid inequality*, if there exists an *exposing point* $z' \in \text{bd}(\mathcal{C})$ such that $\lambda \cdot z' = \sigma_{\mathcal{C}}(\lambda)$ and, for all $\lambda' \in \text{dom}(\sigma_{\mathcal{C}}) \setminus \{\rho\lambda\}_{\rho>0}$, $\lambda' \cdot z' < \sigma_{\mathcal{C}}(\lambda')$.

Observation 2. *An exposed valid inequality must be a supported valid inequality. Conversely, a supported valid inequality is an exposed valid inequality if the manifold $\text{bd}(\mathcal{C})$ is smooth at its supporting point. For example, $\mathcal{C}_1 := \{(x, y) \in \mathbb{R}^2 : y = x^2\}$ is a smooth manifold, so any supported valid inequality of \mathcal{C}_1 is exposed; $\mathcal{C}_2 := \{(x, y) \in \mathbb{R}^2 : y = |x|\}$ is smooth at $x \in [1, 2]$, so any supported valid inequality of \mathcal{C}_2 with support point (x, y) ($x \in [1, 2]$) is also exposed by the same point; however, a supported valid inequality of \mathcal{C}_2 with supporting point (x, y) ($x = 0$) cannot be exposed, since there are infinitely many supported valid inequalities at the same point.*

The first theorem we present holds for full-dimensional nonconvex sets \mathcal{S} . As shown in Fig. 1, we have observed the geometric equivalence between the closed convex inner approximation of $\text{cl}(\mathcal{S}^c)$ and \mathcal{S} -free sets. The theorem provides a sufficient condition for the maximality of closed convex inner approximations.

Theorem 2. *Let \mathcal{F} be a full-dimensional closed set in \mathbb{R}^p , and let $\mathcal{C} \subseteq \mathcal{F}$ be a full-dimensional closed convex set. If, for any $z^* \in \text{int}(\mathcal{F} \setminus \mathcal{C})$ and any $\lambda \in \text{dom}(\sigma_{\mathcal{C}})$ such that $\lambda \cdot z^* > \sigma_{\mathcal{C}}(\lambda)$, there exists a point $z' \in \text{bd}(\mathcal{F}) \cap \text{bd}(\mathcal{C})$ exposing $\lambda \cdot z \leq \sigma_{\mathcal{C}}(\lambda)$, then \mathcal{C} is a maximal convex inner approximation of \mathcal{F} .*

We call z^* in Thm. 2 an *outlier point*, by which we try to enlarge an \mathcal{S} -free set, and let $L(z^*) := \{\lambda \in \text{dom}(\sigma_{\mathcal{C}}) : \lambda \cdot z^* > \sigma_{\mathcal{C}}(\lambda)\}$. The proof of Thm. 2 (in the appendix) was adapted from [53, Thm. 2], which excludes the presence of the outlier point and requires a stronger assumption, namely that for any $\lambda \in \text{dom}(\sigma_{\mathcal{C}})$ there exists a point $z' \in \text{bd}(\mathcal{F}) \cap \text{bd}(\mathcal{C})$ exposing $\lambda \cdot z \leq \sigma_{\mathcal{C}}(\lambda)$. As we will see in the proof of Thm. 3, $\bigcup_{z^* \in \text{int}(\mathcal{F} \setminus \mathcal{C})} L(z^*)$ can be a proper subset of $\text{dom}(\sigma_{\mathcal{C}})$, so we do not need to check that all $\lambda \in \text{dom}(\sigma_{\mathcal{C}})$ are exposed.

We next focus on a specific type of function, namely *positive homogeneous functions*. We summarize their properties as follows.

Lemma 3. *Let f be a positive homogeneous function of degree $d \in \mathbb{R}$, such that, for any $z \in \text{dom}(f) \subseteq \mathbb{R}^p$ and any $\rho \in \mathbb{R}_{++}$, $f(\rho z) = \rho^d f(z)$. Then: (i) $\text{int}(\text{dom}(f))$ is a cone, and (ii) if $d = 1$, then for any $\check{z} \in \text{dom}(f)$, $\Xi_{\check{z}}^f(z) = \nabla f(\check{z}) \cdot z$ for $z \in \text{dom}(f)$ and $\Xi_{\check{z}}^f(z) = f(z)$ for $z = \rho \check{z}$ with $\rho \in \mathbb{R}_{++}$.*

The proof is in the appendix. We recall that $\Xi_{\check{z}}^f$ in the above lemma is defined in Eq. (5). Moreover, $\text{dom}(f)$ is embedded in \mathbb{R}^p , so we call \mathbb{R}^p the *ambient space* of f .

The second theorem we present offers a more structured result, specifically related to nonconvex DCC sets \mathcal{S} . [60, Thm. 5.48] provides a sufficient condition for the maximality of the \mathcal{S} -free set described in Lemma 1. However, to clearly distinguish it from our result below, we translate the condition into our setting as follows: (i) the functions f_1 and f_2 are superlinear, i.e. they are positive homogeneous of degree 1 and superadditive (note that superlinear functions are concave), (ii) they are separable and act independently on different variables u and v , (iii)

f_1 is negative everywhere except at 0, (iv) the linearization point \tilde{v} of f_2 is nonzero, and (v) the domains $\text{dom}(f_1)$ and $\text{dom}(f_2)$ are Euclidean spaces.

Our second theorem provides an alternative condition for maximality that relaxes condition (i) by requiring only that one of f_1 or f_2 be positive homogeneous of degree 1, while imposing mild regularity conditions. Moreover, the domains can be full-dimensional convex cones.

Theorem 3. *For every $i \in \{1, 2\}$, let f_i be concave. Let $\mathcal{S} := \{(u, v) \in \text{dom}(f_1) \times \text{dom}(f_2) : f_1(u) - f_2(v) \leq 0\}$. Suppose that: (i) at least one of f_1, f_2 is positive homogeneous of degree 1, (ii) f_1, f_2 are both positive/negative over the interiors of their domains, (iii) f_1 is continuously differentiable over $\text{int}(\text{dom}(f_1))$, and (iv) $\text{dom}(f_1), \text{dom}(f_2)$ are full-dimensional in the ambient spaces of f_1, f_2 , respectively. Then, for any $\tilde{v} \in \text{int}(\text{dom}(f_2))$, $\mathcal{C} := \{(u, v) \in \text{dom}(f_1) \times \text{dom}(f_2) : f_1(u) - \Xi_{\tilde{v}}^{f_2}(v) \geq 0\}$ is maximally \mathcal{S} -free in $\text{dom}(f_1) \times \text{dom}(f_2)$.*

Proof. We first adapt Lemma 1 by restricting the domain of z to the convex ground set $\mathcal{G} := \text{dom}(f_1) \times \text{dom}(f_2)$. It follows from Lemma 1 that \mathcal{C} is an \mathcal{S} -free set in \mathcal{G} . Since $\text{dom}(f_1) \times \text{dom}(f_2)$ are full-dimensional, $\mathcal{S}, \mathcal{C}, \mathcal{G}$ are full-dimensional. As $\mathcal{S}, \mathcal{C} \subseteq \mathcal{G}$, the maximality of \mathcal{C} in \mathcal{G} is equivalent to that \mathcal{C} is a maximal convex inner approximation of $\mathcal{F} := \text{cl}(\mathcal{S}^c) \cap \mathcal{G} = \{(u, v) \in \mathcal{G} : f_1(u) - f_2(v) \geq 0\}$. Note that \mathcal{F} is full-dimensional. We then apply Thm. 2 to prove that \mathcal{C} is a maximal convex inner approximation of \mathcal{F} . Let $z^* \in \text{int}(\mathcal{F} \setminus \mathcal{C})$ be any outlier point. It follows from the separating hyperplane theorem that there exists a supported valid inequality $\lambda \cdot z \leq \sigma_{\mathcal{C}}(\lambda)$ of \mathcal{C} such that $\lambda \cdot z^* > \sigma_{\mathcal{C}}(\lambda)$. Since $\mathcal{F} \setminus \mathcal{C} \subseteq \mathcal{G}$, $\text{int}(\mathcal{F} \setminus \mathcal{C}) \subseteq \mathcal{G}$. Since $\mathcal{C} \subseteq \mathcal{G}$, the inequality cannot be supported by a valid inequality at $\text{bd}(\mathcal{G})$, so the inequality must be a valid inequality supported at $\mathcal{C} \setminus \text{bd}(\mathcal{G})$. It follows from the concavity of f_1 that the inequality must admit the form $\Xi_{\tilde{u}}^{f_1}(u) - \Xi_{\tilde{v}}^{f_2}(v) \geq 0$ for some $\tilde{u} \in \text{dom}(f_1)$ (identical up to a positive multiplier). By the smoothness of f_1 , w.l.o.g, we can perturb \tilde{u} such that it is in $\text{int}(\text{dom}(f_1))$. Let $\tilde{v} := \tilde{v}$. We now have that $\tilde{u} \in \text{int}(\text{dom}(f_1)), \tilde{v} \in \text{int}(\text{dom}(f_2))$. We will prove that $\Xi_{\tilde{u}}^{f_1}(u) - \Xi_{\tilde{v}}^{f_2}(v) \geq 0$ is exposed by a point $(u', v') \in (\text{bd}(\mathcal{F}) \cap \text{bd}(\mathcal{C})) \cap \text{int}(\mathcal{G})$. It suffices to show that the following three equations hold:

$$\begin{aligned} \Xi_{\tilde{u}}^{f_1}(u') - \Xi_{\tilde{v}}^{f_2}(v') &= 0 \quad (\text{i.e., supported at } (u', v')), \\ f_1(u') - \Xi_{\tilde{v}}^{f_2}(v') &= 0 \quad (\text{i.e., } (u', v') \in \mathcal{C}), \\ f_1(u') - f_2(v') &= 0 \quad (\text{i.e., } (u', v') \in \mathcal{F}). \end{aligned} \tag{8}$$

Since $\mathcal{C} \subseteq \mathcal{F}$ and they are both full-dimensional, the last two equations imply that $(u', v') \in \text{bd}(\mathcal{C}) \cap \text{bd}(\mathcal{F})$. As f_1 is continuously differentiable and concave in the interior of its domain, the graph of $f_1(u) - \Xi_{\tilde{v}}^{f_2}(v)$ over $\text{int}(\mathcal{G})$ is a smooth manifold embedded in $\text{int}(\mathcal{G}) \times \mathbb{R}$. The intersection of a smooth manifold with a hyperplane yields another lower-dimensional smooth manifold. This implies that the level set \mathcal{C} of $f_1(u) - \Xi_{\tilde{v}}^{f_2}(v)$ is also smooth at any point $(u, v) \in \text{int}(\mathcal{G}) \cap \mathcal{C}$. By Obs 2, (u, v) is an exposing point. Since $(u', v') \in \mathcal{C} \cap \text{int}(\mathcal{G})$, (u', v') is an exposing point, and the maximality of \mathcal{C} is verified. We now proceed to construct (u', v') from (\tilde{u}, \tilde{v}) and prove (8). Let $\rho := f_2(\tilde{v})/f_1(\tilde{u})$. Since $\tilde{u} \in \text{int}(\text{dom}(f_1)), \tilde{v} \in \text{int}(\text{dom}(f_2))$, by the assumption, $\rho > 0$. We consider the following two cases separately.

Case i. We first suppose that f_1 is positive homogeneous of degree 1. Let $(u', v') := (\rho\tilde{u}, \tilde{v})$, which, by Lemma 3, is in $\text{int}(\mathcal{G})$. We have that:

$$f_1(u') \stackrel{(i.1)}{=} \Xi_{\tilde{u}}^{f_1}(u') \stackrel{(i.2)}{=} \rho f_1(\tilde{u}) \stackrel{(i.3)}{=} f_2(\tilde{v}) \stackrel{(i.4)}{=} f_2(v') \stackrel{(i.5)}{=} \Xi_{\tilde{v}}^{f_2}(v'),$$

where equations (i.1), (i.2) follow from Lemma 3, (i.3) follows from the definition of ρ , and (i.4), (i.5) follow from $v' = \tilde{v}$.

Case ii. We then suppose that f_2 is positive homogeneous of degree 1. Let $(u', v') := (\check{u}, \check{v}/\rho) \in \text{int}(\mathcal{G})$. We have that:

$$\Xi_{\check{u}}^{f_1}(u') \stackrel{(ii.1)}{=} f_1(u') \stackrel{(ii.2)}{=} f_1(\check{u}) \stackrel{(ii.3)}{=} f_2(\check{v})/\rho \stackrel{(ii.4)}{=} f_2(v') \stackrel{(ii.5)}{=} \Xi_{\check{v}}^{f_2}(v'),$$

where equations (ii.1), (ii.2) follow from $\check{u} = u'$, (ii.3) follows from the definition of ρ , and (ii.4), (ii.5) follow from Lemma 3. Therefore, (8) are satisfied in both cases. \square

We present the motivation for restricting the maximality of the set \mathcal{C} within the ground set $\text{dom}(f_1) \times \text{dom}(f_2)$. The main reason for this restriction arises from the difficulty of finding a nontrivial concave extension of f_1 over its ambient space such that for all $u \notin \text{dom}(f_1)$, $f_1(u) > -\infty$. While such an extension can exist geometrically, the construction of a closed expression remains unclear. In the next section, we will examine a specific example to illustrate this point.

Moreover, we will apply the above theorem to develop DCC formulations for a nonconvex set. In particular, the functions f_1 and f_2 must not simultaneously have positive homogeneity of degree 1, and their domains are non-negative orthants. Consequently, the relaxed condition for homogeneous degrees and domains in Thm. 3 becomes necessary. We give two examples for verification Thm. 3.

Example 2. Let $f_1(u) := u$ with $\text{dom}(f_1) \in \mathbb{R}$, and let $f_2(v) := \sum_{i \in [n]} \sqrt{v_i}$ with $\text{dom}(f_2) = \mathbb{R}_+^n$. Note that f_1, f_2 are concave, $\text{dom}(f_2)$ is a non-negative orthant, and f_1 is positive homogeneous of degree 1. Let $\mathcal{G} := \mathbb{R} \times \mathbb{R}_+^n$. One can verify that the presupposition of Thm. 3 is satisfied. Then, $\mathcal{S} := \{(u, v) \in \mathcal{G} : u - \sum_{i \in [n]} \sqrt{v_i} \leq 0\}$ is a convex set. It is easy to see that $\mathcal{C} := \{(u, v) \in \mathcal{G} : u - \sum_{i \in [n]} (\sqrt{\tilde{v}_i} + (v_i - \tilde{v}_i)/\sqrt{\tilde{v}_i}) \geq 0\}$ is maximally \mathcal{S} -free in \mathcal{G} with $\tilde{v} > 0$.

Example 3. Exchange the functions f_1, f_2 in the previous examples. Then, $\mathcal{S} := \{(u, v) \in \mathcal{G} : \sum_{i \in [n]} \sqrt{v_i} - u \leq 0\}$ is a reverse-convex set. It is easy to see that $\mathcal{C} := \{(u, v) \in \mathcal{G} : \sum_{i \in [n]} \sqrt{v_i} - u \geq 0\}$ is the unique maximal \mathcal{S} -free set in \mathcal{G} .

4 Signomial-lift-free sets and intersection cuts

In this section, we construct (maximal) signomial-lift-free sets and generate intersection cuts for SP.

4.1 Signomial-lift-free and signomial-term-free sets

We introduce and study new formulations of signomial term sets. We transform signomial term sets into DCC sets. We also construct signomial term-free sets and lift them to signomial term-lift-free sets. The maximality of these sets is studied, and a comparison is made between signomial term-free sets derived from different DCC formulations.

We consider an n -variate signomial term $\psi_\alpha(x)$ arising in the extended formulation (2) of SP. The exponent vector α may contain negative/zero/positive entries. We extract two sub-vectors α^- and α^+ from α such that $\alpha^- \in \mathbb{R}_{--}^{h'}$ and $\alpha^+ \in \mathbb{R}_{++}^{\ell'}$, and let $x_- \in \mathbb{R}^{h'}$ and $x_+ \in \mathbb{R}^{\ell'}$ be the corresponding sub-vectors of x . Entries x_j with $\alpha_j = 0$ are excluded from consideration, and so $h' + \ell'$ may be smaller than n . Since $\psi_\alpha(x)$ only depends on x_- and x_+ , it can be represented in the form of $x_-^{\alpha^-} x_+^{\alpha^+}$ of lower order. We consider the *signomial term set* as epigraph or hypograph of $x_-^{\alpha^-} x_+^{\alpha^+}$:

$$\mathcal{S}_{\text{st}} = \{(x_-, x_+, t) \in \mathbb{R}_{++}^{h'+\ell'+1} : t \leq x_-^{\alpha^-} x_+^{\alpha^+}\}. \quad (9)$$

We first give DCC reformulations of signomial term sets. Let \lesseqgtr denote $<$ or $>$. The interior of \mathcal{S}_{st} in (9) is

$$\text{int}(\mathcal{S}_{\text{st}}) = \{(x_-, x_+, t) \in \mathbb{R}_{++}^{h'+\ell'+1} : t \lesseqgtr x_-^{\alpha^-} x_+^{\alpha^+}\}.$$

Reorganizing the signomial terms and taking the closure of the set, we recover

$$\mathcal{S}_{\text{st}} = \{(x_-, x_+, t) \in \mathbb{R}_{++}^{h'+\ell'+1} : tx_-^{\alpha^-} \lesseqgtr x_+^{\alpha^+}\}.$$

Notably, the exponents associated with signomial terms on both sides are now strictly positive. Let $u := (t, x_-)$, $v := x_+$, let $h := h' + 1$, and let $\ell := \ell'$. Then, $\psi_{\beta'}(u) = tx_-^{\alpha^-}$ and $\psi_{\gamma'}(v) = x_+^{\alpha^+}$, where $\beta' := (1, -\alpha^-) \in \mathbb{R}_{++}^h$ and $\gamma' := \alpha^+ \in \mathbb{R}_{++}^\ell$. After the change of variables, the set admits the following form:

$$\mathcal{S}_{\text{st}} = \{(u, v) \in \mathbb{R}_{++}^{h+\ell} : \psi_{\beta'}(u) \lesseqgtr \psi_{\gamma'}(v)\}. \quad (10)$$

The formulation (10) exhibits symmetry between u and v . We can therefore consider w.l.o.g. the inequality “ \leq ” throughout the subsequent analysis. Since the signomial terms $\psi_{\beta'}(u), \psi_{\gamma'}(v)$ are non-negative over $\mathbb{R}_{++}^h, \mathbb{R}_{++}^\ell$, we can take any positive power $\eta \in \mathbb{R}_{++}$ on both sides of (10). Finally, the signomial term set in (9) admits the following form:

$$\mathcal{S}_{\text{st}} = \{(u, v) \in \mathbb{R}_{++}^{h+\ell} : \psi_\beta(u) - \psi_\gamma(v) \leq 0\}, \quad (11)$$

where $\beta := \eta\beta'$, and $\gamma := \eta\gamma'$.

A signomial term $\psi_\alpha(x)$ is said to be a *power function* if $\alpha \geq 0$, and $\|\alpha\|_1 \leq 1$. According to [4, 16], power functions are concave over the non-negative orthant; if additionally $\|\alpha\|_1 = 1$, $\psi_\alpha(x)$ is positive homogeneous of degree 1. Through an appropriate scaling of the parameter η , we obtain a family of DCC reformulations (11) of signomial term sets. We let $\mathcal{G} := \mathbb{R}_{++}^{h+\ell}$, and use the reverse-linearization technique to construct signomial-term-free sets. We recall that the definition of the operator Ξ is given in Eq. (5).

Proposition 1. *Let $\max(\|\beta\|_1, \|\gamma\|_1) \leq 1$. For any $\tilde{v} \in \mathbb{R}_{++}^\ell$,*

$$\mathcal{C} := \{(u, v) \in \mathbb{R}_{++}^h \times \mathbb{R}^\ell : \psi_\beta(u) - \Xi_{\tilde{v}}^{\psi_\gamma}(v) \geq 0\} \quad (12)$$

is a signomial-term-free (\mathcal{S}_{st} -free) set. If $\max(\|\beta\|_1, \|\gamma\|_1) = 1$, then \mathcal{C} is a maximal signomial-term-free set in \mathcal{G} .

Proof. Since $\max(\|\beta\|_1, \|\gamma\|_1) \leq 1$, $\psi_\beta(u), \psi_\gamma(v)$ are concave. By Lemma 1, \mathcal{C} is signomial-term-free. If $\max(\|\beta\|_1, \|\gamma\|_1) = 1$, then at least one of $\|\beta\|_1, \|\gamma\|_1$ is 1. Therefore, one of $\psi_\beta(u), \psi_\gamma(v)$ is positive homogeneous of degree 1. Moreover, $\psi_\beta(u), \psi_\gamma(v)$ are both continuously differentiable and positive over positive orthants $\mathbb{R}_{++}^h, \mathbb{R}_{++}^\ell$ (the interiors of their domains). Since $\mathcal{G} = \text{dom}(\psi_\beta) \times \text{dom}(\psi_\gamma)$, by Thm. 3, $\mathcal{C} \cap \mathcal{G} = \{(u, v) \in \mathcal{G} : \psi_\beta(u) - \Xi_{\tilde{v}}^{\psi_\gamma}(v) \geq 0\}$ is a maximal signomial-term-free set in \mathcal{G} . Therefore, \mathcal{C} is also a maximal signomial-term-free set in \mathcal{G} . \square

Given that $\max(\|\beta\|_1, \|\gamma\|_1) = 1$ results in a desirable DDC formulation for the signomial term set, we refer to this formulation as its *normalized DCC formulation*. Comparing Prop. 1 to Thm. 3, we extend the domain of $\Xi_{\tilde{v}}^{\psi_\gamma}(v)$ from \mathbb{R}_{++}^ℓ to \mathbb{R}^ℓ , since it is an affine function. However, the further extension requires a non-trivial concave extension of the power function ψ_β , which we are unaware of.

We have reduced the n -variate signomial term $\psi_\alpha(x)$ to a signomial term $x_-^{\alpha^-} x_+^{\alpha^+}$ of lower order and constructed the corresponding signomial-term-free sets. A similar reduction is observed for g_i to g'_i in Subsec. 3.1, where we demonstrate the relationship between $\mathcal{S}_{\text{lift}}$ -free sets and $\text{epi}(g'_i)$ -free/hypo(g'_i)-free sets.

Next, we let the lifted set $\mathcal{S}_{\text{lift}}$ be the signomial lift, where all g_i are signomial terms. Each equality constraint $y_i = g_i(x)$ defining the signomial lift is equivalent to two inequality constraints $y_i \leq g_i(x)$ and $y_i \geq g_i(x)$. Applying the normalized DDC reformulation to these inequality constraints, we thus obtain a reformulation of the signomial lift, which we call its *normalized DCC reformulation*.

Corollary 1. *Let \mathcal{C} be as in (12), where $\psi_\alpha = g_i$ for some $i \in [k]$ and $\max(\|\beta\|_1, \|\gamma\|_1) = 1$. Then the orthogonal lifting of \mathcal{C} w.r.t. g_i is a maximal signomial-lift-free ($\mathcal{S}_{\text{lift}}$ -free) set in the non-negative orthant.*

Proof. We verify that the conditions of Thm. 2 are satisfied by the signomial lift. For any $i \in [k]$, the signomial term g_i is continuous, and its domain and range are \mathbb{R}_{++} . Let J_i be the index set of variables of its reduced signomial term g'_i . Let $\mathcal{X} := \times_{j \in [n]} \mathbb{R}_{++}$, $\mathcal{Y} := \times_{j \in [k]} \mathbb{R}_{++}$. For all $j \in [n+k]$, let $\mathcal{D}_j := \mathbb{R}_{++}$. For all $i \in [k]$, let $\mathcal{X}^i := \times_{j \in J_i} \mathbb{R}_{++}$, $\mathcal{Y}^i := \mathbb{R}_{++}$. The underlying sets of the signomial lift are $\mathcal{X}, \mathcal{Y}, \{\mathcal{X}^i, \mathcal{Y}^i\}_{i \in [k]}$ that are 1d-convex decomposable by $\{\mathcal{D}_j\}_{j \in [n+k]}$. By Prop. 1, \mathcal{C} is a maximal hypo(g'_i)-free set in $\mathcal{X} \times \mathcal{Y}$. By Thm. 2, its orthogonal lifting w.r.t. g_i is a maximal signomial-lift-free set in positive orthant. By continuity of ψ_β, ψ_γ , we change the ground set (the positive orthant) to its closure, i.e., non-negative orthant. \square

The following examples show signomial term-free sets from different DDC formulations.

Example 4 (Comparison of DCC formulations). *Consider $\mathcal{S}_{\text{st}} = \{(u, v) \in \mathbb{R}_+^2 : u \leq v\}$, which is already in normalized DCC formulation. It is easy to see that $\mathcal{C}_1 := \{(u, v) \in \mathbb{R}_+ \times \mathbb{R} : u \geq v\}$ is a maximal \mathcal{S}_{st} -free set in \mathbb{R}_+^2 given by Prop. 1. Let $\tilde{v} \in \mathbb{R}_{++}$ be a linearization point. As $\text{int}(\mathcal{S}_{\text{st}}) = \{(u, v) \in \mathbb{R}_+^2 : \log(u) \leq \log(v)\}$ admits a DCC formulation, applying the reverse-linearization technique at \tilde{v} yields $\mathcal{C}_2 := \{(u, v) \in \mathbb{R}_+^2 : \log(u) - (\log(\tilde{v}) + (v - \tilde{v})/\tilde{v}) \geq 0\}$, which is also an \mathcal{S}_{st} -free set. For any $0 < \eta < 1$, $\mathcal{S}_{\text{st}} = \{(u, v) \in \mathbb{R}_+^2 : u^\eta \leq v^\eta\}$ is a DDC set, applying the reverse-linearization technique at \tilde{v} yields $\mathcal{C}_3 := \{(u, v) \in \mathbb{R}_+^2 : u^\eta - ((1-\eta)\tilde{v}^\eta + \eta\tilde{v}^{\eta-1}v) \geq 0\}$, which is also an \mathcal{S}_{st} -free set. However, $\mathcal{C}_2, \mathcal{C}_3$ cannot be maximal in \mathbb{R}_+^2 , because their intersections with \mathbb{R}_+^2 are not polyhedral. These sets are visualized in Fig. 2 with a linearization point $\tilde{v} = 0.5$ and scaling parameter $\eta = 0.7$.*




Figure 2: \mathcal{S}_{st} -free sets from Example 4.

Example 5. *Consider the hypograph of signomial term $x_1^{-2}x_2^2$ and $\mathcal{S}_{\text{st}} = \{(x, y) \in \mathbb{R}_+^3 : y \leq x_1^{-2}x_2^2\}$. For $(x, y) \in \mathbb{R}_{++}^3$, $y \leq x_1^{-2}x_2^2$ if and only if $y^{1/3}x_1^{2/3} \leq x_2^{2/3}$. The following set is maximal \mathcal{S}_{st} -free in $\mathcal{G} = \mathbb{R}_+^3$: $\mathcal{C}_4 := \{(x, y) \in \mathbb{R}_+^3 : y^{1/3}x_1^{2/3} \geq \tilde{x}_2^{2/3} + \frac{2}{3}\tilde{x}_2^{-1/3}(x_2 - \tilde{x}_2)\}$, where $\tilde{x}_2 \in \mathbb{R}_{++}$. See Fig. 3a for $\tilde{x}_2 = 0.2$.*

Example 6. Consider the epigraph of signomial term $x_1^3 x_2$ and $\mathcal{S}_{\text{st}} = \{(x, y) \in \mathbb{R}_+^3 : y \geq x_1^3 x_2\}$. For $(x, y) \in \mathbb{R}_{++}^3$, $y \geq x_1^3 x_2$ if and only if $y^{1/4} \geq x_1^{3/4} x_2^{1/4}$. The following set is maximal \mathcal{S}_{st} -free in $\mathcal{G} = \mathbb{R}_+^3$: $\mathcal{C}_5 := \{(x, y) \in \mathbb{R}_+^3 : \tilde{y}^{1/4} + \frac{1}{4}\tilde{y}^{-3/4}(y - \tilde{y}) \leq x_1^{3/4} x_2^{1/4}\}$, where $\tilde{y} \in \mathbb{R}_{++}$. See Fig. 3b for $\tilde{y} = 0.2$.




Figure 3: \mathcal{S}_{st} and \mathcal{S}_{st} -free sets from Examples 5 and 6.

4.2 Intersection cuts

We focus on the separation of intersection cuts for the extended formulation of SP. In Sec. 2 we presented a method to construct a simplicial cone \mathcal{R} from an LP relaxation. The vertex of this cone is a relaxation solution $\tilde{z} = (\tilde{x}, \tilde{y})$.

We assume that the LP relaxation includes all linear constraints from (2). If \tilde{z} is infeasible for (2), then \tilde{z} does not belong to the signomial lift. Thus, there is a signomial term g_i such that $\tilde{y}_i \neq g_i(\tilde{x})$. Given the reduced form g'_i , we obtain a set of signomial terms \mathcal{S}_{st} : If $g_i(\tilde{x}) > \tilde{y}_i$, we choose \mathcal{S}_{st} to be the epigraph of g'_i ; otherwise, we choose it to be the hypograph of g'_i . This signomial term set yields a signomial term-free set \mathcal{C} in (12) containing (\tilde{u}, \tilde{v}) in its interior (Lemma 1). Using orthogonal lifting of Cor. 1, we can transform \mathcal{C} into a signomial-lift-free set $\bar{\mathcal{C}}$.

We next show how to construct an intersection cut in (7). It suffices to compute step lengths η_j^* in (6) along extreme rays r^j of \mathcal{R} . Each step length η_j^* corresponds to a boundary point $\tilde{z} + \eta_j^* r^j$ in $\text{bd}(\bar{\mathcal{C}})$. The left-hand-side $\psi_\beta(u) - \Xi_v^{\psi_\gamma}(v)$ of the inequality in (12) is a concave function over $(u, v) \in \mathbb{R}_+^h \times \mathbb{R}^\ell$. Its restriction along the ray $\tilde{z} + \eta_j r^j$ ($\eta_j \in \mathbb{R}_+$) is a univariate concave function:

$$\tau_j : \mathbb{R}_+ \rightarrow \mathbb{R}, \eta_j \mapsto \tau_j(\eta_j) := \psi_\beta(\tilde{u} + r_u^j \eta_j) - \Xi_v^{\psi_\gamma}(\tilde{v} + r_v^j \eta_j),$$

where r_u^j and r_v^j are the projections of r^j on u and v respectively. Let $\bar{\eta}_j := \sup_{\eta_j \geq 0} \{\eta_j : \tilde{u} + r_u^j \eta_j \geq 0\}$. Therefore, η_j^* is the first point in $[0, \bar{\eta}_j]$ satisfying the boundary condition: either $\tau_j(\eta_j^*) = 0$ or $\eta_j^* = \bar{\eta}_j$. Since τ_j is a univariate concave function and $\tau_j(0) > 0$, there is at most one positive point in \mathbb{R}_+ where τ_j is zero. We employ the bisection search method [56] to find such η_j^* .

5 Convex outer approximation

In this section we propose a convex nonlinear relaxation for the extended formulation (2) of SP. This relaxation is easy to derive and allows us to generate valid linear inequalities, called outer approximation cuts, for SP. Unlike intersection cuts, outer approximation cuts do not require an LP relaxation *a priori*, so solvers can employ them to generate an initial LP relaxation of (2).

To ensure the convergence of the sBB algorithm, the feasible region of the extended formulation (2) should be compact. Therefore, we assume in the sequel that the signomial lift is in a hypercube.

We construct the convex nonlinear relaxation by approximating each signomial term set of the signomial lift within the hypercube. W.l.o.g., we consider a signomial term set in hypergraph or epigraph form. As Subsec. 4.1, we can convert it into normalized DDC formulation:

$$\mathcal{S}_{\text{st}} = \{(u, v) \in \mathcal{U} \times \mathcal{V} : \psi_\beta(u) - \psi_\gamma(v) \leq 0\}, \quad (13)$$

where $\max(\|\beta\|_1, \|\gamma\|_1) = 1$, and \mathcal{U}, \mathcal{V} are two hypercubes in $\mathbb{R}_+^h, \mathbb{R}_+^\ell$ respectively. Due to Subsec. 5.3, the signomial term set is usually nonconvex, so our construction involves convexifying the concave function ψ_β in (13). This procedure yields a convex outer approximation of \mathcal{S}_{st} , which is non-polyhedral. Consequently, replacing \mathcal{S}_{st} by its convex outer approximation, we obtain the convex nonlinear relaxation of (2).

Next, we introduce the procedure of convexification. We should import the formal concepts of convex underestimators and convex envelopes. Given a function f and a closed set $\mathcal{D} \subseteq \mathbb{R}^p$, a convex function $f' : \text{conv}(\mathcal{D}) \rightarrow \mathbb{R}$ is called a convex underestimator of f over \mathcal{D} , if for all $x \in \mathcal{D}$ $f'(x) \leq f(x)$. The convex envelope $\text{conv}_{\mathcal{D}}(f)$ of f is defined as the pointwise maximum convex underestimator of f over \mathcal{D} .

The following lemma gives an extended formulation of the convex envelope of a concave function over a polytope, where the formulation is uniquely determined by the function values at the vertices of the polytope.

Lemma 4. [26, Thm. 3] *Let P be a polytope in \mathbb{R}^n , let $f : P \rightarrow \mathbb{R}$ be a concave function over P , and let Q be vertices of P . Then, $\text{conv}_{\mathcal{D}}(f)(x) = \max\{\sum_{q \in Q} \lambda_q f(q) : \exists \lambda \in \mathbb{R}_+^Q, \sum_{q \in Q} \lambda_q = 1, x = \sum_{q \in Q} \lambda_q q\}$.*

Based on the lemma above, we observe that the concave function f is convex-extensible from its vertices (i.e., $\text{conv}_{\mathcal{D}}(f)(x) = \text{conv}_{\mathcal{Q}}(f)(x)$ for $x \in P$), and $\text{conv}_{\mathcal{D}}(f)$ is a polyhedral function.

For the case of $P = \mathcal{U} := \prod_{j \in [h]} [\underline{u}_j, \bar{u}_j]$ and $f = \psi_\beta$, $Q = \{q \in \mathbb{R}^h : \forall j \in [h] q_j = \underline{u}_j \vee q_j = \bar{u}_j\}$ is the set of vertices of the hypercube \mathcal{U} . The lemma yields an extended formulation of $\text{conv}_{\mathcal{U}}(\psi_\beta)$. Replacing ψ_β by its convex envelope $\text{conv}_{\mathcal{U}}(\psi_\beta)$, we obtain the convex outer approximation of \mathcal{S}_{st} in (13):

$$\bar{\mathcal{S}}_{\text{st}} := \{(u, v) \in \mathcal{U} \times \mathcal{V} : \text{conv}_{\mathcal{U}}(\psi_\beta)(u) \leq \psi_\gamma(v)\}.$$

By using this extended formulation, our convex nonlinear relaxation of SP contains additional auxiliary variables. In particular, we need 2^h variables λ_q to represent each convex envelope. For most SP problems in MINLPLib where the degrees of the signomial terms are less than 6 and h is less than 3, the convex nonlinear relaxation is computationally tractable.

5.1 Outer approximation cuts

To increase efficiency, we propose a cutting plane algorithm to separate valid linear inequalities in (u, v) -space from the extended formulation of the convex outer approximation. This algorithm

generates a low-dimensional projected approximation of $\overline{\mathcal{S}}_{\text{st}}$. Moreover, the projection procedure converts the convex nonlinear relaxation into an LP relaxation, which is suitable for many solvers.

Given a point $(\tilde{u}, \tilde{v}) \in \mathcal{U} \times \mathcal{V}$, the algorithm determines whether it belongs to $\overline{\mathcal{S}}_{\text{st}}$. This verification can be done by checking the sign of $\text{conv}_{\mathcal{U}}(\psi_{\beta})(\tilde{u}) - \psi_{\gamma}(\tilde{v})$. If $\text{conv}_{\mathcal{U}}(\psi_{\beta})(\tilde{u}) - \psi_{\gamma}(\tilde{v}) \leq 0$, then $(\tilde{u}, \tilde{v}) \in \overline{\mathcal{S}}_{\text{st}}$.

Since $\text{conv}_{\mathcal{U}}(\psi_{\beta})$ is a convex polyhedral function, our cutting plane algorithm evaluates the function by searching for an affine underestimator $a \cdot u + b$ of $\text{conv}_{\mathcal{U}}(\psi_{\beta})$ such that $a \cdot \tilde{u} + b = \text{conv}_{\mathcal{U}}(\psi_{\beta})(\tilde{u})$. If $(\tilde{u}, \tilde{v}) \notin \overline{\mathcal{S}}_{\text{st}}$, then $a \cdot u + b \leq \psi_{\gamma}(v)$ is a valid nonlinear inequality of $\overline{\mathcal{S}}_{\text{st}}$. Subsequently, our cutting plane algorithm linearizes this inequality, resulting in an outer approximation cut $a \cdot u + b \leq \Xi_{\tilde{v}}^{\psi_{\gamma}}(v)$: we recall that $\Xi_{\tilde{v}}^{\psi_{\gamma}}(v)$ is the linearization of $\psi_{\gamma}(v)$ at \tilde{v} defined in Eq. (5).

Due to Lemma 4, we can solve the following LP to find the affine underestimator:

$$\max_{a \in \mathbb{R}^h, b \in \mathbb{R}} a \cdot \tilde{u} + b \quad \text{s.t. } \forall q \in Q \quad a \cdot q + b \leq \psi_{\gamma}(q), \quad (14)$$

where we omit the linear constraints that bound (a, b) . The maximum value resulting from this LP is exactly $\text{conv}_{\mathcal{U}}(\psi_{\beta})(\tilde{u})$. The affine underestimator $a \cdot u + b$ is called a *facet* of the envelope $\text{conv}_{\mathcal{U}}(\psi_{\beta})$, if $a \cdot u + b \leq t$ is a facet of $\text{epi}(\text{conv}_{\mathcal{U}}(\psi_{\beta}))$. It should be noted that the solution of the LP is not necessarily a facet.

For $h = 1, 2$ we can provide explicit projected formulations of convex envelopes of power functions. This allows us to obtain facets of $\text{conv}_{\mathcal{U}}(\psi_{\beta})$ without having to solve LPs. As a result, our cutting plane algorithm can efficiently separate outer approximation cuts for low-order problems.

To simplify our representation, we translate and scale the domain of ψ_{β} to $[0, 1]^h$. This leads to a new function $s(w) := \psi_{\beta}(u)$, where for all $j \in [h]$, $u_j := \bar{u}_j + (\bar{u}_j - \underline{u}_j)w_j$. After these transformations, we have $\mathcal{U} = [0, 1]^h$ and $Q = \{0, 1\}^h$. W.l.o.g., we focus on the study and computation of facets of $\text{conv}_{\mathcal{U}}(s)$. For $h = 1$, the only facet is $s(0) + (s(1) - s(0))w_1$.

A set $D \subseteq \mathbb{R}^h$ is called a *product set*, if $D = \times_{j \in [h]} D_j$ for $D_j \subseteq \mathbb{R}$. Moreover, a function $f : D \rightarrow \mathbb{R}$ is *supermodular* over D ([66, Sec. 2.6.1]), if the increasing difference condition holds: for all $w^1, w^2 \in D, d \in \mathbb{R}_+^h$ such that $w^1 \leq w^2$ and $w^1 + d, w^2 + d \in D$, $f(w^1 + d) - f(w^1) \leq f(w^2 + d) - f(w^2)$. We find that the following operations preserve supermodularity.

Lemma 5. *Let $w' \in \mathbb{R}^h, \rho \in \mathbb{R}_{++}^h$, and let D' be a product subset of D . The following results hold: (restriction) f is supermodular over D' ; (translation) $f(w + w')$ is supermodular over $D - d$; (scaling) $f(\rho * w)$ is supermodular over D/ρ , where $+, -, *, /$ are taken entry-wise.*

Proof. The results follow from the definition. □

We note that when $D = Q$, d is in Q . We observe a useful property of s .

Proposition 2. *s is supermodular over Q and $\text{conv}_{\mathcal{U}}(s) = \text{conv}_Q(s)$.*

Proof. According to [66, Example 2.6.2], the signomial term ψ_{α} with $\alpha > 0$ is supermodular over \mathbb{R}_+^h . This implies that the power function ψ_{β} is supermodular over \mathbb{R}_+^h . By Lemma 5, s is supermodular over $\mathcal{U} = [0, 1]^h$. As $Q = \{0, 1\}^h$ is a product subset of \mathcal{U} , s is supermodular over Q . After the scaling and translation, s is still concave. By Lemma 4, $\text{conv}_{\mathcal{U}}(s) = \text{conv}_Q(s)$. □

Finding facets of s can be reduced to a more general problem of finding facets of supermodular functions over Boolean hypercubes. We note that a similar argument can show that both power functions and multilinear terms over any product subset of \mathbb{R}_+^h are supermodular.

5.2 Projected convex envelopes of bivariate supermodular functions

We present a general characterization of projected convex envelopes of supermodular functions and give a closed-form expression for bivariate cases. Let f be a supermodular function over Q . We can use a bit representation to denote Boolean points in Q . For example, 10 denotes the point w that $w_1 = 1$ and $w_2 = 0$. For an affine function $a \cdot w + b$, we call Boolean points in Q where $a \cdot w + b$ equals $f(w)$ its *supporting points*.

Lemma 6. *Each facet of f is supported by $h + 1$ affinely independent points in Q .*

See the proof in the appendix. We can enumerate all possible subsets of $h + 1$ affinely independent points of Q . Each such subset $S := \{w^1, \dots, w^{h+1}\}$ determines a function over \mathbb{R}^h via the following affine combination:

$$f_S(w) := \left\{ \sum_{j \in [h+1]} \lambda_j f(w^j) : \exists \lambda \in \mathbb{R}^{h+1} \sum_{j \in [h+1]} \lambda_j = 1 \wedge \sum_{j \in [h+1]} \lambda_j w^j = w \right\}.$$

Because of the affine independence of S , the Barycentric coordinate λ is unique for any w in the above affine combination. We can consider f_S as a single-valued affine function and call it the *supported function* of S . Since $f_S(w) = f(w)$ solves the linear system $a \cdot w + b = f(w)$ (for $w \in S$), we can compute a, b which defines f_S . If the supported function f_S underestimates f , we call the subset S *facet-inducing*.

Assuming we have a collection of facets of f , we then determine whether these facets define the convex envelope of f . We recall that the convex hull of $h + 1$ affinely independent points is an h -*simplex*. A finite collection of h -simplices $\{P_k\}_k$ is called a *triangulation* of the unit cube \mathcal{U} , if the following conditions hold: (i) $\bigcup_k P_k = \mathcal{U}$, (ii) $P_k \cap P_{k'}$ is empty or a face of both $P_k, P_{k'}$ for any k, k' , (iii) the vertices of P_k are contained in Q for all k .

Proposition 3. *Let $\{S_k\}_k$ be a collection of facet-inducing subsets of Q . For each k , let f_{S_k} be the supported function induced by S_k , and let $P_k := \text{conv}(S_k)$ be the simplex spanned by S_k . If $\{P_k\}_k$ is a triangulation of \mathcal{U} , then $\text{conv}_Q(f)(w) = \max_k f_{S_k}(w)$ for all $w \in \mathcal{U}$.*

Proof. Since $\bigcup_k P_k = \mathcal{U}$, for any $w \in Q$, there exists P_k such that $w \in P_k$. Since the vertices S_k of P_k are contained in Q , w must be in S_k . Therefore, $f_{S_k}(w) = f(w)$. This implies that $f(w) = \max_k f_{S_k}(w)$ for all $w \in Q$, i.e., $\max_k f_{S_k}$ is an exact convex underestimator. Suppose, to aim at a contradiction, that there exists another convex underestimator f' of f such that $f'(w) = f(w)$ for all $w \in Q$, and $f'(w') > \max_k f_{S_k}(w')$ for some $w' \in \mathcal{U}$. Again, since $\bigcup_k P_k = \mathcal{U}$, there exists P_k such that $w' \in P_k$, i.e., $w' \in \text{conv}(S_k)$. Let $S_k = \{w^1, \dots, w^{h+1}\}$. It follows from $w' \in \text{conv}(S_k)$ that there exists $\lambda \in [0, 1]^{h+1}$ such that $\sum_{j \in [h+1]} \lambda_j = 1, \sum_{j \in [h+1]} \lambda_j w^j = w'$. Note that S_k induces the supported function f_{S_k} , which has supporting points S_k . This implies that $f_{S_k}(w') = \sum_{j \in [h+1]} \lambda_j f_{S_k}(w^j) = \sum_{j \in [h+1]} \lambda_j f(w^j)$. By the convexity of f' , $f'(w') \leq \sum_{j \in [h+1]} \lambda_j f'(w^j) = \sum_{j \in [h+1]} \lambda_j f(w^j) \leq f_{S_k}(w') \leq \max_k f_{S_k}(w')$, which leads to a contradiction. Thus, $\max_k f_{S_k}$ is the convex envelope of f over Q . \square

The collection $\{S_k\}_k$ is called *envelope-inducing family* in Q , if the presupposition “ $\{P_k\}_k$ is a triangulation” in Prop. 3 is satisfied. Using the above result, we can construct an envelope-inducing family for bivariate supermodular functions. Let

$$S_1^2 := \{00, 10, 01\}, S_2^2 := \{11, 10, 01\}. \quad (15)$$

One can find that $\text{conv}(S_1^2) = \{(w_1, w_2) \in [0, 1]^2 : w_1 + w_2 \leq 1\}$, $\text{conv}(S_2^2) = \{(w_1, w_2) \in [0, 1]^2 : w_1 + w_2 \geq 1\}$ are two triangles in $[0, 1]^2$. We have that

$$\begin{aligned} f_{S_1^2}(w) &= f(00) + (f(10) - f(00))w_1 + (f(01) - f(00))w_2, \\ f_{S_2^2}(w) &= f(11) + (f(01) - f(11))(1 - w_1) + (f(10) - f(11))(1 - w_2). \end{aligned}$$

We show that these two affine functions define the convex envelope of f .

Theorem 4. For $h = 2$, $\{S_k^2\}_{k \in [2]}$ as in (15) is the envelope-inducing family in Q .

Proof. It is easy to see that, for all $k \in [2]$, S_k^2 is affinely independent and $\{\text{conv}(S_k^2)\}_{k \in [2]}$ is a triangulation of \mathcal{U} . Therefore, it suffices to show that $\{S_k^2\}_{k \in [2]}$ is facet-inducing, i.e., $f_{S_1^2}, f_{S_2^2}$ are affine underestimators of f .

Case i. We note that, for all $w \in S_1^2 = \{00, 10, 01\}$, $f_{S_1^2}(w) = f(w)$. Note that $Q \setminus S_1^2 = \{11\}$. It follows from the definition of the affine function $f_{S_1^2}$ that

$$f_{S_1^2}(11) = f_{S_1^2}(10) + (f_{S_1^2}(01) - f_{S_1^2}(00)) = f(10) + (f(01) - f(00)).$$

It follows from the supermodularity of f that

$$f(10) + (f(01) - f(00)) \leq f(10) + (f(11) - f(10)) = f(11).$$

Thereby, $f_{S_1^2}$ underestimates f .

Case ii. We note that, for all $w \in S_2^2 = \{11, 10, 01\}$, $f_{S_2^2}(w) = f(w)$. Note that $Q \setminus S_2^2 = \{00\}$. It follows from the definition of the affine function $f_{S_2^2}$ that

$$f_{S_2^2}(00) = f_{S_2^2}(10) - (f_{S_2^2}(11) - f_{S_2^2}(01)) = f(10) + (f(11) - f(01)).$$

It follows from the supermodularity of f that

$$f(10) - (f(11) - f(01)) \leq f(10) - (f(10) - f(00)) = f(00),$$

which concludes the proof. \square

5.3 Convexity and reverse-convexity

Our cutting plane algorithm can detect convexity/reverse-convexity of signomial term sets. The detection is easily done by normalized DDC formulations.

Denote by e_j^ℓ and e_j^h the j -th unit vector in \mathbb{R}^h and \mathbb{R}^ℓ , respectively. Then, we have the following observations:

- i) if $\|\beta\|_1 = 1, \gamma = 0$, i.e., ψ_β is concave and ψ_γ is 1, then \mathcal{S}_{st} is reverse-convex;
- ii) if $\|\beta\|_1 \leq 1, \gamma = e_j^\ell$ for some $j \in [\ell]$, i.e., ψ_β is concave and ψ_γ is a linear univariate function, then \mathcal{S}_{st} is reverse-convex;
- iii) if $\beta = e_j^h, \|\gamma\|_1 \leq 1$ for some $j \in [h]$, i.e., ψ_β is a linear univariate function and ψ_γ is concave, then \mathcal{S}_{st} is convex;
- iv) if $\|\beta\|_1 = 0, \|\gamma\|_1 = 1$, i.e., ψ_β is 1 and ψ_γ is concave, then \mathcal{S}_{st} is convex.

We note that similar results are found in [17, 45]. The results in [17] are proved by checking the negative/positive-semidefiniteness of the Hessian matrix of a signomial term. According to the normalized DCC formulation, the results are evident.

6 Computational results

In this section, we conduct computational experiments to assess the efficiency of the proposed valid inequalities.

The MINLPLib dataset includes instances of MINLP problems containing signomial terms, and some of these instances are SP problems. To construct our benchmark, we select instances from MINLPLib that satisfy the following criteria: (i) the instance contains signomial functions or polynomial functions, (ii) the continuous relaxation of the instance is nonconvex. Our benchmark consists of a diverse set of 251 instances in which nonlinear functions consist of signomial and other functions. These problems come from practical applications and can be solved by general purpose solvers.

Experiments are performed on a server with Intel Xeon W-2245 CPU @ 3.90GHz, 126GB main memory and Ubuntu 18.04 system. We use SCIP 8.0.3 [11] as a framework for reading and solving problems as well as performing cut separation. SCIP is integrated with CPLEX 22.1 as LP solver and IPOPT 3.14.7 as NLP solver.

We evaluate the efficiency of the proposed valid inequalities in four different settings. In the first setting, denoted `disable`, none of the proposed valid inequalities is applied. In the second setting, denoted `oc`, only the outer approximation cuts are applied. The third setting, denoted `ic`, applies only to the intersection cuts. The fourth setting combines both the `oc` and `ic` settings by applying both cuts. We let SCIP’s default internal cuts handle univariate signomial terms and multilinear terms. Our valid inequalities only handle the other high-order signomial terms. The source code, data, and detailed results can be found in our online repository: github.com/lidingxu/ESPCuts.

Each test run uses SCIP with a particular setting to resolve an instance. To solve the instances, we use the SCIP solver with its sBB algorithm and set a time limit of 3600 seconds. In our benchmark, there are 150 instances classified as *affected* in which at least one of the settings `oc`, `ic`, and `oic` settings adds cuts. Among the affected instances, there are 86 instances where the default SCIP configuration (i.e., `disable` setting) runs for at least 500 seconds. Such instances are classified as *affected-hard*. For each test run, we measure the runtime, the number of sBB search nodes, and the relative open duality gap.

To aggregate the performance metrics for a given setting, we compute shifted geometric means (SGMs) over our test set. The SGM for runtime includes a shift of 1 second. The SGM for the number of nodes includes a shift of 100 nodes. The SGM for relative distance includes a shift of 1%. We also compute the SGMs of the performance metrics over the subset of affected and affected-hard instances. The performance results are shown in Table 1, where we also compute the relative values of the SGMs of the performance metrics compared to the `disable` setting. Our following analysis is based on the results of the affected and affected-hard instances.

Setting		All				Affected				Affected-hard			
		solved	nodes	time	gap	solved	nodes	time	gap	solved	nodes	time	gap
<code>disable</code>	absolute	138/251	6510.5	122.0	4.7%	71/150	15592.4	253.6	5.7%	7/86	175973.8	3600.0	26.7%
	relative		1.0	1.0	1.0		1.0	1.0	1.0		1.0	1.0	1.0
<code>oc</code>	absolute	140/251	5954.1	118.0	4.5%	73/150	13443.9	241.4	5.4%	10/86	115262.3	2872.7	23.3%
	relative		0.91	0.97	0.97		0.86	0.95	0.95		0.65	0.8	0.87
<code>ic</code>	absolute	140/251	6144.3	122.4	4.4%	73/150	14081.5	252.1	5.2%	10/86	128072.7	2994.1	22.0%
	relative		0.94	1.0	0.95		0.9	0.99	0.91		0.73	0.83	0.82
<code>oic</code>	absolute	139/251	5934.6	117.7	4.6%	72/150	13275.6	236.8	5.6%	10/86	118054.1	2758.3	23.0%
	relative		0.91	0.96	0.99		0.85	0.93	0.98		0.67	0.77	0.86

Table 1: Summary of performance metrics on MINLPLib instances.

First, we note that the proposed valid inequalities lead to the successful solution of 2 additional instances compared to the `disable` setting. The `oc` setting solves 2 more instances than the `disable` setting.

The reductions in runtime and relative gap achieved by the `oc` setting are 5% and 5%, respectively, for affected instances and 20% and 13%, respectively, for affected-hard instances. The `ic` setting solves 2 more instances than the `disable` setting. The reduction in runtime and relative gap achieved by the `ic` setting is 1% and 9% for affected instances and 17% and 14% for affected-hard instances, respectively. The `oic` setting resolves 1 additional instance compared to the `disable` setting. The reduction in runtime and relative distance achieved by the `oic` setting is 7% and 2%, respectively, for affected instances and 23% and 14%, respectively, for affected-hard instances.

We note that the runtime does not provide much information about affected-hard instances, since only 10 instances can be solved within 3600 seconds. For these instances, the gap reduction is more useful to measure the reduction of the search space by the proposed valid inequalities. However, for all affected instances, the runtime is still important because it measures the speedup due to the valid inequalities.

Second, we find that all cut settings have a positive effect on SCIP performance, although the magnitude of the reduction varies. When we compare the `oc` and `ic` settings, we find that the `oc` setting leads to a larger reduction in runtime. This difference in runtime is due to the fact that computing intersection cuts requires extracting a simplified cone from the LP relaxation and applying bisection search along each ray of the cone. These procedures require more computational resources compared to the construction of outer approximation cuts.

On the other hand, the `ic` setting shows better performance in terms of reducing gaps. Intersection cuts approximate the intersection of a signomial term set with the simplicial cone, while outer approximation cuts approximate the intersection of a signomial term set with a hypercube. Around the relaxation point, the simplicial cone usually provides a better approximation than the hypercube. Therefore, `ic` achieves a greater reduction in the relative gap. However, the better simplicial conic approximation does yield a significant improvement compared to the hypercubic approximation.

Finally, the `oic` setting combines both the `oc` and `ic` settings and achieves the best reduction in runtime. However, for affected and affected-hard instances, the setting shows different gap reduction results. In fact, the results for affected-hard instances give more insight, since the goal of the valid inequalities is to speed up convergence for hard instances. In this sense, the `oic` setting achieves almost the best result, so it carries the best of both valid inequalities. However, the improvement compared to each setting is not significant.

In summary, the performances of the `oc` and `ic` settings are comparable. They can lead to smaller duality gaps with less computation time, which is desirable for solvers, and one can use either of them. Moreover, they do not hurt each other.

7 Conclusion

In this paper we study valid inequalities for SP problems and propose two types of valid linear inequalities: intersection cuts and outer approximation cuts. Both are derived from normalized DCC formulations of signomial term sets. First, we study general conditions for maximal \mathcal{S} -free sets. We construct maximal signomial term-free sets from which we generate intersection cuts. Second, we construct convex outer approximations of signomial term sets within hypercubes. We provide extended formulations for the convex envelopes of concave functions in the normalized DCC formulations. Then we separate valid inequalities for the convex outer approximations by projection. Moreover, when $h = 2$, we use supermodularity to derive a closed-form expression

for the convex envelopes.

We present a comparative analysis of the computational results obtained with the MINLPLib instances. This analysis demonstrates the effectiveness of the proposed valid inequalities. The results show that intersection cuts and outer approximation cuts have similar performance and their combination takes the best of each setting. In particular, it is easy to implement outer approximation cuts in general purpose solvers. In the future, we plan to carefully optimize outer approximation cuts and develop them as an easy-to-use plugin.

We currently deal with signomial terms explicitly present in the signomial terms, but our results can be extended to deal with multiple signomial terms. In the future, the proposed valid inequalities can approximate nonlinear aggregations of constraints that define the signomial lift. Specifically, given signomial constraints $\{\psi_{\alpha^i}(x) = y_i\}_{i \in [r]}$ with any exponent vector $\zeta \in \mathbb{R}^r$, we can employ *signomial aggregation* to generate a new signomial constraint: $\psi_{(\sum_{i \in [r]} \zeta_i \alpha^i)}(x) = \psi_{\zeta}(y)$. This constraint is valid for the signomial lift and encodes more variables and terms. Next, we can apply the DCC reformulation to the constraints $\psi_{(\sum_{i \in [r]} \zeta_i \alpha^i)}(x) \leq \psi_{\zeta}(y)$ and $\psi_{(\sum_{i \in [r]} \zeta_i \alpha^i)}(x) \geq \psi_{\zeta}(y)$. Finally, we can separate the proposed valid inequalities. As far as we know, the signomial aggregation operator is not yet used for polynomial programming, since it outputs a signomial constraint.

Acknowledgments

We thank Felipe Serrano for sharing his Ph.D. thesis with us [60]. We appreciate the discussion with Antonia Chmiela, Mathieu Besançon, Stefan Vigerske, and Ksenia Bestuzheva on SCIP. This publication was supported by the “Integrated Urban Mobility” chair, which is supported by L’X - École Polytechnique and La Fondation de l’École Polytechnique. Under no circumstances do the partners of the Chair assume any liability for the content of this publication, for which the author alone is responsible.

Declarations

No conflicts of interest with the journal or funders.

Appendix

Proof of Thm. 1. It suffices to consider the case that \mathcal{C} is a maximal $\text{epi}(g'_i)$ -free set in \mathcal{G}^i . W.l.o.g., we can assume that $\mathcal{C}, \mathcal{G}^i$ are full-dimensional in \mathbb{R}^{I_i} . Since $\text{epi}(g'_i)$ includes $\text{graph}(g'_i)$, \mathcal{C} , as an $\text{epi}(g'_i)$ -free set, is also $\text{graph}(g'_i)$ -free. First, we prove that \mathcal{C} is a maximal $\text{graph}(g'_i)$ -free set in \mathcal{G}^i . Assume, to aim at a contradiction, that \mathcal{C}' is a $\text{graph}(g'_i)$ -free set that $\mathcal{C} \cap \mathcal{G}^i \subsetneq \mathcal{C}' \cap \mathcal{G}^i$. Suppose that $\text{epi}(g'_i) \cap \text{int}(\mathcal{C}' \cap \mathcal{G}^i)$ is not empty and contains (x'_{J_i}, y'_i) . As \mathcal{C} is $\text{epi}(g'_i)$ -free, there exists a point $(x_{J_i}, y_i) \in \text{int}(\mathcal{C} \cap \mathcal{G}^i) \subseteq \text{int}(\mathcal{C}' \cap \mathcal{G}^i)$ such that $(x_{J_i}, y_i) \in \text{hypo}(g'_i)$. It follows from the continuity of g'_i that there exists a point $(x^*_{J_i}, y^*_i) \in \text{graph}(g'_i)$ in the line segment joining (x_{J_i}, y_i) and (x'_{J_i}, y'_i) . As $\text{int}(\mathcal{C}' \cap \mathcal{G}^i)$ is convex, we have that $(x^*_{J_i}, y^*_i) \in \text{int}(\mathcal{C}' \cap \mathcal{G}^i)$, which leads to a contradiction to $\text{graph}(g'_i)$ -freeness of \mathcal{C}' . Therefore, $\text{epi}(g'_i) \cap \text{int}(\mathcal{C}' \cap \mathcal{G}^i)$ must be empty, so $\mathcal{C}' \cap \mathcal{G}^i \subseteq \text{hypo}(g'_i)$. This means that \mathcal{C}' is also $\text{epi}(g'_i)$ -free. However, note that $\mathcal{C} \cap \mathcal{G}^i \subsetneq \mathcal{C}' \cap \mathcal{G}^i$, this contradicts with the fact that \mathcal{C} is a maximal $\text{epi}(g'_i)$ -free set in \mathcal{G}^i . Therefore, \mathcal{C} is a maximal $\text{graph}(g'_i)$ -free set in \mathcal{G}^i . Secondly, we prove that $\bar{\mathcal{C}}$ is a maximal $\mathcal{S}_{\text{lift}}$ -free set in \mathcal{G} . Assume, to aim at a contradiction, that there exists an $\mathcal{S}_{\text{lift}}$ -free set $\bar{\mathcal{D}}$ in \mathcal{G} such that $\bar{\mathcal{C}} \cap \mathcal{G} \subsetneq \bar{\mathcal{D}} \cap \mathcal{G}$. We look at their orthogonal projections on \mathbb{R}^{I_i} . It follows from the decomposability

that $\mathcal{C} \cap \mathcal{G}^i = \mathcal{C} \cap \text{proj}_{\mathbb{R}^{I_i}}(\mathcal{G}) = \text{proj}_{\mathbb{R}^{I_i}}(\bar{\mathcal{C}} \cap \mathcal{G}) \subseteq \text{proj}_{\mathbb{R}^{I_i}}(\bar{\mathcal{D}} \cap \mathcal{G})$. Denote $\mathcal{D} := \text{cl}(\text{proj}_{\mathbb{R}^{I_i}}(\bar{\mathcal{D}} \cap \mathcal{G}))$, which is a closed convex set in \mathcal{G}^i . Since $\bar{\mathcal{C}} = \mathcal{C} \times \mathbb{R}^{I_i^c}$, \mathcal{D} must strictly include $\mathcal{C} \cap \mathcal{G}^i$. Note that \mathcal{D} is $\text{graph}(g'_i)$ -free. Since \mathcal{C} is a maximal $\text{graph}(g'_i)$ -free set in \mathcal{G}^i , this implies that $\mathcal{C} \cap \mathcal{G}^i = \mathcal{D}$, which leads to a contradiction. \square

Proof of Thm. 3. Let \mathcal{C} be a set satisfying the hypothesis. Suppose, to aim at a contradiction, that there exists a closed convex set \mathcal{C}^* such that $\mathcal{C} \subsetneq \mathcal{C}^*$ and \mathcal{C}^* is an inner approximation of \mathcal{F} . Then, there must exist an open ball B such that $B \subseteq \mathcal{F} \setminus \mathcal{C}$ and $B \subseteq \mathcal{C}^*$. Let z^* be the center of B , so $z^* \in \text{int}(\mathcal{F} \setminus \mathcal{C})$. W.l.o.g., we let $\mathcal{C}^* = \text{conv}(\mathcal{C} \cup \{z^*\})$, which is a closed convex inner approximation of \mathcal{F} . Since $z^* \notin \mathcal{C}$, by the hyperplane separation theorem, there exists $\lambda \in \text{dom}(\sigma_{\mathcal{C}})$ such that

$$\lambda \cdot z^* > \sigma_{\mathcal{C}}(\lambda). \quad (16)$$

For any such λ , by the hypothesis, there exists a point $z' \in \text{bd}(\mathcal{F}) \cap \text{bd}(\mathcal{C})$ such that

$$\lambda \cdot z' = \sigma_{\mathcal{C}}(\lambda), \quad (17)$$

and z' is an exposing point of \mathcal{C} . We want to show that, for any $\lambda' \in \text{dom}(\sigma_{\mathcal{C}^*})$, $\lambda' \cdot z' < \sigma_{\mathcal{C}^*}(\lambda')$. We consider the following three cases. First, we consider the case $\lambda' = \lambda$. Because $z^* \in \mathcal{C}^*$, by the definition of support functions, we have that

$$\lambda \cdot z^* \leq \sup_{z \in \mathcal{C}^*} \lambda \cdot z = \sigma_{\mathcal{C}^*}(\lambda). \quad (18)$$

It follows from (16), (17), and (18) that

$$\lambda \cdot z' = \sigma_{\mathcal{C}}(\lambda) < \lambda \cdot z^* \leq \sigma_{\mathcal{C}^*}(\lambda) = \sigma_{\mathcal{C}^*}(\lambda'). \quad (19)$$

Second, we consider the case $\lambda' = \rho\lambda$ for some $\rho > 0$. Since $\sigma_{\mathcal{C}^*}$ is positively homogeneous of degree 1, it follows from (19) that $\lambda' \cdot z' = \rho\lambda \cdot z' < \rho\sigma_{\mathcal{C}^*}(\lambda) = \sigma_{\mathcal{C}^*}(\lambda')$. Last, we consider the case $\lambda' \in \text{dom}(\sigma_{\mathcal{C}^*}) \setminus \{\rho\lambda\}_{\rho>0}$. By Lemma 2, $\sigma_{\mathcal{C}} \leq \sigma_{\mathcal{C}^*}$. By the hypothesis that z' is an exposing point of \mathcal{C} , provided that $\lambda' \neq \rho\lambda$, we have that $\lambda' \cdot z' < \sigma_{\mathcal{C}}(\lambda') \leq \sigma_{\mathcal{C}^*}(\lambda')$. In summary, we have proved that for any $\lambda' \in \text{dom}(\sigma_{\mathcal{C}^*})$, $\lambda' \cdot z' < \sigma_{\mathcal{C}^*}(\lambda')$. So by Lemma 2, $z' \in \text{int}(\mathcal{C}^*)$. We find that $z' \in \text{bd}(\mathcal{F}) \cap \text{int}(\mathcal{C}^*)$. This finding means a point near z' exists, which is in \mathcal{C}^* , but not in \mathcal{F} . Hence, \mathcal{C}^* is not an inner approximation of \mathcal{F} , which leads to a contradiction. \square

Proof of Lemma 3. Given $z \in \text{dom}(f)$, $f(\rho z) = \rho^d f(z)$ is a real number for any $\rho \in \mathbb{R}_{++}$, so $\text{int}(\text{dom}(f))$ is a cone. Suppose that f is positive homogeneous of degree 1. For any $z \in \text{dom}(f)$, $\Xi_{\check{z}}^f(z) = f(\check{z}) + \nabla f(\check{z}) \cdot (z - \check{z}) = \nabla f(\check{z}) \cdot z$, where the second equation follows from Euler's homogeneous function theorem: $f(\check{z}) = \nabla f(\check{z}) \cdot \check{z}$. For any $z = \rho\check{z}$ with $\rho \in \mathbb{R}_{++}$, $\Xi_{\check{z}}^f(z) = \nabla f(\check{z}) \cdot \rho\check{z} = \rho \nabla f(\check{z}) \cdot \check{z} = \rho f(\check{z}) = f(\rho\check{z})$, where the first and second equations follow from the previous result, the third follows from that $\Xi_{\check{z}}^f$ has the same value as f at \check{z} , and the last equation follows from the homogeneity. \square

Proof of Lemma 6. We note that any hyperplane in \mathbb{R}^{h+1} is uniquely determined by $h+1$ affinely independent points. By definition, an affine underestimator $a \cdot w + b$ of f is a facet if and only if $a \cdot w + b \leq t$ is a facet of the epigraph $\text{epi}_Q(f)$. Then, the affine underestimator is a facet, if and only if, there exists $h+1$ affinely independent points $\{(w^i, t^i)\}_{i \in [h+1]} \subseteq \text{epi}_Q(f) \subseteq \mathbb{R}^{h+1}$ such that for all $i \in [h+1]$, $a \cdot w^i + b = t^i$. Moreover, t^i must equal $f(w^i)$, otherwise $a \cdot w^i + b = t^i < f(w^i)$, which implies that $a \cdot w + b \leq t$ does not underestimate f at w^i . Therefore, w^i is the supporting point of $a \cdot w + b$. Note that $\{(w^i, t^i)\}$ are affinely independent, if and only if, $\{w^i\}_{i \in [h+1]}$ are affinely independent. \square

References

- [1] F. A. AL-KHAYYAL AND J. E. FALK, *Jointly constrained biconvex programming*, Mathematics of Operations Research, 8 (1983), pp. 273–286.
- [2] K. ANDERSEN, Q. LOUVEAUX, AND R. WEISMANTEL, *An analysis of mixed integer linear sets based on lattice point free convex sets*, Mathematics of Operations Research, 35 (2010), pp. 233–256.
- [3] K. ANDERSEN, Q. LOUVEAUX, R. WEISMANTEL, AND L. A. WOLSEY, *Inequalities from two rows of a simplex tableau*, in Integer Programming and Combinatorial Optimization, M. Fischetti and D. P. Williamson, eds., Berlin, Heidelberg, 2007, Springer Berlin Heidelberg, pp. 1–15.
- [4] M. APS, *Mosek modeling cookbook*, 2018.
- [5] E. BALAS, *Intersection cuts—a new type of cutting planes for integer programming*, Operations Research, 19 (1971), pp. 19–39.
- [6] X. BAO, A. KHAJAVIRAD, N. V. SAHINIDIS, AND M. TAWARMALANI, *Global optimization of nonconvex problems with multilinear intermediates*, Mathematical Programming Computation, 7 (2015), pp. 1–37.
- [7] A. BASU, S. S. DEY, AND J. PAAT, *Nonunique lifting of integer variables in minimal inequalities*, SIAM Journal on Discrete Mathematics, 33 (2019), pp. 755–783.
- [8] P. BELOTTI, J. C. GÓEZ, I. PÓLIK, T. K. RALPHS, AND T. TERLAKY, *A conic representation of the convex hull of disjunctive sets and conic cuts for integer second order cone optimization*, in Numerical Analysis and Optimization, Springer, 2015, pp. 1–35.
- [9] P. BELOTTI, C. KIRCHES, S. LEYFFER, J. LINDEROTH, J. LUEDTKE, AND A. MAHAJAN, *Mixed-integer nonlinear optimization*, Acta Numerica, 22 (2013), pp. 1–131.
- [10] P. BELOTTI, J. LEE, L. LIBERTI, F. MARGOT, AND A. WÄCHTER, *Branching and bounds tightening techniques for non-convex minlp*, Optimization Methods & Software, 24 (2009), pp. 597–634.
- [11] K. BESTUZHEVA, A. CHMIELA, B. MÜLLER, F. SERRANO, S. VIGERSKE, AND F. WEGSCHEIDER, *Global optimization of mixed-integer nonlinear programs with scip 8*, arXiv preprint arXiv:2301.00587, (2023).
- [12] D. BIENSTOCK, C. CHEN, AND G. MUÑOZ, *Intersection cuts for polynomial optimization*, in Integer Programming and Combinatorial Optimization, A. Lodi and V. Nagarajan, eds., Cham, 2019, Springer International Publishing, pp. 72–87.
- [13] M. R. BUSSIECK, A. S. DRUD, AND A. MEERAUS, *Minplib—a collection of test models for mixed-integer nonlinear programming*, INFORMS Journal on Computing, 15 (2003), pp. 114–119.
- [14] S. CAFIERI, J. LEE, AND L. LIBERTI, *On convex relaxations of quadrilinear terms*, Journal of Global Optimization, 47 (2010), pp. 661–685.
- [15] V. CHANDRASEKARAN AND P. SHAH, *Relative entropy relaxations for signomial optimization*, SIAM Journal on Optimization, 26 (2016), pp. 1147–1173.

- [16] R. CHARES, *Cones and interior-point algorithms for structured convex optimization involving powers and exponentials*, PhD thesis, UCL-Université Catholique de Louvain Louvain-la-Neuve, Belgium, 2009.
- [17] J.-S. CHEN AND C.-H. HUANG, *A note on convexity of two signomial functions*, *Journal of Nonlinear and Convex Analysis*, 10 (2009), pp. 429–435.
- [18] M. CONFORTI, G. CORNUÉJOLS, AND G. ZAMBELLI, *Integer programming*, Springer International Publishing, Cham, 2014.
- [19] M. CONFORTI, G. CORNUÉJOLS, AND G. ZAMBELLI, *Corner polyhedron and intersection cuts*, *Surveys in Operations Research and Management Science*, 16 (2011), pp. 105–120.
- [20] G. CORNUÉJOLS, L. WOLSEY, AND S. YILDIZ, *Sufficiency of cut-generating functions*, *Mathematical Programming*, 152 (2015), pp. 643–651.
- [21] A. COSTA AND L. LIBERTI, *Relaxations of multilinear convex envelopes: dual is better than primal*, in *International Symposium on Experimental Algorithms*, Springer, 2012, pp. 87–98.
- [22] A. DEL PIA AND R. WEISMANTEL, *Relaxations of mixed integer sets from lattice-free polyhedra*, *4OR*, 10 (2012), pp. 221–244.
- [23] S. S. DEY AND L. A. WOLSEY, *Lifting integer variables in minimal inequalities corresponding to lattice-free triangles*, in *Integer Programming and Combinatorial Optimization*, A. Lodi, A. Panconesi, and G. Rinaldi, eds., Berlin, Heidelberg, 2008, Springer Berlin Heidelberg, pp. 463–475.
- [24] M. DRESSLER AND R. MURRAY, *Algebraic perspectives on signomial optimization*, *SIAM Journal on Applied Algebra and Geometry*, 6 (2022), pp. 650–684.
- [25] R. J. DUFFIN, *Linearizing geometric programs*, *SIAM review*, 12 (1970), pp. 211–227.
- [26] J. E. FALK AND K. R. HOFFMAN, *A successive underestimation method for concave minimization problems*, *Mathematics of Operations Research*, 1 (1976), pp. 251–259.
- [27] M. FISCHETTI, I. LJUBIĆ, M. MONACI, AND M. SINNL, *On the use of intersection cuts for bilevel optimization*, *Mathematical Programming*, 172 (2018), pp. 77–103.
- [28] M. FISCHETTI AND M. MONACI, *A branch-and-cut algorithm for mixed-integer bilinear programming*, *European Journal of Operational Research*, 282 (2020), pp. 506–514.
- [29] F. GLOVER, *Convexity cuts and cut search*, *Operations Research*, 21 (1973), pp. 123–134.
- [30] R. E. GOMORY, *Some polyhedra related to combinatorial problems*, *Linear algebra and its applications*, 2 (1969), pp. 451–558.
- [31] T. HE AND M. TAWARMALANI, *A new framework to relax composite functions in nonlinear programs*, *Mathematical Programming*, 190 (2021), pp. 427–466.
- [32] T. HE AND M. TAWARMALANI, *Tractable relaxations of composite functions*, *Mathematics of Operations Research*, 47 (2022), pp. 1110–1140.
- [33] J.-B. HIRIART-URRUTY AND C. LEMARÉCHAL, *Fundamentals of convex analysis*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2001.
- [34] F. KILINÇ-KARZAN, *On minimal valid inequalities for mixed integer conic programs*, *Mathematics of Operations Research*, 41 (2016), pp. 477–510.

- [35] L. LIBERTI, *Writing global optimization software*, in Liberti and Maculan [36], pp. 211–262.
- [36] L. LIBERTI AND N. MACULAN, eds., *Global Optimization: from Theory to Implementation*, Springer, Berlin, 2006.
- [37] L. LIBERTI AND C. C. PANTELIDES, *Convex envelopes of monomials of odd degree*, Journal of Global Optimization, 25 (2003), pp. 157–168.
- [38] M.-H. LIN AND J.-F. TSAI, *Range reduction techniques for improving computational efficiency in global optimization of signomial geometric programming problems*, European Journal of Operational Research, 216 (2012), pp. 17–25.
- [39] M. LOCATELLI, *Polyhedral subdivisions and functional forms for the convex envelopes of bilinear, fractional and other bivariate functions over general polytopes*, Journal of Global Optimization, 66 (2016), pp. 629–668.
- [40] M. LOCATELLI, *Convex envelopes of bivariate functions through the solution of KKT systems*, Journal of Global Optimization, 72 (2018), pp. 277–303.
- [41] M. LOCATELLI AND F. SCHOEN, *On convex envelopes for bivariate functions over polytopes*, Mathematical Programming, 144 (2014), pp. 65–91.
- [42] J. LUEDTKE, M. NAMAZIFAR, AND J. LINDEROTH, *Some results on the strength of relaxations of multilinear functions*, Mathematical programming, 136 (2012), pp. 325–351.
- [43] A. LUNDELL, A. SKJÄL, AND T. WESTERLUND, *A reformulation framework for global optimization*, Journal of Global Optimization, 57 (2013), pp. 115–141.
- [44] A. LUNDELL AND T. WESTERLUND, *Solving global optimization problems using reformulations and signomial transformations*, Computers & Chemical Engineering, 116 (2018), pp. 122–134.
- [45] C. D. MARANAS AND C. A. FLOUDAS, *Finding all solutions of nonlinearly constrained systems of equations*, Journal of Global Optimization, 7 (1995), pp. 143–182.
- [46] C. D. MARANAS AND C. A. FLOUDAS, *Global optimization in generalized geometric programming*, Computers & Chemical Engineering, 21 (1997), pp. 351–369.
- [47] G. P. MCCORMICK, *Computability of global solutions to factorable nonconvex programs: Part I—Convex underestimating problems*, Mathematical programming, 10 (1976), pp. 147–175.
- [48] C. A. MEYER AND C. A. FLOUDAS, *Trilinear monomials with mixed sign domains: Facets of the convex and concave envelopes*, Journal of Global Optimization, 29 (2004), pp. 125–155.
- [49] C. A. MEYER AND C. A. FLOUDAS, *Trilinear monomials with positive or negative domains: Facets of the convex and concave envelopes*, in *Frontiers in Global Optimization*, C. A. Floudas and P. Pardalos, eds., Boston, MA, 2004, Springer US, pp. 327–352.
- [50] R. MISENER AND C. A. FLOUDAS, *ANTIGONE: Algorithms for coNTinuous / Integer Global Optimization of Nonlinear Equations*, Journal of Global Optimization, 59 (2014), pp. 503–526.
- [51] R. MISENER AND C. A. FLOUDAS, *A framework for globally optimizing mixed-integer signomial programs*, Journal of Optimization Theory and Applications, 161 (2014), pp. 905–932.

- [52] S. MODARESI, M. R. KILINÇ, AND J. P. VIELMA, *Intersection cuts for nonlinear integer programming: convexification techniques for structured sets*, Mathematical Programming, 155 (2016), pp. 575–611, <https://arxiv.org/abs/1302.2556>.
- [53] G. MUÑOZ AND F. SERRANO, *Maximal quadratic-free sets*, Mathematical Programming, 192 (2022), pp. 229–270.
- [54] R. MURRAY, V. CHANDRASEKARAN, AND A. WIERMAN, *Signomial and polynomial optimization via relative entropy and partial dualization*, Mathematical Programming Computation, 13 (2021), pp. 257–295.
- [55] T. T. NGUYEN, J.-P. P. RICHARD, AND M. TAWARMALANI, *Deriving convex hulls through lifting and projection*, Mathematical Programming, 169 (2018), pp. 377–415.
- [56] W. H. PRESS, S. A. TEUKOLSKY, W. T. VETTERLING, AND B. FLANNERY, *Chapter 9: Root finding and nonlinear sets of equations*, Book: Numerical Recipes: The Art of Scientific Computing, New York, Cambridge University Press, 10 (2007).
- [57] J.-P. P. RICHARD AND S. S. DEY, *The group-theoretic approach in mixed integer programming*, in 50 Years of Integer Programming 1958-2008, Springer, 2010, pp. 727–801.
- [58] A. D. RIKUN, *A convex envelope formula for multilinear functions*, Journal of Global Optimization, 10 (1997), pp. 425–437.
- [59] F. SERRANO, *Intersection cuts for factorable MINLP*, in Integer Programming and Combinatorial Optimization, A. Lodi and V. Nagarajan, eds., Cham, 2019, Springer International Publishing, pp. 385–398.
- [60] F. SERRANO MUSALEM, *On cutting planes for mixed-integer nonlinear programming*, doctoral thesis, Technische Universität Berlin, Berlin, 2021.
- [61] P. SHEN, *Linearization method of global optimization for generalized geometric programming*, Applied Mathematics and Computation, 162 (2005), pp. 353–370.
- [62] H. D. SHERALI, *Convex envelopes of multilinear functions over a unit hypercube and over special discrete sets*, Acta mathematica vietnamica, 22 (1997), pp. 245–270.
- [63] E. SPEAKMAN AND J. LEE, *Quantifying double mccormick*, Mathematics of Operations Research, 42 (2017), pp. 1230–1253.
- [64] M. TAWARMALANI, J.-P. P. RICHARD, AND C. XIONG, *Explicit convex and concave envelopes through polyhedral subdivisions*, Mathematical Programming, 138 (2013), pp. 531–577.
- [65] M. TAWARMALANI AND N. V. SAHINIDIS, *A polyhedral branch-and-cut approach to global optimization*, Mathematical programming, 103 (2005), pp. 225–249.
- [66] D. M. TOPKIS, *Supermodularity and complementarity*, Princeton university press, Princeton, 2011.
- [67] H. TUY, *Concave minimization under linear constraints with special structure*, Optimization, 16 (1985), pp. 335–352.
- [68] G. XU, *Global optimization of signomial geometric programming problems*, European journal of operational research, 233 (2014), pp. 500–510.