



HAL
open science

Bola-Amphiphilic Glycodendrimers: New Carbohydrate Mimicking Scaffolds to Target Carbohydrate-Binding Proteins

Wenzheng Zhang, Dinesh Dhumal, Xiaolei Zhu, Brigino Ralahy, Aleksandra Ellert-Miklaszewska, Jing Wu, Erik Laurini, Yi-Wen W Yao, Chai-Lin L Kao, Juan L Iovanna, et al.

► **To cite this version:**

Wenzheng Zhang, Dinesh Dhumal, Xiaolei Zhu, Brigino Ralahy, Aleksandra Ellert-Miklaszewska, et al.. Bola-Amphiphilic Glycodendrimers: New Carbohydrate Mimicking Scaffolds to Target Carbohydrate-Binding Proteins. Chemistry - A European Journal, In press, 10.1002/chem.202201400 . hal-03725167

HAL Id: hal-03725167

<https://hal.science/hal-03725167v1>

Submitted on 15 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chemistry A European Journal

 **Chemistry
Europe**
European Chemical
Societies Publishing

Accepted Article

Title: Bola-Amphiphilic Glycodendrimers: New
Carbohydrate Mimicking Scaffolds to Target Carbohydrate-
Binding Proteins

Authors: Wenzheng Zhang, Dinesh Dhumal, Xiaolei Zhu, Brigino
Ralahy, Aleksandra Ellert-Miklaszewska, Jing Wu, Erik
Laurini, Yi-Wen Yao, Chai-Lin Kao, Juan L. Iovanna, Sabrina
Priel, Bozena Kaminska, Yi Xia, and Ling Peng

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Chem. Eur. J.* **2022**, e202201400

Link to VoR: <https://doi.org/10.1002/chem.202201400>

WILEY-VCH

RESEARCH ARTICLE

Bola-Amphiphilic Glycodendrimers: New Carbohydrate Mimicking Scaffolds to Target Carbohydrate-Binding Proteins

Wenzheng Zhang,^[a, b] Dinesh Dhumal,^[b] Xiaolei Zhu,^[b] Brigino Ralahy,^[b] Aleksandra Ellert-Miklaszewska,^[c] Jing Wu,^[b] Erik Laurini,^[d] Yi-Wen Yao,^[b, e] Chai-Lin Kao,^[e] Juan L. Iovanna,^[f] Sabrina Pricl,^[d, g] Bozena Kaminska,^[c] Yi Xia*^[a] and Ling Peng*^[b]

- [a] W. Zhang, Prof. Dr. Y. Xia
Chongqing Key Laboratory of Natural Product Synthesis and Drug Research, Innovative Drug Research Center, School of Pharmaceutical Sciences
Chongqing University
Chongqing 401331, China
E-mail: yixia@cqu.edu.cn; <http://sps.cqu.edu.cn/info/1051/1040.htm>
- [b] W. Zhang, Dr. D. Dhumal, Dr. X. Zhu, B. Ralahy, J. Wu, Y. W. Yao, Prof. Dr. L. Peng
Centre Interdisciplinaire de Nanoscience de Marseille (CINaM), UMR 7325, Equipe Labellisé par La Ligue
Aix-Marseille Université, CNRS
Marseille 13288, France
E-mail: ling.peng@univ-amu.fr; <http://www.cinam.univ-mrs.fr/cinam/en/team/immf/equipe-peng>
- [c] Dr. A. E. Miklaszewska, Prof. Dr. B. Kaminska
Laboratory of Molecular Neurobiology, Neurobiology Center
Nencki Institute of Experimental Biology of the Polish Academy of Sciences
Warsaw 02-093, Poland
- [d] Prof. Dr. E. Laurini, Prof. Dr. S. Pricl
Molecular Biology and Nanotechnology (MolBNL@UniTS) Laboratory, DEA
University of Trieste
Piazzale Europa 1, 34127 Trieste, Italy
- [e] Y. W. Yao, Prof. Dr. C. L. Kao
Department of Medicinal and Applied Chemistry
Kaohsiung Medical University
Kaohsiung, Taiwan
- [f] Prof. Dr. J. L. Iovanna
Centre de Recherche en Cancérologie de Marseille (CRCM), INSERM U1068, CNRS UMR 7258
Aix-Marseille Université and Institut Paoli-Calmettes
Marseille 13288, France
- [g] Prof. Dr. S. Pricl
Department of General Biophysics, Faculty of Biology and Environmental Protection
University of Lodz
Lodz, Poland

Supporting information for this article is given via a link at the end of the document.

Abstract: Dendrimers are appealing scaffolds for creating carbohydrate mimics with unique multivalent cooperativity. We report here novel bola-amphiphilic glycodendrimers bearing mannose and glucose terminals, and a hydrophobic thioacetal core responsive to reactive oxygen species. The peculiar bola-amphiphilic feature enabled stronger binding to lectin compared to the conventional amphiphile. In addition, these dendrimers are able to target mannose receptors and glucose transporters expressed at the surface of cells, allowing effective and specific cellular uptake. This highlights their great promise for targeted delivery.

Introduction

Carbohydrates are the most abundant biomacromolecules in living organisms and play important roles in energy production, as structural materials and as signaling moieties.^[1-5] Carbohydrates mediate cell-cell and cell-matrix interactions via carbohydrate-carbohydrate and carbohydrate-protein binding.^[1-5] Although single carbohydrate units often show low binding affinity, the glycoside cluster effect, which exploits the cooperative interactions issued from the branched and multivalent structure of

carbohydrates, improves strength and specificity of binding.^[6-9] This particular effect is essential in cell-cell interactions and in regulating various biological events, such as cell differentiation, proliferation and adhesion, inflammation, cancer metastasis and immune responses.^[1-5]

Dendrimers, by virtue of their unique ramified structure and multivalent cooperativity, are particularly appealing scaffolds for constructing carbohydrate mimics.^[10-14] Such dendrimers, also referred to as glycodendrimers, can structurally and functionally mimic natural polysaccharides, glycoproteins and mucins, and hence constitute a useful means for studying the glycoside cluster effect and for targeting specific carbohydrate-binding proteins. Recently, Percec et al developed Janus amphiphilic glycodendrimers which can self-assemble into dendrimersomes, mimicking the recognition structures of glycans and glycoproteins on the cell.^[15-17] In addition, amphiphilic dendrimers bearing carbohydrate modifications have been developed for drug delivery.^[18, 19] For example, a photo-responsive polypeptide-glycosylated dendron amphiphile was documented to dynamically bind to concanavalin A (ConA) for ovalbumin (OVA) delivery.^[18] Also, a lactose-bearing amphiphilic poly(amidoamine) (PAMAM) dendrimer was reported to effectively interact with RCA₁₂₀ lectin and deliver the anticancer drug doxorubicin.^[19] Compared with

RESEARCH ARTICLE

many reported conventional glycodendrimer systems, amphiphilic glycodendrimers are able to self-assemble and form dynamic systems, offering modular and adaptive interactions with carbohydrate-binding proteins. Amphiphilic glycodendrimers therefore hold great potential in targeted drug delivery.

In this study, we report the synthesis and the use of novel bola-amphiphilic glycodendrimers (Figure 1) for binding and targeting specific carbohydrate-binding proteins. Different from traditional amphiphilic molecules, the dumbbell-like shape of bola-amphiphiles offers a unique bipolar amphiphilic structure,^[20, 21] which is expected to reinforce molecular recognition, binding and stability towards carbohydrate-binding proteins. This approach should allow the exploitation of the glycoside cluster effect and the implementation of multivalent glycoconjugates for targeted delivery. In addition, the core of the bola-amphiphilic glycodendrimers harbors a thioacetal functionality, which is responsive to reactive oxygen species (ROS), thus allowing redox-responsive delivery.^[22]




Figure 1. Bola-amphiphilic glycodendrimers bearing mannose (**1a**) and glucose (**1b**) terminals.

Specifically, in this work we sought to introduce mannose and glucose entities to the bola-amphiphilic dendrimer terminals (Figure 1) and study their capacity to interact with carbohydrate-binding proteins such as lectin ConA, the mannose receptor (MR) and the glucose transporter (GLUT). ConA is a tetramer lectin with a strong affinity for mannose residues, forming a cross-linked network or aggregations with mannosyl carbohydrates via multivalent and cooperative interactions.^[23] The MR can effectively bind mannose ligand and is mainly expressed on the surface of macrophages (including microglia - brain-resident myeloid cells), dendritic cells, endothelial cells and some cancer cells^[24, 25], while the GLUT is widely expressed in the blood-brain barrier (BBB), astrocytes and cancer cells.^[26, 27] The GLUT is mainly responsible for the "two-way" transport of glucose that maintains a stable concentration of this carbohydrate in cells, and plays an important role in normal physiological balance.^[26] As such, the GLUT is often found overexpressed in cancer cells owing to their requirement of large quantities of glucose for rapid proliferation.^[27] Based on these considerations, we designed the bola-amphiphilic glycodendrimers **1a** and **1b**, both carbohydrate mimics bearing respectively mannose and glucose terminals, studied their binding with ConA and their capacity to target specific cells overexpressing MRs and GLUTs.

Results and Discussion

Dendrimer synthesis

For synthesizing the bola-amphiphilic dendrimers **1a** and **1b** we envisaged two strategies (Figure 2), mainly exploiting "click" reactions to generate the bola-amphiphilic dendrimer scaffold and to introduce the carbohydrate terminal units. Specifically, the ester-bearing bola-amphiphilic dendrimer would be first constructed via click reaction by coupling the azido-bearing hydrophobic chain with the alkyne-containing hydrophilic PAMAM dendron; the obtained dendrimer would then be transformed to bear the azido or alkynyl terminals which would, via a second click reaction with the corresponding alkyne- and azide-bearing carbohydrate precursors, to deliver the bola-amphiphilic glycodendrimers.




Figure 2. Strategies proposed for synthesizing the bola-amphiphilic glycodendrimers.

Accordingly, we first prepared the ester terminating dendrimer **3** using Cu(I)-catalyzed 1,3-dipolar cycloaddition to couple the hydrophobic chain **1** bearing two azido terminals with the alkyne-containing PAMAM dendron **2** as reported (Figure 3).^[22] We then attempted strategy A, which started well with the successful conversion of **3** to the amine-terminating dendrimer **4** upon treatment with ethylene diamine. However, the reaction to transform the terminal amino groups in **4** to azido functions was unsuccessful despite attempting various different conditions when using imidazole-1-sulfonyl azide as the amine-azide transfer reagent.^[28] This reaction gave many side products, likely due to incomplete multi-site reactions. In addition, purification of the azido-terminating dendrimer **5** was very difficult.

We then turned to strategy B to synthesize alkynyl-terminating dendrimer **7** (Figure 3). We first hydrolyzed the methyl ester terminated dendrimer **3** using LiOH to yield the carboxylic acid terminated dendrimer **6** within a short time (3 hours). Then dendrimer **6** was conjugated with propargylamine using the coupling reagents EDCI and HOBt in DMF to generate the alkynyl-terminating dendrimer **7**. Remarkably, this conjugation

RESEARCH ARTICLE




Figure 3. Synthesis of the intermediate dendrimers **5** and **7** required for the preparation of the glycodendrimers

went very well, and thin layer chromatography (TLC) showed only one main product formed even though there were eight reaction sites in **6**. Dendrimer **7** was easily purified using column chromatography and was obtained at a high yield of 81%.




Figure 4. Synthesis of the bola-amphiphilic glycodendrimers **1a** and **1b**.

Starting with **7**, the click reactions with the azido-conjugated mannose and glucose precursors **8a-b**, respectively, were extremely successful (Figure 4) despite the existence of multi-site reactions and steric hindrance among the carbohydrate terminals. The desired dendrimers **9a-b** were obtained with yields up to 80% after column chromatography.

NMR analysis of the transformation from dendrimer **7** to dendrimer **9a** is presented in Figure 5. The disappearance of the




Figure 5. ^1H NMR spectral analysis of (A) **8a**, (B) **7** and (C) **9a** in CDCl_3 to confirm the successful synthesis of **9a** via click chemistry. The peaks marked with \blacklozenge , \blacktriangledown and \blackstar belong to the characteristic NMR signals from protons of methylene linked to azide group in **8a**, alkyne terminals in **7** and triazole rings in the terminal part of **9a**, respectively.

RESEARCH ARTICLE

NMR peak at 2.30 ppm (labelled as blue triangle, corresponding to the alkynyl terminals in **7**) revealed that all the alkynyl terminals were fully consumed in the click reaction. Also, the newly generated triazole functions were indicated with the NMR peak at 7.68 ppm (labelled as pink star, corresponding to the protons of the eight triazole rings in the terminal part of **9a**). These NMR data confirmed the successful conjugation of the carbohydrate units at the dendrimer terminals (Figure 5B and 5C). It is noteworthy that the protons on the linker (labelled with green square) between monosaccharide and triazole moved to the low-field owing to the deshielding effect of newly formed triazoles (Figure 5A and 5C).

Upon treatment with LiOH, the 32 acetyl groups in dendrimers **9a** and **9b** were successfully removed within 3 hours, indicating that the large steric hindrance had little impact on the deprotection process. Both **1a** and **1b** were obtained at yields above 90% after purification using dialysis. The structural integrity and purity of all the synthesized dendrimer final products and intermediates were confirmed using ^1H -, ^{13}C -, and ^{19}F -NMR spectral analysis as well as high-resolution mass spectroscopy (HRMS) (See Supporting information).

Biocompatibility with no notable cytotoxicity

The bola-amphiphilic dendrimers **1a** and **1b** are well solubilized in water at concentrations up to 30 mg/mL. In addition, both **1a** and **1b** showed good biocompatibility with no notable cytotoxicity, as revealed by the cytotoxicity data obtained with human embryonic kidney cells (HEK293), mouse fibroblast cells (L929), and Madin-Darby canine kidney cells (MDCK) using PrestoBlue assay and lactate dehydrogenase (LDH) test (Figure 6). The PrestoBlue assay evaluates metabolic toxicity by measuring the cell viability, and the LDH test examines toxicity associated with cell membrane damage by determining the release of lactate dehydrogenase. Results from both the PrestoBlue and LDH assays (Figure 6) indicated that both **1a** and **1b** were devoid of any notable toxicity.




Figure 6. Cytotoxicity evaluation of **1a** (A, C) and **1b** (B, D) on HEK293, L929 and MDCK cells using PrestoBlue assay and lactate dehydrogenase (LDH) test. Cells treated with lysis buffer served as control with 100% LDH release. (mean \pm SD, $n = 3$).

Effective and strong binding to lectin

As ConA has a strong affinity for α -D-mannosyl residues but not for β -D-glucosyl residues,^[23, 29] we studied the binding of **1a** with ConA using **1b** as the negative control. Adding **1a** to ConA immediately generated a white precipitate owing to the strong and multivalent interaction between **1a** and ConA, whereas the ConA solution remained clear upon **1b** addition, even at concentrations up to 500 μM . Using turbidimetric assay at 450 nm, we demonstrated that the binding between **1a** and ConA was concentration dependent, and remained effective even at concentrations as low as 0.10 μM (Figure 7A). Moreover, **1a** showed significantly stronger binding compared to the conventional amphiphilic dendrimer **II** bearing the same number of mannose terminals (Figure 7B, 7C and S1). These results highlight the important contribution of the unique bola-amphiphilic structural feature of glycodendrimer **1a** towards the observed stronger protein binding and higher glycoside cluster effect.




Figure 7. Binding of **1a** towards ConA using turbidimetric assay and isothermal titration calorimetry. Turbidity curves recorded at 450 nm for the precipitation of ConA with (A) **1a** at different concentrations, and (B) **1a** and **II**, respectively, at 0.10 μM , in HEPES buffer at room temperature. (C) Chemical structure of the conventional amphiphilic dendrimer counterpart **II**. (D) ITC Thermodynamics parameters for ConA binding at 300 K. Errors are less than 8% of the value. [a] Number of ligands/number of protein monomers.

Further evaluation of the binding affinity and thermodynamics of the bola-amphiphilic dendrimer **1a** towards ConA was carried out using isothermal titration calorimetry (ITC), with the conventional amphiphilic dendrimer **II** serving as the control for comparison (Figure 7D). All the thermodynamics results are reported in Table 1 while the raw thermograms and the integrated heats for both systems can be found in Supporting Information (Figure S2). The thermodynamic signature of binding and the K_d values obtained for the single mannose molecule, used as control, agree with literature data.^[30] Furthermore, the number of detected binding sites (n) of 1.0, is consistent with the monovalent nature of the interaction of this carbohydrate with ConA. Regarding the newly synthesized glycodendrimers, the values of $n < 1$ for both dendrimers corroborate the hypothesis of a multivalent nature in the binding between their terminal carbohydrates and lectin.

RESEARCH ARTICLE

Moreover, the high entropic penalty ($-T\Delta S$) detected for both systems is in line with the possibility of a cross-linking mechanism between each dendrimer and the protein. According to the data reported in Figure 7D, **1a** seems to exploit multivalency in a more effective way upon protein binding, as the corresponding observed value of n is closer to the ideal value of 0.125, whilst that obtained for the **II/ConA** complex is more than twofold larger. Finally, a further confirmation of the lectin binding assay results comes from the analysis of the thermodynamics patterns; indeed, **1a** outperforms **II** with a more favorable ΔG value of -9.55 kcal/mol (vs. -8.44 kcal/mol, Table 1), corresponding to an affinity for ConA of 0.112 μM .

Cell-specific targeting

We also assessed the targeting ability of **1a** and **1b** towards MRs and GLUTs by examining their cellular uptake in primary microglia and astrocyte cultures, respectively chosen for their overexpression of MRs and high levels of GLUTs. Notably, microglia internalized **1a** more efficiently compared to **1b** (Figure 8A), whereas astrocytes showed prominent cellular uptake of **1b** but little of **1a** (Figure 8B). These results demonstrate the effective and specific targeting shown by **1a** to MRs on microglia and by **1b** to GLUTs on astrocytes.

Considering the high cell surface expression of MRs and GLUTs shown by many cancers,^[27] we also evaluated the targeting ability of **1a** and **1b** towards human pancreatic cancer Panc-1 cells and mouse glioma GL261 cells. Both **1a** and **1b** were effectively internalized by both cells (Figure 8C and 8D) in a concentration-dependent manner (Figure S3). Collectively, these data highlight the cancer cell-targeting ability of **1a-b** and their potential use for targeted delivery of anticancer drugs.




Figure 8. Fluorescent imaging on the cellular uptake of Cy5-labelled glycodendrimers **1a** and **1b** in (A) primary microglia cultures, (B) primary astrocyte cultures, and of the Cy3-labelled glycodendrimers **1a** and **1b** in (C) human pancreatic cancer Panc-1 cells and (D) mouse glioblastoma GL261 cells.

Conclusion

In conclusion, we established the novel bola-amphiphilic glycodendrimers bearing mannose and glucose terminals, and studied their properties with regards to binding and targeting specific carbohydrate-binding proteins. These dendrimers were readily synthesized using sequential click reactions and showed good water solubility and biocompatibility. The unique bola-amphiphilic structure of **1a** allowed its stronger binding to the lectin

ConA compared to the conventional amphiphilic glycodendrimer **II**. In addition, **1a** specifically targeted the MR while **1b** specifically targeted the GLUT, and demonstrated effective cellular uptake in microglia and astrocytes, respectively, as well as cancer cells. Collectively, these bola-amphiphilic glycodendrimers hold great promise for targeted drug delivery towards cells expressing carbohydrate-binding proteins; we are actively working in this direction.

Experimental Section

General information

Compounds **1** and **2** were synthesized according to the protocol previously established.^[22, 31] The carbohydrate derivatives **8a** and **8b** were synthesized using the reported methods.^[32, 33] Chemicals were purchased from Sigma Aldrich or Alfa Aesar. Methyl acrylate, *N,N*-dimethylethylenediamine, ethylenediamine, dichloromethane and methanol were dried using the described methods and distilled prior to use. The other chemicals were used without further purification. Concanavalin A (Con A) for carbohydrate binding studies was purchased from Sigma-Aldrich. Dialysis tubing was purchased from Sigma Aldrich (St. Quentin Fallavier, France). Analytical thin layer chromatography (TLC) was performed using silica gel 60 F254 plates 0.2 mm thick with UV light (254 and 364 nm) as revelator. Chromatography was prepared on silica gel (Merck 200-300 mesh). ^1H -, ^{13}C - and ^{19}F -NMR spectra were recorded on a Bruker Avance III 400 at 400, 100 and 376 MHz, respectively. Chemical shifts (δ) are presented in parts per million (ppm) using the residual peak of CHCl_3 at 7.26 ppm or the residual peak of CH_3OH at 3.34 ppm as internal reference. The high-resolution mass spectra (HRMS) were recorded on a SYNAPT G2 HDMS (Waters) mass spectrometer equipped with an electrospray ionization (ESI) source. The sample was ionized in positive electrospray mode under the following conditions: electrospray voltage: 2.8 kV; orifice voltage: 20 V; and nebulizer gas flow (nitrogen): 100 L/h.

Synthesis of 6. To a solution of compound **3** (180 mg, 0.10 mmol) in MeOH (10 mL) was added aqueous solution of LiOH (39 mg, 1.6 mmol) while stirring under the protection of argon. The reaction mixture was stirred at 25 °C for 3 h. The MeOH was removed by rotavapor and followed by the addition of 1 M HCl solution until the pH = 4. The product was purified by dialysis using dialysis tube of MWCO 2000, followed by lyophilization. After repeating 4 times the operation of dialysis and lyophilization, the product was lyophilized to give compound **6** as a white solid (164 mg, 93%).

Synthesis of 7. To the compound **6** (170 mg, 0.10 mmol) were added HOBt (250 mg, 1.6 mmol), EDCI (310 mg, 1.6 mmol) and propargylamine (88 mg, 1.6 mmol) in 15 mL anhydrous DMF. The vessel was sealed and purged with argon for 5 min. The reaction mixture was stirred at 25 °C for 72 h until the reaction was complete, as indicated by TLC analysis. The DMF was evaporated under reduced pressure and the resulting residue was suspended in NaHCO_3 solution (30 mL). The water phase was extracted with CH_2Cl_2 (30 mL \times 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The resulting crude product was purified by column chromatography on silica gel to yield **7** as a slight yellow oil (190 mg, 81%); $R_f = 0.35$ ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 6:1).

Synthesis of 9a. To the compound **7** (100 mg, 0.050 mmol) were added $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$ (10 mg, 0.040 mmol), sodium ascorbate (16 mg, 0.080 mmol) and a solution of **8a** (250 mg, 0.60 mmol) in 12 mL THF. The vessel was sealed and purged with argon for 5 min and 3.0 mL H_2O was then added into the mixture. The reaction mixture was stirred at 60 °C for 6 h until the reaction was complete, as indicated by TLC analysis. The THF was evaporated under reduced pressure and the resulting residue was suspended in EDTA solution (30 mL). The water phase was extracted with

RESEARCH ARTICLE

CH_2Cl_2 (30 mL \times 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The resulting crude product was purified by column chromatography on silica gel to yield **9a** as a slight yellow oil (210 mg, 79%); $R_f = 0.30$ ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 5:1).

Synthesis of 9b. To the compound **7** (100 mg, 0.050 mmol) were added $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$ (10 mg, 0.040 mmol), sodium ascorbate (16 mg, 0.080 mmol) and a solution of **8b** (250 mg, 0.60 mmol) in 12 mL THF. The vessel was sealed and purged with argon for 5 min and 3.0 mL H_2O was then added into the mixture. The reaction mixture was stirred at 60 °C for 6 h until the reaction was complete, as indicated by TLC analysis. The THF was evaporated under reduced pressure and the resulting residue was suspended in EDTA solution (30 mL). The water phase was extracted with CH_2Cl_2 (30 mL \times 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The resulting crude product was purified by column chromatography on silica gel to yield **9b** as a slight yellow oil (220 mg, 82%); $R_f = 0.30$ ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 5:1).

Synthesis of 1a. To a solution of compound **9a** (110 mg, 0.020 mmol) in MeOH (12 mL) was added aqueous solution of LiOH (31 mg, 1.3 mmol) while stirring under the protection of argon. The reaction mixture was stirred at 25 °C for 3 h. The MeOH was removed by rotavapor and followed by the addition of 1 M HCl solution until the pH = 7. The product was purified by dialysis using dialysis tube of MWCO 2000, followed by lyophilization. After repeating 4 times the operation of dialysis and lyophilization, the product was lyophilized to give compound **1a** as a white solid (72 mg, 90%).

Synthesis of 1b. To a solution of compound **9b** (110 mg, 0.020 mmol) in MeOH (12 mL) was added aqueous solution of LiOH (31 mg, 1.3 mmol) while stirring under the protection of argon. The reaction mixture was stirred at 25 °C for 3 h. The MeOH was removed by rotavapor and followed by the addition of 1 M HCl solution until the pH = 7. The product was purified by dialysis using dialysis tube of MWCO 2000, followed by lyophilization. After repeating 4 times the operation of dialysis and lyophilization, the product was lyophilized to give compound **1b** as a white solid (75 mg, 94%).

Turbidimetric assay

Mannose dendrimers and Concanavalin A (Con A) binding was studied using turbidity assay by measuring the increasing absorbance at $\lambda = 450$ nm after addition of mannose dendrimers to Con A solution. 1.0 mg/mL Con A solution in HEPES-buffer (pH 7.0) was stirred for 2 h at room temperature and filtered through a 4 μm syringe filter. 1.0 mM mannose dendrimers stock solution prepared in HEPES-buffer. HEPES-buffer solution was transferred into a quartz cuvette (1 cm cell length) and used as blank solution to set the absorbance as zero. Later Con A solution (1.9 mL) transferred into a quartz cuvette and absorbance readings were collected every second at 450 nm. After 2 min a specific volume from the respective mannose dendrimer stock solution added in Con A solution mixed vigorously for 2s using a micropipette to get desired dendrimer final concentration. The absorbance was recorded at 450 nm for 18 min and then the inhibition assay was performed by adding α -methyl mannose solution (20 μL at 10 mg/mL). Absorbance data were recorded at 450 nm for 5 min.

Isothermal titration calorimetry measurements

ITC experiments were conducted using a MicroCal PEAQ-ITC calorimeter (Malvern Ltd.) at 300 K. The cell volume was 208 μL . All experiments were conducted by step-by-step injections of a constant volume of ligand (Mannose, **1a** or **1b**) solution into the calorimetric cell containing a solution of ConA in 0.10 M HEPES buffer containing NaCl (0.90 %), CaCl_2 (0.10 mM) and MnCl_2 (0.10 mM) at pH 7.2. Specifically, after a pre-rinse injection of 0.40 L, the ligand solution was injected in 18 portions of 2.0 μL at 150 s intervals. All solutions were degassed for 30 min at room temperature under stirring at 750 r.p.m. prior to each experiment. After washing, the cell

was prerinse with a portion of the simple buffer solution and upon filling the cell and syringe, stirring was turned on and each system was allowed to thermally equilibrate for 30 min. The heat signals resulting from mixing, dilution effects and liquid friction were further confirmed by control experiments (data not shown); accordingly, they were subtracted from the relevant datasets to yield the corrected integrated data. All experiments were run in triplicate.

Cells

Primary microglia cultures were prepared from the cerebral cortices of P0-P2 C57BL6J mouse pups as previously described.^[34] Microglial cells were isolated from glial culture monolayers by gentle shaking, then collected by centrifugation, checked for viability and seeded at required density in high-glucose culture medium (DMEM supplemented with Glutamax, 10% fetal bovine serum [ThermoScientific; CA, USA] and antibiotics. The remaining glial cell culture monolayer was left with strong shaking at 180 rpm overnight to remove sparse oligodendrocytes and receive >97% pure astrocyte culture.

Human human pancreatic cancer cells Panc-1, human embryonic kidney 293 cells (HEK 293 cells), mouse fibroblast cells (L929 cells), Madin-Darby canine kidney cells (MDCK cells) and Murine glioma GL261 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Gibco, Invitrogen) containing 10% fetal bovine serum (FBS) (Biosera). Cells were maintained at 37 °C with 5% CO_2 in a humidified chamber.

PrestoBlue assay

HEK 293, L929/MDCK and Panc-1/GL261 cells were seeded at 1.0×10^4 cells/well, 4.0×10^3 cells/well and 5.0×10^3 cells/well in 96-well plates, and all the cells were allowed to grow overnight. Cells were then treated with various concentration (1.0 - 100 μM) of the bola-amphiphilic dendrimer for 72 h. Then 10 μL PrestoBlue reagent was added to each well containing 100 μL of blank, control, or treated cells in culture and incubated for another 1 - 3 h at 37 °C. The changes in cell viability were detected using fluorescence spectroscopy (excitation 570 nm; emission 610 nm). The cell viability was expressed as a percentage relative to the cells untreated with dendrimers. All samples were run in triplicate.

Lactate dehydrogenase (LDH) assay

The different cell lines were seeded in 96-well plates and cultured overnight. The density of HEK293 cells was 1.0×10^4 cells per well. The density of MDCK cells and L929 cells was 4.0×10^3 cells per well. Then, HEK293, MDCK and L929 cells were respectively treated with various concentrations (1.0 - 100 μM) of dendrimer for 24 h. Afterwards, cell membrane damage toxicity was determined using CytoTox-ONE™ Homogeneous Membrane Integrity Assay (Promega). The LDH reaction mixture was freshly prepared according to the manufacturer's protocol, 100 μL added to each well of a 96-well plate containing 100 μL of blank, control or treated cells. The cells were incubated for 30 min at 25 °C followed by adding 50 μL of stop solution. The optical density (OD) of these solutions were measured at 490 nm via fluorescence spectroscopy. Control was performed with lysis buffer and medium, and set as 100% and 0% LDH release, respectively. Each assay was performed in triplicate.

$$\text{LDH}\% = \left[\frac{(\text{the absorbance of sample} - \text{the absorbance of negative control})}{(\text{the absorbance of positive control} - \text{the absorbance of negative control})} \right] \times 100\%$$

Cellular uptake

Visualization of cells for analyzing cellular uptake was performed using fluorescence microscopes.

RESEARCH ARTICLE

Microglia were seeded at a density of $2.0 - 3.0 \times 10^5$ cells/cm² onto glass culture slides and astrocytes were plated on poly-L-lysine-coated glass cover slips in 24-well culture plates at a density of $1.0 - 1.5 \times 10^5$ cells/cm² two days before transfection. The cells were incubated with Cy5-labeled bola-amphiphilic glycodendrimers (4.0 μ M) for 3 h at 37 °C. Next the medium was removed, the cells were washed twice with PBS and the microphotographs were taken using Olympus IX70 microscope.

Panc-1 and GL261 cells were respectively seeded in 12-well plate (1.0×10^5 cells/well) one day before transfection. The cells were incubated with Cy3-labeled bola-amphiphilic glycodendrimers (2.0 μ M and 4.0 μ M) for 3 h at 37 °C. The cells were washed with cold PBS buffer, and then Hoechst33342 (1.0 μ g/mL) was added, and stained for 30 min at 37 °C. A Zeiss Apotome.2 microscope (Carl Zeiss Microscopy GmbH, Goettingen, Germany) was used for visualization. Images were acquired using ZEN 3.1 pro software (Carl Zeiss GmbH)

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available in the supplementary material of this article.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (81502920, YX), Graduate Research and Innovation Foundation of Chongqing (CYB19066, WZ), Campus France PHC CAI YUAN PEI program (YX, JLI) and Orchid program (LP,C-LK), China Scholarship Council (WZ), Bourse d'Excellence Eiffel (Y-WY), Ligue Nationale Contre le Cancer (EL2021 LNCCLiP, LP), Ministry of Science and Technology of Taiwan (Y-WY, C-LK), French National Research Agency and the National Centre for Research and Development in Poland under the frame of H2020 Era-Net EURONANOMED European Research project 'NANOGLIO' (LP, BK), 'iNANOGLUN' (LP, BK), 'TARBRAINFEFC' (LP), 'NAN4TUM' (LP, C-LK). This project has received funding from EU H2020 Research and Innovation program NMBP "SAFE-N-MEDTECH" (2019-2023) (BR, LP; under grant agreement No. 814607) and H2020 Marie Skłodowska-Curie Innovative Training Network "OLIGOMED" (2021-2024) (JW, LP; under grant agreement No 956070).


Keywords: Bola-amphiphiles • glycodendrimers • Glucose transporters • Mannose receptors • Targeted delivery

Reference

- [1] A. Varki, R. D. Cummings, J. D. Esko, P. Stanley, G. W. Hart, M. Aebi, D. Mohnen, T. Kinoshita, N. H. Packer, J. H. Prestegard, R. L. Schnaar, P. H. Seeberger, *Essentials of glycobiology*, 4th ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, **2022**.
- [2] M. E. Taylor, K. Drickamer, *Introduction to glycobiology*, 3rd ed., Oxford University Press, Oxford ; New York, **2011**.
- [3] T. K. Lindhorst, *Essentials of carbohydrate chemistry and biochemistry*, 2nd, rev. and updated ed., Wiley-VCH, Weinheim, **2003**.
- [4] C. R. Bertozzi, L. L. Kiessling, *Science* **2001**, *291*, 2357-2364.
- [5] N. Sharon, H. Lis, *Science* **1989**, *246*, 227-234.
- [6] Y. C. Lee, R. T. Lee, *Acc. Chem. Res.* **1995**, *28*, 321-327.
- [7] J. J. Lundquist, E. J. Toone, *Chem. Rev.* **2002**, *102*, 555-578.
- [8] M. G. Cuesta, C. O. Mellet, J. M. G. Fernández, *Chem. Commun.* **2020**, *56*, 5207-5222.
- [9] A. P. Davis, *Chem. Soc. Rev.* **2020**, *49*, 2531-2545.
- [10] R. Roy, T. C. Shiao, *Chem. Soc. Rev.* **2015**, *44*, 3924-3941.
- [11] W. B. Turnbull, J. F. Stoddart, *Rev. Mol. Biotechnol.* **2002**, *90*, 231-255.
- [12] L. Mousavifar, R. Roy, *Molecules* **2021**, *26*, 2428.
- [13] C. Müller, G. Despras, T. K. Lindhorst, *Chem. Soc. Rev.* **2016**, *45*, 3275-3302.
- [14] L. Su, Y. Feng, K. Wei, X. Xu, R. Liu, G. Chen, *Chem. Rev.* **2021**, *121*, 10950-11029.
- [15] S. Zhang, R.-O. Moussodia, H.-J. Sun, P. Leowanawat, A. Muncan, C. D. Nusbaum, K. M. Chelling, P. A. Heiney, M. L. Klein, S. André, R. Roy, H.-J. Gabius, V. Percec, *Angew. Chem. Int. Ed.* **2014**, *53*, 10899-10903.
- [16] Q. Xiao, M. Delbianco, S. E. Sherman, A. M. R. Perez, P. Bharate, A. P. Vargas, C. R. Emmenegger, N. Y. Kostina, K. Rahimi, D. Söder, M. Möller, M. L. Klein, P. H. Seeberger, V. Percec, *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117*, 11931-11939.
- [17] Q. Xiao, S. Zhang, Z. Wang, S. E. Sherman, R. O. Moussodia, M. Peterca, A. Muncan, D. R. Williams, D. A. Hammer, S. Vértessy, S. André, H. J. Gabius, M. L. Klein, V. Percec, *Proc. Natl. Acad. Sci. U. S. A.* **2016**, *113*, 1162-1167.
- [18] Y. Song, Y. Chen, P. Li, C. M. Dong, *Biomacromolecules* **2020**, *21*, 5345-5357.
- [19] L. Sun, X. Ma, C. M. Dong, B. Zhu, X. Zhu, *Biomacromolecules* **2012**, *13*, 3581-3591.
- [20] J.-H. Fuhrhop, T. Wang, *Chem. Rev.* **2004**, *104*, 2901-2937.
- [21] J. R. Hughes, A. S. Miller, C. E. Wallace, G. N. Vemuri, P. M. Iovine, *Front. Chem.* **2021**, *8*.
- [22] X. Liu, Y. Wang, C. Chen, A. Tintaru, Y. Cao, J. Liu, F. Ziarelli, J. Tang, H. Guo, R. Rosas, S. Giorgio, L. Charles, P. Rocchi, L. Peng, *Adv. Funct. Mater.* **2016**, *26*, 8594-8603.
- [23] B. S. Cavada, V. J. S. Osterne, C. F. Lossio, V. R. Pinto-Junior, M. V. Oliveira, M. T. L. Silva, R. B. Leal, K. S. Nascimento, *Int. J. Biol. Macromol.* **2019**, *134*, 901-911.
- [24] H. J. P. v. d. Zande, D. Nitsche, L. Schlautmann, B. Guigas, S. Burgdorf, *Front. Immunol.* **2021**, *12*, 765034.
- [25] H. Feinberg, S. A. F. Jégouzo, Y. Lasanajak, D. F. Smith, K. Drickamer, W. I. Weis, M. E. Taylor, *J. Biol. Chem.* **2021**, *296*, 100368.
- [26] A. M. Navale, A. N. Paranjape, *Biophys. Rev.* **2016**, *8*, 5-9.
- [27] P. B. Ancey, C. Contat, E. Meylan, *FEBS J.* **2018**, *285*, 2926-2943.
- [28] P. T. Nyffeler, C. H. Liang, K. M. Koeller, C. H. Wong, *J. Am. Chem. Soc.* **2002**, *124*, 10773-10778.
- [29] M. Liu, X. Wang, D. Miao, C. Wang, W. Deng, *Polym. Chem.* **2020**, *11*, 3054-3065.
- [30] T. K. Dam, R. Roy, S. K. Das, S. Oscarson, C. F. Brewer, *J. Biol. Chem.* **2000**, *275*, 14223-14230.
- [31] T. Yu, X. Liu, A. L. Bolcato-Bellemin, Y. Wang, C. Liu, P. Erbacher, F. Qu, P. Rocchi, J. P. Behr, L. Peng, *Angew. Chem. Int. Ed.* **2012**, *51*, 8478-8484.
- [32] Q. Zhao, Y. Zou, J. Guo, S. Yu, X. Chai, H. Hu, Q. Wu, *Tetrahedron* **2014**, *70*, 7780-7787.
- [33] S. Combemale, J. N. Assam-Evoung, S. Houaidji, R. Bibi, V. Barragan-Montero, *Molecules* **2014**, *19*, 1120-1149.
- [34] K. A. Walentynowicz, N. Ochocka, M. Pasierbinska, K. Wojnicki, K. Stepniak, J. Mieczkowski, I. A. Ciechomska, B. Kaminska, *Front. Immunol.* **2018**, *9*, 1329.

RESEARCH ARTICLE

Entry for the Table of Contents



Innovative bola-amphiphilic glycodendrimers bearing mannose and glucose terminals were designed and synthesized. They demonstrated stronger binding to carbohydrate-binding protein ConA than the conventional amphiphilic glycodendrimer thanks to the unique dumbbell-like structure. In addition, these bola-dendrimers targeted respectively mannose receptors and glucose transporters for effective cellular uptake, and thereby hold great promise in carbohydrate-binding protein-targeted drug delivery.