



**HAL**  
open science

## Crop residues differ in their decomposition dynamics : Review of available data from world literature

S. Ntonta, I. Mathew, R. Zengeni, P. Muchaonyerwa, Vincent Chaplot

### ► To cite this version:

S. Ntonta, I. Mathew, R. Zengeni, P. Muchaonyerwa, Vincent Chaplot. Crop residues differ in their decomposition dynamics : Review of available data from world literature. *Geoderma*, 2022, 419, pp.115855. 10.1016/j.geoderma.2022.115855 . hal-03726502

**HAL Id: hal-03726502**

**<https://hal.science/hal-03726502v1>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# 1 **Crop residues differ in their decomposition dynamics: review of available** 2 **data from world literature**

3 **S. Ntonta<sup>a</sup>, I. Mathew<sup>a</sup>, R. Zengeni<sup>a</sup>, P. Muchaonyerwa<sup>a</sup> and V. Chaplot<sup>a, b</sup>**

4  
5 <sup>a</sup>University of Kwa-Zulu Natal, School of Agricultural, Earth & Environmental Sciences, Scottsville,  
6 3209 Pietermaritzburg, South Africa

7 <sup>b</sup>Laboratoire d'Océanographie et du Climat: Expérimentations et approches numériques (LOCEAN),  
8 UMR 7159, IRD/C NRS/UPMC/MNHN, IPSL, 4, Place Jussieu, 75252 Paris, France

## 9 10 **ABSTRACT**

11 Decomposition of crop residues may affect soil organic carbon (C) stocks, which are key for  
12 soil fertility improvement and mitigation of climate change. Numerous independent studies  
13 across the world point to contradictory results but their existence provides an opportunity to  
14 conduct a comprehensive analysis of the impact of crop type on residue decomposition. In the  
15 present study, data from 394 trials from across the world were used to assess cumulative  
16 CO<sub>2</sub> emissions from residues of 17 crops during 0-30, 0-90 and 0-120 days (i.e. C<sub>R30</sub>, C<sub>R90</sub>  
17 and C<sub>R120</sub>; 1-[C<sub>R30</sub>/C<sub>R120</sub>] ratio as a stability index of C emissions) and to relate the results  
18 with residue quality (C, N and lignin concentrations) and selected soil properties (texture, pH,  
19 soil organic carbon concentration). At all durations, legumes exhibited the  
20 highest CO<sub>2</sub> emissions per gram of C added (1003 mg CO<sub>2</sub>-C g<sup>-1</sup>C after 120 days) followed  
21 by grasses (947), oilseed crops (944) and cereals (846), with the legumes and grasses  
22 showing the lowest temporal stability of C emission as pointed out by a 1-[C<sub>R30</sub>/C<sub>R120</sub>] of  
23 0.78 and 0.79, respectively, versus 0.82 and 0.83 for cereals and oilseed crops. At all  
24 durations, maize residues emitted the least C-CO<sub>2</sub> (86, 275 and 495 mg CO<sub>2</sub>-C g<sup>-1</sup>C),  
25 followed by two other lignin rich crops (cotton and sunflower), while the highest emissions

26 were from Alfalfa residues that produced about 4 times more CO<sub>2</sub> (e.g. 359 at C<sub>R</sub>30 and 1319  
27 at C<sub>R</sub>120) than maize. Overall, CO<sub>2</sub> emissions were positively correlated with soil clay  
28 concentration ( $r>0.22$ ), residue C concentration (e.g.  $r=0.46$  at C<sub>R</sub>90 and  $r=0.37$  with  
29 emission stability,  $P<0.05$ ) but negatively to residue N concentration ( $r=-0.26$  at C<sub>R</sub>120,  
30  $P<0.05$ ). The global trend pointed to decreased CO<sub>2</sub> emissions with increasing residue lignin.  
31 Contrary to what is generally believed, providing the soil with high lignin and high N  
32 concentration may foster C stabilization into soils by soil microbes.

33 *Keywords:* Crop residue decomposition, C emissions, Soil carbon stabilisation, Soil C stocks

---

34

## 35        **1. Introduction**

36    The carbon (C) cycle has received considerable attention in recent years, due to concerns  
37    over the continued increase in atmospheric carbon dioxide (CO<sub>2</sub>) concentration. Annual  
38    increases of CO<sub>2</sub> concentration in the atmosphere were observed on a global scale for the  
39    years 2018 (2.7%) and 2019 (0.6%) (Ritchie and Roser, 2020). Global warming, as a result of  
40    high atmospheric CO<sub>2</sub> causes climate change, characterized by increase in mean temperature  
41    and higher variability in precipitation. Consequently, the need to sequester carbon in  
42    agricultural soils has been identified as a sustainable strategy to mitigate climate change and  
43    promote agricultural sustainability.

44    Soil is the second largest C reservoir with 11% (4655 Pg C), after the 87% of global carbon  
45    stocks being oceanic (38,000 Pg C) and has higher C than the 2% found in the atmosphere  
46    (860 Pg C) (Xiao, 2015; Venter et al., 2021). Carbon transfer from the atmosphere to soil is  
47    achieved by plants through photosynthesis, leading to exudation of C compounds from roots  
48    during the growth cycle, and by the retention of plant root and shoot residues. Some of the C  
49    from plant residues or exudates can be mineralized and emitted back to the atmosphere as  
50    CO<sub>2</sub> while a significant proportion can be stabilized as soil organic matter (Ontl and Schulte,  
51    2012; Dignac, 2017). Therefore, the rate of residue decomposition and associated CO<sub>2</sub>  
52    emissions is often used as a proxy for evaluating the potential of plant residues to become soil  
53    organic matter (e.g. Johnson et al., 2007).

54    The potential to increase soil C stocks is mostly on managed agricultural land where  
55    numerous studies exist on the fate of C from different crop residues. Curtin et al. (2008) for  
56    instance, observed that barley straw emitted significantly higher ( $p < 0.05$ ) amount of CO<sub>2</sub>  
57    (55 g CO<sub>2</sub>-C m<sup>-2</sup>) compared to wheat straw (47 g CO<sub>2</sub>-C m<sup>-2</sup>) after 158 days. Ajwa and  
58    Tabatabai (1994) found 58% of organic C evolved as CO<sub>2</sub>-C from alfalfa-treated soils in 30  
59    days which was higher compared to maize residues (30%), which was attributed to the higher

60 C: N of maize residues. Zeng et al. (2010) attributed the higher CO<sub>2</sub> emissions from peanut  
61 root (maximum of 60 %) compared to maize root residues (35%) to biological nitrogen  
62 fixation, which increases N in leguminous plants, and thus lower their C: N ratio, and overall  
63 quality of the crop residues. Not only does crop residue decomposition depend on the type  
64 and quality of crop residues but also on the internal soil conditions (Mathew, et al., 2017;  
65 Stewart et al., 2015).

66 Mathew et al. (2017) concluded that higher plant C stocks and C transfer to soils occurred in  
67 carbon rich clayey soils of tropical humid areas due to higher biomass production potential  
68 compared to sandy soils. Clayey soils also support high C stocks through their aggregation  
69 and ability to provide physical protection as well as mineral adsorption of C constituents  
70 (Elliott, 1986; Ajwa and Tabatabai, 1994; Martens, 2000; Clark, 2007; Mathew, et al., 2017).

71 Several studies have also reported disparities in CO<sub>2</sub> emissions from residues of different  
72 crops under different soil types and climates, which has led to a lack of consensus on the  
73 impact of these factors on CO<sub>2</sub> emissions from soils (Li et al., 1994; Paustian et al., 2000;  
74 Gregorich et al., 2001; Alvarez, 2005; Abdalla et al., 2016; McClelland et al., 2021; Shakoor  
75 et al., 2021). However, the existence of multiple individual studies across the world provides  
76 an opportunity to conduct a comprehensive analysis of the main factors that control crop  
77 residue decomposition and CO<sub>2</sub> emissions and the consequences for the building of soil  
78 carbon stocks. Therefore, the objective of the current study was to evaluate the impact of crop  
79 type, soil and environmental factors on CO<sub>2</sub> emissions from residues of selected crops using  
80 available global data from published studies.

81

## 82 **2. Materials and methods**

### 83 *2.1. Study setup: Database preparation*

84 Data on CO<sub>2</sub> emissions from crop residues incubated in soil at laboratories were collated  
85 from studies across the world, published in peer reviewed journals and accessible from public  
86 domains such as Google scholar, Ref-seek, Science Direct, Sci-Finder, Scopus, Springer  
87 Link, Research-Gate and Web of Science. Key words such as “litter decomposition”, “residue  
88 decomposition”, “C mineralisation”, “crop residue CO<sub>2</sub> emissions”, “C gases”, “carbon  
89 dynamics” and “decomposition rate” were used to search for relevant journal articles. The  
90 available papers had to report on CO<sub>2</sub> emissions from crops (sorghum, wheat, maize, among  
91 others) and on crop quality factors affecting residue decomposition (e.g. total C and N, C: N  
92 ratio, lignin, cellulose or hemicellulose). Furthermore, environmental factors such as climatic  
93 information (mean annual precipitation and mean annual temperature), geographic variables  
94 (latitude, longitude) as well as soil variables (physicochemical properties) measured during  
95 both laboratory or/and field experiments were considered as controlling variables. The studies  
96 had to strictly compare CO<sub>2</sub> emissions between soils incubated with and without (considered  
97 as the control) residues under the same conditions. Moreover, the mass of residues used must  
98 have been clearly stated or able to be deduced. The data were used to compile a database with  
99 quantitative and qualitative data on plant litter quality.

100

## 101 *2.2. Variables of CO<sub>2</sub> emission*

102 The effect of adding crop residues on soil CO<sub>2</sub> emissions were calculated as the difference  
103 between CO<sub>2</sub> emitted from the soil containing residues and CO<sub>2</sub> evolved from the control.  
104 The values were converted from their original units and normalized to common units (mg  
105 CO<sub>2</sub>-C/ g C added of soil over 30, 90 and 120 day periods of incubation) (Table 2). The  
106 amounts of total CO<sub>2</sub> emissions were recorded over 0-30; 30-90 and 90-120 day periods and  
107 were cumulatively representing lability of residue decomposition. The studies that did not  
108 cover the 0-30, 30-90 and 90-120 periods were estimated by use of linear trendline equation.

109 In addition, a ratio between CO<sub>2</sub> emissions at 30 relative to 120 days was calculated, which  
110 was used as an index of temporal stability of C-CO<sub>2</sub> emissions = 1- (C<sub>R30</sub>/ C<sub>R120</sub>). The  
111 greater the index, the greater stability of the emissions over time.

112

### 113 *2.3. Crop quality, soil and environmental factors*

114 Crops were categorized into cereals, legumes, grasses and oilseed (Table 3), (i.e., cereals-  
115 barley, maize, oat, rice, rye, sorghum, wheat; grass- grass; legumes-alfalfa, bean, clover, pea,  
116 soyabean, vetch; and oilseed-canola, cotton, and sunflower). Cereals and grasses were  
117 separated due to their functional differences, with cereals having evolved and undergone  
118 selection by farmers, making them different from natural or forage grasses. In this study,  
119 grasses refer to natural or forage grass that are not used for human consumption while cereal  
120 refers to monocot grasses that are used primarily for grain production for human  
121 consumption. Legumes are defined as any crop that has a natural ability to fix nitrogen, while  
122 oilseed crops are those that are primarily used for extraction of vegetable oil from their seed.  
123 Soyabean was considered as a legume crop in this study. Furthermore, crop quality was  
124 defined by residue chemical composition, such as initial C (TC) and lignin concentrations,  
125 C:N ratio, total nitrogen (TN), dissolved carbon (DC) and total phosphorus (TP); cellulose,  
126 polyphenols and lignin: N ratios. In addition, soil properties such as texture (clay, silt and  
127 sand concentration), soil pH (CaCl<sub>2</sub>) and organic carbon (SOC) were considered and  
128 classified into different categories (Table 3) following Abdalla et al. (2016); Mutema et al.  
129 (2015) and Mathew et al. (2017). Water-based pH was converted to CaCl<sub>2</sub> pH following the  
130 equation of Lierop (1981): ( $y = 0.53 + 0.98x$ ). Where y is pH on the CaCl<sub>2</sub> scale and x is the  
131 water-based pH.

132 Climatic factors included a 30-year average rainfall and temperature (mean annual  
133 precipitation: MAP) and (mean annual temperature: MAT). The climate was further classified

134 as tropical (hot and wet), sub-tropical (warm and arid to humid) or temperate (cool to cold  
135 and mild to warm). In cases where climatic characteristics were not present in a particular  
136 study, appropriate data such as annual precipitation and temperature were obtained using the  
137 location coordinates or surrogate data for nearby prominent features (e.g. town) through  
138 Google search. In addition, the geographical positioning system (GPS) using latitude and  
139 longitude coordinates were used to depict the global distribution of the studies used in the  
140 review (Fig. 1).

141 A total of 58 journal articles (Table 1) were used, detailing different studies across the world,  
142 which provided 394 observations. The name of authors, year of paper publication, country  
143 and geographical location of experimental site, nature of experiment, experimental duration  
144 (time periods), crop(s) or crop types used in the experiments, quantitative information on  
145 plant quality, soil properties as well as C variables and environmental conditions were  
146 captured onto a database. The definitions and acronyms adopted in this paper are used to  
147 simplify the terms and definitions of variables for purposes of this analysis.



148

#### 149 *2.4. Data analyses*

150 The data were compiled into a database and tested for normality of variables, linearity and  
151 homoscedasticity prior to statistical analyses. Descriptive summary statistics (minimum,  
152 maximum, median, mean, SEM: standard error of mean, 25<sup>th</sup> and 75<sup>th</sup> percentiles, skewness  
153 (Skew), kurtosis (Kurt) and coefficient of variation (CV%) were calculated for all variables  
154 (Table 6). Furthermore, box plots were constructed to depict the distribution of the data  
155 showing minimum, 25<sup>th</sup> quartile median, mean, 75<sup>th</sup> quartile and maximum values within the  
156 5 and 95<sup>th</sup> percentiles. Mean CO<sub>2</sub> emissions were computed for different crops, with different  
157 crop residue quality, soil properties and environmental factor classes. Because the CO<sub>2</sub>  
158 emission data did not conform to normal distributions for us to use parametric analysis, the  
159 non-parametric Kruskal-Wallis analysis was performed. Significant differences between  
160 factor classes were tested with non-parametric analysis (Kruskal-Wallis), at chi-square  
161 probability of <0.001. In addition, bivariate Pearson correlations coefficients at p<0.05 were  
162 calculated among the variables (Table 7). Subsequently, principal component analysis (PCA),  
163 which converts non-linear factors and variables into linear combinations for visualization  
164 (Jambu, 1991), were conducted to investigate the multiple correlations between the variables  
165 (Fig. 10 and 11). Finally, because of low number of data points the variable of residue  
166 dissolved carbon, total residue phosphorus, cellulose and polyphenols were discarded. SiAll  
167 analyses were performed using Statistica 10.0 software (Weiß, 2007).

168

### 169 **3. Results**

#### 170 *3.1. CO<sub>2</sub> emissions from crop residues*

171 The summary statistics (Table 4) which were computed from all study sites showed that the  
172 30-days cumulative CO<sub>2</sub> emissions (C<sub>R30</sub>) ranged between 0.3 and 920.1 mg CO<sub>2</sub>-C g<sup>-1</sup> C,

173 with a mean value at 196.5 mg CO<sub>2</sub>-C g<sup>-1</sup> C. Cumulative 120 days emissions (C<sub>R</sub>120) were  
174 between 3.2 and 3640 mg CO<sub>2</sub>-C g<sup>-1</sup> C with a mean at 914.2 mg CO<sub>2</sub>-C g<sup>-1</sup> C (Table 4). The  
175 resulting emission stability index showed a mean at 0.8.

176 Cumulative emissions also varied among different crops, with legumes exhibiting the highest  
177 mean cumulative emissions of 228.0 mg CO<sub>2</sub>-C g<sup>-1</sup> C after 30 days, 586.7 mg CO<sub>2</sub>-C g<sup>-1</sup> C  
178 after 90 days and 1003.0 mg CO<sub>2</sub>-C g<sup>-1</sup> C after 120 days (Table 5). Grasses ranked second  
179 with respectively, 217.0, 529.7 and 946.8 mg CO<sub>2</sub>-C g<sup>-1</sup> C, while oilseed crops ranked third  
180 and cereals fourth (Table 5). The average difference between legumes and cereals was 25%  
181 after 30 days, 19% after 90 days and 16% after 120 days, which corresponded for C<sub>R</sub>30 and  
182 C<sub>R</sub>90 to significant differences at P<0.001. On average, legumes had the lowest temporal C  
183 emissions stability (0.78) followed by grass (0.79), cereals (0.82) and oilseed (0.83)  
184 (Table 5).

185 Table 6 shows variations in CO<sub>2</sub> emissions between different crop types, with alfalfa having  
186 the highest emissions of 359.0 mg CO<sub>2</sub>-C g<sup>-1</sup>C, amongst legumes (48% higher than clover  
187 and pea), while canola (293.5) had the highest amongst oilseed crops, and sorghum (261.1)  
188 had the highest amongst cereals (67% more than maize) at C<sub>R</sub>30. Maize emitted the least  
189 cumulative CO<sub>2</sub> at C<sub>R</sub>30 (85.8 mg CO<sub>2</sub>-C g<sup>-1</sup>C), C<sub>R</sub>90 (275.1) and C<sub>R</sub>120 (495.0) but  
190 exhibited with 0.84 the highest CO<sub>2</sub> emission stability over time. In contrast, alfalfa exhibited  
191 the lowest stability of CO<sub>2</sub> emissions with a mean value at 0.76, followed by rye (0.77),  
192 clover and pea (0.78), canola, grass and oat (0.79) (Table 6; Fig. 2D).

### 193 *3.2. The influence of crop residue quality on CO<sub>2</sub> emissions*

194 The quality of crop residues, determined by the initial C, N and lignin concentrations,  
195 significantly ( $p < 0.05$ ) affected CO<sub>2</sub> emissions (Table 7). There was a systematic decrease in  
196 C emissions with the increase in residue nitrogen content with  $r$  from -0.22 at C<sub>R30</sub>, -0.26 at  
197 C<sub>R120</sub> and -0.28 at C<sub>R90</sub>. The  $r$  values with lignin content were from -0.08 at C<sub>R30</sub> to -0.12 at  
198 C<sub>R120</sub>. Emissions also decreased with increasing C:N with the highest  $r$  (-0.14) found for  
199 C<sub>R30</sub>. In contrast C emissions increased with increasing residue C content from 0.40 at  
200 C<sub>R120</sub> to 0.59 at C<sub>R30</sub>. Furthermore, the increase in residue C content significantly enhanced  
201 the temporal stability of C emissions ( $r = 0.37$ ).

### 203 *3.3. The impact of soil and environmental properties on CO<sub>2</sub> emissions from different* 204 *crop residues*

205 There were significant variations in cumulative CO<sub>2</sub> emissions among the three different soil  
206 textural classes throughout the incubation periods (Fig. 4). For instance, clayey soils  
207 exhibited significantly higher cumulative 30-days CO<sub>2</sub> emissions (224 mg CO<sub>2</sub>-C g<sup>-1</sup> C) as  
208 compared to sandy soils (178 mg CO<sub>2</sub>-C g<sup>-1</sup> C), while silty soils had intermediate emissions.  
209 Decreasing emissions with increasing sand content was a trend also found for C<sub>R90</sub> and  
210 C<sub>R120</sub>. Finally, the temporal stability of CO<sub>2</sub> emissions was surprisingly the highest under  
211 silty conditions ( $1 - (C_{R30}/C_{R120}) = 0.81$ ) and decrease to 0.80 under clayey conditions and to  
212 0.79 for sandy soils (Fig. 4).

213 Cumulative CO<sub>2</sub> emissions also varied due to soil pH (Fig. 5). For all incubation durations,  
214 the lowest emissions were observed for strongly acidic soils while the highest emissions were  
215 found for slightly acidic soils. Additionally, there was a trend for emissions to lessen from  
216 slightly acidic to alkaline through neutral while the temporal stability of the emissions  
217 consistently rose from acidic to alkaline (Fig. 5D).

218 Figure 6 which depicts the impact of climate points to a significant decrease of emissions  
219 from tropical to temperate through sub-tropical. In the case of  $C_{R30}$ ,  $CO_2$  emissions  
220 decreased from an average of 252 mg  $CO_2-C\ g^{-1}\ C$  for tropical to 193 mg  $CO_2-C\ g^{-1}\ C$  for  
221 subtropical and to 150 mg  $CO_2-C\ g^{-1}\ C$  for temperate, which corresponded in all cases to  
222 significant differences at  $P < 0.01$  (Fig. 6A). Similar trends (i.e. a decrease of emissions from  
223 tropical to temperate) were also observed for  $C_{R90}$  and  $C_{R120}$  but the temporal stability of  
224 the emissions consistently rose from tropical to temperate, the differences between sub-  
225 tropical and temperate being however non-significant (Fig. 6D).

226 Multivariate correlations between  $CO_2$  emission variables on the one hand and soil and crop  
227 residue variables on the other hand are displayed on Figure 7. The two principal components  
228 of this PCA explained 99% of the total variation in the data with the first principal component  
229 (PC1) accounting for 79% of data variance and PC2 accounting for 20%. Cumulative  
230 emissions to 30, 90 and 120 days showed a positive correlation to PC1 and this axis can be  
231 thus interpreted as an axis of crop residue decomposition. The temporal stability of  $CO_2$   
232 emissions had a negative coordinate on Axis 2, meaning that PC2 could be interpreted as an  
233 axis of decomposition in-stability. Residue content in lignin, N and C showed negative  
234 coordinates on PC1 while the C:N ratio showed a positive coordinate. There was thus a  
235 tendency for emissions to increase as C:N increases but C, N and lignin concentration  
236 decreases. Finally, emission instability decreased with increasing pH as pointed by a negative  
237 coordinate of pH on PC2 (Fig. 7). Amongst crops, canola and alfalfa correlated to the high  
238  $CO_2$  emissions level and low  $CO_2$  emission stability pole, while maize, cotton and sunflower  
239 correlated to the low emissions and high stability pole.

240

## 241 **4. Discussion**

242 *4.1. Causes of variation in residue decomposition and soil C building amongst crop types*

243 The different crop types exhibited large variations in CO<sub>2</sub> emissions with legumes emitting  
244 the highest cumulative CO<sub>2</sub> emissions and with decreasing emissions over time versus cereals  
245 pointing to lower emissions but of higher stability. Such variations were shown to correlate  
246 with the quality of crop residues (Machinet, et al., 2009). Ajwa and Tabatabai (1994)  
247 revealed that the significantly higher amounts of CO<sub>2</sub>-C released from alfalfa were to be  
248 attributed to their ability to fix N, as they had higher initial N concentration, of 12.6% as  
249 compared to the 6.9% of maize and the 1.3% of oilseed crops. The underlying hypothesis of  
250 greater emissions at high initial N concentration was, as suggested by Gezahegn, et al.,  
251 (2016), the enhanced microbial activity leading to high decomposition and CO<sub>2</sub> emissions.  
252 But the present study, which was based on 394 trials from across the world, tends to  
253 contradict this past statement as it pointed to a negative correlation between CO<sub>2</sub> emissions  
254 and residue N concentration. Such a trend might be due to the fact that N availability  
255 enhances C uptake by microbes and thus humus formation versus C emissions to the  
256 atmosphere as CO<sub>2</sub>, a mechanism that was described by authors such as Henriksen and  
257 Breland (1999), Rousk and Bååth (2007), Bai et al. (2016) and Köbke et al. (2018). The fact  
258 that legumes, especially Alfalfa which accounted for the most emissions amongst crops  
259 during the maximum 120 days period, emitted large amounts of CO<sub>2</sub> despite a high N  
260 concentration was most likely due to its low lignin concentration.

261 Carbon stabilization into soils in microbial biomass thus seems to be favoured by the supply  
262 of residues with a C and N stoichiometric ratio close to that of living microorganisms. Results  
263 on maize, sunflower and cotton tend to show that lignin rich residues experience low  
264 decomposition rates which constitutes a second route of carbon stabilization into soils.

265

266 *4.2.The impact of crop residues on the loss of soil carbon through priming*

267 Recent studies have indeed shown that the activity of decomposers and their ability to  
268 decompose soil organic matter for their living can be stimulated by the addition of fresh  
269 organic matter resulting in an increase in soil respiration beyond C addition, which is referred  
270 to as 'priming' (e.g. Fontaine et al. 2003; Kuzyakov, 2010). In the present study and as  
271 pointed out in Table 8, 43% of the respiration data points showed 120 days cumulative  
272 emissions beyond C addition, which points to the existence of significant C losses from soil  
273 organic matter. All crop types experienced priming with the proportion of studies with C  
274 losses over residue C from 25% for maize, sunflower and beans to over 55% for sorghum,  
275 alfalfa and canola and with differences between legumes, cereals and oilseed crops being  
276 non-significant at  $p < 0.05$ .

277 Several authors have suggested that chemically recalcitrant residues, such as those rich in  
278 lignin decompose more slowly than residues with low lignin and high N concentrations, thus  
279 leading to enhanced C stabilization into soils and increased soil organic matter (Johnson et al.  
280 2007, Berg and Mc Clagherty, 2008). However, several other authors such as Stewart et al.  
281 (2015), pointed out that high lignin residues are used inefficiently by the soil microbial  
282 community that decompose SOM (priming effect) to acquire key nutrients resulting in much  
283 greater respiration losses and less C stabilization into soils. The present study which points to  
284 a global tendency for CO<sub>2</sub> emissions to decrease with the increase in residue lignin and N  
285 concentration tend to show that providing the soil with high lignin and high N concentration  
286 may limit priming and foster C stabilization into soils by soil microbes. Moreover, the present  
287 data showed that lignin concentration minimally impacted the temporal stability of the  
288 emissions but that emissions decreased more sharply over time at higher initial residue C  
289 concentration.

290

291 *4.3. The impact of soil properties on CO<sub>2</sub> emissions*

292 Soil texture and pH had significant impact on cumulative CO<sub>2</sub> emissions from crop residues.  
293 The higher cumulative CO<sub>2</sub> emissions from clayey soils could be due to favorable living  
294 conditions for decomposers than under coarser soil conditions. Moreover, Schmatz, et al.,  
295 (2017) also found higher C emissions from clay soils as compared to sandy-loam soils due to  
296 high organic carbon concentration in clayey soils and enhanced water retention capability,  
297 thus favoring the activity of microorganisms responsible for residue break-down. Contrarily,  
298 most previous studies generally had described clay soils to enhance the physical protection  
299 and mineral adsorption of C constituents, which was not confirmed by the present analysis of  
300 world data from 120 days duration in which clay soils experiences higher CO<sub>2</sub> emissions per  
301 gram of residue C added. Addition of labile organic materials (crop residues) to clayey soils,  
302 with higher organic carbon, could result in more CO<sub>2</sub> emissions (priming effect) than from  
303 sandy loam soils. In addition, such an increase in CO<sub>2</sub> emissions with the increase in soil clay  
304 concentration might come as suggested above from higher soil moisture and bacteria  
305 concentration, favoring the rapid turnover of residue C.

306 We had no explanation for slightly acidic soils (pH of 5.5-6.4) to experience consistently  
307 higher CO<sub>2</sub> emissions (>50%) than the other pH levels irrespective of incubation durations as  
308 conditions for fungi and bacteria are not optimal (Hågvar, 1994; Stott and Martin, 1989). A  
309 possible reason for the higher CO<sub>2</sub> emissions could be the liming effect of residues that foster  
310 the priming of soil organic matter (Wang et al. 2017; Yaowu et al. 2016).

311

#### 312 *4.4. Climate impact on CO<sub>2</sub> emissions*

313 The finding of this study pointed to higher residue CO<sub>2</sub> emissions under tropical conditions  
314 than under the other climates throughout the incubation days. Tropical climates tend to  
315 experience high temperature and rainfall conditions that are conducive for production of high  
316 plant biomass and microbial activity for decomposition. On the other hand, soils of sub-

317 tropical or temperate climates have lower rainfall and cooler temperature thus limiting  
318 microbial activity and the biochemical processes involved in residue decomposition (Ontl and  
319 Schulte, 2012). As the data used in the present study come from laboratory experiments with  
320 controlled conditions of temperature and humidity, the higher decomposition rates and  
321 cumulative CO<sub>2</sub> emissions under tropical climates could be due to favourable conditions for  
322 microbial activity.

323



324 **5. Conclusions**

325 The analysis of 394 laboratory trials worldwide revealed that on average, legumes exhibited  
326 significantly higher CO<sub>2</sub> emissions than cereals, oilseed and fiber crops, with for instance  
327 alfalfa emitting 2.7 times more CO<sub>2</sub> after 120 days than maize (Figure 2, Table 6).  
328 Additionally, legumes (especially alfalfa) showed the lowest stability of CO<sub>2</sub> emissions over  
329 time (i.e., the highest decrease from 30 to 120 days) followed by grasses, cereals and oilseed  
330 crops.

331 Amongst the two models for enhanced C sequestration into soils the present analysis points to  
332 a global tendency for CO<sub>2</sub> emissions to decrease with the increase in residue lignin. This  
333 contrasted with the studies showing that high lignin concentration induce enhanced microbial  
334 decomposition and thus C outputs from soils i.e., alfalfa (low lignin and high N  
335 concentration) versus maize residue (high lignin and low N concentration). Providing soils  
336 with residues with high lignin such as through maize residues may improve C uptake by  
337 microbes (a key process in humus formation).

338 A large proportion of the variance in the data remained however unexplained thus calling for  
339 further analysis of variables such as soil nutrients, microbial communities and quality of  
340 lignin, and of other organic matter compounds. More work is also to be done on selecting  
341 crop cultivars for generating superior ones in our fight against land and climate degradation.  
342 Finally, the question whether the carbon remaining into the soil after 120 days of incubation  
343 will be stabilized to contribute to soil carbon stocks calls for further research to understand  
344 the fate and underlying mechanisms of C sequestration into soils.

345

346 **6. Acknowledgements**

347 This study received financial support from the Water Research Commission of the Republic  
348 of South Africa (WRC Project No. K5/2721/4). The authors also extend their gratitude to the“

349 Carbon Crew” members Nozibusiso Mbava and Nhlakanipho Mbambo, both student from the  
350 University of KwaZulu-Natal at the time of this study, for their support during gathering of  
351 the data used in the paper.

352

353 **7. Reference**

- 354 Abdalla, K., Chivenge, P., Ciaia, P. and Chaplot, V., 2016. No-tillage lessens soil CO<sub>2</sub> emissions  
355 the most under arid and sandy soil conditions: results from a meta-analysis. *Biogeosciences*,  
356 13(12), pp.3619-3633.
- 357 Abiven, S. and Recous, S., 2007. Mineralisation of crop residues on the soil surface or  
358 incorporated in the soil under controlled conditions. *Biology and Fertility of Soils*, 43(6),  
359 pp.849-852.
- 360 Abiven, S., Recous, S., Reyes, V. and Oliver, R., 2005. Mineralisation of C and N from root,  
361 stem and leaf residues in soil and role of their biochemical quality. *Biology and fertility of*  
362 *soils*, 42(2), p.119.
- 363 Abro, S.A., Tian, X., Wang, X., Wu, F. and Kuyide, J.E., 2011. Decomposition characteristics  
364 of maize (*Zea mays*. L.) straw with different carbon to nitrogen (C/N) ratios under various  
365 moisture regimes. *African Journal of Biotechnology*, 10(50), pp.10149-10156.
- 366 Ajwa, H.A. and Tabatabai, M.A., 1994. Decomposition of different organic materials in soils.  
367 *Biology and Fertility of Soils*, 18(3), pp.175-182.
- 368 Alvarez, R., 2005. A review of nitrogen fertilizer and conservation tillage effects on soil organic carbon  
369 storage. *Soil Use and Management*, 21(1), pp.38-52.
- 370 Angers, D.A. and Recous, S., 1997. Decomposition of wheat straw and rye residues as affected  
371 by particle size. *Plant and soil*, 189(2), pp.197-203.
- 372 Anguria, P., Chemining'wa, G., Onwonga, R. and Ugen, M., 2017. Decomposition and nutrient  
373 release of selected cereal and legume crop residues. *Journal of Agricultural Science*, 9(6),  
374 p.p108.
- 375 Arunachalam, K., Singh, N.D. and Arunachalam, A., 2003. Decomposition of leguminous crop  
376 residues in a 'jhum' cultivation system in Arunachal Pradesh, India. *Journal of Plant*  
377 *Nutrition and Soil Science*, 166(6), pp.731-736.

378 Aulakh, M.S., Walters, D.T., Doran, J.W., Francis, D.D. and Mosier, A.R., 1991. Crop residue  
379 type and placement effects on denitrification and mineralization. *Soil Science Society of*  
380 *America Journal*, 55(4), pp.1020-1025.

381 Begum, N., Guppy, C., Herridge, D. and Schwenke, G., 2014. Influence of source and quality of  
382 plant residues on emissions of N<sub>2</sub>O and CO<sub>2</sub> from a fertile, acidic Black Vertisol. *Biology*  
383 *and fertility of soils*, 50(3), pp.499-506.

384 Berg, B. and McClaugherty, C., 2008. Decomposition as a process. *Plant Litter: Decomposition,*  
385 *Humus Formation, Carbon Sequestration*, pp.11-33.

386 Bertrand, I., Chabbert, B., Kurek, B. and Recous, S., 2006. Can the biochemical features and  
387 histology of wheat residues explain their decomposition in soil?. *Plant and Soil*, 281(1-2),  
388 pp.291-307.

389 Blaise, D. and Bhaskar, K.S., 2003. Carbon mineralization patterns of cotton leaves and stems in  
390 vertisols and inceptisols. *Archives of Agronomy and Soil Science*, 49(2), pp.171-177.

391 Bontti, E.E., Decant, J.P., Munson, S.M., Gathany, M.A., Przeszlowska, A., Haddix, M.L.,  
392 Owens, S., Burke, I.C., Parton, W.J. and Harmon, M.E., 2009. Litter decomposition in  
393 grasslands of central North America (US Great Plains). *Global Change Biology*, 15(5),  
394 pp.1356-1363.

395 Cayuela, M.L., Sinicco, T. and Mondini, C., 2009. Mineralization dynamics and biochemical  
396 properties during initial decomposition of plant and animal residues in soil. *Applied soil*  
397 *ecology*, 41(1), pp.118-127.

398 Cong, W.F., Hoffland, E., Li, L., Janssen, B.H. and van der Werf, W., 2015. Intercropping  
399 affects the rate of decomposition of soil organic matter and root litter. *Plant and*  
400 *Soil*, 391(1-2), pp.399-411.

401 Clark, G.J., Dodgshun, N., Sale, P.W.G. and Tang, C., 2007. Changes in chemical and  
402 biological properties of a sodic clay subsoil with addition of organic amendments. *Soil*  
403 *Biology and Biochemistry*, 39(11), pp.2806-2817.

404 Corbeels, M., Hofman, G. and Van Cleemput, O., 2000. Nitrogen cycling associated with the  
405 decomposition of sunflower stalks and wheat straw in a Vertisol. *Plant and soil*, 218(1-2),  
406 pp.71-82.

407 Curtin, D., Francis, G.S. and McCallum, F.M., 2008. Decomposition rate of cereal straw as  
408 affected by soil placement. *Soil Research*, 46(2), pp.152-160.

409 Datta, A., Jat, H.S., Yadav, A.K., Choudhary, M., Sharma, P.C., Rai, M., Singh, L.K.,  
410 Majumder, S.P., Choudhary, V. and Jat, M.L., 2019. Carbon mineralization in soil as  
411 influenced by crop residue type and placement in an Alfisols of Northwest India. *Carbon*  
412 *Management*, 10(1), pp.37-50.

413 Davies, B.E., 1971. A statistical comparison of pH values of some English soils after  
414 measurement in both water and 0.01 M calcium chloride. *Soil Science Society of America*  
415 *Journal*, 35(4), pp.551-552.

416 De Neergaard, A., Hauggaard-Nielsen, H., Jensen, L.S. and Magid, J., 2002. Decomposition of  
417 white clover (*Trifolium repens*) and ryegrass (*Lolium perenne*) components: C and N  
418 dynamics simulated with the DAISY soil organic matter submodel. *European Journal of*  
419 *Agronomy*, 16(1), pp.43-55.

420 Dignac, M.F., Derrien, D., Barre, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet,  
421 G.T., Garnier, P., Guenet, B. and Hedde, M., 2017. Increasing soil carbon storage:  
422 mechanisms, effects of agricultural practices and proxies. A review. *Agronomy for*  
423 *sustainable development*, 37(2), p.14.

424 Donahue, R.L., Miller, R.W. and Shickluna, J.C., 1983. *Soils. An introduction to soils and plant*  
425 *growth*.

426 Duong, T.T.T., Baumann, K. and Marschner, P., 2009. Frequent addition of wheat straw  
427 residues to soil enhances carbon mineralization rate. *Soil Biology and Biochemistry*, 41(7),  
428 pp.1475-1482.

429 Dlamini, P., 2014. *Grassland Degradation and Rehabilitation of Soil Organic Carbon and*  
430 *Nitrogen Stocks* (Doctoral dissertation, University of KwaZulu-Natal, Pietermaritzburg).

431 Drury, C.F., Yang, X.M., Reynolds, W.D. and McLaughlin, N.B., 2008. Nitrous oxide and  
432 carbon dioxide emissions from monoculture and rotational cropping of corn, soybean and  
433 winter wheat. *Canadian Journal of Soil Science*, 88(2), pp.163-174.

434 Elliott, E.T., 1986. Aggregate structure and carbon, nitrogen, and phosphorus in native and  
435 cultivated soils. *Soil science society of America journal*, 50(3), pp.627-633.

436 Finn, D., Page, K., Catton, K., Strounina, E., Kienzle, M., Robertson, F., Armstrong, R. and  
437 Dalal, R., 2015. Effect of added nitrogen on plant litter decomposition depends on initial  
438 soil carbon and nitrogen stoichiometry. *Soil Biology and Biochemistry*, 91, pp.160-168.

439 Fontaine, S., Mariotti, A. and Abbadie, L., 2003. The priming effect of organic matter: a  
440 question of microbial competition?. *Soil Biology and Biochemistry*, 35(6), pp.837-843.

441 Fruit, L., Recous, S. and Richard, G., 1999. Plant residue decomposition: effect of soil porosity  
442 and particle size. In *Effect of mineral-organic-microorganism interactions on soil and*  
443 *freshwater environments* (pp. 189-196). Springer, Boston, MA.

444 Geisseler, D., Horwath, W.R. and Scow, K.M., 2011. Soil moisture and plant residue addition  
445 interact in their effect on extracellular enzyme activity. *Pedobiologia*, 54(2), pp.71-78.

446 Gezahegn, A.M., Abd Halim, R., Yusoff, M.M. and Abd Wahid, S., 2016. Decomposition and  
447 Nitrogen mineralization of Individual and Mixed Maize and Soybean Residue. *Journal of*  
448 *Agricultural Science*, 2, pp.28-45.

449 Ghimire, B., Ghimire, R., VanLeeuwen, D. and Mesbah, A., 2017. Cover crop residue amount  
450 and quality effects on soil organic carbon mineralization. *Sustainability*, 9(12), p.2316.

451 Glasser, W.G., 1985. Lignin. In *Fundamentals of thermochemical biomass conversion* (pp.  
452 6176). Springer, Dordrecht.

453 Gregorich, E.G., Drury, C.F. and Baldock, J.A., 2001. Changes in soil carbon under long-term maize in  
454 monoculture and legume-based rotation. *Canadian journal of soil science*, 81(1), pp.21-31.

455 Hågvar, S., 1994. Soil Biology: Decomposition and soil acidity. In *Long-term experiments with*  
456 *acid rain in Norwegian forest ecosystems* (pp. 136-139). Springer, New York, NY.

457 Havstad, L.T., Aamlid, T.S. and Henriksen, T.M., 2010. Decomposition of straw from herbage  
458 seed production: Effects of species, nutrient amendment and straw placement on C and N  
459 net mineralization. *Acta Agriculturae Scandinavica Section B–Soil and Plant*  
460 *Science*, 60(1), pp.57-68.

461 Henriksen, T.M. and Breland, T.A., 2002. Carbon mineralization, fungal and bacterial growth,  
462 and enzyme activities as affected by contact between crop residues and soil. *Biology and*  
463 *Fertility of Soils*, 35(1), pp.41-48.

464 Henriksen, T.M. and Breland, T.A., 1999. Nitrogen availability effects on carbon mineralization,  
465 fungal and bacterial growth, and enzyme activities during decomposition of wheat straw in  
466 soil. *Soil Biology and Biochemistry*, 31(8), pp.1121-1134.

467 Jambu, M., 1991. *Exploratory and multivariate data analysis*. Elsevier.

468 Jha, P., Garg, N., Lakaria, B.L., Biswas, A.K. and Rao, A.S., 2012. Soil and residue carbon  
469 mineralization as affected by soil aggregate size. *Soil and Tillage Research*, 121, pp.57-62.

470 Jin, K., Sleutel, S., De Neve, S., Gabriels, D., Cai, D., Jin, J. and Hofman, G., 2008. Nitrogen  
471 and carbon mineralization of surface-applied and incorporated winter wheat and peanut  
472 residues. *Biology and Fertility of Soils*, 44(4), pp.661-665.

473 Johnson, J.M.F., Barbour, N.W. and Weyers, S.L., 2007. Chemical composition of crop biomass  
474 impacts its decomposition. *Soil Science Society of America Journal*, 71(1), pp.155-162.

475 Juan, L., Yong, H.A.N. and Zu-Cong, C.A.I., 2009. Decomposition and products of wheat and  
476 rice straw from a FACE experiment under flooded conditions. *Pedosphere*, 19(3), pp.389-  
477 397.

478 Kuzyakov, Y., 2010. Priming effects: interactions between living and dead organic matter. *Soil*  
479 *Biology and Biochemistry*, 42(9), pp.1363-1371.

480 Khalil, M.I., Hossain, M.B. and Schmidhalter, U., 2005. Carbon and nitrogen mineralization in  
481 different upland soils of the subtropics treated with organic materials. *Soil Biology and*  
482 *Biochemistry*, 37(8), pp.1507-1518.

483 Li, C., Frohling, S. and Harriss, R., 1994. Modeling carbon biogeochemistry in agricultural  
484 soils. *Global biogeochemical cycles*, 8(3), pp.237-254.

485 Li, L.J., Han, X.Z., You, M.Y., Yuan, Y.R., Ding, X.L. and Qiao, Y.F., 2013. Carbon and  
486 nitrogen mineralization patterns of two contrasting crop residues in a Mollisol: Effects of  
487 residue type and placement in soils. *European journal of soil biology*, 54, pp.1-6.

488 Lierop, W.V., 1981. Conversion of organic soil pH values measured in water, 0.01 M CaCl<sub>2</sub> or 1  
489 N KCl. *Canadian Journal of Soil Science*, 61(4), pp.577-579.

490 Lin, B. and Agyeman, S.D., 2020. Assessing Sub-Saharan Africa's low carbon development  
491 through the dynamics of energy-related carbon dioxide emissions. *Journal of Cleaner*  
492 *Production*, 274, p.122676.

493 Lou, Y., Ren, L., Li, Z., Zhang, T. and Inubushi, K., 2007. Effect of rice residues on carbon  
494 dioxide and nitrous oxide emissions from a paddy soil of subtropical China. *Water, Air, and*  
495 *Soil Pollution*, 178(1-4), pp.157-168.

496 Luxhøi, J., Magid, J., Tschirko, D. and Kandeler, E., 2002. Dynamics of invertase, xylanase and  
497 coupled quality indices of decomposing green and brown plant residues. *Soil Biology and*  
498 *Biochemistry*, 34(4), pp.501-508.



499 Machinet, G.E., Bertrand, I., Chabbert, B. and Recous, S., 2009. Decomposition in soil and  
500 chemical changes of maize roots with genetic variations affecting cell wall quality.  
501 *European journal of soil science*, 60(2), pp.176-185.

502 Machinet, G.E., Bertrand, I., Barrière, Y., Chabbert, B. and Recous, S., 2011. Impact of plant  
503 cell wall network on biodegradation in soil: role of lignin composition and phenolic acids in  
504 roots from 16 maize genotypes. *Soil biology and biochemistry*, 43(7), pp.1544-1552.

505 Magid, J., Luxhøi, J. and Lyshede, O.B., 2004. Decomposition of plant residues at low  
506 temperatures separate turnover of nitrogen and energy rich tissue components in time. *Plant  
507 and Soil*, 258(1), pp.351-365.

508 Marstorp, H. and Kirchmann, H., 1991. Carbon and nitrogen mineralization and crop uptake of  
509 nitrogen from six green manure legumes decomposing in soil. *Acta Agriculturae  
510 Scandinavica*, 41(3), pp.243-252.

511 Martens, D.A., 2000. Plant residue biochemistry regulates soil carbon cycling and carbon  
512 sequestration. *Soil Biology and Biochemistry*, 32(3), pp.361-369.

513 Mathew, I., Shimelis, H., Mutema, M. and Chaplot, V., 2017. What crop type for atmospheric  
514 carbon sequestration: Results from a global data analysis. *Agriculture, ecosystems &  
515 environment*, 243, pp.34-46.

516 Moreno-Cornejo, J., Zornoza, R. and Faz, A., 2014. Carbon and nitrogen mineralization during  
517 decomposition of crop residues in a calcareous soil. *Geoderma*, 230, pp.58-63.

518 McClelland, S.C., Paustian, K. and Schipanski, M.E., 2021. Management of cover crops in  
519 temperate climates influences soil organic carbon stocks: a meta-analysis. *Ecological  
520 Applications*, 31(3), p.e02278.

521 Muhammad, W., Vaughan, S.M., Dalal, R.C. and Menzies, N.W., 2011. Crop residues and  
522 fertilizer nitrogen influence residue decomposition and nitrous oxide emission from a  
523 Vertisol. *Biology and Fertility of Soils*, 47(1), pp.15-23.

524 Müller, T., Magid, J., Jensen, L.S. and Nielsen, N.E., 2003. Decomposition of plant residues of  
525 different quality in soil—DAISY model calibration and simulation based on experimental  
526 data. *Ecological Modelling*, 166(1-2), pp.3-18.

527 Mungai, N.W. and Motavalli, P.P., 2006. Litter quality effects on soil carbon and nitrogen  
528 dynamics in temperate alley cropping systems. *Applied Soil Ecology*, 31(1-2), pp.32-42.

529 Murungu, F.S., Chiduza, C., Muchaonyerwa, P. and Mnkeni, P.N.S., 2011. Decomposition,  
530 nitrogen and phosphorus mineralization from winter-grown cover crop residues and  
531 suitability for a smallholder farming system in South Africa. *Nutrient Cycling in*  
532 *Agroecosystems*, 89(1), pp.115-123.

533 Mutema, M., Chaplot, V., Jewitt, G., Chivenge, P. and Blöschl, G., 2015. Annual water,  
534 sediment, nutrient, and organic carbon fluxes in river basins: A global meta-analysis as a  
535 function of scale. *Water Resources Research*, 51(11), pp.8949-8972.

536 Nourbakhsh, F., 2006. Fate of carbon and nitrogen from plant residue decomposition in a  
537 calcareous soil. *Plant Soil and Environment*, 52(3), p.137.

538 Oda, T., Maksyutov, S. and Andres, R.J., 2018. The Open-source Data Inventory for  
539 Anthropogenic Carbon dioxide (CO<sub>2</sub>), version 2016 (ODIAC2016): A global, monthly  
540 fossil-fuel CO<sub>2</sub> gridded emission data product for tracer transport simulations and surface  
541 flux inversions. *Earth system science data*, 10(1), p.87.

542 Ontl, T.A. and Schulte, L.A., 2012. Soil carbon storage. *Nature Education Knowledge*, 3(10).

543 Pascault, N., Cécillon, L., Mathieu, O., Hénault, C., Sarr, A., Lévêque, J., Farcy, P.,  
544 Ranjard, L. and Maron, P.A., 2010. In situ dynamics of microbial communities during  
545 decomposition of wheat, rape, and alfalfa residues. *Microbial ecology*, 60(4), pp.816-828.

546 Paustian, K., Six, J., Elliott, E.T. and Hunt, H.W., 2000. Management options for reducing CO<sub>2</sub>  
547 emissions from agricultural soils. *Biogeochemistry*, 48(1), pp.147-163.

548 Perez, J., Munoz-Dorado, J., De la Rubia, T.D.L.R. and Martinez, J., 2002. Biodegradation and  
549 biological treatments of cellulose, hemicellulose and lignin: an overview. *International*  
550 *microbiology*, 5(2), pp.53-63.

551 Probert, M.E., Delve, R.J., Kimani, S.K. and Dimes, J.P., 2005. Modelling nitrogen  
552 mineralization from manures: representing quality aspects by varying C: N ratio of sub-  
553 pools. *Soil biology and Biochemistry*, 37(2), pp.279-287.

554 Puyuelo, B., Ponsá, S., Gea, T. and Sánchez, A., 2011. Determining C/N ratios for typical  
555 organic wastes using biodegradable fractions. *Chemosphere*, 85(4), pp.653-659.

556 Quemada, M. and Cabrera, M.L., 1995. Carbon and nitrogen mineralized from leaves and stems  
557 of four cover crops. *Soil Science Society of America Journal*, 59(2), pp.471-477.

558 Raiesi, F., 2006. Carbon and N mineralization as affected by soil cultivation and crop residue in  
559 a calcareous wetland ecosystem in Central Iran. *Agriculture, ecosystems &*  
560 *environment*, 112(1), pp.13-20.

561 Redin, M., Guénon, R., Recous, S., Schmatz, R., de Freitas, L.L., Aita, C. and Giacomini, S.J.,  
562 2014. Carbon mineralization in soil of roots from twenty crop species, as affected by their  
563 chemical composition and botanical family. *Plant and soil*, 378(1-2), pp.205-214.

564 Ritchie, H. and Roser, M., 2020. CO<sub>2</sub> and greenhouse gas emissions. *Our world in data*. Rousk,  
565 J. and Bååth, E., 2007. Fungal and bacterial growth in soil with plant materials of different  
566 C/N ratios. *FEMS microbiology ecology*, 62(3), pp.258-267.

567 Schmatz, R., Recous, S., Aita, C., Tahir, M.M., Schu, A.L., Chaves, B. and Giacomini, S.J.,  
568 2017. Crop residue quality and soil type influence the priming effect but not the fate of crop  
569 residue C. *Plant and soil*, 414(1-2), pp.229-245.

570 Shahandeh, H., Chou, C.Y., Hons, F.M. and Hussey, M.A., 2011. Nutrient partitioning and  
571 carbon and nitrogen mineralization of switchgrass plant parts. *Communications in soil*  
572 *science and plant analysis*, 42(5), pp.599-615.

573 Shakoor, A., Shakoor, S., Rehman, A., Ashraf, F., Abdullah, M., Shahzad, S.M., Farooq, T.H.,  
574 Ashraf, M., Manzoor, M.A., Altaf, M.M. and Altaf, M.A., 2021. Effect of animal manure,  
575 crop type, climate zone, and soil attributes on greenhouse gas emissions from agricultural  
576 soils—A global meta-analysis. *Journal of Cleaner Production*, 278, p.124019.

577 Shi, A., Penfold, C. and Marschner, P., 2013. Decomposition of roots and shoots of perennial  
578 grasses and annual barley—separately or in two residue mixes. *Biology and fertility of soils*,  
579 49(6), pp.673-680.

580 Stewart, C.E., Moturi, P., Follett, R.F. and Halvorson, A.D., 2015. Lignin biochemistry and soil  
581 N determine crop residue decomposition and soil priming. *Biogeochemistry*, 124(1-3),  
582 pp.335-351.

583 Scott, N.A., Cole, C.V., Elliott, E.T. and Huffman, S.A., 1996. Soil textural control on  
584 decomposition and soil organic matter dynamics. *Soil Science Society of America Journal*,  
585 60(4), pp.1102-1109.

586 Stott, D.E. and Martin, J.P., 1989. Organic matter decomposition and retention in arid soils. *Arid*  
587 *Land Research and Management*, 3(2), pp.115-148.

588 Vachon, K. and Oelbermann, M., 2011. Crop residue input and decomposition in a temperate  
589 maize-soybean intercrop system. *Soil Science*, 176(4), pp.157-163.

590 Vahdat, E., Nourbakhsh, F. and Basiri, M., 2010. Estimation of net N mineralization from short-  
591 term C evolution in a plant residue-amended soil: is the accuracy of estimation time-  
592 dependent?. *Soil use and management*, 26(3), pp.340-345.

593 Venter, Z.S., Hawkins, H.J., Cramer, M.D. and Mills, A.J., 2021. Mapping soil organic carbon  
594 stocks and trends with satellite-driven high resolution maps over South Africa. *Science of*  
595 *The Total Environment*, 771, p.145384.

596 Waksman, S.A. and Gerretsen, F.C., 1931. Influence of temperature and moisture upon the  
597 nature and extent of decomposition of plant residues by microorganisms. *Ecology*, 12(1),  
598 pp.33-60.

599 Wang, J.J., Pisani, O., Lin, L.H., Lun, O.O., Bowden, R.D., Lajtha, K., Simpson, A.J. and  
600 Simpson, M.J., 2017. Long-term litter manipulation alters soil organic matter turnover in a  
601 temperate deciduous forest. *Science of the Total Environment*, 607, pp.865-875.

602 Wang, W.J., Baldock, J.A., Dalal, R.C. and Moody, P.W., 2004. Decomposition dynamics of  
603 plant materials in relation to nitrogen availability and biochemistry determined by NMR and  
604 wet-chemical analysis. *Soil Biology and Biochemistry*, 36(12), pp.2045-2058.

605 Weiß, C.H., 2007. Statsoft, inc., tulsa, ok.: Statistica, version 8.

606 Xiao, C., 2015. *Soil organic carbon storage (sequestration) principles and management:*  
607 *Potential role for recycled organic materials in agricultural soils of Washington State.*  
608 Waste Resources Program, Washington Department of Ecology.


609 Xu, J.M., Tang, C. and Chen, Z.L., 2006. Chemical composition controls residue decomposition  
610 in soils differing in initial pH. *Soil Biology and Biochemistry*, 38(3), pp.544-552.

611 Yaowu, T.I.A.N., Ning, W.A.N.G. and Jing, L.I.U., 2016. The Priming Effect of Soil Organic  
612 Carbon Induced by Nustedge. *Journal of Nuclear Agricultural Sciences*, 30(12), p.2418.


613 Zaccheo, P., Cabassi, G., Ricca, G. and Crippa, L., 2002. Decomposition of organic residues in  
614 soil: experimental technique and spectroscopic approach. *Organic geochemistry*, 33(3),  
615 pp.327-345.

616 Zeng, D.H., Mao, R., Chang, S.X., Li, L.J. and Yang, D., 2010. Carbon mineralization of tree  
617 leaf  
618 litter and crop residues from poplar-based agroforestry systems in Northeast China: a  
619 laboratory study. *Applied soil ecology*, 44(2), pp.133-137.

620 Zhang, D., Hui, D., Luo, Y. and Zhou, G., 2008. Rates of litter decomposition in terrestrial  
621 ecosystems: global patterns and controlling factors. *Journal of Plant Ecology*, 1(2), pp.85-  
622 93.




1  
2 **Figure 1:** Global distribution map of the study sites used in the present study.




**Figure 2:** CO<sub>2</sub> emissions from residues of different crops incubated at different time periods (A: cumulative CO<sub>2</sub> emitted from crop residues to 30 days, Cr30; B: to 90 days, Cr90; C: to 120 days, Cr120; and D: 1-[Cr30/Cr120]). The data were sorted by mean (dotted line).







**Figure 3:** CO<sub>2</sub> emissions from residues of different crop types incubated at different time periods (A: C<sub>R</sub>30; B: C<sub>R</sub>90; C: C<sub>R</sub>120; and D: 1-C<sub>R</sub>30/C<sub>R</sub>120). Plots with the same lower-case letters are not significantly different at Chi-square probability of < 0.001 (Chi and H values= 0.16. 5.24 at A; 0.08. 6.72 at B; 0.00. 13.79 at C and 0.00. 21005 at D. respectively).



**Figure 4:** CO<sub>2</sub> emissions from soils of different texture ((%) -clayey, silt and sandy soil) at different day time periods. Plots with the same lower-case letters are not significantly different at Chi-square probability of < 0.001 (Chi and H values= 0.26, 2.67 at A; 0.63, 0.92 at B; 0.10, 4.70 at C; and 0.00, 14.07 at D, respectively).



**Figure 5:** CO<sub>2</sub> emissions from soils of different pH (highly basic. neutral. slightly acidic and highly acidic) at different day time periods. Strongly acidic (pH<5.4); slightly acidic (5.5-6.4); neutral (6.5-7.4); alkaline (>7.5) soil pH. Plots with the same lower-case letters are not significantly different at Chi-square probability of < 0.001 (Chi and H values= 0.01, 10.99 at A; 0.01, 12.64 at B, 0.25, 4.09 at C, and 0.00, 20.92 at D, respectively).



**Figure 6:** CO<sub>2</sub> emissions from climate of different rainfall and temperature level (as in tropical, sub-tropical and temperate (MAT, MAP) at different day time periods. Plots with the same lower-case letters are not significantly different at Chi-square probability of < 0.001 (Chi and H values= 0.84, 0.34 at A; 0.11, 4.40 at B; 0.68, 0.77 at C, and 0.00, 18.13 at D, respectively).

1 **Table 1:** References included in database with author, country, crops type, soil texture and  
 2 climatic zones under which the studies were conducted.

No.	Author	Country	Crop type	Soil texture	Climate
1	Abiven et al. (2005)	Brazil	Brachiaria; Rice; Sorghum; Soyabean; Wheat	Clayey	Tropical
2	Abiven and Recous (2007)	Brazil	Brachiaria; Rice; Sorghum; Soyabean; Wheat	Clayey	Tropical
3	Abro et al. (2011)	China	Maize	Clayey	Subtropical
4	Ajwa and Tabatabai (1994)	USA	Alfalfa; Maize; Sorghum; Soyabean	Silt	Subtropical
5	Angers and Recous (1997)	France	Rye; Wheat	Silt	Temperate
6	Anguria et al. (2017)	Uganda	Cowpea; groundnut; millet; sorghum	(blank)	Subtropical
7	Arunachalam et al. (2003)	India	Bean and pea	Sandy	Tropical
8	Aulakh et al. (1991)	USA	Vetch and wheat	Silt	Tropical
9	Begum et al. (2014)	Australia	Canola; Sorghum; Soyabean; Wheat	Clayey	Subtropical
10	Bertrand, et al. (2006)	France	Wheat	Silt	Subtropical
11	Blaise & Bhaskar (2003)	India	Cotton	clayey,	Subtropical
12	Cayuela et al. (2009)	Italy	Cotton; wheat	Sandy	Subtropical
13	Clark et al. (2007)	Australia	Alfalfa; Wheat	Clayey	Subtropical
14	Cong et al. (2015)	China	Bean; Maize; Wheat	Sandy	Temperate
15	Corbeels et al. (2000)	Morocco	Sunflower; Wheat	Clayey	Temperate
16	Curtin et al. (2008)	New Zealand	Barley; Wheat	Silt	Subtropical
17	Datta et al. (2019)	India	Maize; Rice; Wheat	Silt	Subtropical
18	De Neergaard et al. (2002)	Denmark	Clover; Grass	Silt	Temperate
19	Duong et al. (2009)	Australia	Wheat	Sandy	Subtropical
20	Finn et al. (2015)	Australia	Alfalfa; Grass; Wheat	clayey, Silt& sandy	Subtropical
21	Fruit et al. (1999)	France	Wheat	Silt	Subtropical
22	Gezahegn et al. (2016)	Malaysia	Maize; Soyabean	Sandy	Subtropical
23	Ghimire et al. (2017)	USA	Canola; Oat; Pea	Clayey	Subtropical
24	Havstad et al. (2010)	Norway	Barley; Clover; Grass; Meadow; Wheat	Silt	Subtropical
25	Henriksen and Breland (2002)	Norway	Barley; Clover; Wheat	Silt, sandy	Temperate
26	Jha et al. (2012)	India	Wheat	Clayey	Subtropical
27	Jin et al. (2008)	China	Peanut; Wheat	Silt	Tropical
28	Johnson et al. (2017)	USA	Alfalfa; Cuphea; Maize; Soyabean; Switchgrass	Silt	Subtropical
29	Juan et al. (2009)	China	Rice; Wheat	(blank)	Subtropical
30	Khalil et al. (2005)	Bangladesh	Bean; wheat	clayey, Silt &sandy	Tropical
31	Li et al. (2013)	China	Maize; Soyabean	Silt	Temperate
32	Lou et al. (2007)	China	Rice	Clayey	Tropical
33	Luxhoi et al. (2002)	Denmark	Clover; Grass; Rye	Sandy	Tropical
34	Machinet et al. (2009)	France	Maize	Silt	Temperate
35	Machinet et al. (2011)	France	Maize	Clayey	Temperate
36	Magid et al. (2004)	Denmark	Clover; Radish; Rye; Sugarcane; Vetch	Sandy	Temperate
37	Marstorp and kirchmann (1991)	Sweden	Clover	Sandy	Temperate
38	Martens (2000)	USA	Alfalfa; Canola; Maize; Oat; Prairie	Sandy	Subtropical
39	Moreno-Cornejo et al. (2014)	Spain	Pepper	Silt	Subtropical
40	Muhammad et al. (2011)	Australia	Cotton; Maize; Sorghum, Sugarcane	Clayey	Subtropical
41	Müller et al. (2003)	Denmark	Barley; Clover; Grass; Rape	Sandy	Temperate
42	Mungai and Motavalli (2006)	Kenya	Grass; Maize; Soyabean	clayey, Silt	Temperate
43	Murungu et al. (2011)	South Africa	Oat; Peas; Vetch	Sandy	Temperate
44	Nourbakhsh (2006)	Iran	Alfalfa; Wheat	Clayey	Subtropical
45	Pascual et al. (2010)	France	Alfalfa; Rape; Wheat	Clayey	Temperate
46	Quemada and Cabrera, (1995)	USA	Clover; Oat; Rye; Wheat	Sandy	Subtropical
47	Raiesi, (20065)	Iran	Alfalfa; Wheat	Clayey	Subtropical
48	Redin et al. (2014)	Brazil	Maize; Sorghum; Wheat, Soyabean; Sunflower; Vetch	Sandy	Tropical
49	Schmatz et al. (2017)	Brazil	Pea; Vetch; Wheat	clayey, sandy	Tropical
50	Shahande et al. (2011)	USA	Switchgrass	Silt	Subtropical
51	Shi et al. (2013)	Australia	Barley; Grass	Silt	Subtropical
52	Stewart et al. (20157)	USA	Maize; Sorghum; Soyabean; Sunflower; Wheat	Silt	Temperate
53	Vachon and Oelbermann (2011)	Argentina	Maize; Soyabean	Silt	Subtropical
54	Vahdat et al; (2010)	Iran	Alfalfa; Barley; Clover; Grass; Wheat	Clayey	Subtropical
55	Wang et al. (2004)	Australia	Brigalow; Grass; Sugarcane; Wheat	Sandy	Subtropical
56	Xu et al. (2006)	Australia	Alfalfa; Chickpea; Medic; Wheat	Sandy	Tropical
57	Zaccheo et al. (2002)	Italy	Alfalfa; Maize	Sandy	Temperate
58	Zeng et al. (2010)	China	Maize; Peanut; Poplar	Sandy	Temperate

**Table 3:** Crop quality, soils and environmental variables classification used in the study.

Variable	Variable remarks	Category	Symbol	Class	Ref.
<b>Crop class</b>	All different crop types categorised into four classes	Wheat, sorghum,	Crop class	Cereal,	Mathew, et al. (2017)
		grass,		Grass,	
		vetch bean, soyabean		Legume and	
		Canola, cotton		Oilseed	
<b>Residue lignin concentration (%)</b>	Initial lignin concentration of residues	<10	Lignin	Low	
		>10		High	
<b>Residue lignin: N ratio</b>	Residue lignin to nitrogen ratio	>10	lignin: N	High	
		<10		Low	
<b>Residue C:N ratio</b>	Crop residue carbon to nitrogen ratio	<20	C: N	Low	Puyuelo, (2011) and Probert (2005)
		20-30		Medium	
		>30		High	
<b>Residue N concentration (%)</b>	Initial nitrogen concentration of residues	>10	TN	High	
		<10		Low	
<b>Residue C concentration (%)</b>	Initial carbon concentration of residues	>50	TC	High	Abdalla et al., (2016)
		<50		Low	
<b>Clay concentration (%)</b>	Soil Texture based on clay fraction	>10	Soil texture	Clayey	Mathew et al., (2017) and Abdalla et al., (2016)
		20-32		Silt	
		<20		Sandy	
<b>Soil pH concentration (CaCl<sub>2</sub>)</b>	Soil pH	<5.4	pH	Strongly acidic	Davies, 1971; Mathew et al. (2017)
		5.5-6.4		Slightly acidic	
		6.5-7.4		Neutral	
		>7.5		Alkaline	
<b>Soil bulk density (g cm<sup>-3</sup>)</b>	Average bulk density in soil profile	<1.3	BD	Low BD	Mathew et al., (2017)
		>1.3		High BD	
<b>CEC (cmol<sub>c</sub>kg)</b>	Soil cation exchange capacity	>20	CEC	Low CEC	
		<20		High CEC	
<b>Climate</b>	Based on the average annual temperature and precipitation of the study site	>20	MAT	Tropical	Mutema et al., (2015) and Mathew et al., 2017
		>1500	MAP		
		20-10	MAT	Sub-tropical	
		100-1110	MAP		
		<10	MAT	Temperate	
		120-1000	MAP		

**Table 4:** Summary Statistics of plant, soil and environmental variables along with residue CO<sub>2</sub> emissions in different time period.

	Lignin	C:N	TC	TN	Clay	Sand	Silt	pH	SOC	MAP	MAT	C <sub>R</sub> 30	C <sub>R</sub> 90	C <sub>R</sub> 120	1- [C <sub>R</sub> 30/C <sub>R</sub> 120]
	%				-----g kg <sup>-1</sup> -----				%	mm	°C	-----mg CO <sub>2</sub> -C g <sup>-1</sup> C -----			
N	192	304	394	365	295	259	243	333	293	386	386	394	394	394	394
Mean	9.55	58.45	13.24	9.38	29.53	37.81	32.40	6.73	13.13	906.16	17.01	196.5	519.0	914.2	0.80
Median	7.70	39.16	1.15	4.00	25.30	32.00	28.00	6.80	9.50	656.00	17.00	138.0	415.6	795.0	0.80
Min.	0.02	7.80	0.04	0.01	2.00	1.00	1.00	3.87	0.50	89.40	3.90	0.3	1.1	3.2	0.49
Max.	29.40	409.00	165.76	407.00	77.30	96.00	83.00	8.60	39.00	2500.00	30.00	920.1	3205	3640	0.98
Q1	5.35	20.30	0.40	0.90	14.30	17.90	16.30	5.60	8.41	551.00	10.00	31.0	118.4	200.4	0.75
Q3	12.55	75.40	4.12	11.00	40.00	60.00	47.00	7.66	17.40	1095.00	24.00	304.9	858.9	1431.0	0.86
Variance	36.07	3257.25	1164.81	592.74	370.64	737.14	483.34	1.34	65.33	263814.21	46.19	37353.5	213397	585331	0.01
SD.	6.01	57.07	34.13	24.35	19.25	27.15	21.99	1.16	8.08	513.63	6.80	193.3	461.9	765.1	0.07
SE.	0.43	3.27	1.72	1.27	1.12	1.69	1.41	0.06	0.47	26.14	0.35	7.7	18.4	30.5	0.00
CV	63.30	98.10	13.70	262.70	64.90	75.40	67.60	17.30	61.60	56.50	40.30	98.4	89.0	83.7	9.11
Skewness	1.06	2.52	2.97	12.47	0.92	0.51	0.68	-0.30	1.04	1.15	0.10	1.07	1.10	0.73	-0.25
Kurtosis	0.83	8.30	7.69	197.12	0.13	-0.79	-0.32	-1.03	0.69	0.51	-1.12	0.58	2.11	-0.15	-0.23

Statistics: Min and Max =minimum and maximum, respectively. Q1 and Q3= first and third quartile, SD = standard deviation. C emission variables: C<sub>R</sub>30; C<sub>R</sub>90; C<sub>R</sub>120 for cumulative residue CO<sub>2</sub> emissions to day 30, 90 and 120. Crop quality: lignin; C: N ratio; TC= total residue carbon; TN= total residue nitrogen. Soil variables: Clay concentration; sand concentration; silt concentration; soil pH (CaCl). SOC= soil organic carbon concentration; Climatic variables: MAP= mean annual precipitation; MAT= mean annual temperature. “†” values are not zero. but rounded off to one decimal place.

**Table 5:** Sample sizes of crop quality (n), soil and climatic factor categories in association with carbon emissions from residues.

		C <sub>R</sub> 30			C <sub>R</sub> 90			C <sub>R</sub> 120			1-[C <sub>R</sub> 30/C <sub>R</sub> 120]		
		n	Mean	STDEV	n	Mean	STDEV	n	Mean	STDEV	n	Mean	STDEV
Overall		394	196.3	193.3	394	518.8	461.9	394	914.2	765.1	394	0.81	0.07
Crop type	Cereal	195	171.0	185.9	195	473.9	445.4	195	846.6	777.1	195	0.82	0.07
	Grass	54	217.0	185.8	54	529.7	426.7	54	946.8	741.0	54	0.79	0.06
	Legume	120	228.0	203.2	120	586.7	505.7	120	1003.0	758.3	120	0.78	0.08
	Oilseed	25	196.0	211.9	25	519.2	459.2	25	944.0	742.5	25	0.83	0.07
Residue lignin Concentration	High	73	173.0	165.2	73	457.0	402.3	73	816.9	694.2	73	0.80	0.07
	Low	119	218.0	188.9	119	602.3	524.3	119	1055.0	853.0	119	0.84	0.05
Residue C: N Ratio	High	142	144.9	159.0	142	447.0	424.9	142	837.9	777.4	142	0.78	0.07
	Low	103	273.7	223.4	103	686.9	537.4	103	1163.1	814.7	103	0.80	0.07
	Medium	59	201.1	214.7	59	484.9	449.7	59	856.8	734.8	59	0.80	0.08
TC	High	284	201.8	198.7	284	524.6	450.5	284	927.3	759.9	284	0.80	0.08
	Low	110	181.9	168.9	110	503.6	635.6	110	880.2	803.9	110	0.82	0.07
Soil texture	Clayey	121	224.4	179.0	121	587.5	440.7	121	1060.6	777.5	121	0.81	0.07
	Silt	130	191.1	215.6	130	505.8	523.9	130	894.3	819.4	130	0.82	0.08
	Sandy	137	178.2	184.7	137	454.7	417.8	137	789.9	699.2	137	0.79	0.08
Soil pH	Alkaline	133	178.1	170.0	133	487.2	410.7	133	869.2	712.5	133	0.82	0.07
	Neutral	57	195.4	165.4	57	532.3	362.0	57	919.4	613.8	57	0.79	0.07
	Slightly acidic	86	262.1	203.3	86	634.6	560.3	86	1140.2	874.0	86	0.78	0.07
	Strongly acidic	57	175.3	222.3	57	414.7	443.4	57	709.9	687.7	57	0.79	0.09
Soil organic carbon Concentration	High	39	149.4	148.9	39	356.7	529.6	39	591.4	695.8	39	0.81	0.07
	Low	153	157.5	188.9	153	421.3	430.9	153	746.9	725.8	153	0.81	0.07
	Medium	101	230.3	184.9	101	595.2	432.0	101	1066.4	746.6	101	0.81	0.07
Climate	Subtropical	198	192.5	181.8	198	522.8	427.3	198	912.2	736.5	198	0.81	0.07
	Temperate	101	149.8	213.8	101	430.3	536.5	101	779.7	828.1	101	0.82	0.08
	Tropical	95	253.4	193.7	95	604.5	444.7	95	1061.2	755.8	95	0.78	0.08



**Table 6:** Sample sizes of crop type categories associated with residue carbon emission variables.

Crop type	Crop	C <sub>R</sub> 30			C <sub>R</sub> 90			C <sub>R</sub> 120			1-[C <sub>R</sub> 30/C <sub>R</sub> 120]		
		n	Mean	STDEV	n	Mean	STDEV	n	Mean	STDEV	n	Mean	STDEV
	Overall	394	196.3	193.3	394	518.8	461.95	394	914.2	765.07	394	0.81	0.07
Cereal	Barley	9	177.7	156.8	9	547.2	436.1	9	1005	778.6	9	0.84	0.05
	Maize	59	85.8	124.3	59	275.1	319.8	59	495.0	541.7	59	0.84	0.08
	Oat	8	212.1	190.8	8	501.9	425.5	8	872.7	724.5	8	0.79	0.06
	Rice	10	251.9	255.7	10	657.1	662.3	10	1211	1196.6	10	0.82	0.08
	Rye	12	199.3	190.8	12	501.6	453.5	12	824.2	729.5	12	0.77	0.04
	Sorghum	16	261.1	218.8	16	714	496.7	16	1303	855.9	16	0.83	0.08
	Wheat	81	196.3	194.8	81	533.6	452.3	81	950.7	791.8	81	0.81	0.07
Grass	Grass	54	217	184.9	54	529.7	420.3	54	946.8	734.2	54	0.79	0.06
Legume	Alfalfa	24	359.0	262.7	24	796.0	515.0	24	1319.0	801.0	24	0.76	0.07
	Bean	12	223.7	239.9	12	538.7	471.0	12	962.1	667.0	12	0.80	0.06
	Clover	26	184.2	139.6	26	580.6	661.0	26	940.3	866.5	26	0.78	0.06
	Pea	27	184.2	152.7	27	452	343.8	27	809	554.1	27	0.78	0.07
	Soyabean	23	207.8	157.0	23	577.5	401.7	23	1039	715.6	23	0.81	0.10
	Vetch	8	190.2	182.6	8	531.1	433.4	8	870.4	696.1	8	0.80	0.05
Oilseed	Canola	8	293.5	327.3	8	670.5	717.3	8	1211	1184.5	8	0.79	0.07
	Cotton	9	135.5	98.8	9	442.5	281.0	9	841.1	513.9	9	0.85	0.03
	Sunflower	8	166.5	208.7	8	454.2	451.7	8	792.5	667.4	8	0.84	0.10

C<sub>R</sub>30, C<sub>R</sub>90 and C<sub>R</sub>120=Amount of CO<sub>2</sub> emitted from crop residues to day 30, 90 and 120 of the experiment, respectively.

**Table 7:** Correlation matrix statistic table of plant, soil and climatic influence on residue CO<sub>2</sub> emissions.

	Lignin	C:N	TC	TN	Clay	Sand	Silt	pH	SOC	MAP	MAT	CR30	CR90	CR120	$\frac{1-}{[CR30/CR120]}$
Lignin	1.00														
C:N	0.15	1.00													
TC	-0.06	-0.06	1.00												
TN	0.03	-0.19	-0.02	1.00											
Clay	-0.08	-0.11	<b>0.51*</b>	-0.19	1.00										
Sand	-0.02	-0.15	<b>-0.42</b>	<b>0.32*</b>	<b>-0.64*</b>	1.00									
Silt	0.10	<b>0.30*</b>	-0.02	-0.19	<b>-0.26*</b>	<b>-0.57*</b>	1.00								
pH	0.02	0.15	0.16	<b>-0.63*</b>	<b>0.26*</b>	<b>-0.45*</b>	<b>0.29*</b>	1.00							
SOC	-0.03	0.03	<b>0.23*</b>	-0.10	<b>0.31*</b>	<b>-0.37*</b>	0.14	-0.08	1.00						
MAP	-0.02	-0.03	-0.16	0.19	<b>-0.41*</b>	<b>0.4*</b>	-0.07	<b>-0.58*</b>	-0.20	1.00					
MAT	0.00	-0.20	-0.21	0.16	<b>-0.40*</b>	<b>0.41*</b>	-0.08	<b>-0.40*</b>	0.03	<b>0.43*</b>	1.00				
CR30	<b>-0.08*</b>	<b>-0.14*</b>	<b>0.59*</b>	-0.22	<b>0.34*</b>	<b>-0.38*</b>	0.12	<b>0.38*</b>	0.15	<b>-0.31*</b>	0.07	1.00			
CR90	<b>-0.11*</b>	<b>-0.10*</b>	<b>0.46*</b>	<b>-0.28*</b>	<b>0.26*</b>	<b>-0.32*</b>	0.13	<b>0.44*</b>	0.09	<b>-0.32*</b>	0.10	<b>0.96*</b>	1.00		
CR120	<b>-0.12*</b>	<b>-0.09*</b>	<b>0.40*</b>	<b>-0.26*</b>	0.22	<b>-0.29*</b>	0.14	<b>0.42*</b>	0.07	<b>-0.31*</b>	0.10	<b>0.93*</b>	<b>0.99*</b>	1.00	
$\frac{1-}{[CR30/CR120]}$	0.15	<b>-0.07*</b>	<b>0.37*</b>	0.11	0.10	-0.12	0.05	-0.19	<b>0.27*</b>	0.19	0.21	<b>0.27*</b>	0.07	-0.03	1.00

Residue CO<sub>2</sub> emission to day 30, 90 and 120 of the experiment. lignin: residue lignin concentration; C:N: residue carbon to nitrogen ratio; TC: Initial residue carbon concentration; Clay, sand & silt: soil texture based on clay fraction (%clay); pH: soil pH (KCl); SOC: soil organic carbon concentration; MAP&MAT: climatic factors-mean annual precipitation and mean annual temperature. \* Significant at p<0.05.

**Table 8:** Proportion of data points with “priming”, i.e. with a 120 days cumulative C-CO<sub>2</sub> emissions above the amount of C added to the soil.

<b>Crop</b>	<b>%</b>
Alfalfa	58
Barley	44
Bean	25
Canola	63
Clover	27
Cotton	44
Grass	48
Maize	25
Oat	50
Pea	33
Rice	50
Rye	58
Sorghum	56
Soyabean	48
Sunflower	25
Vetch	38
Wheat	36
<i>Mean</i>	<b>43</b>