



HAL
open science

Variability of fungi detection in the respiratory tract of racehorses according to the sampling site

Louise C Lemonnier, Anne Couroucé, Margot Cessans, Ludovic Petit, Jacqueline Martina Cardwell, Marie-Pierre Toquet, Eric Richard

► To cite this version:

Louise C Lemonnier, Anne Couroucé, Margot Cessans, Ludovic Petit, Jacqueline Martina Cardwell, et al.. Variability of fungi detection in the respiratory tract of racehorses according to the sampling site. 11th International Conference on Equine Exercise Physiology (ICEEP11), Jun 2022, Uppsala, Sweden. , 18 (Supp 1). hal-03727163

HAL Id: hal-03727163

<https://hal.science/hal-03727163v1>

Submitted on 19 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Variability of fungi detection in the respiratory tract of racehorses according to the sampling site

Louise C. LEMONNIER¹, A. COUROUCE^{1,2}, M. CESSANS¹, L. PETIT³, J.M. CARDWELL⁴, M.-P. TOQUET³, E.A. RICHARD^{2,3}

¹Department of Equine Internal Medicine, University Animal Hospital, CISCO-Oniris, Route de Gachet, 44300 Nantes, France

²BIOTARGEN, Normandie Univ, UNICAEN, 14000 Caen, France

³Labeo-Frank Duncombe Laboratory, 1 route de Rosel, 14053 Caen, France

⁴Department of Pathobiology and Population Sciences, Royal Veterinary College, Hatfield, Hertfordshire, UK

INTRODUCTION

- The clinical relevance of colonization of equine airways by fungi is unclear.
- Diagnosis of fungi in the respiratory tract is fastidious and there is no method of reference.

OBJECTIVES

- To characterize fungi from upper and lower airways of racehorses
- To determine the most reliable methodology for fungal detection

HYPOTHESIS

- Fungal cytology of a pooled BAL (BAL_P) is representative of corresponding left (BAL_L) and right (BAL_R) samples.

METHODS

- 30 Standardbred racehorses in active training, aged 4.6 ± 1.6 years, from 3 different training stables
- TW, followed by BAL from each lung separately, collected through a guarded catheter and gastroscope

- Fungi detection, identification and characterization by cytology and culture
 - Positive culture: ≥ 1 mold identified
 - Positive cytology: ≥ 1 fungal element (hyphae, spore) observed

RESULTS

- Fungal culture (Table 1)
 - TW: 28/30 (93%) positive
 - BAL_P: 4/30 (13%) positive → All positive in TW, 3/4 with same fungi
 - 9 genus (alone or concomitant) : *Aspergillus* (5 genus) 13/28 (46%), *Penicillium* 12/28 (43%)

Table 1: Identification of fungi in horses with a positive BAL_P culture

Case number	9		17		18		21	
	TW	BAL _P	TW	BAL _P	TW	BAL _P	TW	BAL _P
<i>A. fumigatus</i>								x
<i>A. nidulans</i>	x		x	x	x			
<i>A. versicolor</i>								x
<i>Penicillium</i>	x		x		x	x	x	x
<i>Aureobasidium</i>		x						
<i>Chrysosporum</i>	x							

Table 2: Cytology of pooled and individual BALF, and of TW

	Cytology positive	Spores	Hyphae	Phagocyted
TW	22/30	18/22	16/22	14/22
BAL _P	7/30 3 bilateral 4 unilateral	7/7	2/7	6/7
BAL _R	8/30	5/8	6/8	5/8
BAL _L	7/30	7/7	0/7	6/7

Comparison of sample sites (Table 3)

- Poor agreement between BAL_L and BAL_R
- Poor agreement between culture and cytology
- BAL_P vs separate lungs:
 - Not equivalent (p = 0.025 by McNemar Test)
 - Sensitivity = 0,58 ; Specificity = 1,00

Fungal cytology (Table 2)

- TW: 22/30 (73%) positive ; 12/22 (55%) with both spores and hyphae, frequently phagocyted
- BAL: Large disparity between lungs

Table 3: Comparison of BALF sample sites, and of BALF methods of analysis

	BAL _L only	BAL _R only	BAL _{L+R}	Agreement (κ)
Cytology	4/7	5/8	3	0,015
	Culture only	Cytology only	Culture + cytology	Agreement (κ)
BAL _P	3/4	6/7	1/7	0,015

DISCUSSION

- Despite a similar prevalence of fungi detection in left and right BAL, a large disparity was found between lungs on cytology.
- Detecting fungi by cytology on either BAL_P or BAL_L + BAL_R was not equivalent

CONCLUSION

Pooled BAL is a highly specific while fairly sensitive alternative to combined cytology of individual left/right samples for mold/fungi detection.