



HAL
open science

Inhomogeneous Poisson processes in the disk and interpolation

Andreas Hartmann, Xavier Massaneda

► **To cite this version:**

Andreas Hartmann, Xavier Massaneda. Inhomogeneous Poisson processes in the disk and interpolation. Proceedings of the Edinburgh Mathematical Society, 2024, 67 (3), pp.749-777. <hal-03728142>

HAL Id: hal-03728142

<https://hal.science/hal-03728142v1>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

INHOMOGENEOUS POISSON PROCESSES IN THE DISK AND INTERPOLATION

ANDREAS HARTMANN, XAVIER MASSANEDA

ABSTRACT. We investigate different geometrical properties of the inhomogeneous Poisson point process Λ_μ associated to a positive, locally finite, σ -finite measure μ on the unit disk. In particular, we characterize the processes Λ_μ such that almost surely: 1) Λ_μ is a Carleson-Newman sequence; 2) Λ_μ is the union of a given number M of separated sequences. We use these results to discuss the measures μ such that the associated process Λ_μ is almost surely an interpolating sequence for the Hardy, Bloch or weighted Dirichlet spaces.

1. INTRODUCTION AND MAIN RESULTS

Important notions in spaces of analytic functions include zero-sets, Carleson measures, interpolation, sampling, frames, etc. Such properties have been studied for many well-known spaces of analytic functions in a deterministic setting. A canonical example is the Hardy space, where all these properties are well established, see [16]. In other spaces such properties admit theoretical characterizations which are not checkable in general (e.g. interpolation in Dirichlet spaces), see e.g. [24] for a general reference. There also exist situations where a general characterization is not available. In these circumstances it is useful to consider a random setting, which allows to see whether certain properties are “generic” in a sense. The random model we are interested in here is the Poisson point process.

A *Poisson point process* in the unit disk \mathbb{D} is a random sequence Λ defined in the following way: for any Borel set $A \subset \mathbb{D}$ the counting random variable $N_A = \#(A \cap \Lambda)$ is well defined and

- (a) N_A is a Poisson random variable, i.e., there exists $\mu(A) \geq 0$ such that the probability distribution of N_A is

$$\mathbb{P}(N_A = k) = e^{-\mu(A)} \frac{(\mu(A))^k}{k!}, k \geq 0.$$

In particular $\mathbb{E}[N_A] = \text{Var}[N_A] = \mu(A)$.

- (b) If $A, B \subset \mathbb{D}$ are disjoint Borel sets then the variables N_A, N_B are independent.

Date: July 20, 2022.

2010 Mathematics Subject Classification. 30E05, 30H10, 30H25, 30H30, 60G55.

Key words and phrases. Poisson point process, Carleson measure, separation, interpolation, spaces of holomorphic functions.

The first author partially supported by the project REPKA (ANR-18-CE40-0035). Second author partially supported by the Generalitat de Catalunya (grant 2017 SGR 359) and the spanish Ministerio de Ciencia e Innovación (project PID2021-123405NB-I00).

It turns out that these two properties uniquely characterize the point process. Also, the values $\mu(A)$ define a σ -finite Borel measure on \mathbb{D} , which is called the *intensity* of the process.

The Poisson process is a well-known statistical model for point distributions with no (or weak) interactions, and it has multiple applications in a great variety of fields [26]. Because of property (b), it is clearly not adequate to describe distributions in which each point is not statistically independent of the other points of the process. For such situations other models have been proposed (e.g. determinantal processes or zeros of Gaussian analytic functions for random sequences with repulsion, or Cox processes for situations with positive correlations and clumping [18]).

It is also possible to create a Poisson process from a given, σ -finite, locally finite, positive Borel measure μ in \mathbb{D} , in the sense that there exists a point process Λ_μ with intensity μ , i.e, whose counting functions satisfy properties (a) and (b) above. This is a well-known, non-trivial fact that can be found, for example, in [19, Theorem 3.6]. Such a Poisson process Λ_μ is sometimes called *inhomogeneous*, or *non-stationary*.

In this paper, given a positive Borel measure μ on \mathbb{D} , we study elementary geometric properties of the inhomogeneous Poisson process of intensity μ , specifically in relation to conditions used to describe interpolating sequences for various spaces of analytic functions in \mathbb{D} . We shall always assume that $\mu(\mathbb{D}) = +\infty$, since otherwise Λ_μ would be finite almost surely.

The probabilistic point of view has already been explored before in connection with interpolation. Here we mention Cochran [10] and Rudowicz [22] who considered the probabilistic model $\Lambda = \{r_n e^{i\theta_n}\}_n$ in which the radii $r_n \subset (0, 1)$ are fixed a priori and the arguments θ_n are chosen uniformly and independently in $[0, 2\pi]$ (a so-called Steinhaus sequence). For this model they established a zero-one condition on $\{r_n\}_n$ so that the resulting random sequence is almost surely interpolating for the Hardy spaces. In [9] similar results, for the same probabilistic model, were proven for the scale of weighted Dirichlet spaces between the Hardy space and the classical Dirichlet space. See also [13] for related results in the unit ball and the polydisk.

We express our results in terms of a dyadic discretization of μ . Consider first the dyadic annuli

$$A_n = \{z \in \mathbb{D} : 2^{-(n+1)} < 1 - |z| \leq 2^{-n}\}, \quad n \geq 0.$$

Each A_n can be split into 2^n boxes of the same size 2^{-n} :

$$T_{n,k} = \left\{ z = r e^{it} \in A_n : \frac{k}{2^n} \leq \frac{t}{2\pi} < \frac{k+1}{2^n} \right\}, \quad k = 0, 1, \dots, 2^n - 1.$$

These boxes can be viewed as the top halves of the Carleson windows

$$Q(I_{n,k}) = \left\{ z = r e^{i\theta} \in \mathbb{D} : r > 1 - 2^{-n}, e^{i\theta} \in I_{n,k} \right\}$$

associated to the dyadic intervals

$$(1) \quad I_{n,k} = \left\{ e^{it} \in \mathbb{T} : \frac{k}{2^n} \leq \frac{t}{2\pi} < \frac{k+1}{2^n} \right\}, \quad n \geq 0, k = 0, 1, \dots, 2^n - 1.$$




FIGURE 1. Carleson window $Q(I_{n,k})$ associated to the dyadic interval $I_{n,k}$ and its top half $T_{n,k}$.

Denote $X_{n,k} = N_{T_{n,k}}$, which by hypothesis is a Poisson random variable of parameter

$$\mu_{n,k} := \mathbb{E}[X_{n,k}] = \text{Var}[X_{n,k}] = \mu(T_{n,k}).$$

In these terms, the assumption $\mu(\mathbb{D}) = +\infty$ is just

$$\mu(\mathbb{D}) = \sum_{n \in \mathbb{N}} \sum_{k=0}^{2^n-1} \mu_{n,k} = \sum_{n,k} \mu_{n,k} = +\infty.$$

A first geometric property on random sequences we are interested in is separation. For this, we recall that the pseudo-hyperbolic distance in \mathbb{D} is given by

$$\rho(z, w) = \left| \frac{z - w}{1 - \bar{w}z} \right| \quad z, w \in \mathbb{D}.$$

Definition 1.1. A sequence $\Lambda = \{\lambda_k\}_{k \geq 1} \subset \mathbb{D}$ is *separated* if there exists $\delta > 0$ such that

$$\rho(\lambda_k, \lambda_l) \geq \delta, \quad k \neq l.$$

When we need to specify the separation constant we say that Λ is δ -separated.

We are now in a position to state our result characterizing those Λ_μ which can (almost surely) be expressed as finite unions of separated sequences.

Theorem 1.2. *Let Λ_μ be the Poisson process associated to a positive, σ -finite, locally finite measure μ and let $M \geq 1$ be an integer. Then*

$$\mathbb{P}(\Lambda_\mu \text{ union of } M \text{ separated sequences}) = \begin{cases} 1 & \text{if } \sum_{n,k} \mu_{n,k}^{M+1} < \infty \\ 0 & \text{if } \sum_{n,k} \mu_{n,k}^{M+1} = \infty. \end{cases}$$

In particular,

$$\mathbb{P}(\Lambda_\mu \text{ separated}) = \begin{cases} 1 & \text{if } \sum_{n,k} \mu_{n,k}^2 < \infty \\ 0 & \text{if } \sum_{n,k} \mu_{n,k}^2 = \infty. \end{cases}$$

The characterization of a.s. separated sequences was first obtained, with a different proof, in [2, Teorema 3.2.1].

Our second result deals with so-called α -Carleson sequences. Given any arc $I \subset \mathbb{T} = \partial\mathbb{D}$ let $|I|$ denote its normalized length and consider the associated Carleson window

$$Q(I) = \{z = re^{i\theta} \in \mathbb{D} : r > 1 - |I|, e^{i\theta} \in I\}.$$

Definition 1.3. Let $\alpha \in (0, 1]$. The sequence Λ satisfies the α -Carleson condition if there exists $C > 0$ such that for all arcs $I \subset \mathbb{T}$

$$\sum_{\lambda \in Q(I)} (1 - |\lambda|)^\alpha \leq C|I|^\alpha.$$

Such sequences will also be called α -Carleson sequences.

Observe that to check the α -Carleson condition it is enough to test on the dyadic intervals $I_{n,k}$ given in (1).

The sequences Λ satisfying the 1-Carleson condition are by far the most studied, because of their rôle in the famous characterization of the interpolating sequences for the algebra H^∞ of bounded holomorphic functions, given by L. Carleson [8] (see Section 3). They are sometimes found in the literature under the name of *Carleson-Newman* sequences.

The α -Carleson property above is a special case of a more general condition: a finite, positive Borel measure σ on \mathbb{D} is a Carleson-measure of order $\alpha \in (0, 1]$ if $\sigma(Q(I)) \leq C|I|^\alpha$ for some $C > 0$ and all intervals I . As shown by L. Carleson (see e.g. [16]), Carleson measures (of order 1) are precisely those for which the embedding $H^2 \subset L^2(\mathbb{D}, \sigma)$ holds; here H^2 is the classical Hardy space (see the definition in Subsection 3.1 below). Carleson measures of order $\alpha < 1$ have been used, for example, in providing sufficient conditions for solvability of the $\bar{\partial}_b$ -equation in L^p , $L^{p,\infty}$ and in Lipschitz spaces of the boundary of strictly pseudoconvex domains [3].

Theorem 1.4. Let Λ_μ be the Poisson process associated to a positive, σ -finite, locally finite measure μ . Then

(a)

$$\mathbb{P}(\Lambda_\mu \text{ is a 1-Carleson sequence}) = \begin{cases} 1 & \text{if there exists } \gamma > 1 \text{ such that } \sum_{n,k} \mu_{n,k}^\gamma < \infty \\ 0 & \text{if for all } \gamma > 1 \quad \sum_{n,k} \mu_{n,k}^\gamma = \infty. \end{cases}$$

(b) Let $\alpha \in (0, 1)$. If there exists $1 < \gamma < \frac{1}{1-\alpha}$ such that $\sum_{n,k} \mu_{n,k}^\gamma < +\infty$, then

$$\mathbb{P}(\Lambda_\mu \text{ is } \alpha\text{-Carleson}) = 1$$

(c) There exists a positive, σ -finite, locally finite measure μ such that $\sum_{n,k} \mu_{n,k}^{1/(1-\alpha)} < +\infty$ and

$$\mathbb{P}(\Lambda_\mu \text{ is } \alpha\text{-Carleson}) = 0.$$

(d) For every $\gamma > 1$ there exists a positive, σ -finite, locally finite measure μ such that $\sum_{n,k} \mu_{n,k}^\gamma = +\infty$ but

$$\mathbb{P}(\Lambda_\mu \text{ is } \alpha\text{-Carleson}) = 1$$

for all $\alpha \in (0, 1)$.

Remarks. 1) The first statement in part (a) is connected with the first part of the statement in Theorem 1.2, since it is a well-known fact that every 1-Carleson (or Carleson-Newman) sequence can be split into a finite number of separated sequences, each of which being of course 1-Carleson [20, Lemma 21] (obviously a finite number of arbitrary separated sequences may not be Carleson-Newman). However Theorem 1.4(a) does not give a precise information on the number of separated sequences involved. It is also mentionable that the condition for a.s. separation from Theorem 1.2 implies automatically the Carleson condition (picking $\gamma = 2 > 1$). This is perhaps more surprising and may be explained by the nature of the process: the independence of the different points allows for big fluctuations, so the probability of finding pairs of points arbitrarily close is quite big unless the number of points in the process is restricted severely (up to $\sum_{n,k} \mu_{n,k}^2 < \infty$).

2) It is interesting to point out that for the inhomogeneous Poisson process we have a characterization of 1-Carleson sequences, while in the a priori simpler random model with fixed radii and random arguments there is only a sufficient – still optimal – condition (see [9, Theorem 1.4]).

3) In the case $\alpha \in (0, 1)$ the results are less precise than when $\alpha = 1$. The value $1/(1 - \alpha)$ turns out to be an optimal breakpoint, but nothing specific can be said beyond this value without additional conditions on the distribution of μ . The example given in (c) is part of a certain parameter dependent scale of measures which will be discussed in Section 4.1, and for which the α -Carleson condition is characterized in terms of the parameter.

4) Our conditions, both here and in Theorem 1.2, are expressed in terms of $\mu_{n,k}$, thus re-distributing continuously μ on $T_{n,k}$ if necessary, we can always assume that μ is absolutely continuous with respect to the Lebesgue measure.

The structure of the paper is as follows. In Section 2 we prove the main Theorems 1.2 and 1.4. Section 3 deals with the consequences of these results in the study of interpolating sequences for various spaces of holomorphic functions. In particular, we find precise conditions so that a Poisson process Λ_μ is almost surely an interpolating sequence for the Hardy spaces H^p , $0 < p \leq \infty$, the Bloch space \mathcal{B} , or the Dirichlet spaces \mathcal{D}_α , $\alpha \in (1/2, 1)$. A final section is devoted to

provide examples of Poisson processes associated to some simple measures and to give integral conditions (non-discrete) on μ which are in some cases equivalent to the discrete versions used in the statements.

We finish this introduction recalling the Borel-Cantelli lemma, which is a central tool in this paper. We refer to [4] for a general source on probability theory. Given a sequence of events A_k let $\limsup A_k = \{\omega : \omega \in A_k \text{ for infinitely many } k\}$.

Lemma 1.5. *Let $(A_k)_k$ be a sequence of events in a probability space. Then*

- (1) *If $\sum \mathbb{P}(A_k) < \infty$, then $\mathbb{P}(\limsup A_k) = 0$,*
- (2) *If the events A_k are independent and $\sum \mathbb{P}(A_k) = \infty$, then $\mathbb{P}(\limsup A_k) = 1$.*

Acknowledgements: The authors would like to thank Joaquim Ortega-Cerdà for suggesting the consideration of Poisson processes and for helpful discussions.

2. PROOF OF THEOREMS 1.2 AND 1.4.

2.1. Proof of Theorem 1.2. Assume first that $\sum_{n,k} \mu_{n,k}^{M+1} < +\infty$, and define the events

$$A_{n,k} = \{X_{n,k} > M\} = \{X_{n,k} \geq M + 1\}.$$

Then

$$\mathbb{P}(A_{n,k}) = 1 - \sum_{j=0}^M \mathbb{P}(X_{n,k} = j) = 1 - e^{-\mu_{n,k}} \left(\sum_{j=0}^M \frac{\mu_{n,k}^j}{j!} \right).$$

By hypothesis $\lim_n (\sup_k \mu_{n,k}) = 0$, so we can use Taylor's formula

$$(2) \quad 1 - e^{-x} \left(\sum_{j=0}^M \frac{x^j}{j!} \right) = \frac{x^{M+1}}{(M+1)!} + o(x^{M+1}) \quad x \rightarrow 0$$

to deduce that

$$\sum_{n,k} \mathbb{P}(A_{n,k}) \lesssim \sum_{n,k} \frac{\mu_{n,k}^{M+1}}{(M+1)!} < +\infty.$$

By the Borel-Cantelli lemma $X_{n,k} \leq M$ for all but at most a finite number of $T_{n,k}$.

In principle this does not imply that Λ_μ can be split into M separated sequences, because it might happen that points of two neighboring $T_{n,k}$ come arbitrarily close. This possibility is excluded by repeating the above arguments to a new dyadic partition, made of shifted boxes $\tilde{T}_{n,k}$ having the ‘‘lower vertices’’ (those closer to \mathbb{T}) at the center of the $T_{n,k}$'s (see Figure 2 below); let

$$\tilde{T}_{n,k} = \left\{ z = r e^{it} : \frac{3}{2} 2^{-(n+2)} < 1 - r \leq \frac{3}{2} 2^{-(n+1)}; \frac{k+1/4}{2^n} \leq \frac{t}{2\pi} < \frac{k+3/4}{2^n} \right\}.$$

Since each $\tilde{T}_{n,k}$ is included in the union of at most four $T_{m,j}$, we still have $\sum_{n,k} \tilde{\mu}_{n,k}^{M+1} < \infty$, and therefore, as before, $\tilde{X}_{n,k} = N_{\tilde{T}_{n,k}}$ is at most M , except for maybe a finite number of indices (n, k) . This prevents that two adjacent $T_{n,k}$ have more than M points getting arbitrarily close. In

conclusion, for all but a finite number of indices $X_{n,k} \leq M$, hence the part of Λ_μ in these boxes can be split into M separated sequences. Adding the remaining finite number of points to any of these sequences may change the separation constant, but not the fact that they are separated.




FIGURE 2. Dyadic partitions: $\{T_{n,k}\}_{n,k}$ in blue, $\{\tilde{T}_{n,k}\}_{n,k}$ in red.

Assume now that $\sum_{n,k} \mu_{n,k}^{M+1} = +\infty$. We shall prove that for every $\delta_{l_0} = 2^{-l_0}$, $l_0 \in \mathbb{N}$,

$$\mathbb{P}(\Lambda \text{ union of } M \text{ } \delta_{l_0}\text{-separated sequences}) = 0.$$

Split each side of $T_{n,k}$ into 2^{l_0} segments of the same length. This defines a partition of $T_{n,k}$ in 2^{2l_0} small boxes of side length $2^{-n}2^{-l_0}$, which we denote by

$$T_{n,k}^{l_0,j} \quad j = 1, \dots, 2^{2l_0}.$$

Let $X_{n,k}^{l_0,j} = N_{T_{n,k}^{l_0,j}}$ denote the corresponding counting variable, which follows a Poisson law of parameter $\mu_{n,k,l_0,j} = \mu(T_{n,k}^{l_0,j})$.

It is enough to show that for any l_0 ,

$$\mathbb{P}(X_{n,k}^{l_0,j} > M \text{ for infinitely many } n, k, j) = 1.$$

By the second part of the Borel-Cantelli lemma, since the $X_{n,k}^{l_0,j}$ are independent, we shall be done as soon as we see that

$$(3) \quad \sum_{n,k} \sum_{j=1}^{2^{2l_0}} \mathbb{P}(X_{n,k}^{l_0,j} \geq M+1) = +\infty.$$

For any Poisson variable X of parameter λ , the probability

$$\mathbb{P}(X \geq M+1) = e^{-\lambda} \left(\sum_{m=M+1}^{\infty} \frac{\lambda^m}{m!} \right) = 1 - e^{-\lambda} \left(\sum_{m=0}^M \frac{\lambda^m}{m!} \right)$$

increases in λ . Hence there is no restriction in assuming that $0 \leq \mu_{n,k,l_0,j} \leq \mu_{n,k} \leq 1/2$ for all n, k, j . Then we can use Taylor's formula (2) to deduce that

$$\mathbb{P}(X_{n,k}^{l_0,j} \geq M+1) \simeq \frac{\mu_{n,k,l_0,j}^{M+1}}{(M+1)!},$$

and therefore (3) is equivalent to

$$\sum_{n,k} \sum_{j=1}^{2^{2l_0}} \mu_{n,k,l_0,j}^{M+1} = +\infty.$$

That this sum is infinite is just a consequence of the hypothesis and the elementary estimate

$$\mu_{n,k}^{M+1} = \left(\sum_{j=1}^{2^{2l_0}} \mu_{n,k,l_0,j} \right)^{M+1} \leq 2^{2l_0(M+1)} \sum_{j=1}^{2^{2l_0}} \mu_{n,k,l_0,j}^{M+1}.$$

2.2. Proof of Theorem 1.4. (a) Assume first that $\sum_n \mu_{n,k}^\gamma < +\infty$ for some $\gamma > 1$. It is enough to check the Carleson condition

$$\sum_{\lambda \in Q(I)} (1 - |\lambda|) \leq C|I|$$

on the dyadic intervals $I_{n,k}$. Let $Q_{n,k} = Q(I_{n,k})$. Decomposing the sum on the different layers A_m , it is enough to show that almost surely there exists $C > 0$ such that for all $n \geq 0, k = 0, \dots, 2^{n-1}$

$$\sum_{\lambda \in Q_{n,k}} (1 - |\lambda|) \simeq \sum_{m \geq n} \sum_{j: T_{m,j} \subset Q_{n,k}} 2^{-m} X_{m,j} \leq C2^{-n}.$$

This is equivalent to

$$(4) \quad \sup_{n,k} 2^n \sum_{m \geq n} \sum_{j: T_{m,j} \subset Q_{n,k}} 2^{-m} X_{m,j} < \infty$$

Denote

$$X_{n,m,k} = N_{Q_{n,k} \cap A_m} = \#(\Lambda \cap Q_{n,k} \cap A_m) = \sum_{j: T_{m,j} \subset Q_{n,k}} X_{m,j},$$

which is a Poisson variable of parameter

$$\mu_{n,m,k} = \mu(Q_{n,k} \cap A_m) = \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}.$$

Set

$$Y_{n,k} = 2^n \sum_{m \geq n} 2^{-m} X_{n,m,k} = \sum_{m \geq n} 2^{n-m} X_{n,k,m},$$

so that (4) becomes $\sup_{n,k} Y_{n,k} < +\infty$.

Let $A > 0$ be a big constant to be fixed later on. Again by the Borel-Cantelli Lemma, it is enough to show that

$$(5) \quad \sum_{n,k} \mathbb{P}(Y_{n,k} > A) < +\infty,$$

since then $Y_{n,k} \leq A$ for all but maybe a finite number of n, k ; in particular $\sup_{n,k} Y_{n,k} < \infty$.

The first step of the following reasoning is an adaptation to the Poisson process of the proof given in [9, Theorem 1.1] and which allowed to improve the result on Carleson sequences for the probabilistic model with fixed radii and random arguments. However, while in the original proof the Carleson boxes $Q_{n,k}$ are decomposed into layers $Q_{n,k} \cap A_m$ ($m \geq n$), in this new situation (as well as for (b)), Carleson boxes are decomposed into top-halves $T_{m,j} \subset Q_{n,k}$, which requires more delicate arguments to reach the convergence needed in the Borel-Cantelli lemma.

Recall that the probability generating function of a Poisson variable X of parameter λ is $\mathbb{E}(s^X) = e^{\lambda(s-1)}$. By the independence of the different $X_{n,k,m}$, $m \geq n$,

$$\mathbb{E}(s^{Y_{n,k}}) = \prod_{m \geq n} \mathbb{E}((s^{2^{n-m}})^{X_{n,m,k}}) = \prod_{m \geq n} e^{\mu_{n,m,k}(s^{2^{n-m}} - 1)}.$$

Thus for any $s > 1$, by Markov's inequality

$$\mathbb{P}(Y_{n,k} > A) = \mathbb{P}(s^{Y_{n,k}} > s^A) \leq \frac{1}{s^A} \mathbb{E}(s^{Y_{n,k}}) = \frac{1}{s^A} \prod_{m \geq n} e^{\mu_{n,m,k}(s^{2^{n-m}} - 1)}.$$

Using the estimate $x(a^{1/x} - 1) \leq a$, for $a, x > 1$, with $a = s$ and $x = 2^{m-n}$,

$$\begin{aligned} \log \mathbb{P}(Y_{n,k} > A) &\leq -A \log s + \sum_{m \geq n} (s^{2^{n-m}} - 1) \mu_{n,m,k} \\ &\leq -A \log s + \sum_{m \geq n} s 2^{n-m} \mu_{n,m,k} \\ &= -A \log s + s \sum_{m \geq n} 2^{n-m} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}. \end{aligned}$$

We want to optimize this estimate for $s > 1$. Set

$$B_{n,k} = \sum_{m \geq n} 2^{-(m-n)} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}$$

and define

$$\phi(s) = -A \log s + s B_{n,k}.$$

Let us observe first that the $B_{n,k}$ are uniformly bounded (they actually tend to 0). Indeed, let β denote the conjugate exponent of γ ($\frac{1}{\gamma} + \frac{1}{\beta} = 1$). Since for $m \geq n$ there are 2^{m-n} boxes $T_{m,j}$ in $Q_{n,k}$, by Hölder's inequality on the sum in the index j we deduce that

$$B_{n,k} \leq \sum_{m \geq n} 2^{-(m-n)} \left(\sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}^\gamma \right)^{1/\gamma} 2^{(m-n)/\beta}$$

$$= \sum_{m \geq n} 2^{-(m-n)/\gamma} \left(\sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}^\gamma \right)^{1/\gamma} < +\infty.$$

Taking A big enough we see that the minimum of ϕ is attained at $s_0 = A/B_{n,k} > 1$. Hence

$$\log \mathbb{P}(Y_{n,k} > A) \leq \phi(s_0) = -A \log \frac{A}{B_{n,k}} + A.$$

Therefore

$$\mathbb{P}(Y_{n,k} > A) \leq \left(\frac{B_{n,k}}{A} \right)^A e^A,$$

and

$$\sum_{n,k} \mathbb{P}(Y_{n,k} > A) \leq \left(\frac{e}{A} \right)^A \sum_{n,k} B_{n,k}^A.$$

The estimate on $B_{n,k}$ obtained previously is not enough to prove that this last sum converges. In order to obtain a better estimate take $p > 1$, to be chosen later on, its conjugate exponent q (i.e. $\frac{1}{p} + \frac{1}{q} = 1$), and apply Hölder's inequality in the following way:

$$\begin{aligned} B_{n,k} &= \sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-(m-n)} \mu_{m,j} = 2^n \sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m}{p}} 2^{-\frac{m}{q}} \mu_{m,j} \\ &\leq 2^n \left(\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\beta}{p}} \right)^{1/\beta} \times \left(\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\gamma}{q}} \mu_{m,j}^\gamma \right)^{1/\gamma}. \end{aligned}$$

Choose now p so that $1 < p < \beta$; then

$$\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\beta}{p}} = \sum_{m=n}^{\infty} 2^{-\frac{m\beta}{p}} 2^{m-n} = 2^{-n} \sum_{m=n}^{\infty} 2^{-m(\frac{\beta}{p}-1)} \simeq 2^{-n} 2^{-n(\frac{\beta}{p}-1)} = 2^{-n\frac{\beta}{p}}.$$

Thus, from the above estimate,

$$B_{n,k} \leq 2^{\frac{n}{q}} \left(\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\gamma}{q}} \mu_{m,j}^\gamma \right)^{1/\gamma}.$$

Choosing $A = \gamma$ yields

$$\sum_{n,k} B_{n,k}^\gamma \leq \sum_{n,k} 2^{\frac{n\gamma}{q}} \sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\gamma}{q}} \mu_{m,j}^\gamma.$$

We now apply Fubini's theorem to exchange the sums. The important observation here is that each $T_{m,j}$ has only one ancestor at each level $n \leq m$ (i.e, one $T_{n,k}$ containing $T_{m,j}$). Hence

$$\sum_{n,k} B_{n,k}^\gamma \leq \sum_{m,j} 2^{-\frac{m\gamma}{q}} \mu_{m,j}^\gamma \sum_{\substack{n \leq m \\ k: Q_{n,k} \supseteq T_{m,j}}} 2^{\frac{n\gamma}{q}} = \sum_{m,j} 2^{-\frac{m\gamma}{q}} \mu_{m,j}^\gamma \sum_{n \leq m} 2^{\frac{n\gamma}{q}}$$

$$\leq 2 \sum_{m,j} 2^{-\frac{m\gamma}{q}} \mu_{m,j}^\gamma 2^{\frac{m\gamma}{q}} = 2 \sum_{m,j} \mu_{m,j}^\gamma.$$

This finishes the proof of (5), hence of this part of the theorem.

Let us now assume that $\sum_{n,k} \mu_{n,k}^\gamma = +\infty$ for every $\gamma > 1$. Suppose $M \geq 1$ is an integer. Since the sum diverges for $\gamma = M + 1$, Theorem 1.2 implies that the sequence Λ_μ is almost surely not a union of M separated sequences. In particular, a.s. there is $\lambda_0 \in \Lambda_\mu$ such that $D_{\lambda_0} = \{z \in \mathbb{D} : \rho(\lambda_0, z) < 1/2\}$ contains at least $M + 1$ points of Λ_μ . Then, letting I_{λ_0} be the interval centered at $\lambda_0/|\lambda_0|$ with length $1 - |\lambda_0|$, we have $\sum_{\lambda \in Q(I_{\lambda_0})} (1 - |\lambda|) \gtrsim M|I_{\lambda_0}|$, where the underlying constant does not depend on M or λ_0 . This being true for every integer $M \geq 1$, the sequence cannot be 1-Carleson.

(b) Proceeding as in the first implication of (a) we see that it is enough to prove that almost surely

$$(6) \quad \sup_{n,k} Y_{n,k} < +\infty,$$

where now

$$(7) \quad Y_{n,k} = 2^{n\alpha} \sum_{m \geq n} 2^{-m\alpha} \sum_{j: T_{m,j} \subset Q_{n,k}} X_{m,j}.$$

The same estimates as in (a) based on the probability generating function yield, for $s > 1$,

$$\log \mathbb{P}(Y_{n,k} \geq A) \leq \phi(s) = -A \log s + B_{n,k},$$

where now

$$B_{n,k} = \sum_{m \geq n} 2^{-(m-n)\alpha} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}.$$

As in (a), the hypotheses imply that $B_{n,k}$ is uniformly bounded: letting β denote the conjugate exponent to γ ($\frac{1}{\gamma} + \frac{1}{\beta} = 1$) and noticing that $\alpha - 1/\beta = 1/\gamma - (1 - \alpha) > 0$,

$$\begin{aligned} B_{n,k} &\leq \sum_{m \geq n} 2^{-(m-n)\alpha} \left(\sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}^\gamma \right)^{1/\gamma} 2^{(m-n)/\beta} \\ &= \sum_{m \geq n} 2^{-(m-n)(\alpha-1/\beta)} \left(\sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}^\gamma \right)^{1/\gamma}. \end{aligned}$$

Therefore, optimizing the estimate for $s > 1$ exactly as we did in (a), we obtain $\mathbb{P}(Y_{n,k} \geq A) \lesssim B_{n,k}^A$, and we are lead to prove that for some $A > 0$

$$(8) \quad \sum_{n,k} \mathbb{P}(Y_{n,k} \geq A) \lesssim \sum_{n,k} B_{n,k}^A < \infty.$$

Again, we introduce an auxiliary weight p – to be determined later – and its conjugate exponent q . Split $2^{-m\alpha} = 2^{-\frac{m\alpha}{p}} 2^{-\frac{m\alpha}{q}}$ and use Hölder's inequality to obtain

$$B_{n,k} \leq 2^{n\alpha} \left(\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\alpha\beta}{p}} \right)^{1/\beta} \times \left(\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\alpha\gamma}{q}} \mu_{m,j}^\gamma \right)^{1/\gamma}.$$

The first sum is finite: since by hypothesis $\alpha\beta = \frac{\alpha\gamma}{\gamma-1} > 1$ there exists $1 < p < \frac{\alpha\gamma}{\gamma-1}$ and,

$$\sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\alpha\beta}{p}} = \sum_{m \geq n} 2^{-\frac{m\alpha\beta}{p}} 2^{m-n} = 2^{-n} \sum_{m \geq n} 2^{-m(\frac{\alpha\beta}{p}-1)} \simeq 2^{-n \frac{\alpha\beta}{p}}.$$

This implies that

$$B_{n,k}^\gamma \lesssim 2^{n\alpha \frac{\gamma}{q}} \sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\alpha\gamma}{q}} \mu_{m,j}^\gamma$$

and we can conclude the proof of (8) as before:

$$\begin{aligned} \sum_{n,k} B_{n,k}^\gamma &\lesssim \sum_{n,k} 2^{n\alpha \frac{\gamma}{q}} \sum_{\substack{m \geq n \\ j: T_{m,j} \subset Q_{n,k}}} 2^{-\frac{m\alpha\gamma}{q}} \mu_{m,j}^\gamma = \sum_{m,j} \mu_{m,j}^\gamma 2^{-m\alpha \frac{\gamma}{q}} \sum_{\substack{n \leq m \\ k: Q_{n,k} \supseteq T_{m,j}}} 2^{n\alpha \frac{\gamma}{q}} \\ &= \sum_{m,j} \mu_{m,j}^\gamma 2^{-m\alpha \frac{\gamma}{q}} \sum_{n \leq m} 2^{-n\alpha \frac{\gamma}{q}} \simeq \sum_{m,j} \mu_{m,j}^\gamma < +\infty. \end{aligned}$$

(c) Here we give a measure μ for which $\sum_{n,k} \mu_{n,k}^{1/(1-\alpha)} < +\infty$ but $\mathbb{P}(\Lambda_\mu \text{ is } \alpha\text{-Carleson}) = 0$. Let

$$d\mu(z) = \frac{dm(z)}{(1-|z|^2)^{1+\alpha} \log\left(\frac{e}{1-|z|^2}\right)},$$

which is the measure $\mu = \mu(1+\alpha, 1)$ given in the family of examples of Section 4.1. By a simple computation (see (10))

$$\mu_{n,k} \simeq \frac{2^{-n(1-\alpha)}}{n} \quad n \geq 1, \quad k = 0, \dots, 2^n - 1$$

and therefore, since k ranges over 2^n terms,

$$\sum_{n,k} \mu_{n,k}^{1/(1-\alpha)} \simeq \sum_{n \geq 1} \frac{1}{n^{1/(1-\alpha)}} < +\infty.$$

On the other hand, letting $Y_{n,k}$ be as in the proof of part (b) (see (7)) we get

$$\begin{aligned} \mathbb{E}(Y_{n,k}) &= 2^{n\alpha} \sum_{m \geq n} 2^{-m\alpha} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j} \simeq 2^{n\alpha} \sum_{m \geq n} 2^{-m\alpha} 2^{m-n} \frac{2^{-(1-\alpha)m}}{n} \\ &= 2^{-(1-\alpha)n} \sum_{m \geq n} \frac{1}{n} = +\infty \end{aligned}$$

Thus the expected weight of any single Carleson window $Q_{n,k}$ is infinite and Λ_μ cannot be α -Carleson.

(d) One could think of considering a divergent series $\sum_{n,k} \mu_{n,k}^\gamma = +\infty$ such that $\sum_{n,k} \mu_{n,k}^{\gamma'} < +\infty$ for every $\gamma' > \gamma$, and then apply (b), showing that Λ_μ is α -Carleson when $\gamma' < \frac{1}{1-\alpha}$, i.e. when $\alpha > 1 - \frac{1}{\gamma'} = \frac{\gamma'-1}{\gamma'}$. However, this does not yield the whole range $\alpha \in (0, 1)$ for a fixed measure, as required by the statement.

In order to construct an example working for all $\alpha \in (0, 1)$, we pick a measure μ supported in a Stolz angle of vertex 1, i.e. let, for $n \geq 1$,

$$\mu_{n,k} = \begin{cases} \frac{1}{n^{1/\gamma}} & \text{if } k = 0 \\ 0 & \text{if } k > 1. \end{cases}$$

(We could equivalently take the measure $\tau(2, 1/\gamma)$ given in Subsection 4.1, Example 3). Then

$$(9) \quad \sum_{n,k} \mu_{n,k}^\gamma = \sum_n \frac{1}{n} = \infty$$

but for every $\gamma' > \gamma$,

$$\sum_{n,k} \mu_{n,k}^{\gamma'} = \sum_n \frac{1}{n^{\gamma'/\gamma}} < +\infty.$$

To prove that Λ_μ is almost surely α -Carleson we will argue as before. Set $Y_{n,k}$ as in the proof of (b) (see (7)) and follow the same steps to prove that

$$\mathbb{P}(Y_{n,k} \geq A) \lesssim B_{n,k}^A,$$

where

$$B_{n,k} = \sum_{m \geq n} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j} 2^{-(m-n)\alpha}.$$

By construction $B_{n,k} = 0$ for all $k > 0$. On the other hand

$$B_{n,0} = 2^{n\alpha} \sum_{m \geq n} 2^{-m\alpha} \mu_{m,0} = 2^{n\alpha} \sum_{m \geq n} \frac{2^{-m\alpha}}{m^{1/\gamma}} \leq \frac{1}{n^{1/\gamma}}.$$

(Observe that this last expression is independent of α .) Hence

$$\sum_{n,k} B_{n,k}^{\gamma'} = \sum_n B_{n,0}^{\gamma'} \leq \sum_n \frac{1}{n^{\gamma'/\gamma}} < +\infty,$$

and as in the proof of (b) the Borel-Cantelli lemma allows to conclude that Λ is almost surely α -Carleson.

3. RANDOM INTERPOLATING SEQUENCES

In this section we discuss several consequences of Theorems 1.2 and 1.4 on random interpolating sequences Λ_μ for various spaces of holomorphic functions in \mathbb{D} . The results are rather straightforward consequences of the aforementioned theorems and the known conditions for such sequences.

3.1. Hardy (and Bergman) spaces. In this section we completely characterize the measures μ for which the associated Poisson process Λ_μ is almost surely an interpolating sequence for the Hardy spaces.

Recall that a sequence $\Lambda = \{\lambda_n\}_{n \in \mathbb{N}} \subset \mathbb{D}$ is interpolating for

$$H^\infty = \left\{ f \in H(\mathbb{D}) : \|f\|_\infty = \sup_{z \in D} |f(z)| < \infty \right\}$$

whenever for every sequence of bounded values $\{w_n\}_{n \in \mathbb{N}} \subset \mathbb{C}$ there exists $f \in H^\infty$ such that $f(\lambda_n) = w_n$, $n \in \mathbb{N}$. According to a famous theorem by L. Carleson, Λ is H^∞ -interpolating if and only if it is separated and 1-Carleson [8]. This characterization extends to all Hardy spaces

$$H^p = \left\{ f \in H(\mathbb{D}) : \|f\|_p = \sup_{r < 1} \left(\int_0^{2\pi} |f(re^{it})|^p \frac{dt}{2\pi} \right)^{1/p} < +\infty \right\} \quad 0 < p < \infty,$$

for which the interpolation problem is defined in a similar manner (the data w_n to be interpolated should satisfy $\sum_n (1 - |\lambda_n|^2) |w_n|^p < +\infty$, see e.g. [14, Chapter 9]).

The separation condition given in Theorem 1.2 implies immediately that Λ_μ is 1-Carleson, by Theorem 1.4, hence the following result follows.

Theorem 3.1. *Let Λ_μ be the Poisson process associated to a positive, σ -finite, locally finite measure μ . Then, for any $0 < p \leq \infty$,*

$$\mathbb{P}(\Lambda_\mu \text{ is } H^p\text{-interpolating}) = \begin{cases} 1 & \text{if } \sum_{n,k} \mu_{n,k}^2 < \infty \\ 0 & \text{if } \sum_{n,k} \mu_{n,k}^2 = \infty. \end{cases}$$

To complete the picture we discuss zero sequences Λ for H^p , $0 < p \leq \infty$. These are deterministically characterized by the Blaschke condition $\sum_{\lambda \in \Lambda} (1 - |\lambda|) < \infty$. Noticing that $\{\sum_{\lambda \in \Lambda_\mu} (1 - |\lambda|) < \infty\}$ is a tail event and using Kolmogorov's 0-1 law we get:

Proposition 3.2. *Let Λ_μ be the Poisson process associated to a positive, σ -finite, locally finite measure μ . Then, for any $0 < p \leq \infty$,*

$$\mathbb{P}(\Lambda_\mu \text{ is a zero set for } H^p) = \begin{cases} 1 & \text{if } \sum_{n,k} 2^{-n} \mu_{n,k} < \infty \\ 0 & \text{if } \sum_{n,k} 2^{-n} \mu_{n,k} = \infty. \end{cases}$$

Observe that the condition is just

$$\mathbb{E}\left[\sum_{\lambda \in \Lambda_\mu} (1 - |\lambda|)\right] = \mathbb{E}\left[\sum_{n,k} \sum_{\lambda \in T_{n,k}} (1 - |\lambda|)\right] \simeq \sum_{n,k} 2^{-n} \mathbb{E}[X_{n,k}] = \sum_{n,k} 2^{-n} \mu_{n,k} < \infty.$$

Observe also that $\sum_{k=0}^{2^n-1} \mu_{n,k} = \mu(A_n)$ for all $n \in \mathbb{N}$, hence

$$\sum_{n,k} 2^{-n} \mu_{n,k} = \sum_n 2^{-n} \mu(A_n).$$

Proof of Proposition 3.2. Denote $X_n = N_{A_n} = \sum_{k=0}^{2^n-1} X_{n,k}$ and denote $\mu_n = \mathbb{E}[X_n] = \mu(A_n)$.

Assume first that $\sum_n 2^{-n} \mu_n < +\infty$. Set $Y = \sum_n 2^{-n} X_n$ and observe that, by the independence of the different X_n ,

$$\mathbb{E}[Y] = \sum_n 2^{-n} \mu_n < +\infty, \quad \text{Var}(Y) = \sum_n 2^{-2n} \mu_n < +\infty.$$

Then, by Markov's inequality

$$\mathbb{P}(Y \geq 2\mathbb{E}(Y)) \leq \frac{1}{2}.$$

Since $\{Y = \infty\}$ is a tail event, Kolmogorov's 0-1 law implies that $\mathbb{P}(Y = +\infty) = 0$, and in particular the Blaschke sum is finite almost surely.

Assume now that $\sum_n 2^{-n} \mu_n = +\infty$. Split the sum in two parts:

$$\sum_n 2^{-n} \mu_n = \sum_{n: \mu_n \leq 2^n/n^2} 2^{-n} \mu_n + \sum_{n: \mu_n > 2^n/n^2} 2^{-n} \mu_n.$$

It is enough to consider the second sum, since the first one obviously converges. Since $\text{Var}[X_n] = \mu_n$, Chebyshev's inequality yields,

$$\mathbb{P}(X_n \leq \frac{1}{2}\mu_n) = \mathbb{P}(X_n \leq \mu_n - \frac{\mu_n}{2}) \leq \mathbb{P}(|X_n - \mu_n| \geq \frac{\mu_n}{2}) \leq \frac{4}{\mu_n}.$$

Hence

$$\sum_{n: \mu_n > 2^n/n^2} \mathbb{P}(X_n \leq \frac{1}{2}\mu_n) \leq \sum_{n: \mu_n > 2^n/n^2} \frac{4}{\mu_n} \leq \sum_{n: \mu_n > 2^n/n^2} \frac{4n^2}{2^n} < +\infty.$$

Now, by the Borel-Cantelli lemma, $X_n > \frac{1}{2}\mu_n$ for all but maybe a finite number of the n with $\mu_n > 2^n/n^2$; hence

$$\sum_{n: \mu_n > 2^n/n^2} 2^{-n} X_n \gtrsim \frac{1}{2} \sum_{n: \mu_n > 2^n/n^2} 2^{-n} \mu_n,$$

which diverges, by hypothesis. ■

3.1.1. *Remark. Interpolation in Bergman spaces.* Interpolating sequences Λ for the (weighted) Bergman spaces

$$B_\alpha^p = \left\{ f \in H(\mathbb{D}) : \|f\|_{\alpha,p}^p = \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^{\alpha p - 1} dm(z) < \infty \right\},$$

with $0 < \alpha, 0 < p \leq \infty$ are characterized by the separation together with the upper density condition

$$D_+(\Lambda) := \limsup_{r \rightarrow 1^-} \sup_{z \in \mathbb{D}} \frac{\sum_{1/2 < \rho(z,\lambda) \leq r} \log \frac{1}{\rho(z,\lambda)}}{\log\left(\frac{1}{1-r}\right)} < \alpha$$

(see [23] and [17, Chapter 5] for both the definitions and the results).

Since every 1-Carleson sequence has density $D_+(\Lambda) = 0$, the same conditions of Theorem 3.1 also characterize a.s. Bergman interpolating sequences, regardless of the indices α and p . Again, because of the big fluctuations of the Poisson process, the conditions required to have separation a.s. are so strong that they can only produce sequences of zero upper density.

Another indication of the big fluctuations of the Poisson process is the following. For the invariant measure $d\nu(z) = \frac{dm(z)}{(1-|z|^2)^2}$, which obviously satisfies $\nu_{n,k} \simeq 1$ for all n, k , it is not difficult to see that almost surely,

$$D_+(\Lambda_\nu) = +\infty \quad \text{and} \quad D_-(\Lambda_\nu) := \liminf_{r \rightarrow 1^-} \inf_{z \in \mathbb{D}} \frac{\sum_{1/2 < \rho(z,\lambda) \leq r} \log \frac{1}{\rho(z,\lambda)}}{\log\left(\frac{1}{1-r}\right)} = 0.$$

Therefore there are way too many points for Λ_ν to be interpolating for any B_α^p , but there are too few for it to be sampling, since these sets must have strictly positive lower density $D_-(\Lambda)$ (see [17, Chapter 5]).

3.2. Interpolation in the Bloch space. We consider now interpolation in the Bloch space \mathcal{B} , consisting of functions f holomorphic in \mathbb{D} such that

$$\|f\|_{\mathcal{B}} := |f(0)| + \sup_{z \in \mathbb{D}} |f'(z)|(1 - |z|^2) < +\infty.$$

Since Bloch functions satisfy the Lipschitz condition $|f(z) - f(w)| \leq \|f\|_{\mathcal{B}} \delta(z, w)$, where $\delta(z, w) = \frac{1}{2} \log \frac{1 + \rho(z, w)}{1 - \rho(z, w)}$ denotes the hyperbolic distance, A. Nicolau and B. Bøe defined interpolating sequences for \mathcal{B} as those $\Lambda = \{\lambda_n\}_{n \in \mathbb{N}}$ such that for every sequence of values $\{v_n\}_{n \in \mathbb{N}}$ with $\sup_{n \neq m} \frac{|v_n - v_m|}{\delta(\lambda_n, \lambda_m)} < \infty$ there exists $f \in \mathcal{B}$ with $f(\lambda_n) = v_n$, $n \in \mathbb{N}$ [5].

Theorem ([5, pag.172], [24, Theorem 7]). *A sequence Λ of distinct points in \mathbb{D} is an interpolating sequence for \mathcal{B} if and only if:*

- (a) Λ can be expressed as the union of at most two separated sequences,
- (b) for some $0 < \gamma < 1$ and $C > 0$,

$$\#\{\lambda \in \Lambda : \rho(z, \lambda) < r\} \leq \frac{C}{(1-r)^\gamma}$$

independently on $z \in \mathbb{D}$.

As explained in [5], condition (b) can be replaced by:

$$(b)' \text{ for some } 0 < \gamma < 1 \text{ and } C > 0, \text{ and for all Carleson windows } Q(I), \\ \#\{\lambda \in Q(I) : 2^{-(l+1)}|I| < 1 - |\lambda| < 2^{-l}|I|\} \leq C2^{\gamma l}, \quad l \geq 0.$$

In [24, Corollary 2] it is mentioned that it can also be replaced by:

(b)'' there exist $0 < \gamma < 1$ and such that Λ is γ -Carleson.

In view of conditions (a) and (b)'' the following characterization of Poisson processes which are a.s. Bloch interpolating sequences follows from Theorems 1.2 and 1.4(b) (with $\gamma \in (2/3, 1)$).

Theorem 3.3. *Let Λ_μ be the Poisson process associated to a positive, σ -finite, locally finite measure μ . Then,*

$$\mathbb{P}(\Lambda_\mu \text{ is } \mathcal{B}\text{-interpolating}) = \begin{cases} 1 & \text{if } \sum_{n,k} \mu_{n,k}^3 < \infty \\ 0 & \text{if } \sum_{n,k} \mu_{n,k}^3 = \infty. \end{cases}$$

Note. In case $\sum_{n,k} \mu_{n,k}^3 < \infty$ it is also possible to prove (b)' directly, with the same methods employed in the proof of Theorem 1.4. It is enough to prove the estimate for dyadic arcs $I_{n,k}$, and for those

$$\#\{\lambda \in Q(I_{n,k}) : 2^{-(l+1)}|I_{n,k}| < 1 - |\lambda| < 2^{-l}|I_{n,k}|\} \simeq \sum_{j: T_{n+l,j} \subset Q_{n,k}} X_{n+l,j}.$$

In the above, the left hand side corresponds essentially to the number of points in the layer $Q(I_{n,k}) \cap A_{n+l}$. Thus, with $m = n + l$, (b)' is equivalent to

$$\sup_{n,k} \sup_{m \geq n} 2^{-\gamma(m-n)} \sum_{j: T_{m,j} \subset Q_{n,k}} X_{m,j} < +\infty.$$

Letting

$$Y_{n,k,m} = 2^{-\gamma(m-n)} \sum_{j: T_{m,j} \subset Q_{n,k}} X_{m,j}, \quad \mathbb{E}[Y_{n,k,m}] = 2^{-\gamma(m-n)} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}$$

and proceeding as in the first part of the proof of Theorem 1.4(a) we get (taking $A = 3$):

$$\begin{aligned} \sum_{n,k} \sum_{m \geq n} \mathbb{P}(Y_{n,k,m} \geq 3) &\lesssim \sum_{n,k} \sum_{m \geq n} \left[2^{-\gamma(m-n)} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j} \right]^3 \\ &\leq \sum_{n,k} \sum_{m \geq n} 2^{-3\gamma(m-n)} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}^3 2^{2(m-n)} \\ &= \sum_{m,j} \mu_{m,j}^3 \sum_{n \leq m} \sum_{k: Q_{n,k} \supseteq T_{m,j}} 2^{-(3\gamma-2)(m-n)}. \end{aligned}$$

For any $\gamma > 2/3$ this sum is bounded by $\sum_{m,j} \mu_{m,j}^3$, so we can conclude with the Borel-Cantelli lemma.

3.3. Interpolation in Dirichlet spaces. Our last set of results concerns interpolation in the Dirichlet spaces,

$$\mathcal{D}_\alpha = \left\{ f \in H(\mathbb{D}) : \|f\|_{\mathcal{D}_\alpha}^2 = |f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 (1 - |z|^2)^\alpha dm(z) < \infty \right\},$$

with $\alpha \in (0, 1)$. The limiting case $\alpha = 1$ can be identified with the Hardy space H^2 .

In these spaces, interpolating sequences are characterized by the separation and a Carleson type condition. This was initially considered by W.S. Cohn, see [11]; we refer also to the general result [1]. While separation is a simple condition, that in our random setting is completely characterized by Theorem 1.2, the characterization of Carleson measures in these spaces is much more delicate. This was achieved by D. Stegenga using the so-called α -capacity [25]. In our setting it is however possible to use an easier sufficient one-box condition that can be found in K. Seip's book, see [24, Theorem 4, p.38], which we recall here for the reader's convenience.

Theorem 3.4 (Seip). *A separated sequence Λ in \mathbb{D} is interpolating for \mathcal{D}_α , $0 < \alpha < 1$ if there exist $0 < \alpha' < \alpha$ such that Λ is α' -Carleson.*

The reader should be alerted that in Seip's book the space \mathcal{D}_α is defined in a slightly different way, and that the above statement is adapted to our definition.

For these spaces Theorems 1.2 and 1.4 lead to less precise conclusions. Indeed, in view of Theorem 1.4(c),(d) we cannot hope for complete characterizations if we do not impose additional conditions on the measure μ .

Theorem 3.5. *Let Λ_μ be the Poisson process associated to a positive, σ -finite, locally finite measure μ .*

(a) *If $1/2 < \alpha < 1$, then*

$$P(\Lambda_\mu \text{ is interpolating for } \mathcal{D}_\alpha) = \begin{cases} 1 & \text{if } \sum_{n,k} \mu_{n,k}^2 < +\infty \\ 0 & \text{if } \sum_{n,k} \mu_{n,k}^2 = +\infty. \end{cases}$$

(b) *If $0 \leq \alpha < 1/2$ and there exists $1 < \gamma < \frac{1}{1-\alpha}$ such that $\sum_{n,k} \mu_{n,k}^\gamma < +\infty$, then*

$$P(\Lambda_\mu \text{ is interpolating for } \mathcal{D}_\alpha) = 1.$$

Clearly, the condition $\sum_{n,k} \mu_{n,k}^2 < +\infty$ is also necessary in the case (b) (if the sum diverges, then Λ_μ is almost surely not separated).

Proof. (a) If $\sum_{n,k} \mu_{n,k}^2 = +\infty$, then Λ_μ is almost surely not separated by Theorem 1.2, hence it is almost surely not interpolating.

If $\sum_{n,k} \mu_{n,k}^2 < +\infty$, Theorem 1.2 shows again that the sequence Λ_μ is almost surely separated. By Seip's theorem, it remains to show that Λ_μ is almost surely α' -Carleson for some $\alpha' < \alpha$. Pick $1/2 < \alpha' < \alpha < 1$, so that $1/(1 - \alpha') > 2$. Choosing $\gamma \in (2, 1/(1 - \alpha'))$ we get

$$\sum_{n,k} \mu_{n,k}^\gamma \lesssim \sum_{n,k} \mu_{n,k}^2 < +\infty,$$

and by Theorem 1.4(b) we conclude that Λ_μ is almost surely α' -Carleson.

(b) If $\alpha < 1/2$ then $1/(1 - \alpha) < 2$ and the value γ given by the hypothesis satisfies $1 < \gamma < 2$. Therefore

$$\sum_{n,k} \mu_{n,k}^2 \lesssim \sum_{n,k} \mu_{n,k}^\gamma < +\infty,$$

which allows to deduce from Theorem 1.2 that Λ_μ is almost surely separated.

Since the inequality $\gamma < 1/(1 - \alpha)$ is strict, we also have $\gamma < 1/(1 - \alpha')$ for some $\alpha' < \alpha$ sufficiently close to α . Again, Theorem 1.4(b) shows that Λ_μ is almost surely α' -Carleson, and Seip's theorem implies that Λ_μ is almost surely interpolating. ■

3.4. Additional remarks and comments. The above results show several applications of our Theorems 1.2 and 1.4, but they also give rise to many challenging questions. Is it possible to get a necessary counterpart of Theorem 1.4(b) under reasonable conditions on μ (more general than the class considered in Section 4.1 below)? Is it possible to get precise statements when $\alpha = 1/2$? Also, the case of the classical Dirichlet space seems to be largely unexplored for Poisson point processes, while the situation regarding interpolation, separation and zero-sets for the radial probabilistic model is completely known for all $\alpha \in [0, 1]$ (see [9, 6]).

4. EXAMPLES AND INTEGRAL CONDITIONS FOR THE MEASURE μ

In the first part of this final section we illustrate the above results with three simple families of measures on \mathbb{D} . In the second part we briefly discuss alternative, non-discrete, formulations of the conditions given in the previous statements.

4.1. Examples. *1. Radial measures.* Let dm denote the normalized Lebesgue measure and let $d\nu(z) = \frac{dm(z)}{(1-|z|^2)^2}$ be the invariant measure in \mathbb{D} . Define

$$d\mu(a, b)(z) = \frac{dm(z)}{(1 - |z|^2)^a \log^b\left(\frac{e}{1-|z|^2}\right)} = \frac{d\nu(z)}{(1 - |z|^2)^{a-2} \log^b\left(\frac{e}{1-|z|^2}\right)},$$

where either $a > 1$, $b \in \mathbb{R}$, or $a = 1$ and $b \leq 1$ (so that $\mu(a, b)(\mathbb{D}) = +\infty$).

Observe that

$$\mu(a, b)_{n,k} \simeq \frac{2^{-n(2-a)}}{n^b} \quad n \geq 1, \quad k = 0, \dots, 2^n - 1,$$

and therefore, for $\gamma > 0$,

$$(10) \quad \sum_{n,k} \mu(a, b)_{n,k}^\gamma \simeq \sum_n 2^n \frac{2^{-n(2-a)\gamma}}{n^{b\gamma}} = \sum_n \frac{2^{-n[(2-a)\gamma-1]}}{n^{b\gamma}}.$$

Proposition 4.1. *Consider the Poisson process $\Lambda_{a,b}$ associated to the measure $\mu(a, b)$, with either $a > 1$ or $a = 1$ and $b \leq 1$.*

- (a) $\Lambda_{a,b}$ can a.s. be expressed as a union of M separated sequences if and only if either $a < 2 - \frac{1}{M+1}$ and $b \in \mathbb{R}$, or $a = 2 - \frac{1}{M+1}$ and $b > \frac{1}{M+1}$.
- (b) In particular, $\Lambda_{a,b}$ is a.s. separated if and only if either $a < 3/2$ and $b \in \mathbb{R}$, or $a = 3/2$ and $b > 1/2$.
- (c) $\Lambda_{a,b}$ is a.s. a 1-Carleson sequence if and only if $a < 2$, $b \in \mathbb{R}$.
- (d) Let $\alpha \in (0, 1)$. Then $\Lambda_{a,b}$ is a.s. an α -Carleson sequence if and only if $a < 1 + \alpha$ or $a = 1 + \alpha$ and $b > 1$.

Proof. (a) is immediate from Theorem 1.2 and (10) with $\gamma = M + 1$.

(c) If $a \geq 2$ the series in (10) diverges for all $\gamma > 1$, thus by Theorem 1.4(a) $\Lambda_{a,b}$ is a.s. not 1-Carleson.

On the other hand, if $a < 2$ there exists γ such that $(2 - a)\gamma - 1 > 0$ (i.e., such that $\gamma > \frac{1}{2-a}$). For that γ the series in (10) converges, and we can conclude again by Theorem 1.4(a).

(d) Suppose first that $a < 1 + \alpha$. As in the previous case, since $2 - a > 1 - \alpha$ there exists $\gamma \in (\frac{1}{2-a}, \frac{1}{1-\alpha})$. For this γ the series in (10) converges and we can apply Theorem 1.4(b).

If $a > 1 + \alpha$ and $b \in \mathbb{R}$, then $\Lambda_{\mu(a,b)}$ contains in the mean more points than $\Lambda_{\mu(1+\alpha,1)}$ for which we have shown in Theorem 1.4(c) that it is almost surely not α -Carleson.

It thus remains the case $a = 1 + \alpha$. Again, when $b = 1$ — and thus also when $b < 1$ since then we have more points in the mean — the proof of Theorem 1.4(c) shows that the corresponding sequence is almost surely not α -Carleson.

Finally, suppose that $a = 1 + \alpha$ and $b > 1$. Recall from (7) the notation

$$Y_{n,k} = 2^{n\alpha} \sum_{m \geq n} 2^{-m\alpha} \sum_{j: T_{m,j} \subset Q_{n,k}} X_{m,j}.$$

In the proof of Theorem 1.4(b) we have shown that

$$P(Y_{n,k} \geq A) \leq B_{n,k}^A,$$

where

$$B_{n,k} = \sum_{m \geq n} 2^{-(m-n)\alpha} \sum_{j: T_{m,j} \subset Q_{n,k}} \mu_{m,j}.$$

From the explicit form of $\mu_{m,j}$ we get

$$B_{n,k} \simeq \sum_{m \geq n} 2^{-(m-n)\alpha} \times 2^{m-n} \times \frac{2^{-m(2-a)}}{m^b} = 2^{n(\alpha-1)} \sum_{m \geq n} \frac{2^{-m(\alpha+1-a)}}{m^b}$$

which converges exactly when $a < 1 + \alpha$ or when $a = 1 + \alpha$ and $b > 1$ which is the case we are interested in here. In this situation, we get

$$B_{n,k} \simeq \frac{2^{-n(2-a)}}{n^b}$$

Clearly, when $A \geq 1/(2-a) = 1/(1-\alpha)$, then $\sum_{n,k} B_{n,k}^A$ converges, and the Borel-Cantelli lemma shows that $Y_{n,k} \geq A$ can happen for an at most finite number of Carleson windows $Q_{n,k}$. Hence $\Lambda_{\mu(a,b)}$ is a.s. α -Carleson. \blacksquare

2. *Measures with a singularity on \mathbb{T} .* Define now

$$d\sigma(a,b)(z) = \frac{dm(z)}{|1-z|^a \log^b\left(\frac{e}{|1-z|}\right)},$$

where either $a > 2$, $b \in \mathbb{R}$, or $a = 2$ and $b \leq 1$ (so that $\sigma(a,b)(\mathbb{D}) = +\infty$). Here

$$\sigma(a,b)_{n,k} = \sigma(a,b)(T_{n,k}) \simeq \frac{2^{-2n}}{[(k+1)2^{-n}]^a \log^b\left(\frac{e}{(k+1)2^{-n}}\right)}, \quad n \in \mathbb{N}, k = 0, \dots, 2^n - 1.$$

Hence for $\gamma > 1$,

$$(11) \quad \sum_{n,k} \sigma(a,b)_{n,k}^\gamma \simeq \sum_n 2^{-n\gamma(2-a)} \sum_{k=1}^{2^n} \frac{1}{\left(k^a \log^b\left(\frac{e}{k2^{-n}}\right)\right)^\gamma}.$$

Let us examine the growth of the sum in k . For that, set

$$S_n(a,b,\gamma) = \sum_{k=1}^{2^n} \frac{1}{k^{a\gamma} \log^{b\gamma}\left(\frac{e}{k2^{-n}}\right)} \simeq \int_1^{2^n} \frac{dx}{x^{a\gamma} \log^{b\gamma}\left(\frac{e}{x2^{-n}}\right)}$$

The change of variable $t = \log\left(\frac{e}{x2^{-n}}\right)$ leads to

$$S_n(a,b,\gamma) \simeq \int_{\log(2^n e)}^1 \left(\frac{e^t}{e2^n}\right)^{a\gamma-1} \frac{-dt}{t^{b\gamma}} = \frac{2^{-n(a\gamma-1)}}{e^{a\gamma-1}} \int_1^{\log(2^n e)} e^{t(a\gamma-1)} \frac{dt}{t^{b\gamma}}$$

Our standing assumption being $a > 2$ or $a = 2$ and $b \leq 1$, we only need to consider these two cases. In both cases, $e^{t(a\gamma-1)}/t^{b\gamma} \rightarrow +\infty$ when $t \rightarrow +\infty$, and the last integral behaves essentially as the value in the upper bound of the integration interval

$$\int_1^{\log(2^n e)} e^{t(a\gamma-1)} \frac{dt}{t^{b\gamma}} \simeq \frac{2^{n(a\gamma-1)}}{n^{b\gamma}}.$$

Hence

$$S_n(a,b,\gamma) \simeq \frac{1}{n^{b\gamma}},$$

and

$$(12) \quad \sum_{n,k} \sigma(a,b)_{n,k}^\gamma \simeq \sum_n 2^{-n\gamma(2-a)} \times \frac{1}{n^{b\gamma}} = \sum_n \frac{2^{-n\gamma(2-a)}}{n^{b\gamma}}.$$

We are now in a position to prove the following result.

Proposition 4.2. *Consider the Poisson process $\tilde{\Lambda}_{a,b}$ associated to the measure $\sigma(a, b)$, with either $a > 2$ or $a = 2$ and $b \leq 1$.*

- (a) *For $a > 2$ the process $\tilde{\Lambda}_{a,b}$ is a.s. neither a finite union of separated sequences nor an α -Carleson, for any $\alpha \in (0, 1]$.*
- (b) *For $a = 2$ the process $\tilde{\Lambda}_{2,b}$ is*
 - (i) *the union of M separated sequences if and only if $b > \frac{1}{M+1}$,*
 - (ii) *α -Carleson for $\alpha \in (0, 1)$ if $b > 1 - \alpha$.*

Proof. (a) is immediate from Theorems 1.2 and 1.4, since (12) diverges for all $\gamma > 0$.

(b) In this case the series (12) is just $\sum_n 1/n^{b\gamma}$.

The case (i) follows from Theorem 1.2 with $\gamma = M + 1$.

For (ii), by the hypothesis $1/b < 1/(1 - \alpha)$, there exists $1/b < \gamma < 1/(1 - \alpha)$, for which the series (12) converges. We can conclude by Theorem 1.4. \blacksquare

3. *Measures in a cone.* Given a point $\zeta \in \mathbb{T}$, consider a Stolz region

$$\Gamma(\zeta) = \left\{ z \in \mathbb{D} : \frac{|\zeta - z|}{1 - |z|} < 2 \right\}.$$

We discuss the previous measures restricted to $\Gamma(\zeta)$. With no restriction of generality we can assume that $\zeta = 1$. Let thus

$$d\tau(a, b)(z) = \chi_{\Gamma(1)}(z) d\mu(a, b)(z) = \chi_{\Gamma(1)}(z) \frac{dm(z)}{(1 - |z|^2)^a \log^b\left(\frac{e}{1 - |z|^2}\right)},$$

where now either $a > 2$, $b \in \mathbb{R}$, or $a = 2$ and $b \leq 1$ (so that $\nu(a, b)(\mathbb{D}) = +\infty$). Since in Γ the measures $d\mu(a, b)$ and $d\sigma(a, b)$ behave similarly, we could replace $d\mu(a, b)$ by $d\sigma(a, b)$ in the definition of $d\tau(a, b)$.

Observe that $\nu(a, b)_{n,k}$ is non-zero only for a finite number N of k at each level n , and that for those k

$$\tau(a, b)_{n,k} \simeq \frac{2^{-n(2-a)}}{n^b} \quad n \geq 1, \quad k = 0, \dots, N.$$

Hence

$$(13) \quad \sum_{n,k} \tau(a, b)_{n,k}^\gamma \simeq \sum_n \frac{2^{-n(2-a)\gamma}}{n^{b\gamma}},$$

which is exactly the same estimate as in (12) and thus immediately leads to the same result as Proposition 4.2. This might look surprising since $\sigma(a, b)$ (and a fortiori $\mu(a, b)$) puts infinite mass outside $\Gamma(\zeta)$ (actually outside Stolz angles at ζ with arbitrary opening).

Proposition 4.3. *Consider the Poisson process $\hat{\Lambda}_{a,b}$ associated to the measure $\tau(a, b)$, with either $a > 2$ or $a = 2$ and $b \leq 1$.*

- (a) For $a > 2$ the process $\hat{\Lambda}_{a,b}$ is a.s. not a finite union of separated sequences separated or α -Carleson for any $\alpha \in (0, 1]$.
- (b) For $a = 2$ the process $\hat{\Lambda}_{2,b}$ is
 - (1) the union of M separated sequences if and only if $b > \frac{1}{M+1}$,
 - (2) α -Carleson for $\alpha \in (0, 1)$ if $b > 1 - \alpha$.

4.2. Integral conditions on μ . Given a locally finite, σ -finite measure μ , a natural question is whether the discretized conditions appearing in Theorems 1.2 and 1.4 can be reformulated in terms of integrals. Let us assume that μ is absolutely continuous with respect to the Lebesgue (or the invariant) measure on \mathbb{D} . In view of the aforementioned discrete conditions this is not really restrictive, since in case μ had a singular part we could just redistribute its mass continuously on each $T_{n,k}$. Assume thus that $d\mu = h d\nu$, where $d\nu = \frac{dm(z)}{(1-|z|^2)^2}$ is the invariant measure, $h \geq 0$ and $h \in L^1_{loc}(\mathbb{D}; \nu)$ (but $h \notin L^1(\mathbb{D}; \nu)$, so that $\mu(\mathbb{D}) = \infty$).

As a result of Jensen’s inequality applied to ν on $T_{n,k}$, we deduce the following general observation.

Proposition 4.4. *Let $\mu = h d\nu$, where $h \geq 0$ and $h \in L^1_{loc}(\mathbb{D})$. For every $\gamma > 1$ there exists $C > 0$ such that*

$$(14) \quad \sum_{n,k} \mu_{n,k}^\gamma \leq C \int_{\mathbb{D}} h^\gamma(z) d\nu(z).$$

Of course without additional conditions on h the conditions on the sum and on the integral cannot be equivalent, and there are standard construction methods to find measures for which the sum is convergent while the integral diverges. We do not go into details of that here.

One is obviously more interested in situations where the sum and the integral conditions are equivalent. This is for instance the case when h is radial with some regularity conditions (as in the first class of examples in the previous section) or when h is subharmonic (as in the second class of examples).

Another mildly regular case in which an equivalent reformulation in terms of integrals is possible is when μ is doubling, meaning that there exists $C > 0$ such that $\mu(2D) \leq C\mu(D)$ for all open disc $D \subset \mathbb{D}$. Here $2D$ denotes the disk with the same center as D but with double radius (in the pseudohyperbolic metric). Fixing any $c \in (0, 1)$ and defining $F_\mu(z) = \mu(D(z, c))$ one immediately sees that for any $\gamma > 1$

$$\sum_{n,k} \mu_{n,k}^\gamma \simeq \int_{\mathbb{D}} F_\mu^\gamma(z) d\nu(z).$$

REFERENCES

[1] Aleman, A.; Hartz, M.; McCarthy, J.; Richter, S., *Interpolating Sequences in Spaces with the Complete Pick Property*, Int. Math. Res. Not. 2019, no. 12, 3832–3854. pages 18

- [2] Aparicio Monforte, A. *Successions aleatòries a $\mathbb{B}_n \subset \mathbb{C}^n$* . Master's thesis. Universitat de Barcelona (2005). pages 4
- [3] Amar, É. ; Bonami, A. *Mesures de Carleson d'ordre α et solutions au bord de l'équation $\bar{\partial}_b$* . Bull. Soc. Math. France 107 (1979), no. 1, 23–48. pages 4
- [4] Billingsley, P. *Probability and measure*. Wiley, New York, 1979. pages 6
- [5] Bøe, B. ; Nicolau, A. *Interpolation by functions in the Bloch space*. J. Anal. Math. 94 (2004), 171–194. pages 16, 17
- [6] Bogdan, K. *On the zeros of functions with finite Dirichlet integral*. Kodai Math. J. 19 (1996), no. 1, 7–16. pages 19
- [7] Bomash, G.; *A Blaschke-type product and random zero sets for Bergman spaces*. Ark. Mat. 30 (1992), no. 1, 45–60. pages
- [8] Carleson, L; *An interpolation problem for bounded analytic functions*. Amer. J. Math. 80 (1958), 921–930. pages 4, 14
- [9] Chalmoukis, N. ; Hartmann, A.; Kellay, K. ; Wick, B.D. *Random Interpolating Sequences in Dirichlet Spaces*. Int. Math. Res. Not. IMRN (2021), rnab110, <https://doi-org.sire.ub.edu/10.1093/imrn/rnab110> pages 2, 5, 9, 19
- [10] Cochran, W. G.; *Random Blaschke products*. Trans. Amer. Math. Soc. 322 (1990), no. 2, 731–755. pages 2
- [11] Cohn, W.S. ; *Interpolation and multipliers on Besov and Sobolev spaces*. Complex Variables Theory Appl. 22 (1993), no. 1-2, 35–45. pages 18
- [12] Daley, D. J.; Vere-Jones, D. *An introduction to the theory of point processes*. Vol. I. Elementary theory and methods. Second edition. Probability and its Applications (New York). Springer-Verlag, New York, 2003. pages
- [13] Dayan, A.; Wick, B.D.; Wu, Sh.; *Random Interpolating Sequences in the Polydisc and the Unit Ball*, preprint, <http://arxiv.org/abs/2012.05381v4> pages 2
- [14] Duren, P. L. ; *Theory of H^p spaces*. Pure and Applied Mathematics, Vol. 38 Academic Press, New York–London 1970. pages 14
- [15] El Fallah, O.; Kellay, K.; Mashreghi, J. ; Ransford, T. *One-box conditions for Carleson measures for the Dirichlet space*. Proc. Amer. Math. Soc. 143 (2015), no. 2, 679–684. pages
- [16] Garnett, J.B.; *Bounded analytic functions*. Pure and Applied Mathematics, 96. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1981. xvi+467 pp. pages 1, 4
- [17] Hedenmalm, H.; Korenblum, B.; Zhu, K.; *Theory of Bergman spaces*. Graduate Texts in Mathematics, 199. Springer-Verlag, New York, 2000. pages 16
- [18] Hough, J. B. ; Krishnapur, M. ; Peres, Y. ; Virág, B. *Zeros of Gaussian analytic functions and determinantal point processes*. University Lecture Series, 51. American Mathematical Society, Providence, RI, 2009. pages 2
- [19] Last, G.; Penrose, M.; *Lectures on the Poisson process*. Institute of Mathematical Statistics Textbooks, 7. Cambridge University Press, Cambridge, 2018. pages 2
- [20] McDonald, G.; Sundberg, C. *Toeplitz operators on the disc*. Indiana Univ. Math. J. 28 (1979), no. 4, 595–611 pages 5
- [21] Rudin, W.; *Function theory in the unit ball of \mathbb{C}^n* . Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 241. Springer-Verlag, New York-Berlin, 1980. xiii+436 pp. pages
- [22] Rudowicz, R.; *Random sequences interpolating with probability one*. Bull. London Math. Soc. 26 (1994), no. 2, 160–164. pages 2
- [23] Seip, K.; *Beurling type density theorems in the unit disk*. Invent. Math. 113 (1993), no. 1, 21–39. pages 16
- [24] Seip, K. *Interpolation and Sampling in Spaces of Analytic Functions*. University Lecture Series, vol. 33. American Mathematical Society, 2004. pages 1, 16, 17, 18
- [25] D. Stegenga, *Multipliers of the Dirichlet space*. Illinois J. Math. 24 (1980), no. 1, 113–139. pages 18

- [26] Wikipedia contributors, *Poisson point process*, Wikipedia, The Free Encyclopedia, 19 April 2022 19:38 UTC, (https://en.wikipedia.org/w/index.php?title=Poisson_point_process&oldid=1083617092) [accessed 19 May 2022] pages 2

UNIV. BORDEAUX, CNRS, BORDEAUX INP, IMB, UMR 5251, F-33400, TALENCE, FRANCE

DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA 585,
08007-BARCELONA, CATALONIA