



**HAL**  
open science

# On the existence of two solutions for a boundary value problem on the half-line

Ibrahim Soufi

► **To cite this version:**

Ibrahim Soufi. On the existence of two solutions for a boundary value problem on the half-line. 2022.  
hal-03731350

**HAL Id: hal-03731350**

**<https://hal.science/hal-03731350>**

Preprint submitted on 21 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On the existence of two solutions for a boundary value problem on the half-line

Ibrahim Soufi<sup>1</sup>

<sup>1</sup>ESSA. Algeria, Algiers

\*E-mail : [soufimath@gmail.com](mailto:soufimath@gmail.com)

---

## Abstract

In this note we consider Dirichlet boundary value problem on a half line. Using critical point theory and Mountain Pass Theorem, we prove the existence of two nontrivial solutions.

## Keywords

Dirichlet boundary value problem, half line, critical point, mountain pass.

---

## I INTRODUCTION

In this work is concerned with the following boundary value problem (BVP for short) on the half-line

$$\begin{cases} -(p(t)u'(t))' = \lambda f(t, u(t)), & \text{a.e. } t > 0, \\ u(0) = u(+\infty) = 0, \end{cases} \quad (1)$$

where  $\lambda > 0$  be a numerical parameter and  $f : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function. The coefficient  $p : [0, +\infty) \rightarrow (0, +\infty)$  satisfies  $\frac{1}{p} \in L^1(0, +\infty)$ , and

$$\int_0^{+\infty} \left( \int_t^{+\infty} \frac{1}{p(s)} ds \right) dt < +\infty.$$

This paper is devoted to the generalization of problem in [9], when we consider the problem in the half-line.

Define the space

$$H_{0,p}^1(0, +\infty) = \{u \in AC([0, +\infty), \mathbb{R}) \mid u(0) = u(+\infty) = 0, \sqrt{p}u' \in L^2(0, +\infty)\}.$$

Take  $v \in H_{0,p}^1(0, +\infty)$ , multiply the equation in problem (1) by  $v$ , and then integrate over  $(0, +\infty)$ , we get

$$\int_0^{+\infty} p(t)u'(t)v'(t)dt = \lambda \int_0^{+\infty} f(t, u(t))v(t)dt.$$

This leads to the natural concept of weak solution for problem (1).

**Definition 1:** We say that a function  $u \in H_{0,p}^1(0, +\infty)$  is a weak solution of problem (1) if

$$\int_0^{+\infty} p(t)u'(t)v'(t)dt - \lambda \int_0^{+\infty} f(t, u(t))v(t)dt = 0,$$

for all  $v \in H_{0,p}^1(0, +\infty)$ .

In order to study problem (1), we consider the functional  $J : H_{0,p}^1(0, +\infty) \rightarrow R$  defined by

$$J(u) = \frac{1}{2} \int_0^{+\infty} p(t)u'^2(t)dt - \lambda \int_0^{+\infty} F(t, u(t))dt, \quad (2)$$

where

$$F(t, u) = \int_0^u f(t, s)ds.$$

**Lemma 1.**  $H_{0,p}^1(0, +\infty)$  embeds in  $L^2(0, +\infty)$ .

**Proof.** For  $u \in H_{0,p}^1(0, +\infty)$ , we have

$$|u(t)| = \left| \int_t^{+\infty} u'(s)ds \right| = \left| \int_t^{+\infty} \sqrt{p(s)}u'(s) \frac{1}{\sqrt{p(s)}}ds \right|.$$

Then, by the Cauchy-Schwartz inequality

$$\begin{aligned} |u(t)|^2 &\leq \left( \int_t^{+\infty} p(s)u'^2(s)ds \right) \left( \int_t^{+\infty} \frac{1}{p(s)}ds \right) \\ &\leq \left( \int_0^{+\infty} p(s)u'^2(s)ds \right) \left( \int_t^{+\infty} \frac{1}{p(s)}ds \right). \end{aligned}$$

Hence

$$\int_0^{+\infty} |u(t)|^2 dt \leq \left( \int_0^{+\infty} \left( \int_t^{+\infty} \frac{1}{p(s)}ds \right) dt \right) \left( \int_0^{+\infty} p(s)|u'(s)|^2 ds \right),$$

that is

$$\|u\|_{L^2} \leq \sqrt{M} \|\sqrt{p}u'\|_{L^2}.$$

**Theorem 1.** (a) *The operator*

$$\begin{aligned} T : H_{0,p}^1(0, +\infty) &\rightarrow T(H_{0,p}^1(0, +\infty)) \subset L^2(0, +\infty) \times L^2(0, +\infty) : = L_2^2(0, +\infty) \\ u &\rightarrow T(u) = (u, \sqrt{p}u') \end{aligned}$$

*is an isometric isomorphism.*

(b)  $H_{0,p}^1(0, +\infty)$  is a reflexive space.

**Proof.** (a) It is clear that  $T$  is a linear operator and that  $T$  conserves norms, i.e.,

$$\forall u \in H_{0,p}^1(0, +\infty), \quad \|Tu\|_{L_2^2} = \|u\|_p.$$

Indeed

$$\begin{aligned} \|Tu\|_{L_2^2} &= \|(u, \sqrt{p}u')\|_{L_2^2} \\ &= \|u\|_{L^2} + \|\sqrt{p}u'\|_{L^2} \\ &= \|u\|_p. \end{aligned}$$

(b) Since  $L^2(0, +\infty)$  is a reflexive Banach space, the cartesian product  $L^2_2(0, +\infty)$  is also a reflexive Banach space with respect to the norm

$$\|u\|_{L^2_2} = \|u_1\|_{L^2} + \|u_2\|_{L^2}, \quad \text{where } u = (u_1, u_2) \in L^2_2(0, +\infty).$$

From part (a),  $T(H^1_{0,p}(0, +\infty))$  is a closed subspace of  $L^2_2(0, +\infty)$ , the space  $T(H^1_{0,p}(0, +\infty))$  is reflexive. Consequently  $H^1_{0,p}(0, +\infty)$  is also reflexive.

Notice that  $H^1_{0,p}(0, +\infty)$  is a Banach space equipped with the norm

$$\|u\|_{0,p} = \sqrt{\int_0^{+\infty} p(t)u'^2(t)dt + \int_0^{+\infty} u^2(t)dt},$$

or the equivalent norm

$$\|u\|_p = \|u\|_{L^2} + \|\sqrt{p}u'\|_{L^2}.$$

Moreover the space  $H^1_{0,p}(0, +\infty)$  is reflexive.

**Theorem 2.** [8] On  $H^1_{0,p}(0, +\infty)$ , the quantity  $\|u\| = \sqrt{\int_0^{+\infty} p(t)u'^2(t)dt}$  is a norm which is equivalent to the  $H^1_{0,p}(0, +\infty)$ -norm.

**Lemma 2.**  $(H^1_{0,p}(0, +\infty), \|\cdot\|)$  embeds in  $(C_0[0, +\infty), \|u\|_\infty)$ , where  $C_0[0, +\infty) = \{u \in C([0, +\infty), R) \mid \lim_{t \rightarrow +\infty} u(t) = 0\}$  and  $\|u\|_\infty = \sup_{t \in [0, +\infty)} |u(t)|$ , and we have

$$\|u\|_\infty \leq M\|u\| \quad \text{where } M = \sqrt{\left\| \frac{1}{p} \right\|_{L^1}}. \quad (3)$$

**Theorem 3.** The embedding

$$H^1_{0,p}(0, +\infty) \hookrightarrow C_0[0, +\infty) \quad (4)$$

is compact.

Let  $E$  be a real reflexive Banach space, and write  $B_\rho$  for the closed ball centered at 0 with radius  $\rho$  in  $E$  and  $\partial B_\rho = \{u \in E \mid \|u\| = \rho\}$  for the boundary of  $B_\rho$ .

We will employ the following conditions.

(H<sub>1</sub>)  $\Phi : E \rightarrow R$  is a convex, continuously Fréchet differentiable functional

with derivative  $\varphi : E \rightarrow E^*$ ;

(H<sub>2</sub>)  $H : E \rightarrow R$  is a continuously Fréchet differentiable functional with

derivative  $h : E \rightarrow E^*$ ;

(H<sub>3</sub>) the operator  $h : E \rightarrow E^*$  is compact;

(H<sub>4</sub>) there exist constants  $\alpha, \alpha_1 > 1, \gamma > 0$  such that

$$\gamma\|v\|_E^\alpha \leq \langle \varphi(v), v \rangle \quad \text{for all } v \in E$$

and

$$\lim_{\|u\|_E \rightarrow +\infty} \sup \frac{\Phi(u)}{\|u\|_E^{\alpha_1}} = +\infty;$$

(H<sub>5</sub>)  $H$  is a convex functional.

$\langle \cdot, \cdot \rangle$  denotes the action of a derivative on a suitable element or else a duality pairing. We will determine such a value  $\lambda^* > 0$  that for each  $\lambda \in (0, \lambda^*]$  the corresponding Euler action functional  $J : E \rightarrow R$

$$J(u) = \Phi(u) - \lambda H(u),$$

has a critical point on  $B_\rho$ . This implies the solvability of

$$\varphi(u) = \lambda h(u), \quad u \in E. \quad (5)$$

**Theorem 4.** [9] Assume that  $\mathbf{H}_1, \mathbf{H}_2, \mathbf{H}_5$  are satisfied. Fix some  $\lambda^* > 0$  and let  $u_0, v \in E$  be such that

$$J(u_0) \leq J(v) \quad \text{and} \quad \varphi(v) = \lambda^* h(u_0). \quad (6)$$

Then  $u_0$  is a critical point to  $J$ , and thus it solves (5).

**Theorem 5.** [9] Let  $E$  be an infinite dimensional reflexive Banach space and let  $B_\rho$  be fixed. Assume that  $\mathbf{H}_1 - \mathbf{H}_5$  are satisfied. Then there exists  $\lambda^* > 0$  such that for each  $\lambda \in (0, \lambda^*]$  there exists  $u_0 \in B_\rho$  with

$$J(u_0) = \inf_{u \in B_\rho} J(u) \quad (7)$$

and such that  $u_0$  is a critical point to  $J$ , and thus it solves (5).

**Theorem 6.** [9] Let  $E$  be an infinite dimensional reflexive Banach space. Assume that  $\mathbf{H}_1 - \mathbf{H}_5$  are satisfied. Take some  $\rho > 0$ . Then there exists  $\lambda^* > 0$  such that for each  $\lambda \in (0, \lambda^*]$  there exists  $u_0 \in B_\rho$  with

$$J(u_0) = \inf_{u \in B_\rho} J(u) \quad (8)$$

and such that  $u_0$  is a critical point of  $J$ , and thus it solves (5). If for some  $v \in B_\rho$  it holds that  $J(v) < 0$  and  $J(0) = 0$  or else  $h(0) \neq 0$  and  $J(0) = 0$ , then  $u_0$  is non-trivial.

Assume additionally that there exists  $\lambda_1^* \leq \lambda^*$  such that for all  $\lambda \in (0, \lambda_1^*)$

(a)  $J$  satisfies the PS-condition,

(b)  $J(0) < \inf_{u \in \partial B_{\rho_1}} J(u)$  for some  $\rho_1 > \|u_0\|_E$ ,

(c) there exists  $w \in E \setminus B_{\rho_1}$  with  $J(w) \leq 0$ .

Then for all  $\lambda \in (0, \lambda_1^*)$  the functional  $J$  has two nontrivial critical points, namely  $u_0$  and another non-zero critical point  $z_0$  different from  $u_0$ .

## II APPLICATIONS

We will employ the following hypotheses.

(A<sub>1</sub>) There exist positive functions  $a, b$  with  $a, b \in L^1(0, +\infty)$  and  $\sigma > 1$  such that

$$|f(t, u)| \leq a(t)|u|^\sigma + b(t), \quad \text{for a.e. } t \in [0, +\infty) \text{ and all } u \in R,$$

(A<sub>2</sub>) there exists functions  $c_1, c_2 : [0, +\infty) \rightarrow (0, +\infty)$  with  $c_1, c_2 \in L^1(0, +\infty)$ , and  $\theta > 2$  such that

(a)  $\theta F(t, u) \leq uf(t, u)$ , for a.e.  $t \in [0, +\infty)$  and all  $u \in R \setminus \{0\}$ ,

(b)  $F(t, u) \geq c_1(t)|u|^\theta - c_2(t)$ , for a.e.  $t \geq 0$  and all  $u \in R$ ,

(A<sub>3</sub>) the function  $u \mapsto F(t, u)$  is convex on  $R$  for a.e.  $t \in [0, +\infty)$ .

**Theorem 7.** Let  $\lambda > 0$  be fixed. The functional  $J$  is well-defined and continuously differentiable on  $H_{0,p}^1(0, +\infty)$ . The derivative of  $J$  at any  $u \in H_{0,p}^1(0, +\infty)$  has the following form

$$\langle J'(u), v \rangle = \int_0^{+\infty} p(t)u'(t)v'(t)dt - \lambda \int_0^{+\infty} f(t, u(t))v(t)dt, \quad \forall v \in H_{0,p}^1(0, +\infty).$$

In particular,  $u \in H_{0,p}^1(0, +\infty)$  is a solution of problem (1) if and only if  $u$  is a critical point of  $J$ .

## 2.1 Existence of solution by the mountain pass lemma

**Lemma 3.** Suppose that  $(A_1)$ ,  $(A_2(a))$  hold. Then for any  $\lambda > 0$ , the functional  $J$  given by (2) satisfies the PS-condition.

**proof.** Let us take a sequence  $\{u_k\}_{k=1}^{\infty} \subset H_{0,p}^1(0, +\infty)$  such that  $\{J(u_k)\}_{k=1}^{\infty}$  is bounded and  $J'(u_k) \rightarrow 0$ , as  $k \rightarrow \infty$ . We shall show that  $\{u_k\}_{k=1}^{\infty}$  has a convergent subsequence.

Since  $J'(u_k) \rightarrow 0$ , we see that for some  $\epsilon > 0$ , there exists  $k_0$  with  $\|J'(u_k)\| \leq \epsilon$  for  $k \geq k_0$ . Note that for  $k \geq k_0$

$$|\langle J'(u_k), u_k \rangle| \leq \epsilon \|u_k\|$$

and because we have

$$\langle J'(u_k), u_k \rangle = \int_0^{+\infty} p(t) (u_k'(t))^2 dt - \lambda \int_0^{+\infty} f(t, u_k(t))u_k(t)dt,$$

then

$$-\lambda \int_0^{+\infty} f(t, u_k(t))dt \geq \frac{-\epsilon}{\theta} \|u_k\| - \frac{1}{\theta} \|u_k\|^2.$$

Since  $\{J(u_k)\}_{k=1}^{\infty}$  is bounded, there exists a constant  $C$  such that  $|J(u_k)| \leq C$ ,  $\forall k \in N$ . Using the above estimates, we obtain

$$C \geq \left( \frac{\theta - 2}{2\theta} \right) \|u_k\|^2 - \frac{\epsilon}{2} \|u_k\|.$$

Since  $\theta > 2$ , then the sequence  $(u_k)$  is bounded in  $H_{0,p}^1(0, +\infty)$  i.e.,  $\exists M_2 > 0$  such that  $\|u_k\| \leq M_2$ ,  $\forall k \in N$ . Next, we prove that  $(u_k)$  converges strongly to some  $\bar{u}$  in  $H_{0,p}^1(0, +\infty)$ . Since  $(u_k)$  is bounded in the reflexive Banach space  $H_{0,p}^1(0, +\infty)$ , there exists a subsequence of  $(u_k)$  still denoted  $(u_k)$  such that  $(u_k)$  converges weakly to some  $\bar{u}$  in  $H_{0,p}^1(0, +\infty)$ , and  $\|\bar{u}\| \leq M_2$ . Then  $(u_k)$  converges to  $\bar{u}$  in  $C_0[0, +\infty)$  by Lemma 2.

Since  $\lim_{k \rightarrow +\infty} J'(u_k) = 0$  and  $(u_k)$  converges weakly to some  $\bar{u}$ , we get

$$\lim_{k \rightarrow +\infty} \langle J'(u_k) - J'(\bar{u}), u_k - \bar{u} \rangle \rightarrow 0 \quad (9)$$

and

$$\langle J'(u_k) - J'(\bar{u}), u_k - \bar{u} \rangle = \|u_k - \bar{u}\|^2 - \lambda \int_0^{+\infty} (f(t, u_k(t)) - f(t, \bar{u}(t))) (u_k(t) - \bar{u}(t)) dt.$$

Since  $u_k \rightarrow \bar{u}$  in  $C_0[0, +\infty)$ ,  $\forall t \in [0, +\infty)$ , then  $u_k(t) \rightarrow \bar{u}(t)$  for  $t \in [0, +\infty)$ , and since  $f$  is Carathéodory, we have  $f(t, u_k(t)) \rightarrow f(t, \bar{u}(t))$  as  $k \rightarrow +\infty$  p.p.  $t \in [0, +\infty)$ , and using  $(\mathbf{A}_1)$  we have

$$|f(t, u_k(t))| \leq M_2^\sigma M^\sigma a(t) + b(t). \quad (10)$$

Since:  $a \in L^1(0, +\infty)$ ,  $b \in L^1(0, +\infty)$ , then

$$M_2^\sigma M^\sigma a + b \in L^1(0, +\infty). \quad (11)$$

By the Lebesgue dominated convergence theorem, we obtain

$$\lim_{k \rightarrow +\infty} \int_0^{+\infty} f(t, u_k(t)) dt = \int_0^{+\infty} f(t, \bar{u}(t)) dt. \quad (12)$$

Then (9) and (12) imply that  $\{u_k\}_{n=1}^\infty$  is strongly convergent.

**Lemma 4.** *Suppose that  $(\mathbf{A}_1)$ ,  $(\mathbf{A}_2(\mathbf{b}))$  hold. There exists  $\lambda_1^* > 0$  such that for any  $\lambda \leq \lambda_1^*$  there exist numbers  $\rho_1, \alpha > 0$  such that*

- (1) *there are positive constants  $\rho_1, \alpha$  such that  $J(u) \geq \alpha$  for all  $u \in H_{0,p}^1(0, +\infty)$  with  $\|u\| = \rho_1$ .*
- (2) *There exists an element  $w \in H_{0,p}^1(0, +\infty)$  with  $\|w\| > \rho_1$*

*and such that  $J(w) < 0$ .*

**proof.** Thus in (2)

$$J(u) = \frac{1}{2} \|u\|^2 - \lambda \int_0^{+\infty} F(t, u(t)) dt.$$

We first verify that  $J$  satisfies assumption (1) in Lemma 4.

Define  $\beta_1 := \frac{1}{\sigma + 1} M^\sigma \rho_1^\sigma \|a\|_{L^1} + \|b\|_{L^1}$ . Then we see that for any  $u \in \partial B(0, \rho_1)$  with  $\|u\| = \rho_1 = 2M$ , (2) and  $(\mathbf{A}_1)$  give

$$\left| \int_0^{+\infty} F(t, u(t)) dt \right| \leq M \rho_1 \beta_1.$$

Put  $\lambda_1^* = \frac{1}{\beta_1}$  and fix  $\lambda \in (0, \lambda_1^*)$ .

Moreover,

$$J(u) \geq 2M^2(1 - \lambda\beta_1),$$

this implies that assumption (1) holds if we take  $\rho_1 = 2M$  and  $\alpha = 2M^2(1 - \lambda\beta_1)$ .

Now **(A2)** guarantees that for some  $w_0 \in H_{0,p}^1(0, +\infty)$  with  $w_0 \neq 0$  and  $s \in R_+$ , we have the following estimation

$$J(sw_0) \leq \frac{1}{2}s^2\|w_0\|^2 - \lambda s^\theta \int_0^{+\infty} c_1(t)|w_0(t)|^\theta dt + \lambda \int_0^{+\infty} c_2(t)dt.$$

Since  $\theta > 2$  we see that  $J(sw_0) \rightarrow -\infty$  as  $s \rightarrow +\infty$ . Thus there is some  $s_0$  such that  $w = s_0w_0$  we have  $J(w) < 0$ .

Therefore assumption (2) in Lemma 4 is also satisfied.

Now the mountain pass lemma allows us to formulate the following existence result.

**Theorem 8.** *Suppose also that **(A<sub>1</sub>)**, **(A<sub>2</sub>)** hold. Then there exists  $\lambda_1^* > 0$  such that for all  $\lambda \in (0, \lambda_1^*)$ , problem (1) has at least one nontrivial solution.*

### Example 1

Consider the boundary value problem

$$\begin{cases} -(e^t u'(t))' & = \lambda \left( \frac{3u^3 + 2u}{(20 + t^2)\sqrt{u^2 + 1}} \right), \\ u(0) = u(+\infty) & = 0. \end{cases} \quad (13)$$

It can be easily checked that all conditions of Theorem 8 are satisfied with

$$f(t, u) = \frac{3u^3 + 2u}{(20 + t^2)\sqrt{u^2 + 1}}, \quad \sigma = 3, \quad \theta = 2.1,$$

$$a(t) = \frac{1}{1 + t^2}, \quad b(t) = \frac{5}{t^2 + 1}, \quad p(t) = e^t,$$

$$c_1(t) = \frac{1}{20 + t^2}, \quad c_2(t) = \frac{4}{10}e^{-t},$$

and

$$F(t, u) = \frac{u^2\sqrt{u^2 + 1}}{20 + t^2} - \frac{1}{10}e^{-t}$$

Then there is some  $\lambda_1^* > 0$  such that for all  $\lambda \in (0, \lambda_1^*)$ , problem (13) has at least one nontrivial solutions.

## 2.2 Existence of solution by the critical point theorem on a closed ball

**Theorem 9.** *Let  $B(0, \rho)$ , fixed. Assume that **A<sub>1</sub>**, **A<sub>3</sub>** are satisfied and that  $f(t, 0) \neq 0$ . Then there exist  $\lambda^* > 0$  such that for all  $\lambda \in (0, \lambda^*)$ , there exists  $u_0 \in B(0, \rho)$  with*

$$J(u_0) = \inf_{u \in B_\rho} J(u) \quad (14)$$

and such that  $u_0$  is a critical point, and thus it solves (1).



### 2.3 Existence of two solutions

Concerning the multiple solutions, we have the main result of this section where we need only that  $F$  is convex in addition to assumptions leading to a mountain pass solution.

**Theorem 10.** *Assume that  $A_1 - A_3$  are satisfied and that  $f(t, 0) \neq 0$ . Then there exist  $0 < \lambda_1^* \leq \lambda^*$  such that for all  $0 < \lambda \leq \lambda_1^*$ , problem (1) has at least two nontrivial solutions.*

### REFERENCES

- [1] A.M. Arthurs. *Complementary Variational Principles*. Clarendon Press, Oxford (1980).
- [2] M. Badiale, E. Serra, *Semilinear Elliptic Equations for Beginners*. Springer-Verlag, London, 2011
- [3] G. Bonanno, D. O'Regan, *A boundary value problem on the half-line via critical point methods*. Dynam. Systems Appl. 15 (2006), no. 3-4, 395–407.
- [4] B. E. Breckner, D. Repovas and C. Varga, *On the existence of three solutions for the Dirichlet problem on the Sierpinski gasket*, Nonlinear Anal. 73 (2010), no. 9, 2980–2990
- [5] H. BRÉZIS, *Analyse Fonctionnelle, Théorie et Applications*. Masson, Paris, 1983.
- [6] H. BRÉZIS, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2010.
- [7] C. Corduneanu. *Integral Equations and Stability of Feedback Systems*. Cambridge University Press, Cambridge (1973). Iranian Math. Society, 43(1) 2017, 223-237.
- [8] M. Galewski., T. Moussaoui., I. Soufi, *On the existence of solutions for a boundary value problem on the half-line*, ejtde 2018, No. 12, 1 – 12.
- [9] M. Galewski, *Multiple Solutions to a Dirichlet Problem on the Sierpinski Gasket*. Taiwanese journal of mathematics 20(5) October 2016.
- [10] M. Galewski, *On a new multiple critical point theorem and some applications to anisotropic problems*. Taiwanese journal of mathematics 19(5) 2015. .
- [11] J. Mawhin, *Problèmes de Dirichlet variationnels non linéaires*, Les Presses de l'Université de Montréal, 1987.