

The Role of Motor Inhibition During Covert Speech Production

Ladislas Nalborczyk, Ursula Debarnot, Marieke Longcamp, Aymeric Guillot,

F.-Xavier Alario

► To cite this version:

Ladislas Nalborczyk, Ursula Debarnot, Marieke Longcamp, Aymeric Guillot, F.-Xavier Alario. The Role of Motor Inhibition During Covert Speech Production. Frontiers in Human Neuroscience, 2022, 16, 10.3389/fnhum.2022.804832 . hal-03737946

HAL Id: hal-03737946 https://hal.science/hal-03737946

Submitted on 12 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	The role of motor inhibition during covert speech production
2	Ladislas Nalborczyk ^{1,2} , Ursula Debarnot ^{3,4} , Marieke Longcamp ² , Aymeric Guillot ^{3,4} , and
3	and FXavier Alario ¹
4	¹ Aix Marseille Univ, CNRS, LPC, Marseille, France
5	² Aix Marseille Univ, CNRS, LNC, Marseille, France
6	³ Inter-University Laboratory of Human Movement Biology-EA 7424, University of Lyon,
7	University Claude Bernard Lyon 1, Villeurbanne, France
8	⁴ Institut Universitaire de France, Paris, France

Author Note

¹⁰ Correspondence concerning this article should be addressed to Ladislas

11 Nalborczyk. E-mail: ladislas.nalborczyk@univ-amu.fr.

9

Abstract

Covert speech is accompanied by a subjective multisensory experience with auditory and 13 kinaesthetic components. An influential hypothesis states that these sensory percepts 14 result from a simulation of the corresponding motor action that relies on the same 15 internal models recruited for the control of overt speech. This simulationnist view raises 16 the question of how it is possible to imagine speech without executing it. In this 17 perspective, we discuss the possible role(s) played by motor inhibition during covert 18 speech production. We suggest that considering covert speech as an inhibited form of 19 overt speech maps naturally to the purported progressive internalisation of overt speech 20 during childhood. However, we argue that the role of motor inhibition may differ widely 21 across different forms of covert speech (e.g., condensed vs. expanded covert speech) and 22 that considering this variety helps reconciling seemingly contradictory findings from the 23 neuroimaging literature. 24

Keywords: covert speech, inner speech, motor imagery, motor simulation, motor
 control, motor inhibition

Introduction

1

The ability to mentally examine our verbal thoughts is central to our subjective 28 experience. This covert (internal) production of speech typically accompanies everyday 29 activities such as problem solving (Baldo et al., 2005; Sokolov, 1972), future planning 30 (D'Argembeau et al., 2011), reading (e.g., Lœvenbruck et al., 2005; Perrone-Bertolotti 31 et al., 2012), or writing (Frith, 1979). Because overt speech production results from 32 sequences of motor commands that are assembled to reach a given communication goal, it 33 belongs to the broader category of motor actions (Jeannerod, 2006a). Therefore, a 34 parallel can be drawn between covert speech, also known as *inner speech* or *speech* 35 *imagery* (for reviews, see Alderson-Day & Fernyhough, 2015; Lœvenbruck et al., 2018; 36 Perrone-Bertolotti et al., 2014), and other imagined actions (i.e., motor imagery). The 37 motor simulation theory of motor imagery (Jeannerod, 1994, 2001, 2006b) postulates a 38 continuum between the covert and the overt execution of an action and that action 39 representations can operate off-line via a simulation mechanism. 40

However, the proposal that overt and covert actions share common processes and 41 neural circuits is faced with a serious problem. If the neural circuits used for the control 42 of overt actions are also used for covert actions, how can covert actions not lead to 43 execution? This puzzle was coined as the problem of inhibition by Jeannerod (2001). It 44 can be rephrased as follows: given the putative role of the motor system in providing the 45 multisensory content of motor imagery, how is it possible for motor imagery not to lead 46 to motor execution? In this perspective, we examine some theoretical and experimental 47 consequences that emerge from considering covert speech as inhibited overt speech. First, 48 we explore the role and plausible neural implementation of inhibitory mechanisms during 49 covert speech production. Second, we relate the maturation of inhibitory control during 50 childhood with the progressive internalisation of overt speech. Third, we consider how 51 inhibitory mechanisms may play different roles across different forms of covert speech. By 52 bridging recent results from the covert speech, motor imagery, and motor inhibition 53 literature, we highlight some novel and possibly fruitful lines of research. 54

2 Covert speech production as inhibited overt speech production

⁵⁶ 2.1 Cognitive and neural mechanisms supporting motor inhibition

First and foremost, we need to make a distinction between at least two different 57 types of inhibition. First, cognitive inhibition, defined as the stopping or overriding of a 58 mental process, with or without intention (MacLeod, 2007). Second, the inhibition of 59 physical response, or motor inhibition, defined broadly as the withholding, suppression, 60 or overriding of an inappropriate, prepotent, or unwanted motor response (Aron, 2007; 61 O'Shea & Moran, 2018). Here, we are concerned with the latter. Ridderinkhof et al. 62 (2014) further described the concept of response inhibition on three continuous 63 dimensions: intentionality, premeditation, and specificity. Inhibition can be employed 64 with more or less intentionality, planned ahead or employed in the moment, and applied 65 to a specific action and effector, or more globally, to all actions and/or effectors. 66

Within Ridderinkhof et al.'s classification of response inhibitions, we hypothesise 67 that covert speech involves an intentional (we know we want to produce these actions 68 covertly rather than overtly) but implicit/automatic (we do not explicitly think about 69 not producing movements) and planned ahead form of response inhibition. The 70 distinction between implicit and explicit inhibition seems important to highlight. Indeed, 71 the type of motor inhibition that may be at play during motor imagery is still different 72 from the "proactive inhibition" in the motor inhibition literature. Indeed, in behavioural 73 tasks aiming to assess proactive inhibition, participants are instructed not to execute an 74 action. In contrast, while doing motor imagery, participants are asked to imagine the 75 action, which implies that it should not be executed overtly (Guillot et al., 2012). 76 Moreover, the type of motor inhibition that is implemented during covert speech 77 necessarily has to be planned ahead, otherwise speech acts would sometimes be (at least 78 partially) executed. Regarding the level at which motor inhibition may be applied during 79 covert speech, we hypothesise that inhibition may apply both globally and in an 80 effector-specific manner. Supporting this hypothesis, Rieger et al. (2017) and Bart et al. 81 (2021) have shown, using an action mode (overt vs. covert) switching paradigm, that the 82 motor imagery of hand movements is accompanied by both global and effector-specific 83

⁸⁴ inhibition (these results were also replicated in Scheil & Liefooghe, 2018).

Based on evidence from electrophysiological, neuroimaging, and clinical studies, 85 Guillot et al. (2012) suggested three possible routes whereby motor commands can be 86 inhibited during motor imagery. First, motor inhibition can be integrated within the 87 representation of the action to be produced internally. Second, cerebral regions such as 88 the supplementary motor area (SMA) (Kasess et al., 2008) or the right inferior frontal 89 gyrus (rIFG) may weaken the motor commands that are emitted during motor imagery 90 (e.g., Angelini et al., 2015, 2016). More precisely, the pre-SMA and the rIFG may work 91 together to intercept the action process via the basal ganglia (subthalamic nucleus, STN), 92 hence suppressing the output from the basal ganglia which in turn might inhibit the 93 primary motor cortex (Aron, 2011) (cf. Figure 1). In addition to inter-regional inhibitory 94 influences, motor imagery has been shown to be associated with modulations of 95 short-interval intracortical inhibition within the primary motor cortex itself (Neige et al., 96 2020). Third, downstream regions in the cerebellum (e.g., Lotze et al., 1999), in the 97 brainstem (e.g., Jeannerod, 2001, 2006a), or at the spinal level may contribute to motor 98 inhibition at a later stage. 99

Whether motor commands are emitted and subsequently inhibited during covert 100 speech production is an issue that has been debated (e.g., Geva, 2018). It has been 101 suggested that during covert speech production, motor commands would be "simply 102 specified in sub-threshold way, requiring no active inhibition" (Geva, 2018). Alternatively, 103 motor commands may be emitted but subsequently inhibited by frontal regions (e.g., 104 Grandchamp et al., 2019; Lœvenbruck et al., 2018). However, stating that covert speech 105 (or motor imagery, more generally) only involves subthreshold activity (and therefore is 106 not accompanied by the emission of motor commands that are inhibited) simply shifts 107 the problem from "how and where motor commands are subsequently inhibited" to "how 108 and where the magnitude of activity in the motor system is planned or monitored" (see 109 also Scheil & Liefooghe, 2018). In other words, we still need to explain how (in a 110 mechanistic and/or developmental way) this activity is maintained at a subthreshold 111 level. What cognitive and neural mechanisms operate to maintain this activity under the 112



Figure 1

Plausible implementation of the cortical and subcortical inhibitory mechanisms responsible for the "proactive" (but implicit) response inhibition at play during covert speech production. The pre-SMA, posterior rIFC, and STN together form an inhibitory network known as the inhibitory triangle, which may be responsible for braking motor commands during covert speech production. Figure created with BioRender.com.

threshold? In this section, we provided empirical arguments in favour of the "active
inhibition hypothesis". Proponents of the "subliminal level hypothesis" need to clarify
how this activity is maintained at a subthreshold level during covert speech production,
thus preventing execution.

The putative involvement and functional role of inhibitory mechanisms during 117 covert speech could be assessed in several ways. First, it could be assessed by 118 experimentally manipulating the activity of the inhibitory network responsible for 119 preventing execution during motor imagery. For instance, transcranial magnetic 120 stimulation (TMS) could be used to interrupt these inhibitory mechanisms and thus 121 facilitate execution during motor imagery. Second, the role of inhibitory mechanisms 122 during covert verbal actions could be examined in populations with well-identified 123 inhibitory deficits. For instance, Tourette syndrome is a childhood-onset neurological 124

disorder affecting approximately 1% of children and characterised by chronic motor and 125 phonic tics (Jackson et al., 2015). Verbal tics can consist of repeating sounds, words, or 126 utterances (palilalia), producing inappropriate or obscene utterances (coprolalia), or the 127 repetition of another's words (echolalia). In their review, Jackson et al. (2015) suggested 128 that increased control over motor outputs, acquired by repeatedly trying to suppress tics, 129 is brought about by local increases in GABAergic "tonic" inhibition within regions such 130 as the SMA, leading to localised reductions in the gain of motor excitability. For these 131 reasons, comparing the neural implementation of inhibitory mechanisms during covert 132 speech in patients with Tourette syndrome and healthy controls may shed light on the 133 role and flexibility of these mechanisms. 134

¹³⁵ 2.2 Covert speech development: Learning not to produce speech

Watson (1919) suggested that thought was rooted in overt speech. In his 136 terminology, thought referred to covert speech. Hence, his view was that covert speech 137 matures from overt speech. Vygotsky (1934) further formulated the idea that covert 138 speech is internalised during childhood from private egocentric speech, that is, from 139 self-addressed overt speech. Fernyhough (2004) extended these ideas by proposing four 140 levels of internalisation: external dialogue, private speech, expanded inner speech, and 141 condensed inner speech. These levels represent stages of development but also define 142 movements between levels, that is, how a speaker may go from overt to covert speech and 143 conversely. The level at which speech is expressed may depend on inhibitory control 144 applied at different levels in the production flow, such as the formulation or the 145 articulatory planning level (Grandchamp et al., 2019). Therefore, producing covert 146 speech crucially depends on successfully inhibiting speech production at several levels. 147

We formulate the hypothesis that the progressive internalisation of speech during childhood may be related to the development of inhibitory abilities. This hypothesis could be tested in several ways. First, the relation between speech internalisation and inhibitory abilities could be assessed during development at the critical ages (i.e., between 6 and 8 years). We would expect the ability to imagine actions, and speech specifically, to

be positively correlated with inhibitory skills at this age. Wang et al. (2021) provided 153 correlational evidence that motor imagery (as assessed by a hand laterality judgement 154 task and alphanumeric rotation task) and motor inhibition performance (as assessed by a 155 stop-signal task) improve together between 7 and 11 years old, and that these two 156 abilities correlated at 7 years old but did not correlate at 11 years old. This suggests that 157 inhibitory control may play a peculiar role when speech is being internalised, but its role 158 may weaken with expertise. This is consistent with results from training studies 159 suggesting that, with growing expertise, we tend to rely more and more on memory-based 160 processes to produce mental imagery (e.g., Jolicoeur, 1985; Tarr & Pinker, 1989). 161

Second, the hypothesised co-development of motor imagery and response 162 inhibition abilities could be tested by examining how novel actions are internalised in 163 adults. For instance, let's consider the analogy between speaking and playing a music 164 instrument (e.g., playing the piano). The act of producing speech can be paralleled with 165 the act of playing an instrument. Both actions consist in the coordination of complex 166 movements that result in some modifications of the environment, that in turn generate 167 sensory feedback (e.g., kinaesthetic, auditory) for the agent. This analogy suggests that 168 we might be able to study the development of internal models responsible for the sensory 169 experience accompanying imagined actions in the adult mind (e.g., when an individual is 170 learning either a novel music instrument or a new language with speech sounds that are 171 not present in his/her native language). By examining the development of novel imagined 172 actions in the adult mind and by using motor interference (e.g., articulatory suppression) 173 procedures, we might gain new insights about the internalisation of speech during 174 childhood.¹ 175

¹ While keeping in mind the obvious limitation that the child mind is not equivalent to the adult mind, nor is it equivalent to a smaller version of the adult mind. Nevertheless, examining the development of novel imagined actions in adults avoids the contamination of the process of interest (imagined action) by developmental confounds.

176 2.3 Does covert speech always involve motor inhibition?

The production of covert speech is often, although not always and not for everyone, accompanied by the feeling of hearing speech (Hurlburt, 2011). In other words, covert speech is accompanied by a sensory experience that "feels like" hearing speech. However, covert speech may also be accompanied by the feeling of producing speech. These two facets of covert speech are characterised by different phenomenological experiences. In this section, we discuss how these two forms of covert speech may require motor inhibition to a different extent.

The dual stream prediction model (Tian & Poeppel, 2012, 2013; Tian et al., 2016) 184 describes two neural pathways that may provide the auditory content of covert speech. 185 First, the simulation-estimation prediction stream implements a motor-to-sensory 186 transformation via motor simulation, that is, by simulating speech movements and the 187 perceptual changes that would be associated with these movements (see also Lœvenbruck 188 et al., 2018, for a similar proposal). This stream includes cerebral areas involved in 189 speech motor preparation such as the supplementary motor area, the inferior frontal 190 gyrus, the premotor cortex and the insula, as well as brain areas involved in 191 somatosensory estimation and perception such as primary and secondary somatosensory 192 regions, the parietal operculum, and the supramarginal gyrus (Tian et al., 2016). Second, 193 the memory-retrieval prediction stream may be used to provide auditory percepts by 194 "reconstructing stored perceptual information in modality-specific cortices" (Tian et al., 195 2016). This mechanism provides sensory percepts without the need for computing the 196 predicted sensory consequences of (non executed) motor commands. Auditory percepts 197 may be retrieved from various memory sources, relying (amongst others) on the 198 hippocampal structures (Tian et al., 2016). Auditory percepts may also be retrieved from 199 a broad fronto-temporo-parietal lexico-semantic network (for more details, see Tian et al., 200 2016). 201

The balance between these two mechanisms (i.e., simulation vs. memory retrieval) may depend on the circumstances promoting covert speech or, in the lab, on the precise instructions given to the participants, which may cue them to produce different forms of

9

covert speech. For instance, either one of these two streams may be preferentially 205 recruited depending on whether participants are instructed to "imagine speaking" or to 206 "imagine hearing" (see also the distinction between the "inner ear" and the "inner voice", 207 e.g., Smith et al., 1992). In line with this hypothesis, Tian et al. (2016) have shown that 208 inner speaking more strongly recruits brain regions in the simulation stream than inner 209 hearing, which more strongly recruits brain regions in the memory-retrieval stream. Ma 210 and Tian (2019) have shown that inner speaking and inner hearing have distinct 211 magnetoencephalographic (MEG) correlates and distinct effects on a subsequent phonetic 212 categorisation task (discriminating /ba/ vs. /da/). 213

In line with Tian and Poeppel (2012), we suggest that the balance between these 214 two mechanisms may also depend on situational (e.g., surrounding noise) and intrinsic 215 (e.g., expertise) characteristics. We further suggest that a common currency determining 216 the use of either one of these mechanisms is the computational cost of (or equivalently, 217 the computational resources available for) each alternative. To clarify, we borrow the 218 concept of memoisation as applied to cognition and mental imagery by Dasgupta and 219 Gershman (2021) (cf. Box 1). In these authors' view, memory can be considered as a 220 computational resource to facilitate computational reuse through memoisation. In the 221 context of motor and speech imagery, memoisation can be seen in the increasing reliance 222 on memory in the course of learning. 223

In other words, situational (extrinsic) and individual (intrinsic) characteristics 224 jointly determine the computational cost of (or equivalently, the available computational 225 resources for) the task, which in turn determines the balance between the simulation and 226 association mechanisms. For instance, we hypothesise that novel and/or difficult tasks 227 (which are both computationally more expensive, ceteris paribus) may rely more on the 228 simulation mechanism, whereas well known and/or easy tasks may rely more on 229 associative mechanisms. This idea is supported by several studies showing a greater 230 increase in facial EMG activity during the reading of difficult text or while performing 231 difficult mental arithmetic tasks, compared to easier tasks (e.g., Faaborg-Andersen et al., 232 1958; Sokolov, 1972), suggesting a greater involvement of the speech motor system. 233

Box 1: Memoisation

Memoisation is a programming technique used to speed-up algorithms or programs. It avoids redundant computation by storing computational results and reusing them later. When calling a function (where a function can be a motor primitive), the function call is intercepted by a *memoiser* that inspects the previous calls of a function and its outputs. If a function has already been called with the same input, then the previously computed output is retrieved and reused.

In the context of covert speech, memoisation can be postulated as the process by which covert speech percepts produced by motor simulation are stored for later retrieval and use without invoking the motor simulation mechanism.

Alternatively, these results may suggest a lesser involvement of inhibitory mechanisms (see also the discussion in Nalborczyk, 2019, 2020). This is congruent with the greater reliance on associative mechanisms with greater expertise, as discussed previously.

To sum up, whereas inner speaking may involve active inhibition of motor 237 commands, inner hearing may not. These disparities between inner speaking and inner 238 hearing may explain the variety of neural correlates reported for covert speech production 239 (as reviewed for instance in Geva, 2018). More generally, different forms of covert speech, 240 that may vary in condensation (from thinking without words to thinking in words), 241 dialogicality (whether covert speech features monologues or dialogues), or intentionality 242 (for more details, see Grandchamp et al., 2019), may require inhibitory control to a 243 different extent, from no inhibition at all for condensed forms of covert speech to active 244 inhibition of motor commands for fully expanded forms of covert speech. 245

246

3 Conclusions

We explored some of the theoretical and experimental consequences that emerge from considering covert speech production as an inhibited form of overt speech production. To this end, we connected results from the motor imagery, motor inhibition, and covert speech domains. Regarding the role and implementation of general-purpose inhibitory mechanisms during the production of covert speech, we suggested that these may be similar to the inhibitory network responsible for proactive response inhibition and we summarised some propositions from this literature. We related the development of response inhibition abilities in childhood development with the purported internalisation of private speech around the same period. From the response inhibition perspective, the internalisation of speech from overt to covert speech may essentially be considered as "learning not to produce speech".

Regarding the neural origin of the sensory experience of covert speech, we 258 discussed the dual stream prediction model (Tian & Poeppel, 2012, 2013; Tian et al., 259 2016), which suggests that these sensory percepts may be provided either by a 260 motor-simulation process or by a memory-retrieval process. We suggested that the 261 balance between these two mechanisms may be determined by task instructions, which 262 may prompt different forms of covert speech, and also by the computational cost of the 263 task. More precisely, novel or more difficult tasks are expected to rely more on the 264 motor-simulation mechanisms whereas well-known and/or easy tasks may rely more on a 265 "memoised version" of the motor simulation: the memory-retrieval prediction stream. 266 Whereas the former mechanism should involve active inhibitory mechanisms, the latter 267 should not, as there should be no (or less) motor commands to inhibit. 268

These propositions pave the way for several lines of research that should 269 consolidate our understanding of the relations between overt and covert speech 270 production. Several outstanding questions remain. Amongst others, how does the 271 development of inhibitory control relate with the progressive internalisation of speech 272 during childhood? How do individual and situational constraints shape the role of motor 273 inhibition during covert speech production? How is covert speech affected by poor or 274 degraded inhibitory control? Can we experimentally force the externalisation of speech in 275 adults, for example through neurostimulation? The use of neurostimulation and the 276 comparison between healthy controls and patients with well-identified inhibitory deficits 277 could help refine the involvement of these inhibitory mechanisms during covert speech 278 production, which may lead to applied outcomes in the care of motor and verbal tics. 279

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

284

Author Contributions

²⁸⁵ Conceptualisation: all authors; Funding acquisition: LN, ML, FXA; Supervision:
²⁸⁶ UD, ML, AG, FXA; Writing - original draft: LN; Writing - review and editing: all
²⁸⁷ authors.

Funding
 This work, carried out within the Institut Convergence ILCB
 (ANR-16-CONV-0002), has benefited from support from the French government,
 managed by the French National Agency for Research (ANR) and the Excellence

²⁹² Initiative of Aix-Marseille University (A*MIDEX).

293

Data Availability Statement

296	References
297	Alderson-Day, B. & Fernyhough, C. (2015). Inner speech: Development, cognitive
298	functions, phenomenology, and neurobiology. Psychological Bulletin, $141(5)$,
299	931–965. https://doi.org/10.1037/bul0000021
300	Angelini, M., Calbi, M., Ferrari, A., Sbriscia-Fioretti, B., Franca, M., Gallese, V. &
301	Umiltà, M. A. (2015). Motor Inhibition during Overt and Covert Actions: An
302	Electrical Neuroimaging Study (F. Di Russo, Ed.). PLOS ONE, $10(5)$, e0126800.
303	https://doi.org/10.1371/journal.pone.0126800
304	Angelini, M., Calbi, M., Ferrari, A., Sbriscia-Fioretti, B., Franca, M., Gallese, V. &
305	Umiltà, M. A. (2016). Proactive Control Strategies for Overt and Covert
306	Go/NoGo Tasks: An Electrical Neuroimaging Study (L. Jaencke, Ed.). $PLOS$
307	ONE, 11(3), e0152188. https://doi.org/10.1371/journal.pone.0152188
308	Aron, A. R. (2007). The Neural Basis of Inhibition in Cognitive Control: The
309	<i>Neuroscientist.</i> https://doi.org/10.1177/1073858407299288
310	Aron, A. R. (2011). From Reactive to Proactive and Selective Control: Developing a
311	Richer Model for Stopping Inappropriate Responses. Biological Psychiatry, $69(12)$,
312	e55–e68. https://doi.org/10.1016/j.biopsych.2010.07.024
313	Baldo, J. V., Dronkers, N. F., Wilkins, D., Ludy, C., Raskin, P. & Kim, J. (2005). Is
314	problem solving dependent on language? Brain and Language, $92(3)$, 240–250.
315	https://doi.org/10.1016/j.bandl.2004.06.103
316	Bart, V. K. E., Koch, I. & Rieger, M. (2021). Inhibitory mechanisms in motor imagery:
317	Disentangling different forms of inhibition using action mode switching.
318	Psychological Research, 85(4), 1418–1438.
319	https://doi.org/10.1007/s00426-020-01327-y
320	D'Argembeau, A., Renaud, O. & der Linden, M. V. (2011). Frequency, characteristics and
321	functions of future-oriented thoughts in daily life. Applied Cognitive Psychology,
322	25(1), 96-103. https://doi.org/10.1002/acp.1647
323	_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/acp.1647

324	Dasgupta, I. & Gershman, S. J. (2021). Memory as a Computational Resource. Trends in
325	Cognitive Sciences, 25(3), 240–251. https://doi.org/10.1016/j.tics.2020.12.008
326	Faaborg-Andersen, K., Edfeldt, Å. W. & Nykøbing, F. (1958). Electromyography of
327	intrinsic and extrinsic laryngeal muscles during silent speech: Correlation with
328	reading activity: Preliminary report. Acta Oto-Laryngologica, $49(1)$, 478–482.
329	https://doi.org/10.3109/00016485809134778
330	Fernyhough, C. (2004). Alien voices and inner dialogue: Towards a developmental account
331	of auditory verbal hallucinations. New Ideas in Psychology, $22(1)$, 49–68.
332	https://doi.org/10.1016/j.newideapsych.2004.09.001
333	Frith, U. (1979). Reading by Eye and Writing by Ear. In P. A. Kolers, M. E. Wrolstad &
334	H. Bouma (Eds.), Processing of Visible Language (pp. 379–390). Springer US.
335	$https://doi.org/10.1007/978-1-4684-0994-9_23$
336	Geva, S. (2018). Inner speech and mental imagery: A neuroscientific perspective. In
337	P. Langland-Hassan & A. Vicente (Eds.), Inner Speech: New Voices (pp. 150–130).
338	Oxford University Press.
339	Grandchamp, R., Rapin, L., Perrone-Bertolotti, M., Pichat, C., Haldin, C., Cousin, E.,
340	Lachaux, JP., Dohen, M., Perrier, P., Garnier, M., Baciu, M. & Lœvenbruck, H.
341	(2019). The ConDialInt Model: Condensation, Dialogality, and Intentionality
342	Dimensions of Inner Speech Within a Hierarchical Predictive Control Framework.
343	Frontiers in Psychology, 10. https://doi.org/10.3389/fpsyg.2019.02019
344	Guillot, A., Di Rienzo, F., MacIntyre, T., Moran, A. & Collet, C. (2012). Imagining is not
345	doing but involves specific motor commands: A review of experimental data
346	related to motor inhibition. Frontiers in Human Neuroscience, 6.
347	https://doi.org/10.3389/fnhum.2012.00247
348	Hurlburt, R. T. (2011). Investigating pristine inner experience: Moments of truth. C. U.
349	Press, Ed.
350	Jackson, G. M., Draper, A., Dyke, K., Pépés, S. E. & Jackson, S. R. (2015). Inhibition,
351	Disinhibition, and the Control of Action in Tourette Syndrome. Trends in
352	Cognitive Sciences, 19(11), 655–665. https://doi.org/10.1016/j.tics.2015.08.006

353	Jeannerod, M. (1994). The representing brain: Neural correlates of motor intention and
354	imagery. Behavioral and Brain Sciences, 17(02), 187.
355	https://doi.org/10.1017/S0140525X00034026
356	Jeannerod, M. (2001). Neural simulation of action: A unifying mechanism for motor
357	cognition. NeuroImage, 14(1), S103–S109. https://doi.org/10.1006/nimg.2001.0832
358	Jeannerod, M. (2006a). Motor cognition: What actions tell the self. Oxford University
359	Press
360	OCLC: ocm65195268.
361	Jeannerod, M. (2006b). The origin of voluntary action. History of a physiological concept.
362	Comptes Rendus Biologies, 329(5-6), 354–362.
363	https://doi.org/10.1016/j.crvi.2006.03.017
364	Jolicoeur, P. (1985). The time to name disoriented natural objects. Memory & Cognition,
365	13(4), 289-303. https://doi.org/10.3758/BF03202498
366	Kasess, C. H., Windischberger, C., Cunnington, R., Lanzenberger, R., Pezawas, L. &
367	Moser, E. (2008). The suppressive influence of SMA on M1 in motor imagery $% \left({{\rm{M}}_{\rm{m}}} \right)$
368	revealed by fMRI and dynamic causal modeling. NeuroImage, $40(2)$, 828–837.
369	https://doi.org/10.1016/j.neuroimage.2007.11.040
370	Lœvenbruck, H., Baciu, M., Segebarth, C. & Abry, C. (2005). The left inferior frontal
371	gyrus under focus: An fMRI study of the production of deixis via syntactic
372	extraction and prosodic focus. Journal of Neurolinguistics, 18(3), 237–258.
373	https://doi.org/10.1016/j.jneuroling.2004.12.002
374	Lœvenbruck, H., Grandchamp, R., Rapin, L., Nalborczyk, L., Dohen, M., Perrier, P.,
375	Baciu, M. & Perrone-Bertolotti, M. (2018). A cognitive neuroscience view of inner
376	language: To predict and to hear, see, feel. In P. Langland-Hassan & A. Vicente
377	(Eds.), Inner speech: New voices (p. 37). Oxford University Press.
378	Lotze, M., Montoya, P., Erb, M., Hülsmann, E., Flor, H., Klose, U., Birbaumer, N. &
379	Grodd, W. (1999). Activation of Cortical and Cerebellar Motor Areas during
380	Executed and Imagined Hand Movements: An fMRI Study. Journal of Cognitive
381	Neuroscience, 11(5), 491–501. https://doi.org/10.1162/089892999563553

382	Ma, O. & Tian, X. (2019). Distinct mechanisms of imagery differentially influence speech
383	perception. $eNeuro$. https://doi.org/10.1523/ENEURO.0261-19.2019
384	MacLeod, C. M. (2007). The concept of inhibition in cognition. In D. S. Gorfein $\&$
385	C. M. MacLeod (Eds.), Inhibition in cognition. (pp. 3–23). American
386	Psychological Association. https://doi.org/10.1037/11587-001
387	Nalborczyk, L. (2019). Understanding rumination as a form of inner speech: Probing the
388	role of motor processes (PhD Thesis). Univ. Grenoble Alpes & Ghent University.
389	Nalborczyk, L. (2020). Re-analysing the data from Moffatt et al. (2020): A textbook
390	illustration of the absence of evidence fallacy. PsyArXiv.
391	https://doi.org/10.31234/osf.io/9j76v
392	Neige, C., Rannaud Monany, D., Stinear, C. M., Byblow, W. D., Papaxanthis, C. &
393	Lebon, F. (2020). Unravelling the Modulation of Intracortical Inhibition During
394	Motor Imagery: An Adaptive Threshold-Hunting Study. Neuroscience, 434,
395	102–110. https://doi.org/10.1016/j.neuroscience.2020.03.038
396	O'Shea, H. & Moran, A. (2018). To go or not to go? Pupillometry elucidates inhibitory
397	mechanisms in motor imagery. Journal of Cognitive Psychology, $30(4)$, 466–483.
398	https://doi.org/10.1080/20445911.2018.1461104
399	Perrone-Bertolotti, M., Rapin, L., Lachaux, J. P., Baciu, M. & Lœvenbruck, H. (2014).
400	What is that little voice inside my head? Inner speech phenomenology, its role in
401	cognitive performance, and its relation to self-monitoring. Behavioural Brain
402	Research, 261, 220–239. https://doi.org/10.1016/j.bbr.2013.12.034
403	Perrone-Bertolotti, M., Kujala, J., Vidal, J. R., Hamame, C. M., Ossandon, T.,
404	Bertrand, O., Minotti, L., Kahane, P., Jerbi, K. & Lachaux, JP. (2012). How
405	Silent Is Silent Reading? Intracerebral Evidence for Top-Down Activation of
406	Temporal Voice Areas during Reading. Journal of Neuroscience, 32(49),
407	17554–17562. https://doi.org/10.1523/JNEUROSCI.2982-12.2012
408	Ridderinkhof, K. R., van den Wildenberg, W. P. M. & Brass, M. (2014). "Don't" versus
409	"Won't": Principles, mechanisms, and intention in action inhibition.

MOTOR INHIBITION AND COVERT SPEECH

410	Neuropsychologia, 65, 255–262.
411	https://doi.org/10.1016/j.neuropsychologia.2014.09.005
412	Rieger, M., Dahm, S. F. & Koch, I. (2017). Inhibition in motor imagery: A novel action
413	mode switching paradigm. Psychonomic Bulletin & Review, $24(2)$, 459–466.
414	https://doi.org/10.3758/s13423-016-1095-5
415	Scheil, J. & Liefooghe, B. (2018). Motor command inhibition and the representation of
416	response mode during motor imagery. Acta Psychologica, 186, 54–62.
417	https://doi.org/10.1016/j.actpsy.2018.04.008
418	Smith, J. D., Reisberg, D. & Wilson, M. (1992). Subvocalization and auditory imagery:
419	Interactions between the inner ear and inner voice. Auditory imagery (pp. 95–119).
420	Lawrence Erlbaum Associates, Inc.
421	Sokolov, A. (1972). Inner speech and thought. Springer-Verlag
422	OCLC: 948758651.
423	Tarr, M. J. & Pinker, S. (1989). Mental rotation and orientation-dependence in shape
424	recognition. Cognitive Psychology, 21(2), 233–282.
425	https://doi.org/10.1016/0010-0285(89)90009-1
426	Tian, X. & Poeppel, D. (2012). Mental imagery of speech: Linking motor and perceptual
427	systems through internal simulation and estimation. Frontiers in Human
428	<i>Neuroscience</i> , 6. https://doi.org/10.3389/fnhum.2012.00314
429	Tian, X. & Poeppel, D. (2013). The effect of imagination on stimulation: The functional
430	specificity of efference copies in speech processing. Journal of Cognitive
431	Neuroscience, $25(7)$, 1020–1036. https://doi.org/10.1162/jocn_a_00381
432	Tian, X., Zarate, J. M. & Poeppel, D. (2016). Mental imagery of speech implicates two
433	mechanisms of perceptual reactivation. Cortex, 77, 1–12.
434	https://doi.org/10.1016/j.cortex.2016.01.002
435	Vygotsky, L. S. (1934). Thought and language, revised and expanded edition (A. Kozulin,
436	Trans.). The MIT Press.
437	Wang, C., Li, W., Zhou, Y., Nan, F., Zhao, G. & Zhang, Q. (2021). The relationship
438	between internal motor imagery and motor inhibition in school-aged children: A

MOTOR INHIBITION AND COVERT SPEECH

- 439 cross-sectional study. Advances in Cognitive Psychology, 17, 88–98.
- 440 https://doi.org/10.5709/acp-0319-9
- 441 Watson, J. B. (1919). Psychology from the standpoint of a behaviorist. J B Lippincott
- 442 Company. https://doi.org/10.1037/10016-000