



Therapeutic effects of large-field visual virtual immersion on balance control in unilateral vestibular patients

Olivier A.J. Martin, Denis Faure-Vincent, Jean-Dominique Gascuel, Sébastien Schmerber, Alina Voda, Pascal Bellemain

► To cite this version:

Olivier A.J. Martin, Denis Faure-Vincent, Jean-Dominique Gascuel, Sébastien Schmerber, Alina Voda, et al.. Therapeutic effects of large-field visual virtual immersion on balance control in unilateral vestibular patients. ISPGR 2022 - International Society of Posture and Gait Research, Jul 2022, Montréal, Canada. pp.1-1. hal-03737948

HAL Id: hal-03737948

<https://hal.science/hal-03737948>

Submitted on 25 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Therapeutic effects of large-field visual virtual immersion on balance control in unilateral vestibular patients

Martin O.^{1*}, Faure-Vincent D.¹, Gascuel JD.², Schmerber S.³, Voda A.¹, Bellemain P.¹

¹GIPSA-lab, UMR5216 Grenoble-Alpes University & G-INP, Grenoble France

²INRIA Rhône-Alpes, Maverick Team & Jean Kunzman Lab, Grenoble, France


³Grenoble-Alpes University Hospital, NET Clinic, Grenoble, France

(*Contact : olivier.martin@gipsa-lab.grenoble-inp.fr)

INTRODUCTION | VR-based balance rehab. relevance ?



3. OBJECTIVES




VR REHABILITATION



- Balance processes use visual, vestibular and proprioceptive sensory signals in a synergistic way. [1]

- Vestibular deficit increases a visual dependency due to the vestibular inhibition suppression [2].



- Therapeutic effects of virtual reality (VR) based protocols (optical flows) to restore the reciprocal visual-vestibular-motor to reduce the patients' visual dependency.





RESULTS | VR immersion → Visual perturbation → Balance restoration





- INTER-SESSIONS effect (representative patient)**
- Decreases the Cop Trajectory length
 - Reduces the postural reaction
 - Decrease the balance perturbation
 - Improve the Gaze-Balance interactions



- Flows SPEED effect (all patients)**
- Increase the Cop Trajectory length
 - Increase the balance perturbations
 - Stimulate the compensatory postural adjust.




- ISPEED x STM.1 effect on AP-ML adjustments (all patients)**
- [Speed max. + 2/3 stim.] > [Speed min. + 1D stim.]
 - High [Speed x Stim.] increase the CoP Trajectory length → related to the augmented complexity of the virtual stim.
 - Increase the balance perturbations
 - Multiple visual flows parameters "co-stimulate" the compensatory postural strategies



- ISPEED x STM.1 effect on AP-ML adjustments (all patients)**
- [Speed max. + 2/3 stim.] > [Speed min. + 1D stim.]
 - High [Speed x Stim.] increase the CoP Trajectory length → related to the augmented complexity of the virtual stim.
 - Increase the balance perturbations
 - Multiple visual flows parameters "co-stimulate" the compensatory postural strategies

GAZE ANCHORAGE (GA)

- Visuo-spatial reference
- With GA : session 1-4 ; Without GA : session 5-8
- [With] : session 9 ; [Without] : session 10



INTRA-SESSIONS effect (all patients)

- Increase the CoP Trajectory length
- Increase of the postural reaction
- "Positive fatigue effect" ? → Optimal Difficulty Level ?

- GAZE-BALANCE INTERACTION (representative patient)**
- Functionally (neuro-mechanically) coupled and synchronized
 - Synergistic improvement and interactive stabilisation
 - Gaze Balance enhancement ... ocular OR vestibular efficiency thanks to the visual-retinal flows ... Which sub-system drives the improvement ?

PROTOCOL

1 Trial = Visual stimulation

• Rest – Flow – Rest : 25 sec.

1 Serie = 8 stim. (8 trials)

• 5 different 1D/2D-3D visual flows

1 Session = 6 Speed

• 8 trials x 6 speed = 48 trials

2 Gaze Anchorage cond.

• Sess. 1-4 : With GA

• Sess. 5-8 : Without GA

• (48 trials x 8) = 384 trials (from 15 sec.)

→ Total stim. duration = 96 min. (1h36)


[2 Phases = Compar./Valid.]

• 8 VR-based sess. (PIVVIT)

→ "Proof of concept" +++

• 8 Standard OKS sess.

→ "Proof of non-inferiority" +++



DATA RECORDING AND ANALYSIS

- Balance control : Center-of-Pressure (Wiifit board, 100 Hz) : **Cop Trajectory length** → Disequilibrium indicators (lost of balance control, falls)

- Gaze behaviour (Eye-Head-body) stability ; BlueGain EOG, 100Hz) : **EOGxy** trajectory (preliminary results)

- [Questions-Responses] (subjective estimation of the "balance comfort" in every day life tasks)

DISCUSSION – CONCLUSION | Rapid VR-based rehab. benefits

At a Clinical/Rehab. level : Positive effects of the "large visual field" immersion on balance restoration

➢ METH. Patient-related sessions repetition + difficulty-dependent tuneable scenarios + patient follow-up + methodical practice through short period (**two months**)

➢ BENEFITS. Rapid functional restoration of efficient synergistic control between balance and gaze process.

➢ PSYCHOL. Patients' motivation and self-confidence !

At a Neuro-functional level : "visual-vestibular-motor" adaptation occurs despite the vestibular deficits

➢ PROCESS. Reweighting of the visual-vestibular sensory inputs → reciprocal inhibition restoration

➢ ADAPTATION. Reduction of the visual dependence by visual-vestibular reverse compensations

➔ Questions about the neural basis of the "visual-vestibular plasticity"

➢ Sensory adaptive threshold/gain of the VOR and/or OKN ?

➢ Dual sensorimotor plasticity in balance restoration : Interaction MT visual area - vestibular nuclei → visual-vestibular integration for motor command tuning/adjustment.

CONCLUSION

➢ Optimized rehabilitation strategies based on simple virtual visual immersions Vs. standardized rehabilitation protocols (uncontrolled OKS) for unilateral vestibular patient

➢ Limit: No long-term effects (reported by patients after a six months period) requiring periodic booster sessions

➢ Our study corroborate the proof of concept of interactive VR-rehabilitation based on a large visual field stimulation device (experimental and clinical validation).

REFERENCES. [1] Peterka, R. J. J. Neurophysiol, 2002. [2] Roberts et al., Adv. Clin. Neurosci. Rehab. 2016. [3] Voda et al., IFAC PapersOnLine 2019.

ACKNOWLEDGEMENTS. Thanks to all patients, and to the Grenoble University Hospital (CHUGA NET Clinic) – Grant DRCI (Dept. Clinical Research & Innovation) : PIVVIT Project #1045.

