

Cord blood leptin level and a common variant of its receptor as determinants of the BMI trajectory: The EDEN mother-child cohort

Aminata Cissé, Marion Taine, Muriel Tafflet, Blandine de Lauzon-Guillain, Karine Clément, Olfa Khalfallah, Laetitia Davidovic, Sandrine Lioret, Marie Charles, Barbara Heude

▶ To cite this version:

Aminata Cissé, Marion Taine, Muriel Tafflet, Blandine de Lauzon-Guillain, Karine Clément, et al.. Cord blood leptin level and a common variant of its receptor as determinants of the BMI trajectory: The EDEN mother–child cohort. Pediatric Obesity, In press, 17 (11), pp.e12955. 10.1111/ijpo.12955. hal-03739236

HAL Id: hal-03739236 https://hal.science/hal-03739236

Submitted on 29 Jul2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title:

Cord blood leptin level and a common variant of its receptor as determinants of the BMI trajectory: the EDEN mother-child cohort

Running title: Cord leptin as a determinant of peak and rebound

Authors' names:

Aminata Hallimat Cissé¹, Marion Taine¹, Muriel Tafflet¹, Blandine de Lauzon-Guillain¹, Karine Clément², Olfa Khalfallah³, Laetitia Davidovic³, Sandrine Lioret¹, Marie Aline Charles¹, Barbara Heude¹

Affiliations:

¹Université de Paris, Inserm, INRAE, Centre for Research in Epidemiology and StatisticS (CRESS), F-75004 Paris, France

²Sorbonne Université, INSERM, NutriOmics Research Unit, Assistance Publique-Hôpitaux de Paris, Pitié □ Salpêtrière Hopital, Nutrition department Paris, France
³Institut de Pharmacologie Moléculaire et Cellulaire, CNRS, INSERM, Université Nice Côte d'Azur, UMR7275, UMR_S, Valbonne, France

Corresponding author:

Address: Aminata CISSE, INSERM U1153, Team 6 EAROH 16 Avenue Paul Vaillant Couturier, Bâtiment Leriche 94807 Villejuif Cedex, France Phone: + 33 145-595-194 Email: <u>aminata-hallimat.cisse@inserm.fr</u> Abbreviations list AP: Adiposity peak AR: Adiposity rebound BMI: Body mass index EDEN: Etude des Déterminants pré et post natals de la santé de l'Enfant *LEPR*: Leptin receptor SNP: Single nucleotide polymorphism

1 Abstract (200/200)

Background: Cord blood leptin is an indicator of neonatal fat mass and could shape
postnatal adiposity trajectories. Investigating genetic polymorphisms of the leptin receptor
gene (*LEPR*) could help understand the mechanisms involved.

5 Objectives: We aimed to investigate the association of cord blood leptin level and the *LEPR*6 rs9436303 polymorphism, with BMI at adiposity peak (AP) and age at adiposity rebound
7 (AR).

8 **Methods:** In the EDEN cohort, BMI at AP and age at AR were estimated with polynomial 9 mixed models, for 1713 and 1415 children, respectively. Multivariable linear regression 10 models allowed for examining the associations of cord blood leptin level and *LEPR* 11 rs9436303 genotype with BMI at AP and age at AR adjusted for potential confounders 12 including birth size groups. We also tested interactions between cord blood leptin level and 13 rs9436303 genotype.

14 **Results:** Increased leptin level was associated with reduced BMI at AP and early age at AR 15 (comparing the highest quintile of leptin level to the others). Rs9436303 G-allele carriage was 16 associated with increased BMI at AP and later age at AR but did not modulate the association 17 with leptin level.

18 Conclusion: These results illustrate the role of early life body composition and the 19 intrauterine environment in the programming of adiposity in childhood.

Keywords: Cord blood leptin, SNP rs9436303, BMI at adiposity peak, age at adiposity
rebound, EDEN cohort.

3

23 The body mass index (BMI) trajectory during infancy and childhood is one main predictor of 24 later cardio-metabolic health (1,2) and comprises two key periods. Over the first year of life, 25 the BMI increases to a maximum, called adiposity peak (AP), at about 9 months of age. It 26 then decreases to a minimum and starts to increase again between 4 and 6 years of age, on 27 average, which corresponds to the adiposity rebound (AR) period (3,4). In epidemiological study, higher BMI at AP and earlier age at AR seem to be relevant predictors of high-risk 28 29 trajectories to later obesity (5–7). These factors are affected by child sex, parental educational 30 level, genetic susceptibility to obesity and parental BMI (8-11). We also previously showed 31 that being born small for gestational age (SGA) and higher maternal gestational weight gain 32 during pregnancy were additional predictors of age at AR (12). These latter findings highlight the important role of intrauterine life in postnatal growth and these prenatal conditions are 33 34 likely involved in the fetal programming of the adiposity trajectory.

35 Hormones such as leptin are probably involved in this programming phenomenon and 36 measuring their levels in cord blood may help predict later adiposity trajectories (13,14). 37 Leptin is an adipokine secreted mainly by adipocytes but also by enterocytes in the small 38 intestine and several tissues such as placenta (14). The placenta is one of the main sites for 39 leptin production during pregnancy, and part of the leptin produced in the placenta passes into 40 the fetal circulation. As a consequence, cord blood leptin consists of leptin secreted by fetal 41 adipocytes and placenta but in proportions still unknown (15). Children born SGA exhibit a 42 low level of leptin in cord blood, a low percentage of body fat and higher catch-up growth 43 than others (16-18). Leptin measured in cord blood is therefore considered a marker of 44 neonatal fat mass and is possibly involved in early postnatal growth (19,20). In addition, 45 studies have shown that placental leptin production is increased in severe conditions

46 associated with fetal growth abnormalities, such as diabetes and pre-eclampsia (21,22).
47 However, to our knowledge, no study has investigated the role of leptin in the trajectory of
48 BMI, particularly at AP and AR.

49 To regulate satiety, leptin binds centrally to its hypothalamic receptors and activates 50 multiple signaling pathways that control food intake (23). Mutations in the genes encoding 51 leptin or its receptor (LEPR) can induce leptin resistance and disrupt its central and peripheral actions (23). Some very rare mutations are responsible for severe obesity syndrome in 52 53 humans (24), and more frequent variants have been associated with BMI variability and 54 obesity in children and adolescents (Aris et al., 2019; Fonseca et al., 2021; Péneau et al., 55 2017)(23,25). However, the association between individual LEPR variants and the evolution 56 of child BMI has been investigated in few studies and has been suggested to vary according to 57 age (26,27). We hypothesized that cord blood leptin level and carriage of the rare G allele of a 58 common LEPR variant, the rs9436303 single nucleotide polymorphism (SNP), can affect, 59 independently or in interaction, early postnatal growth, and later adiposity trajectories.

To obtain further insights into the role of cord blood leptin level in the development of adiposity in childhood, we aimed to 1) determine whether the cord blood leptin level and the *LEPR* rs9436303 variant are associated with later BMI trajectory, specifically BMI at AP and age at AR, and 2) examine whether *LEPR* rs9436303 modulates the association between cord blood leptin level and its indicators of child BMI trajectory.

65 **2.** Participants and methods

66 **2.1. Study design**

67 The EDEN (Etude des Déterminants pré et post natals de la santé de l'ENfant) mother-child study is a prospective French cohort that aims to assess prenatal and postnatal determinants of 68 69 child growth, health, and development. This cohort recruited 2002 pregnant women (before 70 24 weeks' amenorrhea) older than 18 years old between 2003 to 2006 in two university hospital centers in Nancy and Poitiers. All details of the study protocol were published 71 72 previously (28). Briefly, the exclusion criteria were pre-pregnancy diabetes, multiple 73 pregnancies, inability to read and write in French and planning to move outside the region; 74 53% of mothers were eligible.

Both parents gave their written consent when their child was included. The Data
Protection Authority and the relevant ethical research committee of Kremlin- Bicêtre
Hospital gave their approval for the study.

78

2.2. Cord blood leptin measurement

79 Immediately after birth, venous cord blood was sampled from 1425 newborns and allowed to clot. Blood samples were centrifuged within 24 h post-collection, serum was collected and 80 81 stored at -80°C. Cord blood serum was assessed for leptin level using the electro-82 chemiluminescence (ECL)-based immunoassay V-PLEX Human Leptin Kit (Meso Scale 83 Diagnostics, Rockville, MD, USA). All samples were above the lower limit of detection 84 (LLOD) of 43 pmol/mL. Inter-run coefficients of variations (CV) were below 20%, therefore 85 complying with the recommendations of the US Food and Drug Administration regarding 86 ligand binding assays (29). All assays were completed within 1 week by the same 87 investigator. All assays were performed according to the manufacturer's instructions.

88

2.3. Genotyping of LEPR SNP rs9436303

The SNP rs9436303 was previously studied in relation to adiposity peak and age at rebound in the Alves et *al.* study, in which the EDEN cohort was used for replication (11). Therefore, this polymorphism was the only *LEPR* SNP genotyped in EDEN and available for this study.

92 DNA was extracted from cord blood samples. Individual SNP genotyping involved 93 using amplified DNA with TaqMan probes (Applied Biosystems, Foster City, CA, USA). 94 SNP rs9436303 probes were ordered from Applied Biosystems, and after synthesis, used 95 according to the manufacturer's guidelines. The TaqMan assays were then read on a 7900HT 96 Fast Real-Time PCR System (Applied Biosystems), and the alleles were called by using the SDS software (Applied Biosystems). The call rate was > 98% and the Hardy Weinberg 97 98 Equilibrium P-value was 0.22. These genetic data were available for 1340 children in the 99 cohort. The G allele of this variant is the rare allele. In this study, we considered the SNP 100 variable (classified as 0, 1, and 2 according to the number of G alleles carried) as a continuous 101 trait, assuming a co-dominant effect on the adiposity traits.

102

2.4. AP and AR assessments

103 Children's weight and height were measured during clinical examinations or collected from 104 child's health booklets from birth to early adolescence. Weight was measured by previously 105 trained midwives at birth, 1, 3, and 5 years by using an electronic scale (SecaLtd or Terraillon 106 SL-351). Length was measured at birth and 1 year with a somatometer (Testut,NMMedical), 107 and height at 3 and 5 years with a stadiometer (Seca Ltd). Children had a mean of 10 weight 108 and height measurements (interquartile range 6- 14 and 5-13, respectively) from birth to 13 109 years.

110 The method used to model individual BMI trajectories, described elsewhere (1), was based on that of Sovio et al. and extended by adding a random effect to each parameter of a 111 112 mixed-effects cubic model (equation provided in supplementary files). Anthropometric data collected between age 3 days and 24 months were used to determine AP and those between 113 114 age 18 months and 13 years to determine AR. For each period and each participant, we 115 estimated AP by the first derivative null and second negative derivatives of individual BMI 116 functions and AR by the first derivative null and second positive derivatives of individual 117 BMI function. The modelling was performed over 2 different age periods (between age 3 days 118 and 24 months and between age 18 months and 13 years). The only criterion for inclusion was 119 to have at least three BMI measurements per child over each period. Individual BMI curves 120 were modelled for girls and boys separately for both periods.

121 As a result, AP data were available for 1713 children and AR data for 1415 children of 122 the cohort. We then considered these two distinct populations according to the outcome of 123 interest (Figure 1).

124 **2.5. Covariates**

125 Maternal smoking status during pregnancy (yes/no), maternal and paternal educational level 126 (years) were collected during pregnancy. At birth, we collected maternal age (years); newborn 127 characteristics including sex; prematurity (yes/no); and birth weight-for-gestational age separated into three classes: SGA ($\leq 10^{\text{th}}$ percentile), appropriate for gestational age (AGA, 128 $>10^{th}$ percentile to $\le 90^{th}$ percentile), and large for gestational age (LGA, $>90^{th}$ percentile) by 129 130 using a customized approach calculated according to Gardosi et al. references (30). 131 Gestational weight gain (kg) and gestational diabetes (yes/no) were collected from obstetrics 132 records. Maternal pre-pregnancy weight and paternal weight and height were collected from 133 self-reports at inclusion. Maternal height was measured at inclusion. These measurements

8

- were used to calculate BMI (kg/m²). For simplicity maternal pre-pregnancy BMI and paternal
 BMI were referred to as parental BMI in the following parts of the manuscript.
- 136

2.6. Statistical analysis

Descriptive analyses of leptin level, *LEPR* SNP and covariates were performed before multiple imputations, with mean (SD) or percentage (N). We compared children with estimated BMI at AP and age at AR to non-included children by Student *t* test for continuous variables and chi-square test for categorical variables. We also described the mean cord blood leptin level according to rs9436303 genotype and birth size groups.

142 We used multivariable linear regression models to evaluate the associations of the 143 cord blood leptin level and the rs9436303 genotype with BMI at AP and age at AR. These 144 multivariable linear regression models allow for assessing the association between an 145 exposure of interest (leptin) and our outcome (BMI at AP or age at AR), whatever the level of the other covariates introduced in the models, including birth size group. Analyses were 146 147 performed separately for each outcome. We did not stratify the analyses by sex because we 148 did not detect a significant interaction between sex and leptin (p>0.3). Given the skewed 149 distribution for the leptin variable, it was log-transformed and standardized by reduced 150 centered mean before being introduced into the models. We first analyzed the unadjusted 151 associations between leptin level, the LEPR SNP and outcomes. We then produced models adjusted for confounders (center, child sex, prematurity, birth size groups, maternal age at 152 153 delivery, gestational weight gain, gestational diabetes, parental educational level, and parental 154 BMI) for leptin level and the LEPR SNP separately. Finally, we considered leptin level and 155 the LEPR SNP simultaneously in one model adjusted for all confounders. Multivariable 156 analyses used quintiles of cord blood leptin level to assess whether the association was linear.

To test whether the association between leptin level, the *LEPR* SNP and outcomes did not depend on other variables, we performed two interaction analyses: one between cord blood leptin level and rs9436303 genotype and another between cord blood leptin level and birth size groups.

161 We used multiple imputation (SAS MI and MIANALYZE procedures) techniques to 162 deal with missing values due to leptin level, the LEPR SNP and covariates (Supplementary 163 table S1) to reduce potential selection bias. We generated 40 independent imputed datasets 164 using the fully conditional specification method. For each dataset, we used inverse probability 165 weighting (IPW) to account for potential attrition bias. We assigned to each participant a 166 weight corresponding to the inverse probability of being included in the analysis sample in a 167 logistic regression model adjusted for covariates (31). The results of different imputed 168 datasets were combined by using the SAS MIANALYZE procedure, in which standard errors 169 are calculated using Rubin's rules (32).

As sensitivity analyses, we repeated all analyses on complete cases (children without missing values for covariates) and after excluding mothers with pre-eclampsia because this condition has been associated with elevated placental leptin secretion (21).

We used R v3.5.1 for modelling BMI trajectories and estimating AP and AR (package
NLME). SAS v9.4 (SAS, Cary, NC, on an AIX 7.1 platform) was used for all other analyses.
Nominal p <0.05 was considered statistically significant.

176 **3. Results**

177

3.1. Characteristics of study populations

178 The children included in our two samples (children with estimated BMI at AP and children with estimated age at AR) were in general similar (Table 1). The mean [SD] BMI at AP was 179 17.5 [1.3] kg/m² and mean age at AR 5.5 [1.4] years. As compared with children excluded 180 181 from sample 1, children with estimated BMI at AP, children included had a higher mean cord blood leptin level and the proportion of preterm births was lower (Supplementary Table S2). 182 183 The mothers of the children included in the two samples had a higher level of education, were 184 older, and were less frequently smokers during pregnancy than those of the excluded children. 185 We found no difference in distribution of the LEPR variant. The proportion of children with 186 each variant of the LEPR SNP was similar between children included or not in both samples 187 (p=0.53 and 0.80 for samples 1 and 2, respectively). The allelic frequency of the G allele was 188 26.1% and 26.6% in samples 1 and 2, respectively.

Infants born SGA had a lower mean [SD] cord blood leptin level than those with AGA and LGA (8.3 [0.99] vs 8.9 [0.84] and 9.5 [0.85], p<.0001) (Supplementary figure S1a). We found no significant difference between mean cord blood leptin level according to number of risk alleles (Supplementary figure S1b).

193

3.2. Cord blood leptin level, BMI at AP, and age at AR

194 Cord blood leptin level was negatively related to BMI at AP in both the unadjusted (mean 195 $[SE] = -0.14 \ [0.04] \ \text{kg/m}^2$; p=0.0001) and the fully adjusted (-0.19 $[0.04] \ \text{kg/m}^2$; p<.0001) 196 models (Table 2). Furthermore, investigating quintiles of cord blood leptin revealed a linear 197 shape of the association with BMI at AP (p trend <.0001) (Figure 2A).

198 The negative association between cord blood leptin level and age at AR before any 199 adjustment (Table 2) did not remain significant in the multivariable model. However, when considering quintiles of leptin level in the model (Figure 2B), the highest quintile was negatively associated with age at AR as compared with all the other quintiles combined (mean [SE] = -87.3 [39.4] days or about 2.9 [1.3] months; p=0.03).

After adjusting for cord blood leptin level, as compared with children born AGA, those born SGA had lower BMI at AP (-0.57 [0.10] kg/m²; p<.0001) and earlier age at AR (-101.0 [39.8] days or about -3.2 [1.3] months; p=0.01), whereas those born LGA had higher BMI at AP (0.54 [0.12] kg/m²; p<.0001) (Tables 2). We did not find any significant interaction between birth size groups and cord blood leptin level (p>0.15).

208 **3.3.** *LEPR* polymorphism, BMI at AP, and age at AR

In the fully adjusted model, carrying a rare allele (G) in the *LEPR* SNP was associated with increased BMI at AP by about 0.24 kg/m² (p<0.001) and delayed age at AR by about 67 days (2.2 months; p=0.005) (Tables 2). We did not find any interaction between leptin level and rs9436303 G-allele carriage with the two outcomes considered (p-interaction=0.48 and 0.28 for BMI at AP and age at AR, respectively).

214

3.4. Sensitivity analyses

The results of the complete-case analyses are presented in Supplementary table S3. Results from analyses restricted to children with no missing data were consistent overall with those after multiple imputation and IPW. When mother–child pairs presenting pre-eclampsia pregnancy complications were excluded from the analysis, the results remained unchanged (data not shown).

4. Discussion

In the prospective EDEN mother-child cohort, we demonstrated that whatever the birth size group, increased cord blood leptin level was associated with reduced BMI at AP and earlier age at AR when considering the highest quintile of leptin level versus the other quintiles (Figure 2). Carrying the rare G-allele of the *LEPR* SNP rs9436303 did not modulate these associations but was independently related to higher BMI at AP and later age at AR.

226 High BMI at AP and early age at AR have been associated with increased BMI, risk of 227 cardiovascular disease, and obesity later in childhood and in adulthood (1-3). However, in 228 their work on the ELANCE cohort, Rolland-Cachera and Péneau highlighted two trajectories 229 at risk of later obesity (5): a first one marked by a high BMI at any age and a second 230 characterized by low BMI at AP followed by an early rebound and a subsequent increase in 231 BMI. According to this study, children with early AR had lower BMI before and higher BMI 232 after the rebound, on average, and were at increased risk of overweight in adulthood (5). We 233 interpreted our results on the association between cord blood leptin and the LEPR 234 polymorphism in light of these previously described trajectory.

235 To our knowledge, no previous studies have investigated cord blood leptin in 236 association with such specific features but rather analyzed BMI at specific ages in infancy or 237 childhood. In a prospective study of 642 children born in Greece, Karakosta et al. showed 238 high cord blood leptin level related to reduced BMI from age 6 months to 4 years (33). 239 Likewise, other studies of mother-infant pairs reported a significant inverse association 240 between cord blood leptin level and both BMI standard deviation score and weight gain in the 241 first year of life (34–36). Several analyses conducted in the project VIVA cohort showed low 242 cord blood leptin level related to small birth size (14) and increased BMI z-score at age 3 243 years (13) but low measures of adiposity from childhood to early adolescence (37). A recent 244 meta-analysis including some of these studies concluded that higher cord blood leptin levels 245 were associated with lower adiposity up to 3 years of age but not between 4 and 7 years (38). 246 Our results are fully consistent with this overview because we show high cord blood leptin 247 level associated with reduced BMI at AP but earlier age at AR, recognized as a risk factor for 248 later obesity (1,2,6,7). Furthermore, this trajectory, characterized by a reduced BMI at AP 249 more likely followed by early age at the time of AR (dashed line in Supplementary Figure 250 S2), corresponds precisely to one of the two risk trajectories described by Rolland-Cachera 251 and Peneau (5).

252 Mechanisms underlying such associations are still unclear but deserve to be elucidated 253 and discussed. The leptin hormone regulates satiety, and its circulating level is positively 254 correlated with body fat mass and negatively correlated with satiety (39-41). Reduced 255 circulating leptin level enhances food intake and increases body fat mass and vice versa 256 (19,39). This is consistent with our finding showing lower leptin level at birth associated with 257 higher BMI at AP. Furthermore, animal studies suggest that in addition to its role in satiety, 258 leptin would contribute to the modification of the release of other growth hormones in the 259 hypothalamus (42) and to the regulation of certain neuroendocrine functions (43).

Our findings could explain in part the rapid postnatal weight gain observed in children born SGA as compared with other children (44,45). Indeed, the level of leptin was lower, on average, in children born SGA than their peers. However, in our study, being born SGA and high cord blood leptin level were independent predictors of reduced BMI at AP, which suggests that different physiological mechanisms are involved in catch-up growth. Indeed, infants are born SGA usually due to poor fetal growth and nutrition. Children born SGA 266 exhibit abnormal body composition with lower lean mass and at higher risk for metabolic267 disorders (46,47).

268 Our results also reveal that very high cord blood leptin level was associated with early 269 age at AR. Leptin measured in venous cord blood consists of placental and fetal leptin 270 (19,48). The placenta is one of the main sites for leptin production during pregnancy, and part 271 of the leptin produced in the placenta passes into the fetal circulation, but the proportion of 272 placental leptin concentration measured in cord blood at birth is unknown (15). Thus, cord 273 blood leptin reflects both placental and fetal leptin. According to some studies, placental 274 leptin production is increased during fetal hypoxia in response to an adverse intrauterine 275 condition (21,49). Cord blood leptin level higher than expected for a given amount of fat mass 276 could be a marker of adverse intrauterine conditions, themselves associated with BMI 277 trajectories characterized by early AR (46,47). This suggestion is consistent with the hypothesis of the existence of a fetal programming for BMI trajectory at the time of AR 278 279 suggested by several authors (50–53). Our sensitivity analysis revealed similar results with or 280 without the exclusion of mothers with pre-eclampsia, but the number of pre-eclampsia 281 mothers was very low.

282 To exert its action on satiety in the individual, leptin binds to hypothalamic receptors. 283 Altered *LEPR* gene expression may lead to leptin resistance and thus modulate the satiety effect of leptin during childhood and affect the BMI trajectory and AR. Leptin resistance is 284 285 characterized by the inability of leptin to reach the target cell owing to reduced expression of 286 LEPR or disrupted LEPR signaling. This resistance leads to decreased effectiveness of its 287 action and a major decrease in the effect of leptin on satiety (54). Thus, the effects of leptin on 288 appetite and body mass are impaired and may result in higher leptin secretion as a 289 compensatory mechanism. Its action on satiety occurs in the postnatal period only, so there is

290 no reason to believe that a change in LEPR level could affect prenatal leptin production as a 291 compensatory mechanism, which explains why no association between leptin levels in cord 292 blood and LEPR genotype was expected. Indeed, we did not find any difference in cord blood 293 leptin level between carriers or not of the LEPR rs9436303 rare G-allele. Our hypothesis that 294 the G allele would interact in the association between cord blood leptin level and early growth 295 was not confirmed by our results. However, children carrying the G allele had higher BMI at 296 AP and later age at AR (dotted line in Supplementary figure S2). In a previous genome-wide 297 association study, which included 7215 children from five European cohorts, carrying the G 298 allele was also associated with higher BMI at AP, and the rs9436303 genotype explained 299 0.3% of the variance in BMI at AP (11) as compared with 1.3% in our study (data not shown). 300 To our knowledge, the association between rs9436303 genotype and age at AR, has not been investigated in genetic studies. Another study addressed the link between 28 LEPR 301 302 polymorphisms and BMI in 522 Spanish children aged 6 to 15 years and showed a strong 303 association between rs11804091 carriage and obesity-related traits in children(23). The 304 rs11804091 rare allele frequency was found in linkage disequilibrium with rs9436303 305 $(R^2=0.502)$. Therefore, variants in *LEPR* could determine the overall BMI trajectory, 306 regardless of cord blood leptin level, and future studies should analyze the long-term 307 association with adult obesity. A systematic analysis and meta-analysis of 17 studies in adults 308 examined the association between three other variants of LEPR and obesity-related 309 phenotypes and did not find any association with adult overweight (55). Genetic variation in 310 LEPR may have different effects depending on the variant considered and the age window 311 explored, as with leptin level. Further studies are needed to better understand the 312 physiological mechanisms involved and the long-term consequences of LEPR genetic 313 variation on the BMI trajectory.

314 This study is among the first to examine the links of both cord blood leptin level and 315 of the LEPR SNP rs9436303 genotype with BMI trajectory, especially at the times of AP and 316 AR. Anthropometric data were collected prospectively, which allowed us to estimate AP and 317 AR for many children in the EDEN cohort. However, the generalization of results remains 318 limited because of potential selection and attrition biases that we attempted to address by 319 applying multiple imputation and IPW. The complete-case analyses showed well that the 320 impact of these biases was limited. Moreover, cord blood leptin reflects both placental and 321 fetal leptin, which limits interpretation regarding the origin of leptin variation (placenta, fetal) 322 in our sample. Of note, previous studies have shown that cord blood leptin level was not 323 correlated with maternal leptin level or BMI but with birth weight and length, ponderal index, 324 and weight/length ratio (20,56). These data support that cord blood leptin reflects fetal fat 325 mass, which suggests that leptin measured in cord blood is primarily of fetal origin and 326 strengthens the relevance of relying on cord blood leptin level as a proxy of neonatal 327 adiposity. However, our study was not designed with the aim of developing a biological 328 signature at birth, that could serve as a tool in routine clinical practice for the early detection 329 of children at risk of developing obesity during childhood; further dedicated studies would be 330 necessary for this purpose.

In conclusion, we have shown that cord blood leptin level and *LEPR* SNP rs9436303 genotype interfere with BMI trajectory in children, independent of each other and birth size group. These results illustrate the role of early life body composition in the programming of childhood adiposity. In addition to being a marker of fetal adiposity, cord blood leptin level could also reflect an unfavorable intrauterine environment and may predict a trajectory at risk for subsequent obesity.

Acknowledgements

The authors thank the EDEN mother-child cohort study group, whose members are I. Annesi-Maesano, J.Y. Bernard, M.A. Charles, P. Dargent-Molina, B. de Lauzon-Guillain, P. Ducimetière, M. de Agostini, B. Foliguet, A. Forhan, X. Fritel, A. Germa, V. Goua, R. Hankard, B. Heude, M. Kaminski, B. Larroquey, N. Lelong, J. Lepeule, G. Magnin, L. Marchand, C. Nabet, F Pierre, R. Slama, M.J. Saurel-Cubizolles, M. Schweitzer, and O. Thiebaugeorges.

Author contributions

None of the authors have any financial relationships or conflict of interest to disclose.

AC and BH designed the research, wrote the manuscript and analyzed the data.

BH, MAC, MuT, MaT, SL, KL, OK, LD and BLG were responsible for data collection and genotyping in EDEN. All authors reviewed drafts, provided critical feedback, approved the final manuscript and were responsible for the final content of the paper. BH had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Funding information

The EDEN study was supported by Foundation for medical research (FRM), National Agency for Research (ANR), National Institute for Research in Public health (IRESP: TGIR cohorte santé 2008 program), French Ministry of Health (DGS), French Ministry of Research, INSERM Bone and Joint Diseases National Research (PRO-A), and Human Nutrition National Research Programs, Paris-Sud University, Nestlé, French National Institute for Population Health Surveillance (InVS), French National Institute for Health Education (INPES), the European Union FP7 programmes (FP7/2007–2013, HELIX, ESCAPE, ENRIECO, Medall projects), Diabetes National Research Program (through a collaboration with the French Association of Diabetic Patients [AFD]), French Agency for Environmental Health Safety (now ANSES), Mutuelle Générale de l'Education Nationale a complementary health insurance (MGEN), French national agency for food security, French-speaking association for the study of diabetes and metabolism (ALFEDIAM). KKO is supported by the Medical Research Council (Unit programme: MC_UU_12015/2).

Conflict of interest

The authors declare that they have no conflict of interest.

References

- 1. Aris IM, Rifas-Shiman SL, Li L-J, Kleinman KP, Coull BA, Gold DR, Hivert M-F, Kramer MS, Oken E. Patterns of body mass index milestones in early life and cardiometabolic risk in early adolescence. Int J Epidemiol 2019;0:11.
- 2. Péneau S, Giudici KV, Gusto G, Goxe D, Lantieri O, Hercberg S, Rolland-Cachera M-F. Growth Trajectories of Body Mass Index during Childhood: Associated Factors and Health Outcome at Adulthood. J Pediatr 2017;186:64-71.e1.
- 3. Fonseca MJ, Moreira C, Santos AC. Adiposity rebound and cardiometabolic health in childhood: results from the Generation XXI birth cohort. Int J Epidemiol 2021;dyab002.
- 4. Rolland-Cachera MF, Deheeger M, Bellisle F, Sempé M, Guilloud-Bataille M, Patois E. Adiposity rebound in children: a simple indicator for predicting obesity. Am J Clin Nutr 1984;39:129–35.
- 5. Rolland-Cachera MF, Péneau S. Growth trajectories associated with adult obesity. World Rev Nutr Diet 2013;106:127–34.
- 6. Hof MHP, Vrijkotte TGM, de Hoog MLA, van Eijsden M, Zwinderman AH. Association between Infancy BMI Peak and Body Composition and Blood Pressure at Age 5–6 Years. Wang L, editor. PLoS ONE 2013;8:e80517.
- 7. Silverwood RJ, De Stavola BL, Cole TJ, Leon DA. BMI peak in infancy as a predictor for later BMI in the Uppsala Family Study. Int J Obes 2009;33:929–37.
- 8. Péneau S, González-Carrascosa R, Gusto G, Goxe D, Lantieri O, Fezeu L, Hercberg S, Rolland-Cachera MF. Age at adiposity rebound: determinants and association with nutritional status and the metabolic syndrome at adulthood. Int J Obes 2016;40:1150–6.
- 9. Aris IM, Rifas-Shiman SL, Li L-J, Kleinman K, Coull BA, Gold DR, Hivert M-F, Kramer MS, Oken E. Pre-, Perinatal, and Parental Predictors of Body Mass Index Trajectory Milestones. J Pediatr 2018;201:69-77.e8.
- 10. Ip EH, Marshall SA, Saldana S, Skelton JA, Suerken CK, Arcury TA, Quandt SA. Determinants of Adiposity Rebound Timing in Children. J Pediatr 2017;184:151-156.e2.
- 11. Couto Alves A, De Silva NMG, Karhunen V, Sovio U, Das S, Taal HR, Warrington NM, Lewin AM, Kaakinen M, Cousminer DL, et al. GWAS on longitudinal growth traits reveals different genetic factors influencing infant, child, and adult BMI. Sci Adv 2019;5:eaaw3095.
- Cissé AH, Lioret S, de Lauzon-Guillain B, Forhan A, Ong KK, Charles MA, Heude B. Association between perinatal factors, genetic susceptibility to obesity and age at adiposity rebound in children of the EDEN mother–child cohort. Int J Obes [Internet] 2021 [cited 2021 May 13]; Available from: http://www.nature.com/articles/s41366-021-00847-w

- 13. Boeke CE, Mantzoros CS, Hughes MD, L. Rifas-Shiman S, Villamor E, Zera CA, Gillman MW. Differential associations of leptin with adiposity across early childhood: Leptin and Adiposity Across Early Childhood. Obesity 2013;21:1430–7.
- 14. Mantzoros CS, Rifas-Shiman SL, Williams CJ, Fargnoli JL, Kelesidis T, Gillman MW. Cord Blood Leptin and Adiponectin as Predictors of Adiposity in Children at 3 Years of Age: A Prospective Cohort Study. PEDIATRICS 2009;123:682–9.
- 15. Geary M, Pringle PJ, Persaud M, Wilshin J, Hindmarsh PC, Rodeck CH, Brook CGD. Leptin concentrations in maternal serum and cord blood: relationship to maternal anthropometry and fetal growth. BJOG Int J Obstet Gynaecol 1999;106:1054–60.
- 16. Ong K, Ahmed M, Dunger D. The role of leptin in human growth and puberty. Acta Paediatr 1999;88:95–8.
- 17. Valūniene M, Verkauskiene R, Boguszewski M, Dahlgren J, Lasiene D, Lasas L, Wikland KA. Leptin levels at birth and in early postnatal life in small- and appropriate-for-gestational-age infants. Med Kaunas Lith 2007;43:784–91.
- Martínez-Cordero C, Amador-Licona N, Guízar-Mendoza JM, Hernández-Méndez J, Ruelas-Orozco G. Body Fat at Birth and Cord Blood Levels of Insulin, Adiponectin, Leptin, and Insulin-like Growth Factor-I in Small-for-Gestational-Age Infants. Arch Med Res 2006;37:490–4.
- 19. Taine M, Khalfallah O, Forhan A, Glaichenhaus N, Charles M-A, Heude B. Does cord blood leptin level mediate the association between neonatal body size and postnatal growth? Results from the EDEN mother–child cohort study. Ann Hum Biol 2020;47:159–65.
- 20. Clapp JF, Kiess W. Cord Blood Leptin Reflects Fetal Fat Mass. J Soc Gynecol Investig 1998;5:300–3.
- 21. Mise H, Sagawa N, Matsumoto T, Yura S, Nanno H, Itoh H, Mori T, Masuzaki H, Hosoda K, Ogawa Y, et al. Augmented Placental Production of Leptin in Preeclampsia: Possible Involvement of Placental Hypoxia¹. J Clin Endocrinol Metab 1998;83:3225–9.
- 22. Persson B, Westgren M, Celsi G, Nord E, Örtqvist E. Leptin Concentrations in Cord Blood in Normal Newborn Infants and Offspring of Diabetic Mothers. Horm Metab Res 1999;31:467–71.
- 23. Olza J, Rupérez AI, Gil-Campos M, Leis R, Cañete R, Tojo R, Gil Á, Aguilera CM. Leptin Receptor Gene Variant rs11804091 Is Associated with BMI and Insulin Resistance in Spanish Female Obese Children: A Case-Control Study. Int J Mol Sci 2017;18:1690.
- 24. Clément K, Vaisse C, Lahlou N, Cabrol S, Pelloux V, Cassuto D, Gourmelen M, Dina C, Chambaz J, Lacorte J-M, et al. A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. Nature 1998;392:398–401.

- 25. Wheeler E, Huang N, Bochukova EG, Keogh JM, Lindsay S, Garg S, Henning E, Blackburn H, Loos RJF, Wareham NJ, et al. Genome-wide SNP and CNV analysis identifies common and low-frequency variants associated with severe early-onset obesity. Nat Genet 2013;45:513–7.
- 26. Helgeland Ø, Vaudel M, Juliusson PB, Lingaas Holmen O, Juodakis J, Bacelis J, Jacobsson B, Lindekleiv H, Hveem K, Lie RT, et al. Genome-wide association study reveals dynamic role of genetic variation in infant and early childhood growth. Nat Commun 2019;10:4448.
- 27. Silva NMGD, Sebert S, Alves AC, Sovio U, Das S, Taal R, Warrington NM, Lewin AM, Kaakinen M, Cousminer D, et al. Genetic architecture of early childhood growth phenotypes gives insights into their link with later obesity [Internet]. bioRxiv; 2017 [cited 2022 Mar 21]. p. 150516. Available from: https://www.biorxiv.org/content/10.1101/150516v1
- 28. Heude B, Forhan A, Slama R, Douhaud L, Bedel S, Saurel-Cubizolles M-J, Hankard R, Thiebaugeorges O, De Agostini M, Annesi-Maesano I, et al. Cohort Profile: The EDEN mother-child cohort on the prenatal and early postnatal determinants of child health and development. Int J Epidemiol 2016;45:353–63.
- 29. Barbosa S, Khalfallah O, Forhan A, Galera C, Heude B, Glaichenhaus N, Davidovic L. Immune activity at birth and later psychopathology in childhood. Brain Behav Immun Health 2020;8:100141.
- 30. Gardosi J, Chang A, Kalyan B, Sahota D, Symonds EM. Customised antenatal growth charts. The Lancet 1992;339:283–7.
- 31. Howe CJ, Cole SR, Lau B, Napravnik S, Eron JJ. Selection Bias Due to Loss to Follow Up in Cohort Studies: Epidemiology 2016;27:91–7.
- 32. Bartlett JW, Seaman SR, White IR, Carpenter JR, for the Alzheimer's Disease Neuroimaging Initiative*. Multiple imputation of covariates by fully conditional specification: Accommodating the substantive model. Stat Methods Med Res 2015;24:462–87.
- 33. Karakosta P, Roumeliotaki T, Chalkiadaki G, Sarri K, Vassilaki M, Venihaki M, Malliaraki N, Kampa M, Castanas E, Kogevinas M, et al. Cord blood leptin levels in relation to child growth trajectories. Metabolism 2016;65:874–82.
- 34. Telschow A, Ferrari N, Deibert C, Flöck A, Merz WM, Gembruch U, Ehrhardt C, Dötsch J, Graf C. High Maternal and Low Cord Blood Leptin Are Associated with BMI-SDS Gain in the First Year of Life. Obes Facts 2019;12:575–85.
- 35. Kaar JL, Brinton JT, Crume T, Hamman RF, Glueck DH, Dabelea D. Leptin levels at birth and infant growth: the EPOCH study. J Dev Orig Health Dis 2014;5:214–8.
- Parker M, Rifas-Shiman SL, Belfort MB, Taveras EM, Oken E, Mantzoros C, Gillman MW. Gestational Glucose Tolerance and Cord Blood Leptin Levels Predict Slower Weight Gain in Early Infancy. J Pediatr 2011;158:227–33.

- 37. Li L-J, Rifas-Shiman SL, Aris IM, Young JG, Mantzoros C, Hivert M-F, Oken E. Associations of maternal and cord blood adipokines with offspring adiposity in Project Viva: is there an interaction with child age? Int J Obes 2018;42:608–17.
- 38. Bagias C, Sukumar N, Weldeselassie Y, Oyebode O, Saravanan P. Cord Blood Adipocytokines and Body Composition in Early Childhood: A Systematic Review and Meta-Analysis. Int J Environ Res Public Health 2021;18:1897.
- 39. Friedman JM. Leptin, Leptin Receptors, and the Control of Body Weight. Nutr Rev 2009;56:S38–46.
- 40. Attele AS, Shi ZQ, Yuan C-S. Leptin, gut, and food intake. Biochem Pharmacol 2002;63:1579-83.
- 41. Klok MD, Jakobsdottir S, Drent ML. The role of leptin and ghrelin in the regulation of food intake and body weight in humans: a review. Obes Rev 2007;8:21–34.
- 42. Watanobe H. Leptin directly acts within the hypothalamus to stimulate gonadotropinreleasing hormone secretion in vivo in rats. J Physiol 2002;545:255–68.
- 43. Park H-K, Ahima RS. Physiology of leptin: energy homeostasis, neuroendocrine function and metabolism. Metabolism 2015;64:24–34.
- 44. Fonseca VM, Sichieri R, Moreira MEL, Moura AS. Early Postnatal Growth in Preterm Infants and Cord Blood Leptin. J Perinatol 2004;24:751–6.
- 45. Yeung EH, Sundaram R, Xie Y, Lawrence DA. Newborn adipokines and early childhood growth. Pediatr Obes 2018;13:505–13.
- 46. Ong KKL, Preece MA, Emmett PM, Ahmed ML, Dunger DB. Size at Birth and Early Childhood Growth in Relation to Maternal Smoking, Parity and Infant Breast-Feeding: Longitudinal Birth Cohort Study and Analysis. Pediatr Res 2002;52:863–7.
- 47. Lévy-Marchal C, Czernichow P. Small for Gestational Age and the Metabolic Syndrome: Which Mechanism Is Suggested by Epidemiological and Clinical Studies? Horm Res Paediatr Karger Publishers; 2006;65:123–30.
- 48. West J, Santorelli G, Collings P, Bingham D, Whincup P, Sattar N, Norris T, Wright J, Lawlor DA. Associations of cord leptin and cord insulin with adiposity and blood pressure in White British and Pakistani children aged 4/5 years. Wellcome Open Res 2019;4:157.
- 49. de Knegt VE, Hedley PL, Kanters JK, Thagaard IN, Krebs L, Christiansen M, Lausten-Thomsen U. The Role of Leptin in Fetal Growth during Pre-Eclampsia. Int J Mol Sci 2021;22:4569.
- 50. McMillen IC, Edwards LJ, Duffield J, Muhlhausler BS. Regulation of leptin synthesis and secretion before birth: implications for the early programming of adult obesity. Reproduction Society for Reproduction and Fertility; 2006;131:415–27.

- 51. Breier BH, Vickers MH, Ikenasio BA, Chan KY, Wong WPS. Fetal programming of appetite and obesity. Mol Cell Endocrinol 2001;185:73–9.
- 52. Brynhildsen J, Sydsjö G, Blomberg M, Claesson I-M, Theodorsson E, Nyström F, Sydsjö A, Josefsson A. Leptin and adiponectin in cord blood from children of normal weight, overweight and obese mothers. Acta Paediatr 2013;102:620–4.
- 53. Attig L, Solomon G, Ferezou J, Abdennebi-Najar L, Taouis M, Gertler A, Djiane J. Early postnatal leptin blockage leads to a long-term leptin resistance and susceptibility to diet-induced obesity in rats. Int J Obes 2008;32:1153–60.
- 54. Obradovic M, Sudar-Milovanovic E, Soskic S, Essack M, Arya S, Stewart AJ, Gojobori T, Isenovic ER. Leptin and Obesity: Role and Clinical Implication. Front Endocrinol 2021;12:585887.
- 55. Bender N, Allemann N, Marek D, Vollenweider P, Waeber G, Mooser V, Egger M, Bochud M. Association between Variants of the Leptin Receptor Gene (LEPR) and Overweight: A Systematic Review and an Analysis of the CoLaus Study. Rouet P, editor. PLoS ONE 2011;6:e26157.
- 56. Laml T, Hartmann BW, Ruecklinger E, Preyer O, Soeregi G, Wagenbichler P. Maternal serum leptin concentrations do not correlate with cord blood leptin concentrations in normal pregnancy. J Soc Gynecol Investig 2001;8:43–7.



Figure 1: Flow chart of the children included in the study. Two separate samples based on the outcomes with adiposity peak on the left and adiposity rebound on the right. BMI, body mass index; SNP, single nucleotide polymorphism



Figure 2: Adjusted means^a (SE) for A) BMI at adiposity peak (AP) and B) age at adiposity rebound (AR) according to cord leptin level quintiles^b.

^a Adjusted for center, maternal age, maternal and paternal education, maternal and paternal BMI, gestational weight gain, gestational diabetes, maternal smoking during pregnancy, prematurity, birth size groups and sex.

^b Quintiles 1, 2, 3 and 4 were grouped together and compared to the last quintile in the model

	Children with	Children with
	estimated BMI at AP	estimated age at AR
	N=1713	N=1415
	Mean (SD) or % (N)	Mean (SD) or % (N)
Sex (boys)	51.8 (888)	52.3 (746)
Preterm birth (yes)	5.4 (92)	5.5 (78)
Birth weight (kg)	3.3 (5.1)	3.3 (5.0)
Birth size groups ^a		
SGA	13.1 (218)	13.3 (184)
AGA	78.4 (1309)	77.4 (1070)
LGA	8.6 (143)	9.3 (128)
Gestational weight gain (kg)	13.4 (4.7)	13.3 (4.7)
Gestational diabetes (yes)	6.65 (94)	6.60 (113)
Maternal BMI (kg/m ²)	23.2 (4.6)	23.1 (4.4)
Paternal BMI (kg/m ²)	25.1 (3.7)	25.1 (3.6)
Never smoking during pregnancy	75.8 (1269)	78.6 (1088)
Maternal age at delivery (years)	29.7 (4.8)	29.9 (4.7)
Maternal educational level	13.7 (2.6)	14.0 (2.6)
(years)		
Paternal educational level (years)	13.1 (2.6)	13.3 (2.6)
Age at AR (years)	5.5 (1.4)	5.5 (1.4)
BMI at AP (kg/m ²)	17.5 (1.3)	17.5 (1.3)
Cord blood leptin level (ng/ml)	11.1 (11.0)	11.0 (10.5)
LEPR SNP rs9436303		
A/A	55.4 (668)	55.0 (551)
A/G	36.9 (445)	36.8 (369)
G/G	7.6 (92)	8.2 (82)
G-allele frequency	26.1	26.6

Table 1: Description of study populations

^a Small for gestational age, appropriate for gestational age and large for gestational age classified according to Gardosi references.

AP, adiposity peak; AR, adiposity rebound; BMI, body mass index; SGA, small for gestational age; AGA, appropriate for gestational age; LGA, large for gestational age; SNP, single nucleotide polymorphism

Table 2: Results of association analyses between cord blood leptin level, *LEPR* SNP rs9436303 and BMI at AP (in kg/m², N=1713^a) and age at AR (in days, N=1415^b)

Characteristics	Separate and unadjusted models ^c		Separate adjusted model with leptin level ^d		Separate adjusted model with <i>LEPR</i> SNP ^d		Fully adjusted model ^d	
Outcome = BMI at AP; N=1713 ^a							·	
	β (SE)	p value	β (SE)	p value	β (SE)	p value	B (SE)	p value
Birth size groups ^e (ref=AGA)								
SGA			-0.57 (0.10)	<.0001	-0.48 (0.10)	<.0001	-0.57 (0.10)	<.0001
LGA			0.54 (0.12)	<.0001	0.43 (0.12)	0.0002	0.54 (0.12)	<.0001
Cord blood leptin (log-transformed)	-0.14 (0.04)	0.0001	-0.19 (0.04)	<.0001			-0.19 (0.04)	<.0001
<i>LEPR</i> SNP rs9436303 (number of G alleles carried)	0.25 (0.06)	<.0001			0.24 (0.06)	<.0001	0.24 (0.06)	<.0001
Outcome = Age at AR; N=1415 ^b			·					
Birth size groups ^e (ref=AGA)								
SGA			-101.0 (39.8)	0.01	-94.0 (39.0)	0.02	-104.5 (39.9)	0.009
LGA			20.9 (47.9)	0.66	5.6 (46.8)	0.91	18.1 (47.8)	0.70
Cord blood leptin (log-transformed)	-34.9 (15.0)	0.02	-24.7 (17.6)	0.16			-22.7 (17.6)	0.20
LEPR SNP rs9436303 (number of G alleles carried)	65.3 (24.5)	0.008			67.9 (23.6)	0.004	66.6 (23.7)	0.005

^a Sample of children with estimated BMI at AP, linear regression model after multiple imputation on missing data of leptin, *LEPR* SNP and covariates.

^b Sample of children with estimated age at AR, linear regression model after multiple imputation on missing data of leptin, *LEPR* SNP and covariates.

^c Two separate models for cord blood leptin and rs9436303 without any adjustment.

^d Adjusted for center, maternal age, maternal and paternal education, maternal and paternal BMI, gestational weight gain, gestational diabetes, maternal smoking during pregnancy, prematurity and sex.

^e Small for gestational age, appropriate for gestational age and large for gestational age classified according to Gardosi references.

Supplementary Materials and Methods: Modeling equation for BMI trajectories

The following equation describes the general model applied for the i^{th} individual at the t^{th} time point, whatever the period considered (12).

$$\log (BMI_{it}) = \beta_0 + \sum_{j=1}^{3} \beta_j Age_{it}^{\ j} + b_{0i} + \sum_{j=1}^{3} b_{ji} Age_{it}^{\ j} + \varepsilon_{it}$$

Where:

- $\beta_0, \beta_1, \beta_2, \beta_3$ are the parameters for the fixed effects,
- b_{0i}, b_{1i}, b_{2i}, b_{3i} are the parameter estimates for the random effects on the polynomial coefficients, for each individual,
- ε_{it} are the error terms assumed to be normally distributed and independent

Variables	Type of variable	Model used to predict	Missing
		missing data ^a	value (%)
Age at adiposity peak	Continuous		0
BMI at adiposity peak	Continuous		0
Age at adiposity rebound	Continuous		0
BMI at adiposity rebound	Continuous		0
Center	Binary		0
Sex	Binary		0
Birth weight	Continuous	Linear regression	0.06
Personalized birth weight z-score	Continuous	Linear regression	2.5
Height at birth	Continuous	Linear regression	2.9
Head circumference at birth	Continuous	Linear regression	2.9
Skin folds at birth	Continuous	Linear regression	5.8
Gestational age	Continuous		0
Weight predicted at 5 years by	Continuous		0
Jenss-Bayley modelling ^b			
Height predicted at 5 years by	Continuous		0
Jenss-Bayley modelling ^b			
Maternal age at delivery	Continuous		0
Gestational weight gain	Continuous	Linear regression	1.9
Gestational diabetes	Binary	Logistic regression	0.6
Smoking during pregnancy	Binary	Logistic regression	2.3
Maternal education	Continous		0
Maternal income	Ordinal (3 categories)	Multinomial regression	0.6
Maternal BMI	Continuous	Linear regression	2.0
Paternal education	Continous		0
Paternal BMI	Continuous	Linear regression	8.0
Cord blood leptin level	Continuous	Linear regression	25.0
rs9436303 SNP	Ordinal (3 categories)	Multinomial regression	29.7

Supplementary Table S1: Details of the method used for multiple imputation

From the variables in the table, we generated 40 independent sets of imputed data using the fully conditional specification method (MI procedure of the SAS software). This method assigned data to missing measurements based on the measurement of infants with similar profiles. Then, we computed pooled effect estimates (MIANALYZE procedure of the SAS software).

BMI, body mass index; SNP, single nucleotide polymorphism

^a Fully conditional specification method.

^b Predicted weight and height were calculated by using previously modelled trajectories from the Jenss– Bayley model (1, 2).

1. Carles, S. *et al.* A Novel Method to Describe Early Offspring Body Mass Index (BMI) Trajectories and to Study Its Determinants. *PLOS ONE* **11**, e0157766 (2016).

2. Botton, J. *et al.* Postnatal Weight and Height Growth Modeling and Prediction of Body Mass Index as a Function of Time for the Study of Growth Determinants. *Ann. Nutr. Metab.* **65**, 156–166 (2014)

Table S2: Characteristics of children included or not in each analysis sample

	Children wit	h estimated BM	II at AP	Children with estimated age at AR			
Characteristics	Included	Excluded	P > t or χ2	Included	Excluded	$P > t $ or $\chi 2$	
	N=1713	N=289		N=1415	N=587		
	Mean (SD) or % (N)			Mean (SD) or % (N)			
Sex (boys)	51.8 (888)	59.0 (112)	0.06	52.7 (746)	52.1	0.80	
Preterm birth (yes)	5.4 (92)	9.4 (18)	0.02	5.5 (78)	6.5 (32)	0.41	
Birth weight (kg)	3.3 (0.5)	3.2 (0.5)	0.13	3.3 (0.5)	3.2 (0.5)	0.03	
Birth size groups ^a			0.38			0.04	
SGA	13.1 (218)	15.3 (28)		13.3 (184)	13.2 (62)		
AGA	78.4 (1309)	78.7 (144)		77.4 (1070)	81.3 (383)		
LGA	8.6 (143)	6.0 (11)		9.3 (128)	5.5 (26)		
Gestational weight gain (kg)	13.4 (4.7)	13.7 (5.7)	0.41	13.3 (4.7)	13.8 (5.3)	0.05	
Gestational diabetes (yes)	6.65 (94)	5.92 (29)	0.57	6.60 (113)	5.21 (10)	0.46	
Maternal BMI (kg/m ²)	23.2 (4.6)	23.5 (5.2)	0.43	23.1 (4.4)	23.6 (5.1)	0.10	
Paternal BMI (kg/m ²)	25.1 (3.7)	24.4 (3.4)	0.02	25.1 (3.6)	24.7 (3.7)	0.03	
Never smoking during pregnancy	75.8 (1269)	54.0 (94)	<.0001	78.6 (1088)	59.4 (275)	<.0001	
Maternal age at delivery (years)	29.7 (4.8)	27.9 (5.4)	<.0001	29.9 (4.7)	28.2 (5.2)	<.0001	
Maternal educational level (years)	13.7 (2.6)	12.3 (2.5)	<.0001	14.0 (2.6)	12.5 (2.5)	<.0001	
Paternal educational level (years)	13.1 (2.6)	12.2 (2.2)	<.0001	13.3 (2.6)	12.2 (2.4)	<.0001	

Age at AR (years)				5.5 (1.4)			
BMI at AP (kg/m ²)	17.5 (1.34)						Compari
Cord blood leptin level (ng/ml)	11.1 (11.0)	9.3 (9.2)	0.04	11.0 (10.5)	10.8 (11.9)	0.78	son
LEPR SNP rs9436303			0.53			0.80	between
A/A	55.4 (668)	54.1 (73)		55.0 (551)	56.2 (190)		children with
A/G	36.9 (445)	35.6 (48)		36.8 (369)	36.7 (124)		estimate
G/G	7.6 (92)	10.4 (14)		8.2 (82)	7.1 (24)		d BMI at

age at AR to non-included children from eligible mothers by Student t test for continuous variables and chi- square test for categorical variable

^a Small for gestational age, appropriate for gestational age and large for gestational age classified according to Gardosi references.

BMI, body mass index; AP, adiposity peak; AR, adiposity rebound, SGA, small for gestational age; AGA, appropriate for gestational age; LGA, large for gestational age; SNP, single nucleotide polymorphism

Supplementary Table S3: Results of association analyses of complete cases between cord blood leptin level, *LEPR* SNP rs9436303 SNP and BMI at AP and age at AR

Characteristics	Separate and unadjusted models ^a		Separate adjusted model with leptin level ^b		Separate adjusted model with <i>LEPR</i> SNP ^b		Fully adjusted model ^b	
Outcome = BMI at AP			N=1124		N=1047		N=948	
	β (SE)	p value	β (SE)	p value	β (SE)	p value	B (SE)	p value
Birth size groups ^c (ref=AGA)								
SGA			-0.54 (0.13)	<.0001	-0.48 (0.13)	0.0002	-0.46 (0.14)	0.001
LGA			0.63 (0.14)	<.0001	0.42 (0.14)	0.003	0.59 (0.15)	<.0001
Cord blood leptin (log-transformed)	-0.26 (0.04)	0.0002	-0.21 (0.04)	<.0001			-0.20 (0.05)	<.0001
LEPR SNP rs9436303 (number of G alleles carried)	0.24 (0.06)	<.0001			0.22 (0.07)	0.001	0.24 (0.07)	0.001
Outcome = Age at AR			N=928	N=928 N=880			N=793	
Birth size groups ^c (ref=AGA)								
SGA			-172.46 (52.16)	0.001	-163.13 (53.38)	0.002	-181.82 (56.90)	0.002
LGA			37.12 (56.12)	0.51	3.05 (56.11)	0.96	30.75 (59.72)	0.61
Cord blood leptin (log-transformed)	-26.94	0.08	-28.44 (18.42)	0.12			-25.40 (19.43)	0.19
	(15.34)						× -/	
LEPR SNP rs9436303 (number of G alleles carried)	57.84 (24.50)	0.02			80.04 (26.16)	0.002	85.00 (27.46)	0.002

^a Two separate models for cord blood leptin and rs9436303 without any adjustment.

^b Adjusted for center, maternal age, maternal and paternal education, maternal and paternal BMI, gestational weight gain, gestational diabetes, maternal smoking during pregnancy, prematurity and sex.

^c Small for gestational age, appropriate for gestational age and large for gestational age classified according to Gardosi references



Supplementary Figure S1: Violin plots with quantile bounds and means [95% confidence interval]. Cord blood leptin level by A) birth size groups and B) *LEPR* SNP rs9436303 *G-allele* carriage. The dotted lines represent the quantiles (from bottom to top: 10th, 25th, 75th and 90th percentiles). The symmetrical curves represent the probability densities at different values of leptin. SGA, small for gestational age; AGA, appropriate for gestational age; LGA, large for gestational age.



Supplementary Figure S2: Three fictitious BMI trajectories illustrating the results obtained in this analysis. Dotted line represents higher BMI at AP and later rebound; trajectory associated with carrying a rare G-allele of the rs9436303 SNP. Dashed line represents lower BMI at AP followed by earlier age at AR, trajectory associated with an increase in the cord blood leptin level; trajectory associated with carrying a rare G-allele of the rs9436303 SNP. And solid line a trajectory with normal BMI at AP and age at AR.