



HAL
open science

Conjectures for Invertible Diophantine Equations Of 3-Cyclic and 4-Cyclic Refined Integers

Oliver Von Shtawzen

► **To cite this version:**

Oliver Von Shtawzen. Conjectures for Invertible Diophantine Equations Of 3-Cyclic and 4-Cyclic Refined Integers. *journal of neutrosophic and fuzzy systems*, 2022, 3, pp.32 - 36. 10.54216/jnfs.030203 . hal-03739847

HAL Id: hal-03739847

<https://hal.science/hal-03739847v1>

Submitted on 28 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Conjectures for Invertible Diophantine Equations Of 3-Cyclic and 4-Cyclic Refined Integers

Oliver Von Shtawzen *

University Of Nizwa, Nizwa, Sultanate Of Oman

Email: Vonshtawzen1970abc@gmail.com

Abstract

The objective of this paper is to suggest two new conjectures concerning the invertible elements in 3-cyclic, and 4-cyclic refined neutrosophic rings of integers, where the invertibility condition shows that the solution of some Diophantine equations may determine the classification of the group of units of these algebraic rings.

Keywords: n-cyclic refined neutrosophic ring; n-cyclic refined integer; group of units; Turiyam rings

Introduction

In the theory of rings, it was very useful to build a bigger ring contains another one, which it is called a ring extension [1-2, 11-15]. Recently, it is studied for dealing the four way data analysis using Turiyam ring [16] and its extensive properties [17-19] for solving various decision making problems. However basic proof of some conjecture and equations are required to understand the usability of mathematical algebra [20]. To achieve this goal, the current paper focused on invertibility conditions for some Diophantine equations and its extension to Turiyam rings.

In [3], Abobala has presented a novel way to build a ring extension by using a neutrosophic idea of splitting and indeterminate into some other elements. In our work, we will concentrate the algebraic structure of this extension not the logical one.

If R is a ring, the corresponding n-cyclic refined neutrosophic ring is considered as

$$R_n(I) = \{a_0 + a_1I_1 + \dots + a_nI_n ; a_i \in R\} . \text{ It is clear that } R_n(I) \text{ contains } R.$$

The n-cyclic refined ring is different from the direct product of the ring R with itself $n+1$ times, that is because the multiplication is defined to be distributive on addition with an additional property which is similar to structure of cyclic group of integers, it can be described as follows:

$$I_i I_j = I_{(i+j \bmod n)} .$$

In [4], Sadiq has studied the group of units structure of 2-cyclic refined neutrosophic rings, and provided some interesting open problems. This motivated us to study the invertibility conditions in 3-cyclic and 4-cyclic refined rings of integers.

For more information about neutrosophic algebra [5-10] or four way Turiyam ring author can refer to [16-19] for basic understanding of mathematical proof [20].

For definitions and properties of n-cyclic refined neutrosophic rings see [3].

Main Discussion

Definition:

Let $Z_3(I) = \{a_0 + a_1I_1 + a_2I_2 + a_3I_3; a_i \in Z\}$ be the 3-cyclic refined neutrosophic ring of integers, then $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3$ is called invertible (unit) if and only if there exists $B = b_0 + b_1I_1 + b_2I_2 + b_3I_3$ such that $AB=1$.

Theorem:

Let $Z_3(I) = \{a_0 + a_1I_1 + a_2I_2 + a_3I_3; a_i \in Z\}$ be the 3-cyclic refined neutrosophic ring of integers, then $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3$ is invertible if and only if one of the following two Diophantine equations are true.

$$(a_0 + a_3)^3 + (a_1)^3 + (a_2)^3 - 3a_1a_2(a_0 + a_3) = 1 \text{ or } -1. \text{ With } a_0 \in \{1, -1\}.$$

Proof:

The study of the invertibility of A means that we are forced to study the 3-cyclic refined equation

$$AX=1. \text{ Where } X = x_0 + x_1I_1 + x_2I_2 + x_3I_3.$$

Now, we compute the product AX as follows:

$$AX = a_0x_0 + I_1[x_1(a_0 + a_3) + x_2a_2 + x_3a_1 + x_0a_1] + I_2[x_1a_1 + x_2(a_0 + a_3) + x_3a_2 + x_0a_2] + I_3[x_1a_2 + x_2a_1 + x_3(a_0 + a_3) + x_0a_3] = 1. \text{ This implies that:}$$

$$a_0x_0 = 1, \text{ thus } x_0, a_0 \in \{1, -1\}. \text{ Also,}$$

$$x_1(a_0 + a_3) + x_2a_2 + x_3a_1 = -x_0a_1, x_1a_1 + x_2(a_0 + a_3) + x_3a_2 = -x_0a_2, x_1a_2 + x_2a_1 + x_3(a_0 + a_3) = -x_0a_3.$$

So that, we got three linear equations with three variables x_1, x_2, x_3 . This system is solvable uniquely if and

$$\text{only if the determinant of the coefficients matrix is invertible, i.e. } \begin{vmatrix} a_0 + a_3 & a_2 & a_1 \\ a_1 & a_0 + a_3 & a_2 \\ a_2 & a_1 & a_0 + a_3 \end{vmatrix} = 1 \text{ or } -1.$$

By easy computing, we get the equations

$$(a_0 + a_3)^3 + (a_1)^3 + (a_2)^3 - 3a_1a_2(a_0 + a_3) = 1 \text{ or } -1. \text{ With } a_0 \in \{1, -1\}.$$

Conjecture 1:

The previous two equations has finite number of integer solutions (We mean that the order of the group of units of the 3-cyclic refined ring of integers is finite). Also, this number must be divisible by 6. (That is because the group of units of the integer ring Z has order 2, and the index of the extension is 3).

Definition:

Let $Z_4(I) = \{a_0 + a_1I_1 + a_2I_2 + a_3I_3 + a_4I_4; a_i \in Z\}$ be the 4-cyclic refined neutrosophic ring of integers, then $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3 + a_4I_4$ is called invertible (unit) if and only if there exists $B = b_0 + b_1I_1 + b_2I_2 + b_3I_3 + b_4I_4$ such that $AB=1$.

Theorem:

Let $Z_4(I) = \{a_0 + a_1I_1 + a_2I_2 + a_3I_3 + a_4I_4; a_i \in Z\}$ be the 4-cyclic refined neutrosophic ring of integers, then $A = a_0 + a_1I_1 + a_2I_2 + a_3I_3 + a_4I_4$ is invertible if and only if one of the following two Diophantine equations are true.

$$(a_0 + a_4)^4 - 4a_1a_3(a_0 + a_4)^2 + 4a_2a_3^2(a_0 + a_4) - 4a_1a_3a_2^2 - a_1^4 - a_3^4 + a_2^4 + 4a_2a_1^2(a_0 + a_4) + 2a_1^2a_3^2 - 2a_2^2(a_0 + a_4)^2 = 1 \text{ or } -1, \text{ with } a_0 \in \{1, -1\}.$$

Proof:

The study of the invertibility of A means that we are forced to study the 4-cyclic refined equation

$$AX=1. \text{ Where } X = x_0 + x_1I_1 + x_2I_2 + x_3I_3 + x_4I_4.$$

Now, we compute the product AX as follows:

$$AX = a_0x_0 + I_1[x_1(a_0 + a_4) + x_2a_3 + x_3a_2 + x_0a_1 + x_4a_1] + I_2[x_1a_1 + x_2(a_0 + a_4) + x_3a_3 + x_0a_2 + x_4a_2] + I_3[x_1a_2 + x_2a_1 + x_3(a_0 + a_4) + x_0a_3 + x_4a_3] + I_4[x_1a_3 + x_2a_2 + x_3a_1 + x_0a_4 + x_4(a_0 + a_4)] = 1.$$

This implies that:

$a_0x_0 = 1$, thus $a_0, x_0 = 1$ or -1 . On the other hand, we get:

$$x_1(a_0 + a_4) + x_2a_3 + x_3a_2 + x_4a_1 = -x_0a_1, x_1a_1 + x_2(a_0 + a_4) + x_3a_3 + x_4a_2 = -x_0a_2, x_1a_2 + x_2a_1 + x_3(a_0 + a_4) + x_4a_3 = -x_0a_3, x_1a_3 + x_2a_2 + x_3a_1 + x_4(a_0 + a_4) = -x_0a_4.$$

The previous system of 4 linear equations with 4 variables is solvable if and only if the following determinant is equal to 1 or -1.

$$\begin{vmatrix} a_0 + a_4 & a_3 & a_2 & a_1 \\ a_1 & a_0 + a_4 & a_3 & a_2 \\ a_2 & a_1 & a_0 + a_4 & a_3 \\ a_3 & a_2 & a_1 & a_0 + a_4 \end{vmatrix} = 1 \text{ or } -1. \text{ By computing the value of the determinant, we get the desired equations.}$$

Conjecture 2:

The equations shown in the previous Theorem have finite number of solutions. Also, this number is divisible by 8.

A Discussion Of The Case Of Symbolic Turiyam Ring of Integers

Recently, symbolic Turiyam rings [16] its modules [17] and matrix [18] is studied for dealing the data with Turiyam set [19]. In this section, the invertibility of a symbolic Turiyam integer is discussed as follows:

Let $A = x_0 + x_1T + x_2F + x_3I + x_4Y$, $B = y_0 + y_1T + y_2F + y_3I + y_4Y$ be two symbolic Turiyam integers such that $AB=1$.

We get:

$$x_0, y_0 = 1 \text{ or } -1, y_1(x_0 + x_1) = -x_1y_0, y_1(x_2) + y_2(x_0 + x_1 + x_2) = -x_2y_0,$$

$$y_1(x_3) + y_2(x_3) + y_3(x_0 + x_1 + x_2 + x_3 + x_4) + y_4(x_3) = 0,$$

$$y_1(x_4) + y_2(x_4) + y_3(0) + y_4(x_0 + x_1 + x_2 + x_3 + x_4) = 0.$$

The determinant of the coefficients matrix is:

$$\begin{vmatrix} x_0 + x_1 & 0 & 0 & 0 \\ x_2 & x_0 + x_1 + x_2 & 0 & 0 \\ x_3 & x_3 & x_0 + x_1 + x_2 + x_3 + x_4 & x_3 \\ x_4 & x_4 & 0 & x_0 + x_1 + x_2 + x_3 + x_4 \end{vmatrix} =$$

$$(x_0 + x_1)(x_0 + x_1 + x_2)(x_0 + x_1 + x_2 + x_3 + x_4)(x_0 + x_1 + x_2 + x_3 + x_4) = 1 \text{ or } -1.$$

The previous Diophantine equation can be solved easily, that is because it is written as a product.

From the previous discussion, we get the following result:

A symbolic Turiyam integer x is invertible if and only if $(x_0 + x_1)(x_0 + x_1 + x_2)(x_0 + x_1 + x_2 + x_3 + x_4)(x_0 + x_1 + x_2 + x_3 + x_4) = 1$ or -1 .

For example, the following symbolic Turiyam integer $x = 1 - 2I$ is invertible. In this way the proposed method can be helpful for dealing several decision making process as discussed in[20]. In near future the author will focus on solving some new problems of these conjectures to classify the group of units.

Conclusion

This paper discussed the invertibility of a 3-cyclic\4-cyclic refined neutrosophic integer. Also, we have presented two conjectures concerning the order of the group of units of 3-cyclic\4-cyclic refined neutrosophic ring of integers. Same time condition for the invertibility of a symbolic Turiyam integer is also discussed.

In the future, we aim to look for the solutions of these conjectures and to classify the group of units of these rings.

Acknowledgements: Author thanks the anonymous reviewers and editor for their valuable time and suggestions.

Funding: Author declares that there is no funding for this paper.

Conflict of Interest: Author declares that there is no conflict of interest for this paper.

References

- [1] A. G. Agarg`um, D. D. Anderson, and S. Valdes-Leon, "Factorization in commutative rings with zero divisors III", Rocky Mountain J. Math. 31 (2001), 1–21.
- [2] D. D. Anderson, and R. Markanda, "Unique factorization rings with zero divisors", Houston J. Math. 11 (1985), 15–30.
- [3] Abobala, M, "*n*-Cyclic Refined Neutrosophic Algebraic Systems Of Sub-Indeterminacies, An Application To Rings and Modules", International Journal of Neutrosophic Science, Vol. 12, pp. 81-95 . 2020.
- [4] Sadiq. B., " A Contribution To The group Of Units Problem In Some 2-Cyclic Refined neutrosophic Rings ", International Journal Of Neutrosophic Science, 2022.
- [5] Smarandache, F., " A Unifying Field in Logics: Neutrosophic Logic, Neutrosophy, Neutrosophic Set, Neutrosophic Probability", American Research Press. Rehoboth, 2003.
- [6] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81. 2020.
- [7] Ibrahim, M., and Abobala, M., "An Introduction To Refined Neutrosophic Number Theory", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [8] Olgun, N., Hatip, A., Bal, M., and Abobala, M., " A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", International Journal of Neutrosophic Science, Vol. 16, pp. 72-79, 2021.
- [9] Abobala, M., Bal, M., and Hatip, A., " A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", International Journal of Neutrosophic Science, Vol. 17, 2021.
- [10] Abobala, M., Bal, M., and Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.
- [11] P. Ribenboim, "Rings of generalized power series. II. Units and zero-divisors", J. Algebra 168 (1994), 71–89.
- [12] Z. Pogorzaly, "A generalization of trivial extension algebras", J. Pure Appl. Algebra 203 (2005), 145–165.
- [13] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [14] E. Adeleke. A. Agboola., and F. Smarandache. "Refined Neutrosophic Rings II". International Journal of Neutrosophic Science, Vol. 2, 2020. pp. 89-94.

- [15] D. G. Northcott, "Lessons on Rings, Modules, and Multiplicities", Cambridge Univ. Press, Cambridge, 1968.
- [16] Singh, P. K., Ahmad, K., Bal, M., and Aswad, M., " On The Symbolic Turiyam rings", Journal of Neutrosophic and Fuzzy Systems, Vol. 1, Issue 2, pp. 80-88, 2022.
- [17] Bal M., Singh P. K., and Ahmad K. D., " An Introduction To The Symbolic Turiyam R-Modules and Turiyam Modulo Integers", Journal of Neutrosophic and Fuzzy Systems, Vol. 2 , No. 2 , (2022) : 8-19
- [18] Bal M., Singh P. K., and Ahmad K. D., "A Short Introduction To The Concept Of Symbolic Turiyam Matrix", Journal of Neutrosophic and Fuzzy Systems, Vol. 2 , No. 1 , (2022) : 88-99
- [19] Singh, P.K., " Data With Turiyam Set for Fourth Dimension Quantum Information Processing", Journal of Neutrosophic and Fuzzy Systems, Vol.1, 2022.
- [20] Singh P. K., "A Note on Basic Proof of Some Famous Mathematical Theorem and Its Illustration", Journal of Neutrosophic and Fuzzy Systems, Vol. 3 , No. 1 , (2022) : 39-53