

The renormalized volume of a 4-dimensional Ricci-flat ALE space

Olivier Biquard, Hans-Joachim Hein

▶ To cite this version:

Olivier Biquard, Hans-Joachim Hein. The renormalized volume of a 4-dimensional Ricci-flat ALE space. Journal of Differential Geometry, 2023, 123 (3), pp.411-429. 10.4310/jdg/1683307004. hal-03740923

HAL Id: hal-03740923 https://hal.science/hal-03740923

Submitted on 21 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE RENORMALIZED VOLUME OF A 4-DIMENSIONAL RICCI-FLAT ALE SPACE

Olivier Biquard & Hans-Joachim Hein

Abstract

We introduce a natural definition of the renormalized volume of a 4-dimensional Ricci-flat ALE space. We then prove that the renormalized volume is always less or equal than zero, with equality if and only if the ALE space is isometric to its asymptotic cone. Currently the only known examples of 4-dimensional Ricciflat ALE spaces are Kronheimer's gravitational instantons and their quotients, which are also known to be the only possible examples of special holonomy. We calculate the renormalized volume of these spaces in terms of Kronheimer's period map.

1. Introduction

This article grew out of an attempt to understand 4-dimensional Ricci-flat asymptotically locally Euclidean (ALE) manifolds. The standard example of such a space is the Eguchi-Hanson metric on $T^*\mathbb{S}^2$ [12]. The Eguchi-Hanson metric is actually hyper-Kähler (see Calabi [6]), and a classification of hyper-Kähler ALE 4-manifolds was given by Kronheimer [18, 19]. Finite free quotients of Kronheimer spaces as classified by Şuvaina [25] and Wright [27] are also examples of Ricci-flat ALE 4-manifolds. A fundamental open question due to Bando-Kasue-Nakajima [4] asks whether there exist any other examples. Using Witten's proof of the positive mass theorem [26], Nakajima [21] showed that such examples can never be spin. See Lock-Viaclovsky [20] for some additional restrictions based on the Hitchin-Thorpe inequality of [21]. We did not solve this problem but we have found an interesting new volume property of 4-dimensional Ricci-flat ALE spaces, which we will now explain.

Let (M, g) be a 4-dimensional Ricci-flat ALE manifold, or a 4-dimensional Ricci-flat ALE orbifold with at worst finitely many isolated singularities (we shall call such an orbifold an *ALE space*). By [4], this means that there exist a finite subgroup Γ of SO(4) acting freely on \mathbb{S}^3 and a diffeomorphism $\Phi : (\mathbb{R}^4 \setminus B_1(0))/\Gamma \to M \setminus K$ for some compact subset $K \subset M$ such that for all $k \in \mathbb{N}_0$,

(1.1)
$$|\nabla_{g_0}^k(\Phi^*g - g_0)|_{g_0} = O(r^{-4-k}) \text{ as } r \to \infty,$$

where g_0 denotes the Euclidean metric on \mathbb{R}^4 or on \mathbb{R}^4/Γ . By [7, 8], this behavior already follows from the much weaker assumption that (M, g)is a 4-dimensional complete Ricci-flat manifold (or a complete Ricci-flat orbifold with finite singular set) of maximal volume growth at infinity.

Assume that $\Gamma \neq \{1\}$. By [11] there exist $K \subset M$ compact and a number $\rho_0 > 0$ such that $M \setminus K$ is uniquely foliated by hypersurfaces Σ_{ρ} ($\rho > \rho_0$) of constant mean curvature $3/\rho$ such that Σ_{ρ} is a normal graph of height $O(\rho^{-3})$ over $\Phi(\partial B_{\rho}(0)/\Gamma)$ for any diffeomorphism Φ as in (1.1).

Given these preliminaries, our first main result may be stated as follows.

Theorem A. Let (M, g) be a 4-dimensional Ricci-flat ALE space with $\Gamma \neq \{1\}$.

(1) Let $\Omega_{\rho} \subset M$ denote the domain interior to Σ_{ρ} . Then the quantity

$$\operatorname{Vol}_g(\Omega_{\rho}) - \operatorname{Vol}_{g_0}(B_{\rho}(0)/\Gamma)$$

has a finite limit as $\rho \to \infty$. We refer to this limit as the renormalized volume \mathcal{V} of (M, g).

(2) The renormalized volume satisfies $\mathcal{V} \leq 0$, with equality if and only if $(M,g) \cong (\mathbb{R}^4/\Gamma, g_0)$.

Ozuch [22, Thm 0.7] has recently used this result to prove that spherical or hyperbolic 4-orbifolds cannot be Gromov-Hausdorff limits of smooth Einstein 4-manifolds under certain conditions.

There is a vast literature on renormalized volumes of open manifolds. Our motivation was not so much to extend these ideas to a new setting but to interpret some of our computations for Ricci-flat ALE spaces geometrically. However, as it turns out, the statement of Theorem A is formally similar to several known results. The examples we are aware of are Anderson's comparison theorem [1] for the classical renormalized volume [14] of asymptotically hyperbolic Einstein 4-manifolds, Hu-Ji-Shi's comparison theorem [15] for the Brendle-Chodosh relative volume [5] of asymptotically hyperbolic *n*-manifolds with sharp lower scalar curvature bounds, and the identification [10, 13, 17] of Huisken's isoperimetric mass [16] of asymptotically flat 3-manifolds of nonnegative scalar curvature with the ADM mass [2], combined with the statement of the positive mass theorem [24, 26].

To prove Theorem A, we will first construct a special ALE diffeomorphism Φ as in (1.1) such that $\Phi^*g - g_0$ admits an asymptotic expansion whose leading term has vanishing trace, divergence, and contraction with ∂_r . Due to these properties, the coordinate spheres $\Phi(\partial B_{\rho}(0)/\Gamma)$ are CMC to one higher order than expected. Thus, Σ_{ρ} is a normal graph of height $O(\rho^{-4})$ rather than $O(\rho^{-3})$ over $\Phi(\partial B_{\rho}(0)/\Gamma)$, so by changing Φ very slightly we are able to create a CMC gauge without changing the leading term of the metric. The precise statement is as follows. **Theorem B.** Let (M, g) be a 4-dimensional Ricci-flat ALE space with $\Gamma \neq \{1\}$. Let $\{\Sigma_{\rho}\}_{\rho > \rho_0}$ be the canonical CMC foliation of the end of M constructed in [11]. Then for any $k_0 \in \mathbb{N}$ there exists a diffeomorphism $\Phi : (\mathbb{R}^4 \setminus B_1(0))/\Gamma \to M \setminus K$ of class at least C^{k_0+1} such that $\Phi(\partial B_{\rho}(0)/\Gamma) = \Sigma_{\rho}$ for all $\rho > \rho_0$ and such that there exists a decomposition

(1.2)
$$\Phi^* g - g_0 = h_0 + h',$$

where h_0 is either zero or comparable above and below to r^{-4} in g_0 -norm and satisfies

(1.3)
$$\mathcal{L}_{r\partial_r}h_0 = -2h_0, \\ d_r \,\lrcorner\, h_0 = 0, \ \operatorname{tr}_{g_0}h_0 = 0, \ \operatorname{div}_{g_0}h_0 = 0, \\ \Delta_{g_0}h_0 = 0,$$

and where the remainder h' can be estimated by

(1.4)
$$\sum_{k=0}^{k_0} r^k |\nabla_{g_0}^k h'|_{g_0} = O_{k_0}(r^{-5}) \text{ as } r \to \infty.$$

The space of all symmetric 2-tensors h_0 satisfying (1.3) on $\mathbb{R}^4 \setminus \{0\}$ is isomorphic to $\mathbf{S}^4_+ \oplus \mathbf{S}^4_-$ as a representation of SO(4), where \mathbf{S}^ℓ_\pm denotes the two $(\ell + 1)$ -dimensional fundamental representations of Spin(4). Each element of \mathbf{S}^4_+ arises as the leading term h_0 of a suitable Kronheimer gravitational instanton. All of this will be explained in Section 2. Here we only note that the property $\operatorname{tr}_{g_0} h_0 = 0$ allows us to conclude that the renormalized volume \mathcal{V} exists, which proves Theorem A(1).

The second key step is to construct a function u on M with $\Delta u = 8$ and $\Phi^* u = r^2 + o(1)$, and to integrate by parts in the formula $\operatorname{div}(\operatorname{Hess}_0 u) = \operatorname{Ric} du = 0$. Theorem A(2) follows from an analysis of the boundary terms, relying on the properties $\partial_r \,\lrcorner\, h_0 = 0$, $\operatorname{tr}_{g_0} h_0 = 0$, and $\operatorname{div}_{g_0} h_0 = 0$.

It is worth pointing out that the proof of Theorem A goes through on every 4-dimensional ALE space with $\Gamma \neq \{1\}$ and Ric ≥ 0 that satisfies the conclusion of Theorem B. The property $\Delta_{g_0}h_0 = 0$ is actually not needed for this, although in the Ricci-flat case this property is automatic and allows us to prove the crucial properties $\partial_r \,\lrcorner\, h_0 = 0$, $\operatorname{tr}_{g_0}h_0 = 0$, and $\operatorname{div}_{g_0}h_0 = 0$ because harmonic tensors on $\mathbb{R}^4 \setminus \{0\}$ are easy to classify. Thus, Theorem A certainly still holds under the weaker assumption that Ric ≥ 0 globally and Ric $= O(r^{-6-\varepsilon})$ at infinity. However, it could be that these two conditions together imply that Ric = 0 globally. Using [21], one can show that this is true if M is spin.

Let us also point out that, in the Ricci-flat case, the leading term h_0 of Theorem B can actually be zero even if (M, g) is not isometric to $(\mathbb{R}^4/\Gamma, g_0)$. Indeed, using a computation of Auvray [3], we will see in Sections 2.2.2–2.3 that h_0 vanishes on a Kronheimer space if and only

if the preimages in $H_c^2(M)$ of the three Kähler classes in $H^2(M)$ are orthonormal with respect to the intersection form. Thanks to Kronheimer's Torelli theorem [19], this is possible if and only if $b^2(M) \ge 3$.

In the same spirit, our last main result computes the renormalized volume of a Kronheimer space. Kronheimer's Torelli theorem [19], which we just mentioned, provides a bijection between the set of all ALE hyper-Kähler metrics up to the obvious action of $\text{Diff}_0(M)$ on a fixed smooth manifold M, where M is determined by $\Gamma \subset SU(2)$, and elements of $H_c^2(M) \otimes \mathbb{R}^3$ up to the obvious SO(3)-action. The bijection, which is also known as the *period map*, sends a hyper-Kähler metric to the preimages in $H_c^2(M)$ of the three Kähler classes in $H^2(M)$. The vectors in the image of the period map satisfy a nondegeneracy condition, which is an open dense condition, but this condition can be dropped if we allow for degenerations of smooth ALE hyper-Kähler metrics on Mwith at worst isolated orbifold singularities. There is a further natural identification of $H_c^2(M)$ together with the intersection form with a Cartan subalgebra of a certain simple Lie algebra together with the Killing form.

Theorem C. Let Γ be a finite subgroup of SU(2) acting freely on \mathbb{S}^3 . Let \mathfrak{h} be a Cartan subalgebra of the Lie algebra associated with Γ via the McKay correspondence. Choose any element $\zeta \in \mathfrak{h} \otimes \mathbb{R}^3$. Let (M, g) be the unique Kronheimer gravitational instanton with period point ζ up to the obvious action of SO(3) on $\mathfrak{h} \otimes \mathbb{R}^3$. Then in terms of the Killing form inner product on \mathfrak{h} ,

(1.5)
$$\mathcal{V}(M,g) = -\frac{1}{6}|\zeta|^2.$$

The basic idea behind Theorem A is to determine when the scaling vector field $r\partial_r$ on \mathbb{R}^4/Γ can be extended to a conformal Killing field on (M, g). We conclude this paper with some remarks on the analogous question for Killing fields instead of conformal Killing fields.

Acknowledgments. OB received support from the "Investissements d'Avenir" program ANR-10-IDEX-0001-02 PSL. HJH was partially supported by NSF grant DMS-1745517. We are grateful to the referees for useful comments, and to Michael Anderson for pointing out the similarity between our renormalized volume and Huisken's isoperimetric mass.

2. Asymptotics of Ricci-flat ALE metrics in dimension 4

Recall that the *Bianchi operator* of a Riemannian manifold (M, g) is defined by

$$\mathcal{B}_g: \Gamma(\operatorname{Sym}^2 T^*M) \to \Gamma(T^*M), \ h \mapsto \operatorname{div}_g(h - \frac{1}{2}(\operatorname{tr}_g h)g).$$

Unless stated otherwise, let (M, g) be a 4-dimensional Ricci-flat ALE space with $\Gamma \neq \{1\}$.

2.1. Preparation of an asymptotic expansion. Fix an arbitrary diffeomorphism Φ as in (1.1). By standard Fredholm theory arguments in weighted Hölder spaces, it is possible to modify Φ to satisfy the Bianchi gauge condition relative to g and g_0 , meaning that the tensor

$$h := \Phi^* g - g_0$$

satisfies $\mathcal{B}_{g_0}h = 0$, although a priori Φ will then only be of class C^{k_0+1} for some $k_0 \in \mathbb{N}$, (1.1) will only hold up to k_0 derivatives, and we may need to relax the decay rate in (1.1) slightly. However, we do have the freedom of prescribing an arbitrarily high cutoff k_0 . Moreover, since the Ricci-flat equation is elliptic in any Bianchi gauge, h actually admits an asymptotic expansion.

One can be completely precise about the shape of this expansion [9] but for us the first term will be sufficient. By a standard iteration, the first term is a Γ -equivariant harmonic function

$$h_0: \mathbb{R}^4 \setminus B_1(0) \to \mathrm{Sym}^2 \mathbb{R}^4$$

that decays at infinity. As explained in [8, Thm 5.103], thanks to the Bianchi gauge condition and because $\Gamma \neq \{1\}$, we can assume that each component function of h_0 is homogeneous of degree -4. This then justifies the rate -4 chosen in (1.1). We also learn that the nonlinear terms of the Ricci curvature are at worst $O(r^{-10})$, so we immediately get

(2.1)
$$h = h_0 + h_1 + h_2 + h_3 + h',$$

where the first four terms h_0, h_1, h_2, h_3 are homogeneous,

(2.2)
$$\mathcal{L}_{r\partial_r}h_k = (-2-k)h_k, \ |h_k|_{g_0} \sim r^{-4-k},$$

and satisfy the pair of equations

(2.3)
$$\mathcal{B}_{g_0}h_k = 0, \ \Delta_{g_0}h_k = 0.$$

Moreover, we are free to assume that the remainder h' satisfies

(2.4)
$$\sum_{k=0}^{k_0} r^k |\nabla_{g_0}^k h'|_{g_0} \leqslant C(k_0, \varepsilon) r^{-8+\varepsilon}$$

for any given $k_0 \in \mathbb{N}$ and $\varepsilon \in (0, 1)$. For simplicity, we will from now on absorb h_1, h_2, h_3 into h', so that (2.1) reads $h = h_0 + h'$ and (2.4) holds with $C(k_0, \varepsilon)r^{-8+\varepsilon}$ replaced by $C(k_0)r^{-5}$.

2.2. Two examples of possible leading terms. The remainder of Section 2 is dedicated to a classification of the possible leading terms h_0 . We begin by describing two types of examples.

2.2.1. Harmonic gauge terms. Somewhat surprisingly, the Bianchi gauge condition allows for some residual gauge freedom to leading order. Indeed, recall that for any arbitrary Riemannian manifold (M, g) and any vector field X on M we have the Bochner type formula

(2.5)
$$\mathcal{B}_g(\mathcal{L}_X g) = -\nabla^* \nabla X + \operatorname{Ric} X.$$

Therefore, if g is Ricci-flat and if X is harmonic, then $h := \mathcal{L}_X g$ is in Bianchi gauge $(\mathcal{B}_g h = 0)$ and satisfies the linearized Ricci-flat equation. So if X on $(\mathbb{R}^4 \setminus \{0\})/\Gamma$ satisfies

- 1) X is harmonic: $\Delta_{g_0} X = 0$,
- 2) $\mathcal{L}_{r\partial_r}X = -4X$, so in particular $|X|_{g_0} \sim r^{-3}$,

then $h_0 := \mathcal{L}_X g_0$ will solve equations (2.2) and (2.3) for k = 0, as desired.

On $\mathbb{R}^4 \setminus \{0\}$ the vector fields satisfying (1)–(2) are exactly the ones of the form

(2.6)
$$X = \frac{1}{r^4} \ell_{ij} x^j \frac{\partial}{\partial x^i}$$

for an arbitrary matrix $L = (\ell_{ij}) \in \mathbb{R}^{4 \times 4}$. So the Γ -invariant matrices $L \in (\mathbb{R}^{4 \times 4})^{\Gamma}$ give rise to such a harmonic vector field X on $(\mathbb{R}^4 \setminus \{0\})/\Gamma$ and hence to a possible term $h_0 = \mathcal{L}_X g_0$.

For reference, let us note that thanks to the Bianchi condition, h_0 is trace-free if and only if it is divergence-free, but h_0 may not satisfy either of these conditions. In fact,

(2.7)
$$\operatorname{tr}_{g_0} h_0 = \operatorname{tr}_{g_0}(\mathcal{L}_X g_0) = 2\operatorname{div}_{g_0} X = \frac{2}{r^4} (\delta_{ij} - 4\frac{x^i x^j}{r^2}) \ell_{ij},$$

which vanishes if and only if L is either a multiple of the identity or skew-symmetric. A special case is $L = L_0 = -2$ Id, so that $X = X_0 = \nabla_{g_0}(\frac{1}{r^2})$, which is harmonic because $\frac{1}{r^2}$ is. We then have

(2.8)
$$h_0 = \mathcal{L}_{X_0} g_0 = 2 \operatorname{Hess}_{g_0}(\frac{1}{r^2}) = 2 \nabla_{g_0}(-\frac{2}{r^3}\partial_r) = -\frac{4}{r^4}(g_0 - 4dr^2),$$

which is trace-free by inspection or because X_0 is divergence-free because $\frac{1}{r^2}$ is harmonic.

More abstractly, let **U** denote the space of all harmonic gauge terms $h_0 = \mathcal{L}_X g_0$ on $\mathbb{R}^4 \setminus \{0\}$ with X as in (2.6). As a representation of SO(4) this obviously decomposes into irreducibles as

$\mathbf{U}=\mathbf{U}_1\oplus\mathbf{U}_2\oplus\mathbf{U}_3,$

where $\mathbf{U}_1 = \mathbb{R}$ is spanned by the Hessian of the Green's function in (2.8), $\mathbf{U}_2 = \Lambda^2 \mathbb{R}^4$ corresponds to taking L to be skew-symmetric in (2.6), and $\mathbf{U}_3 = \operatorname{Sym}_0^2 \mathbb{R}^4$ corresponds to taking L to be trace-free symmetric in (2.6). If $h_0 \in \mathbf{U}_3$, then $\operatorname{tr}_{g_0} h_0 = q_0/r^6$ for the harmonic polynomial $q_0(x) := -8\ell_{ij}x^ix^j$ (see (2.7)), so the trace defines an equivariant projection $\mathbf{U} \to \operatorname{Sym}_0^2 \mathbb{R}^4 = \mathbf{U}_3$. **2.2.2. Kronheimer terms.** Let Γ be a finite subgroup of SU(2) that acts freely on \mathbb{S}^3 . Kronheimer's period map [19] is a bijection between hyper-Kähler ALE metrics asymptotic to \mathbb{R}^4/Γ and triples

$$(\zeta_1,\zeta_2,\zeta_3)\in\mathfrak{h}\otimes\mathbb{R}^3$$

up to the obvious SO(3)-action, where \mathfrak{h} is a Cartan subalgebra of the Lie algebra associated with Γ via the McKay correspondence. As explained in [3, Thm 2.1], Kronheimer constructed a special gauge with respect to which the tensor $-(2\pi^2/|\Gamma|)r^6h_0$ takes the form

(2.9)

$$\begin{aligned} |\zeta_1|^2((rdr)^2 + \alpha_1^2 - \alpha_2^2 - \alpha_3^2) + |\zeta_2|^2((rdr)^2 + \alpha_2^2 - \alpha_3^2 - \alpha_1^2) \\ &+ |\zeta_3|^2((rdr)^2 + \alpha_3^2 - \alpha_1^2 - \alpha_2^2) \\ &+ 2\langle\zeta_1, \zeta_2\rangle(\alpha_1 \cdot \alpha_2 - rdr \cdot \alpha_3) + 2\langle\zeta_1, \zeta_3\rangle(\alpha_1 \cdot \alpha_3 - rdr \cdot \alpha_2) \\ &+ 2\langle\zeta_2, \zeta_3\rangle(\alpha_2 \cdot \alpha_3 - rdr \cdot \alpha_1). \end{aligned}$$

Here $\alpha_j = I_j(rdr)$, and (I_1, I_2, I_3) is the standard triple of complex structures on \mathbb{R}^4 given by

(2.10)
$$I_1(x^1, x^2, x^3, x^4) = (-x^2, x^1, -x^4, x^3),$$
$$I_2(x^1, x^2, x^3, x^4) = (-x^3, x^4, x^1, -x^2),$$
$$I_3 = I_1 I_2.$$

Then obviously (2.2) is satisfied, and, by computation,

(2.11)
$$\operatorname{tr}_{g_0} h_0 = 0, \ \operatorname{div}_{g_0} h_0 = 0, \ \Delta_{g_0} h_0 = 0,$$

so (2.3) is satisfied as well.

Let $\operatorname{SU}(2)_{\pm}$ denote the two canonical subgroups of $\operatorname{SO}(4)$, with $\mathfrak{su}(2)_{+} = \langle I_1, I_2, I_3 \rangle$. Then every Kronheimer tensor h_0 as in (2.9) is invariant under $\operatorname{SU}(2)_{-}$. Notice that $\Gamma \subset \operatorname{SU}(2)_{-}$.

The following lemma clarifies the structure of the tensors (2.9) using representation theory. Let \mathbf{S}_{\pm}^{ℓ} be the $(\ell + 1)$ -dimensional irreducible representation of $\mathrm{SU}(2)_{\pm}$. The irreducible representations of $\mathrm{Spin}(4)$ are given by $\mathbf{S}_{\pm}^{\ell} \otimes \mathbf{S}_{\pm}^{m}$ ($\ell, m \in \mathbb{N}_{0}$). This descends to $\mathrm{SO}(4)$ if and only if $\ell + m$ is even.

Lemma 2.1. Let \mathbf{V} denote the space of all symmetric 2-tensors on $\mathbb{R}^4 \setminus \{0\}$ satisfying (2.11) whose component functions are (-4)homogeneous. Recall the SO(4)-invariant subspaces $\mathbf{U}_1, \mathbf{U}_2 \subset \mathbf{V}$ from Section 2.2.1 and let $\mathbf{W} \subset \mathbf{V}$ be the invariant complement of $\mathbf{U}_1 \oplus \mathbf{U}_2$. Let

$$\mathbf{F}: \mathrm{Sym}^2 \mathbb{R}^3 \to \mathbf{V}$$

be the linear map that sends $(\langle \zeta_i, \zeta_j \rangle) \in \text{Sym}^2 \mathbb{R}^3$ to the Kronheimer tensor $h_0 \in \mathbf{V}$ with $-r^6 h_0$ given by (2.9). Then $\operatorname{pr}_{\mathbf{W}} \circ \mathbf{F}$ is a linear isomorphism onto a subrepresentation $\mathbf{S}^4_+ \subset \mathbf{W}$. *Proof.* Even though this is not logically necessary, we begin by constraining the possible irreducible representations contained in **V**. By construction, if $h_0 \in \mathbf{V}$, then all components of h_0 take the form q_0/r^6 , where q_0 is a harmonic quadratic polynomial. Thus, **V** is an invariant subspace of

(2.12) $(\operatorname{Sym}_0^2 \mathbb{R}^4)^{\otimes 2} = (\mathbf{S}_+^2 \otimes \mathbf{S}_-^2)^{\otimes 2} = (\mathbf{S}_+^4 \oplus \mathbf{S}_+^2 \oplus \mathbb{R}) \otimes (\mathbf{S}_-^4 \oplus \mathbf{S}_-^2 \oplus \mathbb{R}).$

The subspace $\mathbf{U}_1 \subset \mathbf{V}$ corresponds to the \mathbb{R} -component in (2.12) and is spanned by

(2.13)
$$\frac{1}{r^6}(3(rdr)^2 - \alpha_1^2 - \alpha_2^2 - \alpha_3^2)$$

according to (2.8). The subspace $\mathbf{U}_2 \subset \mathbf{V}$ corresponds to the $(\mathbf{S}^2_+ \oplus \mathbf{S}^2_-)$ component in (2.12). It is clear that the three tensors $\mathcal{L}_X g_0$, where $X = I(\nabla_{g_0}(\frac{1}{r^2}))$ for $I = I_1, I_2, I_3$, generate the \mathbf{S}^2_+ -part of this space.
Calculating these tensors, we obtain that the \mathbf{S}^2_+ -part of \mathbf{U}_2 is spanned
by

(2.14)
$$\frac{1}{r^6}(rdr) \cdot \alpha_1, \quad \frac{1}{r^6}(rdr) \cdot \alpha_2, \quad \frac{1}{r^6}(rdr) \cdot \alpha_3.$$

By the definition of **F** we have for all $\zeta \in \text{Sym}^2 \mathbb{R}^3$ that

$$-r^{6}\mathbf{F}(\zeta) = \zeta_{11}((rdr)^{2} + \alpha_{1}^{2} - \alpha_{2}^{2} - \alpha_{3}^{2}) + \zeta_{22}((rdr)^{2} + \alpha_{2}^{2} - \alpha_{3}^{2} - \alpha_{1}^{2}) + \zeta_{33}((rdr)^{2} + \alpha_{3}^{2} - \alpha_{1}^{2} - \alpha_{2}^{2}) + 2\zeta_{12}(\alpha_{1} \cdot \alpha_{2} - rdr \cdot \alpha_{3}) + 2\zeta_{13}(\alpha_{1} \cdot \alpha_{3} - rdr \cdot \alpha_{2}) + 2\zeta_{23}(\alpha_{2} \cdot \alpha_{3} - rdr \cdot \alpha_{1}).$$

Define a new linear map **G** on $\text{Sym}^2 \mathbb{R}^3$ by

(2.15)
$$-r^{6}\mathbf{G}(\zeta) := \frac{2}{3}(\zeta_{11}(2\alpha_{1}^{2} - \alpha_{2}^{2} - \alpha_{3}^{2}) + \zeta_{22}(2\alpha_{2}^{2} - \alpha_{3}^{2} - \alpha_{1}^{2}) + \zeta_{33}(2\alpha_{3}^{2} - \alpha_{1}^{2} - \alpha_{2}^{2})) + 2\zeta_{12}\alpha_{1} \cdot \alpha_{2} + 2\zeta_{13}\alpha_{1} \cdot \alpha_{3} + 2\zeta_{23}\alpha_{2} \cdot \alpha_{3}.$$

It is easy to check using (2.13), (2.14) that $\mathbf{F} - \mathbf{G}$ takes values in $\mathbf{U}_1 \oplus \mathbf{U}_2$. Because $(\alpha_1, \alpha_2, \alpha_3)$ is a basis of \mathbf{S}^2_+ and because $\mathbf{S}^4_+ = \mathrm{Sym}^2_0 \mathbf{S}^2_+$, it is clear from (2.15) that \mathbf{G} defines an isomorphism onto the \mathbf{S}^4_+ -component in (2.12), so that necessarily $\mathbf{G} = \mathrm{pr}_{\mathbf{W}} \circ \mathbf{F}$. q.e.d.

Remark 2.2. We can transport the given $SU(2)_+$ -action on $S^4_+ \subset V$ to $Sym^2\mathbb{R}^3$ by using the linear isomorphism $\operatorname{pr}_{\mathbf{W}} \circ \mathbf{F}$ of Lemma 2.1. We obtain an irreducible representation of $SU(2)_+$ on $Sym^2\mathbb{R}^3$ with kernel equal to $\{\pm \operatorname{Id}_{\mathbb{R}^4}\}$, and hence an irreducible representation of SO(3) on $Sym^2\mathbb{R}^3$. This is *not* equal to the canonical SO(3)-action on $Sym^2\mathbb{R}^3$ although it is of course conjugate to it.

Remark 2.3. From (2.14) and (2.15), if some element $h_0 \in \mathbf{V}$ belongs to one of the two invariant subspaces \mathbf{S}^2_+ or \mathbf{S}^4_+ , then $h_0(\partial_r, \partial_r) =$ 0. Since im $\mathbf{F} \subset \mathbf{U}_1 \oplus \mathbf{S}^2_+ \oplus \mathbf{S}^4_+$, it follows that

$$(\mathrm{pr}_{\mathbf{U}_1} \circ \mathbf{F})(\zeta) = \mathbf{F}(\zeta)(\partial_r, \partial_r) \cdot \frac{r^4}{6} \mathrm{Hess}_{g_0}(\frac{1}{r^2}) = -\frac{1}{6} \mathrm{tr}(\zeta) \cdot \mathrm{Hess}_{g_0}(\frac{1}{r^2})$$

for all $\zeta \in \text{Sym}^2 \mathbb{R}^3$. Thus, if $\zeta = (\langle \zeta_i, \zeta_j \rangle)$ with $(\zeta_1, \zeta_2, \zeta_3) \in \mathfrak{h} \otimes \mathbb{R}^3$, then the coefficient has a sign and vanishes if and only if the associated Kronheimer space is isometric to \mathbb{R}^4/Γ . This observation led us to our main theorem and is in fact equivalent to it in the case of a Kronheimer space.

2.3. Classification of the possible leading terms. Lemma 2.1 motivates the following:

Definition 2.4. A reduced Kronheimer term is an element of $\mathbf{S}^4_+ = \operatorname{im}(\operatorname{pr}_{\mathbf{W}} \circ \mathbf{F}) \subset \mathbf{V}$. Explicitly, a reduced Kronheimer term is a symmetric 2-tensor h_0^+ on $\mathbb{R}^4 \setminus \{0\}$ of the form

(2.16)
$$-\frac{3}{2}r^{6}h_{0}^{+} = \zeta_{11}(2\alpha_{1}^{2} - \alpha_{2}^{2} - \alpha_{3}^{2}) + \zeta_{22}(2\alpha_{2}^{2} - \alpha_{3}^{2} - \alpha_{1}^{2}) + \zeta_{33}(2\alpha_{3}^{2} - \alpha_{1}^{2} - \alpha_{2}^{2}) + 3\zeta_{12}(\alpha_{1} \cdot \alpha_{2}) + 3\zeta_{13}(\alpha_{1} \cdot \alpha_{3}) + 3\zeta_{23}(\alpha_{2} \cdot \alpha_{3}),$$

where $\alpha_j = I_j(rdr)$ for I_1, I_2, I_3 as in (2.10) and where (ζ_{ij}) is any symmetric 3×3 matrix.

A reduced Kronheimer term for the opposite orientation is a symmetric 2-tensor h_0^- on $\mathbb{R}^4 \setminus \{0\}$ of the form $h_0^- = R^* h_0^+$ for some reduced Kronheimer term h_0^+ and some $R \in O(4) \setminus SO(4)$.

Remark 2.5. Using this language, the discussion of Section 2.2.2 implies that for $\Gamma \subset SU(2)$ and \mathfrak{h} a Cartan subalgebra of the simple Lie algebra associated with Γ via the McKay correspondence, for $\zeta \in \mathfrak{h} \otimes \mathbb{R}^3$, and for (M, g) the unique Kronheimer gravitational instanton with period point ζ , there exist ALE coordinates for (M, g) such that the leading term h_0 satisfies $h_0 = (|\Gamma|/2\pi^2)h_0^+$, where h_0^+ is the reduced Kronheimer term given by (2.16) for $\zeta_{ij} = \langle \zeta_i, \zeta_j \rangle$. In particular, $h_0 = 0$ if and only if $\langle \zeta_i, \zeta_j \rangle = \delta_{ij}$. However, we should also recall from Remark 2.3 that in Kronheimer's *original* ALE coordinates, the leading term of the metric vanishes if and only if $(M, g) \cong (\mathbb{R}^4/\Gamma, g_0)$.

We are now in position to classify the possible leading terms of Ricciflat ALE metrics.

Proposition 2.6. Let h_0 be a symmetric 2-tensor on $\mathbb{R}^4 \setminus \{0\}$ that satisfies

(2.17)
$$\mathcal{L}_{r\partial_r}h_0 = -2h_0, \ \mathcal{B}_{g_0}h_0 = 0, \ \Delta_{g_0}h_0 = 0.$$

(1) There exists a unique decomposition

(2.18)
$$h_0 = h_0^+ + h_0^- + \mathcal{L}_{X_1} g_0 + \mathcal{L}_{X_2} g_0 + \mathcal{L}_{X_3} g_0,$$

where h_0^{\pm} are reduced Kronheimer terms for the two orientations and where

(2.19)
$$X_k(x) = \frac{1}{r^4} L_k x, \ L_k \in \mathbb{R}^{4 \times 4},$$

with L_1 a multiple of the identity, L_2 skew-symmetric, and L_3 trace-free symmetric. All terms of the decomposition (2.18) again satisfy all the conditions of (2.17).

(2) We have the following characterizations:

(2.20)
$$\operatorname{tr}_{g_0} h_0 = 0 \Longleftrightarrow \operatorname{div}_{g_0} h_0 = 0 \Longleftrightarrow X_3 = 0,$$

$$(2.21) \qquad \qquad \partial_r \,\lrcorner\, h_0 = 0 \Longleftrightarrow X_1 = X_2 = X_3 = 0.$$

(3) Let G be any subgroup of SO(4). If h_0 is G-invariant, then so are h_0^+ , h_0^- , L_1 , L_2 , L_3 .

Proof. Define an equivariant linear map

(2.22)
$$\mathbf{H} : \operatorname{Sym}^{2} \mathbb{R}^{4} \otimes \operatorname{Sym}_{0}^{2} \mathbb{R}^{4} \to \mathbb{R}^{4} \otimes \operatorname{Sym}^{3} \mathbb{R}^{4},$$
$$h \otimes q \mapsto r^{8} \mathcal{B}_{g_{0}}(\frac{q(x)}{r^{6}}h),$$

where $q(x) = q_{ij}x^i x^j$ denotes the harmonic quadratic polynomial associated with $q \in \text{Sym}_0^2 \mathbb{R}^4$. The kernel of **H** is naturally and equivariantly isomorphic to the space of all tensors h_0 satisfying (2.17). By Section 2.2.1, ker **H** contains a 16-dimensional invariant subspace $\mathbb{R} \oplus (\mathbf{S}^2_+ \oplus \mathbf{S}^2_-) \oplus (\mathbf{S}^2_+ \otimes \mathbf{S}^2_-)$ consisting of tensors of the form $\mathcal{L}_{X_1}g_0 + \mathcal{L}_{X_2}g_0 + \mathcal{L}_{X_3}g_0$ as above. By Lemma 2.1 in Section 2.2.2, ker **H** also contains a 10dimensional invariant subspace $\mathbf{S}^4_+ \oplus \mathbf{S}^4_-$ consisting of reduced Kronheimer terms for the two orientations. Thus, to prove item (1) it suffices to compute dim ker $\mathbf{H} = 26$, which can be done on a computer. (See the arXiv version of this paper, arXiv:1901.03647, for a Maxima script.)

To prove item (2), first observe that (2.20) follows from the fact that $\mathcal{B}_{g_0}h_0 = 0$, (2.7), and (2.16). To prove (2.21), notice that the map $h_0 \mapsto \partial_r \,\lrcorner\, h_0$ is equivariant, so by Schur's lemma and (2.16) it suffices to show that it has trivial kernel on tensors of the form $\mathcal{L}_{X_1}g_0$, $\mathcal{L}_{X_2}g_0$, and $\mathcal{L}_{X_3}g_0$. For the first two types this is obvious from (2.13) and (2.14), respectively. For the remaining type, we write $X_3(x) = \frac{1}{r^4}Lx$ for all $x \in \mathbb{R}^4 \setminus \{0\}$ with $L = (\ell_{ij}) \in \mathbb{R}^{4 \times 4}$ trace-free symmetric. Then

$$(\partial_r \,\lrcorner\, \mathcal{L}_{X_3}g_0)(\partial_k) = -\frac{2}{r^7}(r^2Lx + 2q(x)x)^k,$$

where $q(x) = \ell_{ij} x^i x^j$ is the quadratic polynomial associated with L. If this vanishes for all k, then every $v \in \mathbb{R}^4 \setminus \{0\}$ is an eigenvector of Lwith eigenvalue $-2q(\frac{v}{|v|})$, so that L = 0. For item (3), note that the projections from ker **H** onto its SO(4)invariant subspaces are SO(4)-equivariant, so if h_0 is *G*-invariant, then so are h_0^+ , h_0^- , and $\mathcal{L}_{X_k}g_0$ (k = 1, 2, 3). It remains to prove that the vector fields X_k , or equivalently the matrices L_k , are *G*-invariant. But this is clear because the map $X \mapsto \mathcal{L}_X g_0$ for X of the form (2.6) is SO(4)-equivariant and injective. q.e.d.

Remark 2.7. It follows from (2.20) that $\mathbf{W} = \mathbf{S}_{+}^{4} \oplus \mathbf{S}_{-}^{4}$ in Lemma 2.1 is as small as it can be.

3. The volume inequality

3.1. Proof of Theorem B. Let (M, g) be a 4-dimensional Ricci-flat ALE space with $\Gamma \neq \{1\}$. Let $k_0 \in \mathbb{N}$. By Section 2.1 there exists a diffeomorphism

$$\Phi: (\mathbb{R}^4 \setminus B_1(0)) / \Gamma \to M \setminus K$$

of class at least C^{k_0+1} such that

$$\Phi^*g - g_0 = h_0 + h',$$

where the leading term h_0 satisfies

$$\mathcal{L}_{r\partial_r}h_0 = -2h_0, \ \mathcal{B}_{g_0}h_0 = 0, \ \Delta_{g_0}h_0 = 0,$$

and where the remainder h' can be estimated by

$$\sum_{k=0}^{k_0} r^k |\nabla_{g_0}^k h'|_{g_0} \leqslant C(k_0) r^{-5}.$$

Then Proposition 2.6 tells us that after lifting through Γ ,

$$h_0 = h_0^+ + h_0^- + \mathcal{L}_{X_1} g_0 + \mathcal{L}_{X_2} g_0 + \mathcal{L}_{X_3} g_0,$$

where h_0^{\pm} are Γ -invariant reduced Kronheimer terms for the two orientations and X_1 , X_2 , X_3 are Γ -invariant harmonic vector fields of the appropriate types. If we compose Φ with the time-1 flows of these vector fields, the metric changes by the corresponding Lie derivatives of g_0 to leading order. By Taylor expansion, since $|\nabla_{g_0}^j X_k|_{g_0} = O(r^{-3-j})$ for k = 1, 2, 3 and all $j \in \mathbb{N}_0$, all other changes to the metric are certainly $O(r^{-5})$ as $r \to \infty$, with as many derivatives as the regularity of h' allows. Thus, we are free to assume without loss that $X_1 = X_2 = X_3 = 0$, and hence that

(3.1)
$$\partial_r \,\lrcorner\, h_0 = 0, \, \operatorname{tr}_{g_0} h_0 = 0, \, \operatorname{div}_{g_0} h_0 = 0,$$

where the latter two properties are equivalent thanks to the Bianchi gauge condition.

We now modify Φ to become a CMC gauge, preserving all of its other properties except for the number of derivatives of h' that we control, which will drop by a bounded amount. Since this does not affect the statement of Theorem B, we will from now on treat k_0 as a generic constant.

Let us recall how the canonical CMC foliation $\{\Sigma_{\rho}\}_{\rho>\rho_0}$ was constructed in [11]. For all $\rho > 0$ we consider the homothety $h_{\rho}(x) = \rho x$ in \mathbb{R}^4/Γ . Then on any compact set $K \subset (\mathbb{R}^4 \setminus \{0\})/\Gamma$,

(3.2)
$$h_{\rho}^{*}\Phi^{*}g - g_{0} = O_{K}(\rho^{-4}) \text{ as } \rho \to \infty$$

including all derivatives up to order k_0 . The unit sphere \mathbb{S}^3/Γ has constant mean curvature 3, and the linearization of the CMC equation on normal deformations of \mathbb{S}^3/Γ is the Jacobi operator $J = \Delta_{\mathbb{S}^3/\Gamma} + 3$. On \mathbb{S}^3 the kernel of $\Delta_{\mathbb{S}^3} + 3$ is given by the restriction to \mathbb{S}^3 of the linear functions on \mathbb{R}^4 and corresponds to translations of \mathbb{S}^3 inside \mathbb{R}^4 . As $\Gamma \neq \{1\}$, this flexibility disappears in \mathbb{R}^4/Γ , so that J is invertible. It follows that for $\rho \gg 1$ we can deform \mathbb{S}^3/Γ into a hypersurface $\tilde{\Sigma}_{\rho}$ of constant mean curvature 3 with respect to $h_{\rho}^* \Phi^* g$. More precisely, we can write $\tilde{\Sigma}_{\rho}$ as a radial graph

$$\tilde{\Sigma}_{\rho} = \{ (1 + f_{\rho}(x)) x \in \mathbb{R}^4 / \Gamma : x \in \mathbb{S}^3 / \Gamma \},\$$

and (3.2) implies that for all $0 \leq k \leq k_0$ we have

(3.3)
$$\nabla^k f_{\rho} = O(\rho^{-4}) \text{ as } \rho \to \infty$$

Pushing Σ_{ρ} forward by $\Phi \circ h_{\rho}$ now gives the required CMC hypersurfaces Σ_{ρ} . The same argument applied to the obvious family of CMC hypersurfaces around $\tilde{\Sigma}_{\rho}$ with mean curvature between $\frac{3}{2}$ and 6 tells us that the family $\{\Sigma_{\rho}\}_{\rho>\rho_0}$ is actually a foliation.

The key point is that in our setting we are able to improve (3.3) by one order by using (3.1). To see this, we will compute the mean curvature Hof the unit sphere \mathbb{S}^3/Γ with respect to $h_{\rho}^*\Phi^*g$. It is clear from (3.2) that $H = 3 + O(\rho^{-4})$ up to k_0 derivatives, which leads to (3.3). However, we can improve this to $H = 3 + O(\rho^{-5})$, with a corresponding improvement in (3.3), by viewing \mathbb{S}^3/Γ as a level set of the function $u = r^2$ and using (3.1) to exhibit cancellations in the formula

(3.4)
$$H = \frac{1}{|\nabla u|} (\Delta u - \frac{1}{|\nabla u|^2} (\operatorname{Hess} u) (\nabla u, \nabla u)).$$

The details are as follows. First note that the background metric in (3.4) is

$$h_{\rho}^{*}\Phi^{*}g = g_{0} + h_{0}\rho^{-4} + O(\rho^{-5})$$

The Christoffel symbols of this metric are given by

(3.5)
$$\Gamma_{ij}^{k} = \frac{1}{2} (\partial_{i} h_{0,jk} + \partial_{j} h_{0,ik} - \partial_{k} h_{0,ij}) \rho^{-4} + O(\rho^{-5})$$

in the standard Euclidean coordinates of \mathbb{R}^4 in a tubular neighborhood of $\mathbb{S}^3.$ Then

$$\nabla_i u = 2x^i - 2h_{0,ij}x^j \rho^{-4} + O(\rho^{-5}) = 2x^i + O(\rho^{-5}),$$

Hess_{ij} $u = 2\delta_{ij} - (\partial_i h_{0,jk} + \partial_j h_{0,ik} - \partial_k h_{0,ij})x^k \rho^{-4} + O(\rho^{-5}),$

where in the first line we have used the property $\partial_r \,\lrcorner\, h_0 = 0$ from (3.1). Thus,

$$\begin{aligned} |\nabla u|^2 &= 4r^2 + 4h_{0,ij}x^i x^j \rho^{-4} + O(\rho^{-5}) = 4r^2 + O(\rho^{-5}), \\ \Delta u &= 8 - 2(\mathrm{tr}_{g_0}h_0 + (\mathcal{B}_{g_0}h_0)_k x^k)\rho^{-4} + O(\rho^{-5}) = 8 + O(\rho^{-5}), \\ (\mathrm{Hess}\, u)(\nabla u, \nabla u) &= 8r^2 - 4x^i x^j x^k (\partial_i h_{0,jk} + \partial_j h_{0,ik} - \partial_k h_{0,ij})\rho^{-4} \\ &+ O(\rho^{-5}) = 8r^2 + O(\rho^{-5}), \end{aligned}$$

thanks to all three of the properties of (3.1) and by applying Euler's homogeneity relation to h_0 in the last line. Evaluating (3.4) at u = 1, it is then clear that $H = 3 + O(\rho^{-5})$, as desired.

The upshot of all of this is that if we replace Φ by the diffeomorphism

$$\rho x \mapsto \Phi((1+f_{\rho}(x))\rho x) \ (\rho \gg 1, x \in \mathbb{S}^3/\Gamma).$$

then the leading term h_0 of the metric remains unchanged, and the remainder h' satisfies the same estimates as before, but it now holds by construction that $\Phi(\partial B_{\rho}(0)/\Gamma) = \Sigma_{\rho}$.

3.2. Proof of Theorem A. We now use the ALE diffeomorphism Φ provided by Theorem B. One consequence of the property $\operatorname{tr}_{g_0} h_0 = 0$ is that

(3.6)
$$\Phi^*(d\mathrm{Vol}_g) = d\mathrm{Vol}_{\Phi^*g} = d\mathrm{Vol}_{g_0}(1 + O(r^{-5})).$$

This immediately implies that the function

$$g(\rho) := \operatorname{Vol}_g(\Omega_\rho) - \operatorname{Vol}_{g_0}(B_\rho(0)/\Gamma)$$

satisfies for all $\rho_2 > \rho_1 \gg 1$ that

$$g(\rho_2) - g(\rho_1) = \int_{\rho_1}^{\rho_2} \Phi^*(d\operatorname{Vol}_g) - d\operatorname{Vol}_{g_0} = O(\rho_1^{-1}).$$

Thus, $g(\rho)$ has a finite limit as $\rho \to \infty$, which proves Theorem A(1), i.e., the existence of \mathcal{V} .

For Theorem A(2), we first prove that there exists a function u on M such that $\Delta_q u = 8$ and

(3.7)
$$\Phi^* u = r^2 + br^{-2} + O_{\varepsilon}(r^{-3+\varepsilon})$$

for some $b \in \mathbb{R}$ and all $\varepsilon \in (0, 1)$. For this we require the precise expansion

$$\Delta_{\Phi^*g}r^2 = 8 - 2(\operatorname{tr}_{g_0}h_0 + (\mathcal{B}_{g_0}h_0)(r\partial_r)) + O(r^{-5}) = 8 + O(r^{-5}),$$

which follows from the work in Section 3.1 by pushing forward by the homotheties h_{ρ} . Following a standard pattern, we now extend $\Phi_* r^2$ to

a smooth function u_0 on M and let $f = 8 - \Delta_g u_0$. Note that $\Phi^* f = O(r^{-5})$. It suffices to find a function \bar{u} on M such that $\Delta_g \bar{u} = f$ and

$$\Phi^* \bar{u} = br^{-2} + O_{\varepsilon}(r^{-3+\varepsilon})$$

because then $u = u_0 + \bar{u}$ solves the original problem. The existence of \bar{u} is a standard fact and can be proved in many ways. For example, by solving Dirichlet problems on larger and larger balls and using Moser iteration and weighted Sobolev inequalities, one first constructs a solution \bar{u} such that $\Phi^*\bar{u} = O_{\varepsilon}(r^{-2+\varepsilon})$. Then it is clear that $\Delta_{g_0}(\Phi^*\bar{u}) = O(r^{-5})$, so standard properties of the Green's function on \mathbb{R}^4 imply that $\Phi^*\bar{u} = br^{-2} + O_{\varepsilon}(r^{-3+\varepsilon})$, as desired.

Let ν be the exterior unit normal to the domain Ω_{ρ} with respect to g. Integrating the equation $\Delta_g u = 8$ over Ω_{ρ} , integrating by parts, and using (3.6) we get

$$8\operatorname{Vol}_{g}(\Omega_{\rho}) = \int_{\partial\Omega_{\rho}} du(\nu)(\nu \,\lrcorner\, d\operatorname{Vol}_{g})$$
$$= \int_{\partial\Omega_{\rho}} du(\nu)(\nu \,\lrcorner\, \Phi_{*}d\operatorname{Vol}_{g_{0}}) + O(\rho^{-1}).$$

From now on we will ignore the map Φ for convenience. Then, using (3.7) in the first line,

$$du(\nu)|_{\partial\Omega_{\rho}} = 2\rho - 2b\rho^{-3} + 2\rho(dr(\nu) - 1) + O_{\varepsilon}(\rho^{-4+\varepsilon}),$$
$$\nu \,\lrcorner\, d\mathrm{Vol}_{g_0} = dr(\nu)(\partial_r \,\lrcorner\, d\mathrm{Vol}_{g_0}).$$

A priori the error $dr(\nu) - 1$ is $O(\rho^{-4})$ but expanding the equation $g(\nu, \nu) = 1$ yields

$$dr(\nu) - 1 = -\frac{1}{2}h_0(\partial_r, \partial_r) + O(\rho^{-5}) = O(\rho^{-5})$$

thanks to (3.1). Combining these computations, we get

$$8\operatorname{Vol}_g(\Omega_\rho) = 8\operatorname{Vol}_{g_0}(B_\rho(0)/\Gamma) - 2b|\mathbb{S}^3/\Gamma| + O_\varepsilon(\rho^{-1+\varepsilon}),$$

which obviously implies that $b|\mathbb{S}^3/\Gamma| = -4\mathcal{V}$.

We now reinterpret b as the obstruction to ∇u being a conformal Killing field, and the proof will show that $b \ge 0$. Recall that ∇u being conformally Killing means that the trace-free Hessian Hess₀u vanishes, and this is equivalent to ∇u generating a 1-parameter group of conformal diffeomorphisms of (M, g). In this case, the conformal factor is constant because $\Delta u = 8$, so ∇u actually generates a 1-parameter group of homotheties, and by considering the sup norm of the curvature tensor one easily checks that this is equivalent to (M, g) being isometric to $(\mathbb{R}^4/\Gamma, g_0)$ with $u = r^2$.

Using the general identity

$$\Delta \nabla u - \nabla \Delta u = \operatorname{Ric} \nabla u,$$

together with the fact that u has constant Laplacian, we obtain that

(3.8)
$$\operatorname{div}(\operatorname{Hess}_0 u) = \operatorname{Ric} du = 0.$$

Integrating this against du over Ω_{ρ} , we get

$$\int_{\partial\Omega_{\rho}} (\operatorname{Hess}_{0} u)(\nu, \nabla u)(\nu \,\lrcorner\, d\operatorname{Vol}) = \int_{\Omega_{\rho}} |\operatorname{Hess}_{0} u|^{2} + \operatorname{Ric}(\nabla u, \nabla u) \, d\operatorname{Vol} \ge 0.$$

The desired inequality $b \ge 0$, with equality if and only if $\operatorname{Hess}_0 u = 0$, will come out of an expansion of the boundary term. We begin by expanding $\operatorname{Hess}_0 u$. This will turn out to be $O(\rho^{-4})$ to leading order, so there is no need to also expand ν , ∇u , $\nu \lrcorner d$ Vol, and we can simply replace these by their Euclidean approximations. For $\operatorname{Hess}_0 u$, using (3.5) and (3.7) we have

$$\begin{aligned} \operatorname{Hess}_{0,ij} u &= \partial_i \partial_j u - \Gamma_{ij}^k \partial_k u - 2g_{ij} \\ &= -\frac{2b}{r^4} (\delta_{ij} - 4\frac{x^i x^j}{r^2}) - (\partial_i h_{0,jk} + \partial_j h_{0,ik} - \partial_k h_{0,ij}) x^k - 2h_{0,ij} \\ &+ O_{\varepsilon} (r^{-5+\varepsilon}). \end{aligned}$$

This then tells us that, as 3-forms on $\partial \Omega_{\rho}$,

$$(\operatorname{Hess}_{0} u)(\nu, \nabla u)(\nu \lrcorner d\operatorname{Vol})$$

$$= (12b\rho^{-3} - 2x^{i}x^{j}x^{k}(\partial_{i}h_{0,jk} + \partial_{j}h_{0,ik} - \partial_{k}h_{0,ij})\rho^{-1} - 4h_{0}(\partial_{r}, \partial_{r})\rho$$

$$+ O_{\varepsilon}(\rho^{-4+\varepsilon}))(\partial_{r} \lrcorner d\operatorname{Vol}_{g_{0}})$$

$$= (12b\rho^{-3} + O_{\varepsilon}(\rho^{-4+\varepsilon}))(\partial_{r} \lrcorner d\operatorname{Vol}_{g_{0}}),$$

after applying Euler's relation to h_0 and using (3.1) as in Section 3.1.

Combining this with the preceding discussion, we obtain that $b \ge 0$, or equivalently that $\mathcal{V} \le 0$, with equality if and only if (M, g) is isometric to $(\mathbb{R}^4/\Gamma, g_0)$ with $u = r^2$.

Remark 3.1. The last part of the above proof can be interpreted as a reverse Bishop-Gromov type inequality. In fact, similar computations show that if Ω is a bounded domain in an *n*-manifold with Ric ≥ 0 , with boundary mean curvature $\geq \frac{n-1}{\rho}$ for some $\rho > 0$, then $|\Omega| \leq \frac{\rho}{n} |\partial \Omega|$, with equality if and only if Ω is isometric to a ball of radius ρ in \mathbb{R}^n . After completing our work on this paper, we discovered that this result already follows from Ros [**23**, Thm 1], who used a very similar method. Our computation replaces the use of Reilly's formula in [**23**] by (3.8) and (3.4).

3.3. Proof of Theorem C. We now assume that (M, g) is a Kronheimer gravitational instanton with period point $\zeta \in \mathfrak{h} \otimes \mathbb{R}^3$. As mentioned in Section 2.2.2, Kronheimer constructed a particular ALE diffeomorphism Φ : $(\mathbb{R}^4 \setminus B_1(0))/\Gamma \to M \setminus K$ with respect to which

 $-(2\pi^2/|\Gamma|)r^6h_0$ takes the form (2.9). Details can be found in [3, Section 2], including the remarkable property that

(3.9)
$$\operatorname{Vol}_g(U_{\tau}) = \operatorname{Vol}_{g_0}(B_{\tau}(0)/\Gamma) \text{ for all } \tau \gg 1,$$

where $U_{\tau} \subset M$ denotes the domain interior to $S_{\tau} := \Phi(\partial B_{\tau}(0)/\Gamma)$. (See [3, Lemma 2.5] for this. Note that the map F_{ζ} of [3] is a global diffeomorphism from $M \setminus Z$ to $(\mathbb{R}^4 \setminus \{0\})/\Gamma$, where $Z \subset M$ is a finite union of compact 2-dimensional surfaces.) By Lemma 2.1 and its proof, we have

$$h_0 = h_0^+ + \mathcal{L}_{X_1} g_0 + \mathcal{L}_{X_2} g_0$$

where, first of all, h_0^+ is a reduced Kronheimer term, so that

(3.10)
$$\partial_r \lrcorner h_0^+ = 0, \ \operatorname{tr}_{g_0} h_0^+ = 0, \ \operatorname{div}_{g_0} h_0^+ = 0$$

second, the harmonic vector field X_1 is a scalar multiple of $\nabla_{g_0}(\frac{1}{r^2})$; and, third, the harmonic vector field X_2 is a linear combination of $I_j(\nabla_{g_0}(\frac{1}{r^2}))$ for j = 1, 2, 3. In fact, Remark 2.3 tells us that

(3.11)
$$X_1 = -\frac{|\tilde{\zeta}|^2}{12} \nabla_{g_0}(\frac{1}{r^2}), \text{ where } \tilde{\zeta} := \left(\frac{|\Gamma|}{2\pi^2}\right)^{\frac{1}{2}} \zeta.$$

We now replace Kronheimer's gauge Φ by $\Phi' = \Phi \circ \Phi_2 \circ \Phi_1$, where Φ_k denotes the time-1 flow of $-X_k$ for k = 1, 2. In this new gauge, the leading term of the metric is h_0^+ . By Section 3.1, because of (3.10), the canonical CMC hypersurface Σ_{ρ} is a normal graph of height $O(\rho^{-4})$ (rather than the expected $O(\rho^{-3})$) over the coordinate sphere $S'_{\rho} := \Phi'(\partial B_{\rho}(0)/\Gamma)$. Let $U'_{\rho} \subset M$ denote the domain interior to S'_{ρ} . Then it is easy to see that

(3.12)

$$\operatorname{Vol}_{g}(\Omega_{\rho}) = \operatorname{Vol}_{g}(U_{\rho}') + O(\rho^{-1})$$

$$= \operatorname{Vol}_{g}(U_{\tau}) + O(\rho^{-1}) \text{ for } \tau^{4} = \rho^{4} - \frac{2}{3}|\tilde{\zeta}|^{2}.$$

Indeed, $S'_{\rho} = \Phi(\Phi_2(\Phi_1(\partial B_{\rho}(0)/\Gamma)))$ by definition, $\Phi_1(\partial B_{\rho}(0)/\Gamma) = \partial B_{\tau}(0)/\Gamma$ with τ in terms of ρ as in (3.12) by solving a simple ODE (using (3.11)), and Φ_2 obviously preserves each sphere in \mathbb{R}^4/Γ . The desired formula for $\mathcal{V}(M, g)$ now follows from (3.12) and (3.9).

4. Killing fields on 4-dimensional Ricci-flat ALE spaces

The idea of the proof of Theorem A is to search for a conformal Killing vector field asymptotic to $2r\partial_r$. This is done by first producing the harmonic vector field ∇u , where u is asymptotic to r^2 with $\Delta u = 8$, and then integrating by parts in the Bochner type formula div(Hess₀u) = Ric du = 0. Using a similar approach, one can give a criterion for the existence of a Killing field asymptotic to a given $\mathfrak{so}(4)$ symmetry. The necessary computations are very long, and there does not seem to be an application in the spirit of Theorem A, so we will only briefly sketch this result.

Let (M, g) be a 4-dimensional Ricci-flat ALE space asymptotic to \mathbb{R}^4/Γ . Fix an arbitrary ALE diffeomorphism Φ as in (1.1). Then for all $X_0 \in \mathfrak{so}(4)^{\Gamma}$ there exists a unique harmonic vector field X on M such that $(\Phi^{-1})_*X = X_0 + Y_0 + O_{\varepsilon}(r^{-4+\varepsilon})$ for all $\varepsilon \in (0, 1)$, where

$$\mathcal{L}_{r\partial_r} Y_0 = -4Y_0, \ |Y_0|_{q_0} \sim r^{-3}.$$

An asymptotic expansion shows that

$$\Delta_{q_0} Y_0 = 0.$$

The next observation is that the divergence of a harmonic vector field on a Ricci-flat manifold is a harmonic function, and it is clear that div Xgoes to zero at infinity, so div X = 0 by the maximum principle. If the leading term h_0 of the metric with respect to Φ satisfies $\operatorname{tr}_{g_0} h_0 = 0$, then one can deduce from this by asymptotic expansion that

$$\operatorname{div}_{q_0} Y_0 = 0$$

Since Y_0 has to be of the form (2.6), it follows that for some $a \in \mathbb{R}$ and $Z_0 \in \mathfrak{so}(4)$,

$$Y_0 = \frac{1}{r^4} (ar\partial_r + Z_0).$$

If u is the unique solution to $\Delta u = 8$ with $u = r^2 + o(1)$ on M, and if the precise expansion (3.7) holds for u (which we know is true if $\operatorname{tr}_{g_0}h_0 = 0$ and $\mathcal{B}_{g_0}h_0 = 0$), then an integration by parts in the identity $0 = \langle \Delta X, \nabla u \rangle - \langle X, \Delta \nabla u \rangle$ shows after a long computation that a = 0.

If X_0, X'_0 are in $\mathfrak{so}(4)^{\Gamma}$ with harmonic extensions X, X', and if $\partial_r \,\lrcorner\, h_0 = 0$, then another lengthy integration by parts in the identity $0 = \langle \Delta X, X' \rangle - \langle X, \Delta X' \rangle$ shows that $\langle X_0, Z'_0 \rangle = \langle Z_0, X'_0 \rangle$ with respect to the Killing form on $\mathfrak{so}(4)$. Thus, the endomorphism $X_0 \mapsto Z_0$ of $\mathfrak{so}(4)^{\Gamma}$ is selfadjoint.

Integration by parts in the Bochner type formula (2.5) shows that the selfadjoint endomorphism $X_0 \mapsto Z_0$ of $\mathfrak{so}(4)^{\Gamma}$ is nonpositive, and that $Z_0 = 0$ if and only if X is a Killing field. In this case, an asymptotic expansion of the equation $\mathcal{L}_X g = 0$ immediately tells us that $\mathcal{L}_{X_0} h_0 = 0$.

References

- M.T. Anderson, L² curvature and volume renormalization of AHE metrics on 4-manifolds, Math. Res. Lett. 8 (2001), 171–188, MR1825268, Zbl 0999.53034.
- [2] R. Arnowitt, S. Deser, and C.W. Misner, Coordinate invariance and energy expressions in general relativity, Phys. Rev. (2) 122 (1961), 997–1006, MR0127946, Zbl 0094.23003.
- [3] H. Auvray, From ALE to ALF gravitational instantons, Compos. Math. 154 (2018), 1159–1221, MR3826456, Zbl 1401.53066.

- 18 The renormalized volume of a 4-dimensional Ricci-flat ALE space
- [4] S. Bando, A. Kasue, and H. Nakajima, On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth, Invent. Math. 97 (1989), 313–349, MR1001844, Zbl 0682.53045.
- [5] S. Brendle and O. Chodosh, A volume comparison theorem for asymptotically hyperbolic manifolds, Comm. Math. Phys. **332** (2014), 839–846, MR3257665, Zbl 1306.53021.
- [6] E. Calabi, Métriques kählériennes et fibrés holomorphes, Ann. Sci. École Norm. Sup. 12 (1979), 269–294, MR0543218, Zbl 0431.53056.
- [7] J. Cheeger and A. Naber, Regularity of Einstein manifolds and the codimension 4 conjecture, Ann. of Math. 182 (2015), 1093–1165, MR3418535, Zbl 1335.53057.
- [8] J. Cheeger and G. Tian, On the cone structure at infinity of Ricci-flat manifolds with Euclidean volume growth and quadratic curvature decay, Invent. Math. 118 (1994), 493–571, MR1296356, Zbl 0814.53034.
- [9] Y. Chen, On expansions of Ricci flat ALE metrics in harmonic coordinates about the infinity, Comm. Math. Stat. 8 (2020), 63–90, MR4071856, Zbl 1436.32085.
- [10] O. Chodosh, M. Eichmair, Y. Shi, and H. Yu, Isoperimetry, scalar curvature, and mass in asymptotically flat Riemannian 3-manifolds, Comm. Pure Appl. Math. 74 (2021), 865–905, MR4221936, Zbl not yet listed.
- [11] O. Chodosh, M. Eichmair, and A. Volkmann, Isoperimetric structure of asymptotically conical manifolds, J. Differ. Geom. 105 (2017), 1–19, MR3592692, Zbl 1364.53035.
- [12] T. Eguchi and A.J. Hanson, Self-dual solutions to Euclidean gravity, Ann. Physics 120 (1979), 82–106, MR0540896, Zbl 0409.53020.
- [13] X.-Q. Fan, Y. Shi, and L.-F. Tam, Large-sphere and small-sphere limits of the Brown-York mass, Comm. Anal. Geom. 17 (2009), 37–72, MR2495833, Zbl 1175.53083.
- [14] C.R. Graham, Volume and area renormalizations for conformally compact Einstein metrics, The Proceedings of the 19th Winter School "Geometry and Physics" (Srní, 1999), Rend. Circ. Mat. Palermo (2) Suppl. 63, Circ. Mat. Palermo, 2000, pp. 31–42, MR1758076, Zbl 0984.53020.
- [15] X. Hu, D. Ji, and Y. Shi, Volume comparison of conformally compact manifolds with scalar curvature $R \ge -n(n-1)$, Ann. Henri Poincaré **17** (2016), 953–977, MR3472629, Zbl 1338.53071.
- [16] G. Huisken, An isoperimetric concept for mass and quasilocal mass, Report 2/2006–Mathematical aspects of general relativity, Oberwolfach Rep. 3, European Mathematical Society, Zürich, 2006, pp. 87–88, MR2278888, Zbl 1109.83300.
- [17] J.L. Jauregui and D.A. Lee, Lower semicontinuity of mass under C⁰ convergence and Huisken's isoperimetric mass, J. Reine Angew. Math. **756** (2019), 227–257, MR4026453, Zbl 1430.53052.
- [18] P.B. Kronheimer, The construction of ALE spaces as hyper-Kähler quotients, J. Differ. Geom. 29 (1989), 665–683, MR0992334, Zbl 0671.53045.
- [19] _____, A Torelli-type theorem for gravitational instantons, J. Differ. Geom. 29 (1989), 685–697, MR0992335, Zbl 0671.53046.
- [20] M.T. Lock and J.A. Viaclovsky, Quotient singularities, eta invariants, and selfdual metrics, Geom. Topol. 20 (2016), 1773–1806, MR3523069, Zbl 1348.53054.
- [21] H. Nakajima, Self-duality of ALE Ricci-flat 4-manifolds and positive mass theorem, Recent topics in differential and analytic geometry, Adv. Stud. Pure Math.,

vol. 18, Academic Press, Boston, MA, 1990, pp. 385–396, MR1145266, Zbl 0744.53025.

- [22] T. Ozuch, Noncollapsed degeneration of Einstein 4-manifolds II, preprint (2019), arXiv:1909.12960.
- [23] A. Ros, Compact hypersurfaces with constant higher order mean curvatures, Rev. Mat. Iberoamericana 3 (1987), 447–453, MR0996826, Zbl 0673.53003.
- [24] R. Schoen and S.-T. Yau, On the proof of the positive mass conjecture in general relativity, Comm. Math. Phys. 65 (1979), 45–76, MR0526976, Zbl 0405.53045.
- [25] I. Şuvaina, ALE Ricci-flat Kähler metrics and deformations of quotient surface singularities, Ann. Global Anal. Geom. 41 (2012), 109–123, MR2860399, Zbl 1236.53057.
- [26] E. Witten, A new proof of the positive energy theorem, Comm. Math. Phys. 80 (1981), 381–402, MR0626707, Zbl 1051.83532.
- [27] E.P. Wright, Quotients of gravitational instantons, Ann. Global Anal. Geom. 41 (2012), 91–108, MR2860398, Zbl 1236.53060.

Sorbonne Université & Université de Paris, CNRS, IMJ-PRG, 75005 Paris, France

E-mail address: olivier.biquard@sorbonne-universite.fr

Mathematisches Institut, WWU Münster, 48149 Münster, Germany And

DEPARTMENT OF MATHEMATICS, FORDHAM UNIVERSITY, BRONX, NY 10458, USA *E-mail address*: hhein@uni-muenster.de