



# The Hamburg/ESO R-process enhanced star survey (HERES). III. HE 0338-3945 and the formation of the r + s stars

K. Jonsell, P. S. Barklem, B. Gustafsson, Norbert Christlieb, Vanessa Hill,  
Timothy C. Beers, J. Holmberg

## ► To cite this version:

K. Jonsell, P. S. Barklem, B. Gustafsson, Norbert Christlieb, Vanessa Hill, et al.. The Hamburg/ESO R-process enhanced star survey (HERES). III. HE 0338-3945 and the formation of the r + s stars. *Astronomy & Astrophysics - A&A*, 2006, 451, pp.651-670. 10.1051/0004-6361:20054470 . hal-03742443

HAL Id: hal-03742443

<https://hal.science/hal-03742443v1>

Submitted on 21 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The Hamburg/ESO R-process enhanced star survey (HERES)<sup>★,★★</sup>

## III. HE 0338–3945 and the formation of the *r + s* stars

K. Jonsell<sup>1</sup>, P. S. Barklem<sup>1</sup>, B. Gustafsson<sup>1</sup>, N. Christlieb<sup>2</sup>, V. Hill<sup>3</sup>, T. C. Beers<sup>4</sup>, and J. Holmberg<sup>5</sup>

<sup>1</sup> Department of Astronomy and Space Physics, Uppsala Astronomical Observatory, Box 515, 751 20 Uppsala, Sweden  
e-mail: [firstname.lastname@astro.uu.se](mailto:firstname.lastname@astro.uu.se)

<sup>2</sup> Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany

<sup>3</sup> GEPI, Observatoire de Paris-Meudon, 92125 Meudon Cedex, France

<sup>4</sup> Department of Physics and Astronomy and JINA: Joint Institute for Nuclear Astrophysics, Michigan State University, East Lansing, MI 48824, USA

<sup>5</sup> Tuorla Observatory, Väisäläntie 20, 21500 Piikkiö, Finland

Received 4 November 2005 / Accepted 18 January 2006

### ABSTRACT

We have derived abundances of 33 elements and upper limits for 6 additional elements for the metal-poor ( $[Fe/H] = -2.42$ ) turn-off star HE 0338–3945 from high-quality VLT-UVES spectra. The star is heavily enriched, by about a factor of 100 relative to iron and the Sun, in the heavy *s*-elements (Ba, La, ...). It is also heavily enriched in Eu, which is generally considered an *r*-element, and in other similar elements. It is less enriched, by about a factor of 10, in the lighter *s*-elements (Sr, Y and Zr). C is also strongly enhanced and, to a somewhat lesser degree, N and O. These abundance estimates are subject to severe uncertainties due to NLTE and thermal inhomogeneities which are not taken into detailed consideration. However, an interesting result, which is most probably robust in spite of these uncertainties, emerges: the abundances derived for this star are very similar to those of other stars with an overall enhancement of all elements beyond the iron peak.

We have defined criteria for this class of stars, *r + s* stars, and discuss nine different scenarios to explain their origin. None of these explanations is found to be entirely convincing. The most plausible hypotheses involve a binary system in which the primary component goes through its giant branch and asymptotic giant branch phases and produces CNO and *s*-elements which are dumped onto the observed star. Whether the *r*-element Eu is produced by supernovae before the star was formed (perhaps triggering the formation of a low-mass binary), by a companion as it explodes as a supernova (possibly triggered by mass transfer), or whether it is possibly produced in a high-neutron-density version of the *s*-process is still unclear. Several suggestions are made on how to clarify this situation.

**Key words.** stars: population II – stars: fundamental parameters – stars: abundances – Galaxy: halo – Galaxy: abundances – Galaxy: evolution

### 1. Introduction

Elements with atomic numbers  $Z > 30$  are believed to be almost exclusively synthesized in neutron-capture (*n*-capture) reactions. In the most metal-poor stars the overall abundance of these elements varies from star to star, by more than a factor of 100 at a given metallicity (McWilliam et al. 1995; Ryan et al. 1996). Also, the different abundance ratios vary, e.g. the Ba/Eu ratio tends to decline with decreasing  $[Fe/H]$ <sup>1</sup> (McWilliam 1998; Burris et al. 2000). Eu is often thought to be synthesized in conditions of very high neutron fluxes (the

*r*-process) while Ba is most easily made in conditions of much lower neutron fluxes (the *s*-process). The decline of Ba/Eu may show the relative dominance of *r*-process sites in the early history of the Galaxy.

The first Extremely Metal Poor (EMP, Beers & Christlieb 2005) star found to be enriched in *n*-capture elements was the giant HD 115444 with  $[Fe/H] \sim -3$  (Griffin et al. 1982). Westin et al. (2000) found marginal overabundances of Ba and other *s*-elements relative to iron in this star as compared with the Sun. This still makes the star rich in these elements relative to “normal” Population II stars of the same metallicity. However, it has a high abundance of europium,  $[Eu/Fe] = 0.85$ , even compared to the Sun.

The heavy elements of HD 115444 show good agreement in relative abundances with scaled solar *r*-process abundances, as does the EMP giant CS 22892-052 which is even more

<sup>★</sup> Based on observations carried out at the European Southern Observatory, Paranal, Chile. Proposal number (170.D-0010).

<sup>★★</sup> Table 2 and Appendix A are only available in electronic form at <http://www.edpsciences.org>

<sup>1</sup> We use the standard notations  $A/B = N_A/N_B$  and  $[A/B] = \log(N_A/N_B)_* - \log(N_A/N_B)_\odot$  where  $N_X$  are number densities.

*r*-element rich (McWilliam et al. 1995; Sneden et al. 2003a). This agreement suggests that a similar process is responsible for generating the heavy *r*-elements, both for the most metal-poor stars and for the Sun. However, several studies have found a deviation from the scaled solar *r*-process abundance pattern for the light *r*-process elements (e.g. Sneden et al. 2000, 2003a; Cowan et al. 2002; Aoki et al. 2005). Also, similar stars like CS 31082-001 (Hill et al. 2002) and CS 30306-132 (Honda et al. 2004a) seem to have had an “actinide boost”, which has selectively affected the abundances of the most heavy *r*-elements. These variations in behaviour of the *r*-elements suggest that multiple astrophysical sites for the *r*-process may exist.

The discovery of CS 22892-052 was a result of the compilation of metal-poor HK survey objects of Beers et al. (1992). Similarly, the large programme of Cayrel et al. (2004) at ESO, “First Stars”, was founded on that survey. In the First Stars project, high-resolution studies of the very metal-poor giant CS 31082-001 revealed its high Eu abundance. It was possible to identify a U II line in this star, and this opened up the possibility to use the U/Th ratio as a new chronometer. More recently, Honda et al. (2004a,b) reported on two new Eu-rich stars, again taken from the HK survey, namely CS 30306-132 with [Eu/Fe] = +0.85 and CS 22183-031 with [Eu/Fe] = +1.2. According to the classification suggested by Beers & Christlieb (2005) the *latter* of these is an *r*-II star (having [Eu/Fe] > +1.0 and [Ba/Eu] < 0.0) while the *former* is classified as *r*-I (+0.3 ≤ [Eu/Fe] ≤ +1.0 and [Ba/Eu] < 0.0). Another interesting object which is very overabundant in *n*-capture elements, the carbon-enhanced metal-poor (CEMP) star with *s*-element enhancement CS 31062-050, was recently studied by Johnson & Bolte (2004). Interestingly, they concluded that the abundance profile of the star, in spite of its high Eu abundance, suggests that the *r*-process was not responsible in this case. These indications will be further discussed in Sect. 6.

Although the contribution of the *s*-process to the gas in the very early Galaxy may in general be tiny, as one would expect due to the small fraction of iron to start the nucleosynthesis from, some stars with [Fe/H] as low as about -3 have been found to have clear *s*-element signatures (Johnson & Bolte 2002; Simmerer et al. 2004; Sivarani et al. 2004). The implications of the presence of these stars are still not fully understood. It was predicted by Gallino et al. (1998; see also Goriely & Mowlavi 2000), that the fraction of the heaviest *s*-elements should be enhanced in metal-poor environments with respect to the lighter *s*-elements, a result of the scarcity of seed nuclei with respect to neutrons. This prediction was qualitatively confirmed by Aoki et al. (2000) who made the first discovery of Pb in a *s*-element rich very metal-poor star, LP625-44. Somewhat more iron-rich “lead stars” were also discovered by Van Eck et al. (2001; see also Sivarani et al. 2004). We also note that the abundances of the “lighter” *n*-capture elements (Sr, Y, Zr) do not scale very well with those of the heavier elements (Ba etc., Z ≥ 56). This may be due to the fact that the lighter *s*-elements (the so-called “weak component”, see Prantzos et al. 1990) received important contributions from seed nuclei reacting with neutrons created in the  $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$  reaction in massive AGB stars, while the heavy elements were essentially

contributed by neutrons from the  $^{13}\text{C}(\alpha, n)^{16}\text{O}$  reaction in less massive stars.

Another class of stars was discovered by Barbuy et al. (1997) and Hill et al. (2000). Two carbon-rich Population II giants, the CEMP stars CS 22948-027 and CS 29497-034 from the HK survey, were found not only to be rich in the commonly enhanced *s*-elements such as Sr, Y, Ba and La, but also the *r*-element Eu was significantly enhanced. Preston & Sneden (2001) found another star of this class in the HK survey (CS 22898-027), Aoki et al. (2002) added two more (CS 29526-110 and CS 31062-012) and Sivarani et al. (2004) another (CS 29497-030) (see also Ivans et al. 2005). The stars of this type presently known are listed in Table 8. We shall denote these stars *r* + *s* stars, and we postpone a detailed discussion of the definition of this class of stars to Sect. 6.

The origin of the abundance peculiarities of the *r* + *s* stars is not clear, and many scenarios have been proposed (see Sect. 6). However, this is not the only reason for studying them further. In fact, the astrophysical sites of the *r*-process are still unclear. The precise sites of the *s*-process among the most metal-poor stars are also still debatable, as well as the fraction of nuclei provided by neutrons from the  $^{13}\text{C}(\alpha, n)^{16}\text{O}$  reaction as compared with the  $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$  neutron source. One might hope that a clarification of the origin of the *r* + *s* stars may shed some light on the general questions concerning the sites of the *r*- and *s*-processes.

The Hamburg/ESO objective-prism survey for bright quasars (HES; Wisotzki et al. 2000) has been used for a search for metal-poor stars (Christlieb 2003). One of these was the *r* + *s* star HE 2148-1247, found in the Keck pilot programme on EMP stars from the HES (Cohen et al. 2003). In a systematic approach to exploring the *n*-enhanced HES stars, the Hamburg/ESO R-process Enhanced Star survey (HERES) has been carried out (Christlieb et al. 2004, hereafter Paper I). For several hundred candidate stars from HES, “snapshot” spectra at moderate resolving power ( $R \sim 20\,000$ ) and low *S/N* ( $\sim 50$ ) were obtained with the main goal to find *r*-element enriched stars, as disclosed by strong Eu lines. These spectra were analysed with an automatic procedure based on MARCS model atmospheres and synthetic spectra (Barklem et al. 2005, Paper II). As a result of the HERES survey, a number of new detections of *n*-capture enriched EMP stars were made. In particular, 8 new *r*-II stars and 35 *r*-I stars were found.

One of the stars found by the HERES survey is HE 0338-3945. This object is a star located close to the main-sequence turnoff of an old halo population with an overall metallicity of [Fe/H] ∼ -2.4, and enriched in both Ba and Eu. We decided to obtain spectra of higher quality of this star and perform a detailed analysis which is presented here. In Sects. 2 and 3 the observations and analysis respectively are described. In Sect. 4 the abundance results are presented. In Sect. 5 the abundances are compared with those of other stars. This reveals HE 0338-3945 to be very similar to another *r* + *s* star, HE 2148-1247 (Cohen et al. 2003). This result led us to survey the known *r* + *s* stars in the literature, and this survey, along with a comparison with HE 0338-3945 is also presented. In Sect. 6 the results for HE 0338-3945 are discussed, as is both

**Table 1.** Basic data and parameters for the star HE 0338–3945.

Quantity	Value	Uncertainty
RA (J2000)	03h39m54.9s	$\pm 0.1$ s
Dec (J2000)	$-39^\circ 35' 44''$	$\pm 1''$
<i>V</i>	15.333 mag	$\pm 0.007$ mag
<i>B – V</i>	0.420 mag	$\pm 0.016$ mag
<i>V – R</i>	0.235 mag	$\pm 0.011$ mag
<i>V – I</i>	0.546 mag	$\pm 0.010$ mag
<i>J</i>	14.403 mag	$\pm 0.032$ mag
<i>H</i>	14.114 mag	$\pm 0.035$ mag
<i>K</i>	14.084 mag	$\pm 0.059$ mag
<i>E(B – V)<sup>1</sup></i>	0.000 mag	
<i>E(B – V)<sup>2</sup></i>	0.013 mag	
<i>T<sub>eff</sub></i>	6160 K	$\pm 100$ K
[Fe/H]	$-2.42$ dex	$\pm 0.11$ dex
log <i>g</i>	4.13 dex	$\pm 0.33$ dex
$\xi_t$	$1.13 \text{ km s}^{-1}$	$\pm 0.22 \text{ km s}^{-1}$
$V_r$	$177.9 \text{ km s}^{-1}$	$\pm 0.5 \text{ km s}^{-1}$
Class.	turn-off star	

<sup>1</sup> Burstein & Heiles (1982). <sup>2</sup> Schlegel et al. (1998).

the classification and formation of *r + s* stars in general. Finally, in Sect. 7 the conclusions are presented.

## 2. Observations and data reduction

The basic data for the star HE 0338–3945 are presented in Table 1. It was observed in Service Mode at the VLT Unit Telescope 2 (Kueyen) equipped with the spectrograph UVES at the European Southern Observatory at Paranal, Chile, during the nights of 11 December and 23 to 25 December, 2002.

To cover as large a spectral range as possible, UVES was used in dichroic mode with two different settings: Dichroic 1 with central wavelengths of 3460 Å (3055–3850 Å) and 5800 Å (4795–5745 Å and 5855–6755 Å) in the blue and red arm of the spectrograph respectively, and Dichroic 2 with central wavelengths of 4370 Å (3760–4945 Å) and 8600 Å (6720–8425 Å and 8710–10510 Å). The exposure times were in total 6 h for the Dichroic 1 setting and 9.5 h for the Dichroic 2 setting, consisting of single exposure times of 30 to 75 min. The projected slit width was set to 1.2'', yielding a resolving power of  $R = \lambda/\Delta\lambda$  equal to 30 000–40 000. During the observations, the CCD binning was set to  $1 \times 1$ , but after reduction the spectra were rebinned by a factor of two to increase the signal-to-noise ratio (*S/N*) per pixel. Since the pixel scale after rebinning is 0.44''/pixel and 0.32''/pixel in the blue and red arm, respectively, the spectra are appropriately sampled.

The optical photometry data (*BVRI*) for HE 0338-3945 were obtained on the night of 24 October 2002 by J. Holmberg using the Danish 1.5 m telescope with DFOSC on La Silla. Details of the reduction and analysis of these data are provided by Beers et al. (2005, in preparation). These data were supplemented by near-IR *JHK* from the 2MASS Point Source Catalog (Cutri et al. 2003).

The reduction was made in the standard way by using the IDL-package REDUCE, Piskunov & Valenti (2002). There were some problems with discontinuities and other reduction artifacts, cosmic ray hits, and noise in the spectrum, so the usable spectral ranges were finally 3100–5741 Å and 5847–8487 Å, now given in the rest frame of the star. The *S/N* in the spectra was rather poor at the blue end. At 3700 Å it had risen to about 40, and from 4600 Å it was 70 or higher, peaking at 6700 Å at 130. The *S/N* was about 74 in mean for the final spectra.


## 3. Abundance analysis

### 3.1. Fundamental parameters and atmospheric model

In Paper II, a temperature of  $6162 \pm 100$  K was derived from photometry when adopting reddening derived from maps of Schlegel et al. (1998). A subsequent analysis of the snapshot spectrum gave  $\log g = 4.09$  and [Fe/H] =  $-2.41$ . We note that if instead the reddening maps of Burstein & Heiles (1982) were adopted the temperature would be 70 K lower.

The effective temperature for analysis of the higher quality spectrum was redetermined from H $\beta$  and H $\delta$ . The continuum rectification and analysis, including error estimation, were performed following Barklem et al. (2002). These two lines were employed as the continuum rectification method was able to be successfully applied. The gravity and metallicity from Paper II were adopted during the analysis of these Balmer lines. We obtained  $T_{\text{eff}} = 6160 \pm 140$  K, in perfect agreement with the result from photometry. Based on these results, we adopted an effective temperature of  $6160 \pm 100$  K for our analysis. Attempts were also made to derive the effective temperature by comparing the observed colours with calculated colours of the MARCS model atmospheres (Edvardsson et al., in preparation). We found values of  $T_{\text{eff}}$  from *B – V*, *V – R* and *V – I* to range between 6100 K and 6350 K.

A small grid of models in  $\log g$  and [Fe/H] for a fixed  $T_{\text{eff}}$  was calculated with the updated Uppsala model atmosphere code MARCS (Gustafsson et al. in preparation). All models used scaled solar abundances with the exception of the alpha-elements, which were enhanced by 0.4 dex. A microturbulence of  $1 \text{ km s}^{-1}$  was adopted for the line opacities. This grid was used for an initial analysis of the spectrum, deriving  $\log g$  and chemical abundances as described in Sect. 3.3. The parameters from Paper II were used as the initial guess. As C, N and O abundances are significantly enhanced in this star, the grid was then recomputed enhancing these abundances by amounts suggested by the initial analysis. The changes in these abundances led to a warming of the model atmosphere by about 25 K in the typical line forming regions. A final analysis of the spectrum was then made as described in Sect. 3.3, giving final model parameters of  $T_{\text{eff}} = 6160 \pm 100$  K,  $\log g = 4.13 \pm 0.33$  dex, and [Fe/H] =  $-2.42 \pm 0.11$ . The final microturbulence  $\xi_t$  was  $1.13 \pm 0.22 \text{ km s}^{-1}$ . In Fig. 1 we show that the adopted model is consistent with excitation and ionisation equilibrium for Fe and Ti, and that there are no trends in Fe and Ti abundances with line strength or wavelength. We compared these stellar parameters with a [Fe/H] =  $-2.5$  and 12 Gyr



**Fig. 1.** Plots showing trends of abundances found from individual Fe and Ti lines with excitation potential (top), wavelength (middle) and equivalent width (bottom). Circles and crosses indicate lines from the neutral and singly ionised species respectively, and the dashed lines indicate the determined abundances from the simultaneous best fit to all lines of the element in question.

isochrone from Kim et al. (2002), and the result lies near the turnoff but between the positions of the main-sequence and the sub-giant stage. We note that  $\log g$  is expected to be underestimated since overionisation of Fe I will lead to an underestimate of the gravity in our LTE analysis (estimating the pressure in the atmosphere to be lower than it actually is), and thus this star is most likely closer to the main-sequence than the sub-giant branch.

### 3.2. Line selection and data

The lines used in the abundance analysis were selected through a thorough search and selection process. All lines in the observed wavelength regions were extracted from the Vienna Atomic Line Data Base (VALD, Kupka et al. 1999). Predicted

equivalent widths were then calculated based on a MARCS model atmosphere with the stellar parameters and abundances of Paper II. For  $n$ -capture element abundances not derived in Paper II we used solar values appropriately scaled to have similar enhancements to other  $n$ -capture elements. Predicted strengths of C<sub>2</sub>, CH and CN features, in the wavelength ranges were also calculated. For elements with many available lines of reasonable strength, lines with equivalent widths predicted to be smaller than 0.1 mÅ were rejected, and the remaining lines which were predicted not to be overly blended were then scrutinised in the UVES spectra, checking that the line is observed and is indeed unblended. For elements of interest with very few lines, (e.g. Ag, Tb) all possible lines were scrutinised. The lines from the VALD search were complemented with lines of Paper II, and Sneden et al. (1996, 2003a). The preliminary line list was then applied to an initial analysis of the spectrum (see Sect. 3.3), and following visual inspection of the results each line was either adopted in the final line list or rejected. Lines were rejected if the predicted line from the derived abundance did not agree well with the observed line, with respect to the majority lines of the element. This generally indicates a blend or erroneous  $\log gf$ . A large number of lines were rejected in this way. Most blends were found to be due to CH and Nd.

The final line list comprises 621 lines for measuring abundances and 650 blending lines. Blending lines are modelled in the calculated synthetic spectrum using abundances derived from other lines, but are not themselves used in determining the abundances. Atomic data has been compiled for the selected lines. Where several sources of atomic data were available, the data were generally extracted in the following order of priority: Paper II, Sneden et al. (2003a), Sneden et al. (1996), laboratory data from VALD, Jonsell et al. (2005), theoretical data from VALD. In a few cases, we were aware of recent laboratory  $f$ -values which were preferred, e.g. in the case of Nd. In general this reflects our preference for well checked data, and for laboratory data over astrophysical or theoretical data. The original sources of the data were traced where possible, and are presented with the final line list in Table 2, which is only available electronically. The wavelength  $\lambda$ , excitation potential  $\chi$ ,  $\log gf$ -value, collisional broadening due to neutral hydrogen, Stark broadening, and radiation damping constant,  $\gamma_{\text{rad}}$ , are tabulated in Table 2, along with the spectral windows used in the abundance analysis (see the description of the method in Paper II). When applicable and available, data for the isotopic splitting and hyperfine structure (hfs) are given. Blending lines considered are also included in the list. In the Appendix we briefly comment on line selection issues and data sources for each element.

### 3.3. Analysis method

The spectrum has been analysed using the automated spectrum analysis code based on SME (Valenti & Piskunov 1996) which is described in detail in Paper II. Here we summarise the most relevant aspects of the method. The synthetic spectrum was calculated assuming LTE, a 1D plane-parallel geometry for the atmosphere, and Doppler broadening was modelled

through the classical microturbulence and macroturbulence parameters. Model parameters were optimised to minimise the  $\chi^2$  statistic comparing the synthetic and observed spectra. The effective temperature was derived independently as described in Sect. 3.1 and held fixed throughout. The remaining atmospheric parameters ( $\log g$ , [Fe/H],  $\xi_t$ ,  $v_{\text{macro}}$ ) were first determined from an analysis of Fe and Ti lines. These parameters were then adopted, and abundances for all elements determined from appropriate lines considering also blends in the region. As a large number of lines regarded as blends were included in this analysis, a second iteration was performed to ensure that these lines were modelled with appropriate abundances. In some instances where lines were undetectable, we calculated upper limits to the abundances. We computed  $3\sigma$  upper limits using the same code by finding the abundance necessary to produce a line with an equivalent width three times the  $1\sigma$  measurement error in the equivalent width due to noise that a weak line at the relevant location in the spectrum would have.

One significant upgrade of the code was performed, namely the partition functions from the MOOG code (Sneden 1973) have been adopted, which have been significantly updated and corrected in 2002 (see <http://verdi.as.utexas.edu/moog.html>). Some significant corrections to the previously used polynomial fits to the partition functions of Irwin (1981) have been found, e.g. Tb II (Lawler et al. 2001b).

Error estimates in all quantities are computed as detailed in Paper II. The propagation of errors from stellar parameters, oscillator strengths, observational error and continuum placement are considered in detail. In Paper II we also estimated contributions to the error from modelling uncertainties, such as the assumptions of LTE and 1D plane-parallel geometry, as these may contribute to star-to-star scatter in a sample of stars. However, in this case we are examining a single star and this error has not been included, i.e.  $\sigma_e(\text{model}) = 0$ .

As in Paper II, we make a distinction between absolute and relative error estimates. The absolute error,  $\sigma^{\text{abs}}$ , estimates the total uncertainty in a given quantity. The relative error,  $\sigma^{\text{rel}}$ , is an error where mainly the uncertainty in the line  $f$ -values ( $\sigma_{\log gf}$ ) has been neglected. Thus, it is appropriate for star-to-star comparison where the same lines and  $f$ -values are used, as in our homogeneous analysis with HE 2148–1247 (see Sect. 5.3).

Since the procedure globally fits the spectral lines of a given element, an average uncertainty for each element  $\sigma_{\log gf}$  is required when computing  $\sigma^{\text{abs}}$ . As a number of elements studied here were not included in Paper II, in Table 3 we provide an estimated average uncertainty for each element with reference to the original literature, which are adopted for  $\sigma_{\log gf}$ . In the cases of Ho and Hf, we could find no error estimate for the oscillator strengths and adopted 0.2 dex as an estimate.

## 4. Results

### 4.1. Abundance results

We have derived abundances for 33 elements and upper limits for an additional 6 elements. The results of the abundance analysis are given in Table 4 and the results for the  $n$ -capture

**Table 3.** Assigned average values of  $\sigma_{\log gf}$  for each element.

Element	$\sigma_{\log gf}$	Element	$\sigma_{\log gf}$	Element	$\sigma_{\log gf}$
C	0.10	Co	0.10	Eu	0.03
N	0.10	Ni	0.03	Gd	0.05
O	0.02	Cu	0.04	Tb	0.04
Na	0.02	Sr	0.10	Dy	0.04
Mg	0.07	Y	0.03	Ho	0.20
Al	0.11	Zr	0.03	Er	0.12
Ca	0.11	Ag	0.04	Tm	0.02
Sc	0.04	Ba	0.03	Yb	0.05
Ti	0.05	La	0.03	Lu	0.04
V	0.05	Ce	0.10	Hf	0.20
Cr	0.05	Pr	0.04	Pb	0.05
Mn	0.06	Nd	0.03	Th	0.02
Fe	0.03	Sm	0.05	U	0.05

elements are displayed in Fig. 2. In the table we present the derived 1D LTE abundances together with the estimated relative and absolute errors (see Sect. 3.3). In calculating the error estimates, the sensitivity of each abundance to a change in each stellar parameter is calculated, and the results are shown in Table 5.

As has been described above, this analysis is based on the assumptions of 1D plane-parallel geometry and LTE. Such a 1D LTE analysis is still the most usual way to estimate the chemical composition of stars. However, in recent decades, more and more studies of the effects of deviations from these traditional assumptions have been carried out, and these studies have shown that 1D LTE analysis may yield abundances far from the “true” values. This is particularly true for metal-poor stars, which according to 3D calculations are on average much cooler in the line forming layers than predicted by 1D model atmospheres (Asplund 2004). The abundance effects may be as large as –0.5 dex for minority species and low excitation lines, and even more for molecular lines. Also severe NLTE effects may arise due to the steeper temperature structures and temperature inhomogeneities (Asplund 2005).

While this analysis is a 1D LTE analysis, and the 1D LTE abundances will be discussed in the paper, it is of interest to examine the estimated effects of 3D and NLTE, and these should be born in mind in interpretation of the abundances. For the purposes of our following discussion, we define the estimated effects:

$$\Delta = \log \epsilon_{\text{NLTE},3D} - \log \epsilon_{\text{LTE},1D}, \quad (1)$$

$$\Delta_{\text{NLTE}} = \log \epsilon_{\text{NLTE},1D} - \log \epsilon_{\text{LTE},1D}, \quad (2)$$

$$\Delta_{3D} = \log \epsilon_{\text{LTE},3D} - \log \epsilon_{\text{LTE},1D}. \quad (3)$$

In the absence of complete line formation calculations in 3D-NLTE, of which there are presently very few, the effects for the 1D-NLTE and 3D-LTE cases are discussed. Note that  $\Delta$  is generally different from the sum of  $\Delta_{\text{NLTE}}$  and  $\Delta_{3D}$ . In Table 6 we present estimates of the 3D and NLTE effects for this star


**Table 4.** Derived elemental abundances for HE 0338–3945. For each element X the LTE abundance is presented as  $\log \epsilon_X$  and [X/Fe], together with their relative,  $\sigma_{\log \epsilon_X}^{\text{rel}}$ , and absolute,  $\sigma_{\log \epsilon_X}^{\text{abs}}$ , rms error estimates (see text).  $N_{\text{tot}}$  gives the number of lines of the element used for abundance determination for each element, and  $N_{3\sigma}$  gives the number of those features classified as  $3\sigma$  detections.

	Element		$\log \epsilon_X$	$\sigma_{\log \epsilon_X}^{\text{rel}}$	$\sigma_{\log \epsilon_X}^{\text{abs}}$	[X/Fe]	$\sigma_{[\text{X}/\text{Fe}]}^{\text{rel}}$	$\sigma_{[\text{X}/\text{Fe}]}^{\text{abs}}$	$N_{\text{tot}}$	$N_{3\sigma}$
			[dex]	[dex]	[dex]	[dex]	[dex]	[dex]		
6	C	(CH)	8.08	0.09	0.15	2.13	0.06	0.15	—	— <sup>1</sup>
7	N	(CN)	6.92	0.16	0.20	1.55	0.09	0.17	—	— <sup>1</sup>
8	O	(O I)	7.66	0.04	0.09	1.40	0.09	0.11	2	1
11	Na	(Na I)	4.26	0.03	0.06	0.36	0.07	0.10	3	2
12	Mg	(Mg I)	5.45	0.02	0.09	0.30	0.09	0.14	7	5
13	Al	(Al I)	3.16	0.09	0.15	-0.88	0.03	0.13	1	1
20	Ca	(Ca I)	4.31	0.05	0.12	0.38	0.07	0.14	10	7
21	Sc	(Sc II)	1.26	0.13	0.16	0.53	0.05	0.09	5	2
22	Ti	(Ti I+II)	2.96	0.12	0.14	0.37	0.04	0.08	24	11
23	V	(V I+II)	1.76	0.13	0.16	0.19	0.05	0.09	5	0
24	Cr	(Cr I+II)	3.12	0.09	0.11	-0.12	0.03	0.08	8	2
25	Mn	(Mn I+II)	2.47	0.11	0.12	-0.49	0.04	0.08	5	4
26	Fe	(Fe I+II)	5.07	0.10	0.11	0.00	0.03	0.05	69	30
27	Co	(Co I)	2.72	0.09	0.14	0.23	0.03	0.11	4	3
28	Ni	(Ni I)	3.83	0.10	0.11	0.01	0.03	0.06	8	5
29	Cu	(Cu I)	1.02	0.12	0.12	-0.75	0.06	0.08	2	0
38	Sr	(Sr I+II)	1.28	0.08	0.16	0.74	0.03	0.14	5	2
39	Y	(Y II)	0.63	0.13	0.16	0.83	0.05	0.08	11	6
40	Zr	(Zr II)	1.37	0.13	0.15	1.20	0.04	0.08	16	7
47	Ag	(Ag I)	<0.45	0.15	0.16	<1.94	0.11	0.12	2	0
56	Ba	(Ba II)	2.10	0.06	0.07	2.41	0.05	0.08	3	3
57	La	(La II)	1.01	0.13	0.16	2.28	0.04	0.08	8	6
58	Ce	(Ce II)	1.30	0.14	0.18	2.16	0.05	0.12	15	8
59	Pr	(Pr II)	0.44	0.16	0.19	2.16	0.07	0.11	4	0
60	Nd	(Nd II)	1.37	0.14	0.16	2.30	0.05	0.08	34	15
62	Sm	(Sm II)	0.82	0.15	0.18	2.25	0.06	0.10	4	1
63	Eu	(Eu II)	0.02	0.14	0.17	1.94	0.05	0.09	3	3
64	Gd	(Gd II)	0.69	0.13	0.16	2.00	0.05	0.09	8	0
65	Tb	(Tb II)	<0.13	0.16	0.18	<2.66	0.09	0.11	1	0
66	Dy	(Dy II)	0.55	0.14	0.17	1.84	0.05	0.09	16	3
67	Ho	(Ho II)	<1.16	0.15	0.27	<3.33	0.08	0.23	1	0
68	Er	(Er II)	0.74	0.14	0.20	2.24	0.05	0.15	3	1
69	Tm	(Tm II)	0.06	0.13	0.15	2.49	0.05	0.08	4	0
70	Yb	(Yb II)	0.77	0.13	0.15	2.12	0.07	0.09	2	2
71	Lu	(Lu II)	<0.72	0.15	0.17	<3.09	0.09	0.12	1	0
72	Hf	(Hf II)	0.94	0.17	0.28	2.49	0.10	0.23	1	1
82	Pb	(Pb I)	2.62	0.10	0.11	3.10	0.04	0.08	2	1
90	Th	(Th II)	<0.23	0.17	0.19	<2.57	0.08	0.11	1	0
92	U	(U II)	<-0.11	0.38	0.74	<2.82	0.46	0.80	1	0

<sup>1</sup> The detection exceeds  $3\sigma$  for the molecular bands seen as single features.

and the spectral features employed based on results from the literature. The values are in a number of cases uncertain. In particular the estimates of NLTE effects are often plagued by uncertainties in collision rates. Note that a range of 3D effects are often quoted as the effects on individual lines may vary.

Below, we discuss the results and relevant issues in the analysis for each element in turn. Elements not discussed individually have a significant number of unblended lines with adequate  $S/N$  for a clear detection were available, the analysed lines were all well fit by a single abundance using our adopted model, no NLTE or 3D corrections were available, and the



**Fig. 2.** The abundance pattern of HE 0338–3945 (full circles and downward arrows) compared with the solar *r*-process pattern (solid line) scaled to the Eu abundance of HE 0338–3945, and the solar *s*-process pattern (dotted line) scaled to the Ba abundance of the star. The estimated absolute error bars are shown. The *r*- and *s*-process fractions are from Arlandini et al. (1999), except for Th and U which are from Burris et al. (2000). Note, in the case of upper limits (downward arrows) the plotted horizontal bar marks the upper limit *plus* the  $1\sigma$  absolute error estimate in the upper limit.

abundance was unremarkable with respect to “normal” metal-poor stars (see Sect. 5.2). Note that line selection and atomic data for each element are discussed in the Appendix.

**Carbon:** HE 0338–3945 is carbon-rich and the two strong bands and two lines of CH used in the abundance analysis were well fitted. The abundance is, however, uncertain, due to the strong temperature sensitivity of the CH molecule formation (see Table 5). In addition, preliminary calculations suggest that  $\Delta_{3D}$  may amount to approximately –0.5 to –0.3 dex (Asplund & García Pérez 2001; Asplund 2004).

The isotopic ratio  $^{12}\text{C}/^{13}\text{C}$  was determined from reasonably well isolated  $^{13}\text{CH}$  features between 4210 and 4225 Å. The comparisons of observed and synthetic spectra are shown in Fig. 3, where the total derived C abundance derived is adopted. We find  $^{12}\text{C}/^{13}\text{C} \sim 10$ .

**Nitrogen:** The *N* abundance was derived based on bands of the CN molecule. The abundance is uncertain, first because it is both highly temperature sensitive (see Table 5) and dependent on the C abundance, and secondly because it is probably subject to 3D effects. The value of  $\Delta_{3D}$  for CN is expected to be lower than for NH, which may be as much as –0.6 to –0.9 dex for turn-off stars (Asplund 2005). A reasonable estimate of the effect may amount to –0.5 dex. Note, the NH  $\Delta\Pi-\chi\zeta$  band

at 3360 Å gave an abundance consistent with the CN result, but was disregarded in the final analysis due to poor *S/N* and a large degree of blending.

**Oxygen:** Only two of the three lines of the infrared OI triplet at 7773 Å were used in the analysis, as the 7775 Å line was affected by a reduction artefact. The [OI] line at 6300 Å could not be detected. A value of  $\Delta_{\text{NLTE}} \sim -0.1$  dex has been adopted for stars with similar fundamental parameters by Nissen et al. (2002).  $\Delta_{3D}$  is estimated to be approximately –0.1 dex by Asplund (2005).

**Sodium:** The derived Na abundance was dominated by the NaI D lines at 5890 Å, and is determined quite precisely as seen from the small estimated errors in Table 4. However, the true Na abundance may be lower due to NLTE effects. From interpolation in Table 2 of Baumüller et al. (1998) we estimate  $\Delta_{\text{NLTE}} \sim -0.4$  dex. The 3D effect seems not to be severe for sodium (Asplund 2004).

**Magnesium:** The lines of MgI were well fit, and the error in the abundance estimated to be quite small. However, it is affected by both NLTE and 3D effects. According to Asplund (2005) the NLTE effect on the abundance from the MgI line at 5172 Å is around –0.2 dex, while for other

**Table 5.** The change in the logarithmic abundances due to a change of the fundamental parameters by amounts corresponding approximately to their uncertainties.

Element	$T_{\text{eff}} + 100 \text{ K}$	$\log g + 0.3 \text{ dex}$	$\xi_t + 0.2 \text{ km s}^{-1}$
	[dex]	[dex]	[dex]
C	0.15	-0.09	-0.01
N	0.23	-0.09	0.00
O	-0.06	0.10	0.00
Na	0.06	-0.04	-0.03
Mg	0.06	-0.06	-0.01
Al	0.11	-0.03	-0.03
Ca	0.06	-0.02	0.00
Sc	0.05	0.10	-0.03
Ti	0.05	0.08	-0.03
V	0.05	0.09	0.00
Cr	0.09	-0.01	-0.04
Mn	0.10	0.01	-0.01
Fe	0.07	0.03	-0.02
Co	0.09	0.01	-0.01
Ni	0.09	0.01	-0.04
Cu	0.10	0.00	-0.01
Sr	0.08	-0.02	-0.09
Y	0.05	0.09	-0.04
Zr	0.06	0.09	-0.02
Ag <sup>1</sup>	0.11	0.01	-0.01
Ba	0.08	-0.03	-0.02
La	0.05	0.10	-0.01
Ce	0.06	0.10	-0.01
Pr	0.06	0.12	0.00
Nd	0.06	0.10	-0.02
Sm	0.07	0.11	0.00
Eu	0.06	0.11	0.00
Gd	0.05	0.10	0.00
Tb <sup>1</sup>	0.07	0.09	-0.01
Dy	0.07	0.09	-0.02
Ho <sup>1</sup>	0.05	0.11	0.00
Er	0.06	0.10	-0.01
Tm	0.05	0.09	-0.01
Yb	0.07	0.06	-0.03
Lu <sup>1</sup>	0.05	0.09	0.00
Hf	0.08	0.09	-0.06
Pb	0.11	-0.02	-0.01
Th <sup>1</sup>	0.09	0.10	-0.01
U <sup>1</sup>	0.35	-0.80	-0.01

<sup>1</sup> Effects on the upper limits are shown.

Mg I lines the effect is approximately +0.2 dex. We have used a total of 7 lines, and so  $\Delta_{\text{NLTE}}$  should be positive. This is supported by Gehren et al. (2004) who show results for 7 lines, although not exactly the same set of lines as used here. The

**Table 6.** Estimated effects of NLTE and 3D on abundances from literature and adapted to the parameters of HE 0338–3945. The numbers are in many cases highly uncertain; see text for further discussion.

Element	$\Delta_{\text{NLTE}}$	Ref. <sub>NLTE</sub>	$\Delta_{\text{3D}}$	Ref. <sub>3D</sub>
	[dex]		[dex]	
6 C (CH)			-0.5 to -0.3	a, c
7 N (CN)			-0.5	see text
8 O	-0.1	i	-0.1	b
11 Na	-0.4	d	-0.1 to +0.1	a
12 Mg	+0.1	g	-0.5 to -0.3	a
13 Al	+0.5	e	-0.5 to -0.3	a
20 Ca	+0.2	h	-0.3 to -0.1	a
26 Fe (Fe I)	+0.6	f	-0.5 to -0.3	a
26 Fe (Fe II)	+0.2	b	-0.1 to +0.1	a
38 Sr	+0.3	b	-0.5 to -0.3	a
56 Ba	+0.2	b	-0.5 to -0.1	a
63 Eu	+0.1	b		


References: a – Asplund (2004), b – Asplund (2005), c – Asplund & García Pérez (2001), d – Baumüller et al. (1998), e – Baumüller & Gehren (1997), f – Collet et al. (2005), g – Gehren et al. (2004), h – Korn & Mashonkina (2006, in preparation), i – Nissen et al. (2002).

resulting value of  $\Delta_{\text{NLTE}}$  for stars with parameters close to HE 0338–3945 is  $\sim +0.1$  dex. Asplund (2004) estimate that  $\Delta_{\text{3D}}$  may be  $-0.5$  to  $-0.3$  dex for the low excitation lines.

**Aluminium:** The abundance of Al was derived using only the strong resonance line of Al I at 3961 Å, and gives a clearly sub-solar value relative to Fe as is the norm for metal-poor stars (see Sect. 5.2).  $\Delta_{\text{NLTE}}$  could, according to an interpolation made in Table 1 of Baumüller & Gehren (1997), be as high as +0.5 dex. According to Asplund (2004, 2005)  $\Delta_{\text{3D}}$  may amount to a similar value but of opposite sign for low excitation lines.

**Calcium:** The calcium abundance was determined from lines of Ca I. A value of  $\Delta_{\text{NLTE}} \sim +0.2$  dex was adopted from recent work of Korn & Mashonkina (2006, in preparation), and according to Asplund (2004) the  $\Delta_{\text{3D}}$  may amount to  $-0.3$  to  $-0.1$  dex.

**Scandium:** The abundance of scandium was determined using 5 Sc II lines. The abundance seems to be high in comparison to normal metal-poor stars (see Sect. 5.2). The two bluest lines are strongest and dominate the fitting, and are thus well fit. The fit does, however, seem to overestimate the strength of the three weak lines redward of 5000 Å. This may suggest that the blue lines are affected by unknown or poorly modelled blends. We compared the [Sc/Fe] and [C/Fe] abundances from Paper II, and found that while the results are uncorrelated for  $[\text{C}/\text{Fe}] \lesssim 1.5$ , for  $[\text{C}/\text{Fe}] \gtrsim 1.5$  there is a trend to high [Sc/Fe] with high [C/Fe]. Thus, our Sc abundance may be overestimated.



**Fig. 3.** Comparison of observed and synthetic spectra for  $^{13}\text{CH}$  features. The double line shows the observed spectrum. The thin lines show synthetic spectra for isotope ratios  $^{12}\text{C}/^{13}\text{C} = 3, 10$ , and  $30$ .

**Vanadium:** Five weak lines in the blue were used to derive the abundance, the best being only detected at the  $1.8\sigma$  level. However, all lines of both species, VI and V II, are well modelled and we consider this a reliable detection.

**Manganese:** Three lines of Mn II and two of Mn I were analysed. The  $S/N$  was significantly better at the Mn I lines than at the Mn II lines, and thus the Mn I lines dominated the fit. However, the derived abundance did not reproduce the Mn II lines well, the model prediction being significantly weaker than the observed by a consistent amount for all three lines. The discrepancy may be due to overionisation of Mn I. Such a discrepancy has also been noticed by Johnson (2002).

**Iron:** A total of 61 Fe I and 8 Fe II lines were analysed. In Table 6 we give estimates for  $\Delta_{3D}$  and  $\Delta_{NLTE}$  for each species separately. As our Fe abundance is dominated by the numerous Fe I lines the effects on Fe I should dominate. Note, however, that the effects of NLTE are presently highly uncertain due to uncertainty in inelastic collision rates due to hydrogen atoms.

For example, Korn et al. (2003) found much smaller NLTE effects when astrophysically calibrating these collision rates. As noted in Sect. 3.1 NLTE effects on Fe I are expected to lead to underestimation of  $\log g$  in LTE, and this would affect all other elements.

**Copper:** The Cu abundance was derived from two lines in the ultraviolet. Both lines were clearly seen and identified, and well modelled with a single abundance. However, the low  $S/N$  ratio in this spectral region meant that the strongest line was detected only at the  $2.9\sigma$  level, and continuum placement was uncertain. Despite this, we regard it as a definite detection.

**Strontium, yttrium, zirconium:** Sr, Y and Zr, are strongly overabundant compared to normal metal-poor stars (see Sect. 5.2). Interestingly, the LTE abundances of these elements lie below the solar *s*-process abundance pattern as normalised to Ba (see Fig. 2), and curiously enough, at least Y and Zr fit the *r*-process abundance pattern normalised to Eu. Strontium is estimated to have  $\Delta_{NLTE} \sim +0.3$  dex for the lines at 4077, 4161 and 4215 Å in metal-poor turn-off stars (Asplund 2005). We analysed two more weak lines, one with low  $S/N$ . The stronger lines at higher  $S/N$  have high weight in the abundance calculation, and therefore this estimate for  $\Delta_{NLTE}$  is appropriate.  $\Delta_{3D}$  is approximately  $-0.3$  to  $-0.5$  (Asplund 2004).

**Silver:** The upper limit to the silver abundance is estimated from two Ag I lines, both blended and of roughly equal intrinsic strength. The upper limit is sufficiently low to be inconsistent with the scaled solar *r*-process abundance pattern (see Fig. 2). This may reflect overionisation of Ag I (see Sect. 6).

**Ba**rium: Ba is strongly overabundant compared to normal metal-poor stars (see Sect. 5.2). This abundance was derived using three Ba II lines: two weak and the strong resonance line. The resonance line at 4554 Å has an estimated value of  $\Delta_{NLTE}$  of at least  $+0.2$  dex (Asplund 2005).  $\Delta_{3D}$  is of the opposite sign, and is estimated at  $-0.5$  to  $-0.1$  dex (Asplund 2004).

**Lanthanum:** Lanthanum is also strongly overabundant in this star. Eight lines of La II of varying strength were employed, the majority of which were well fit by a single abundance. However, two lines, 4920 and 4921 Å were not well fit, the observed lines being stronger than the predicted lines for the derived abundance. We note that for these lines we had incomplete hfs data (see Appendix), and this could possibly lead to the line strengths being underestimated.

**Praseodymium, Neodymium, Samarium:** Four weak lines were employed to derive the abundance of Pr, the best being only detected at the  $1.6\sigma$  level. However, all 4 lines were well modelled, and thus we consider this a reliable detection. A total of 34 mostly weak lines were used to derive the Nd abundance, only slightly less than half of the lines being detected at the  $3\sigma$  level. For Sm, four weak lines of Sm II were used.

**Europium:** The abundance of Eu is enhanced compared to typical metal-poor stars (see Sect. 5.2). The abundance may be even higher, as  $\Delta_{\text{NLTE}}$  is probably greater than +0.1 (Asplund 2005). No estimate of  $\Delta_{\text{3D}}$  is available, but see the discussion regarding abundance ratios at the end of this section.

**Gadolinium:** Eight weak lines are used, the best being only a  $1.8\sigma$  detection. However, all eight lines are consistently modelled, and we class this as a reliable detection.

**Terbium:** No line was detected, and thus an upper limit was derived from a single line of Tb II.

**Holmium:** No lines are detected, and thus a  $3\sigma$  upper limit was derived. Due to strong blending at the strongest lines of Ho, we were forced to use a rather weak line and thus the derived upper limit is rather high and a weak constraint.

**Erbium, Thulium:** The abundance of Er was based on three weak Er II lines. For Tm, 4 weak lines were employed, the best being only detected at the  $1.6\sigma$  level. However, the lines of these elements were well modelled, and thus we consider these as reliable detections. Both elements have abundances significantly above the scaled solar *s*- and *r*-process patterns (see Fig. 2).

**Lutetium:** No lines were detected, and thus a  $3\sigma$  upper limit was derived. The employed line region is somewhat blended, but we regard the derived upper limit as reliable.

**Hafnium:** Only the resonance line in the ultraviolet was available, but a definite detection was made.

**Lead:** The Pb abundance is significantly above the scaled *s*-process abundance pattern in Fig. 2. Pb has an ionization energy of 7.4 eV, and the two lines observed are both Pb I lines. Here, overionisation is a possibility, which would lead to an underestimated Pb abundance and e.g. Pb/Ba ratio, meaning that this ratio, already remarkably high in the LTE analysis, may be even higher.

**Thorium:** The  $3\sigma$  upper limit was based on a single Th II line at 4019.129 Å. The region contains a number of blends, but a reliable upper limit was possible to derive.

**Uranium:** No lines are detected, so a  $3\sigma$  upper limit has been derived. The region is significantly blended, and the reliability of the derived upper limit is dependent on the correct modelling of these blends.

**Some specific abundance ratios:** We shall finally comment on the accuracy of some important abundance ratios, to be further discussed below. The similarity of Sr, Y, Zr and Ba in terms

of ionisation energies and atomic structure seems to suggest that the effects of NLTE should be limited on Ba/Sr, Ba/Y and Ba/Zr. Also, the most important lines in the analysis have low excitation energies – the Ba abundance determination is dominated by the wings of the 4554 Å resonance line, while the other two lines with 2.5 and 2.7 eV excitation energies are less significant. Thus, the temperature sensitivity of the important line strengths are roughly similar, which is reflected in similar temperature sensitivities of the abundances in Table 5. This should lead to similar sensitivities to 3D effects. Similarly, the effects of NLTE and 3D on abundance ratios of these elements relative to Eu may be expected to be small.

## 4.2. Binarity

Radial velocities were measured from a series of Fe I lines with accurate laboratory wavelengths between 4000 and 5000 Å, using the snapshot spectrum observed on 15 October 2002, and the observations from December 2002 which comprise the high quality spectrum analysed in this work. No difference in radial velocity shift was observed between the different exposures taken in December 2002. The barycentric radial velocities measured from the snapshot spectrum and the co-added high quality spectrum were  $177.7 \pm 0.8$  km s $^{-1}$  and  $177.9 \pm 0.5$  km s $^{-1}$  respectively, where the quoted errors are the standard deviations of the results from different lines; errors due to the wavelength calibration uncertainties are not included. We therefore presently have no evidence for the binarity of HE 0338-3945. A recent study by Lucatello et al. (2005) suggests all CEMP stars are multiple systems.


## 5. Comparisons

### 5.1. Comparison with the results of Paper II


As has been mentioned, a snapshot spectrum of this star was analysed in Paper II. The snapshot spectrum of HE 0338-3945, compared to the spectrum employed in this work, is of relatively low quality with  $S/N \sim 47$  and  $R \sim 20\,000$  and has significantly less wavelength coverage,  $\lambda = 3760\text{--}4980$  Å. The snapshot spectrum was reduced by the UVES pipeline, while the spectrum analysed here was reduced as described in Sect. 2. However, the spectra have been analysed in a similar manner, and it is of interest to check if the results are consistent. We have already seen in Sect. 3 that the results for the stellar parameters are in good agreement. Figure 4 compares the abundances where available from Paper II, and we see that the results are in good agreement.

### 5.2. Comparison with normal metal-poor stars of Paper II

To place HE 0338-3945 in context, it is of interest to compare this star with “normal” Population II stars of similar metallicity. We extracted all stars from Paper II with  $-2.6 < [\text{Fe}/\text{H}] < -2.2$ . It was required that the stars should have  $[\text{Eu}/\text{Fe}] < 1$  and  $[\text{C}/\text{Fe}] < 1$  to be considered as “normal”. The mean value



**Fig. 4.** Difference between the abundances for HE 0338–3945 from this work and those from Paper II.



**Fig. 5.** Difference between abundances for HE 0338–3945 and those for normal Population II stars of similar metallicity from Paper II, derived as described in the text. The vertical bars show the standard deviation among the normal Population II stars.

of [X/Fe] was then computed for the sample where more than five measurements were available.


In Fig. 5 we compare the abundances for HE 0338–3945 with the described mean abundances. For reference, the derived mean abundances and standard deviations are provided in Table 7. In Fig. 5 we clearly see the enhancements of C and the  $n$ -capture elements relative to the normal stars. For the most part the iron-peak element abundances for HE 0338–3945 appear to be normal; however, Sc appears to be significantly enhanced compared to the normal stars. The good agreement found in Sect. 5.1 with results for this star from Paper II means that this cannot be due to a difference in the analysis, e.g. use of different lines or atomic data. However, as discussed in Sect. 4.1 the Sc abundance may be spurious.

### 5.3. Comparisons with the star HE 2148–1247

Soon after examining our results for HE 0338–3945, it became apparent that this star was very similar to the star HE 2148–1247 studied by Cohen et al. (2003). The Keck/HIRES spectrum of this object used by Cohen et al. was kindly provided to us by Judith Cohen. Order merging and continuum placement was performed by us; all other reduction steps had been performed. Thus, using our automated spectrum analysis technique we were easily able to perform an analysis of this spectrum homogeneous with our analysis of HE 0338–3945. Note, as the spectral coverage of the HE 2148–1247 spectrum was not as extensive as for our HE 0338–3945 spectrum, we produced a spectrum of HE 0338–3945 with identical wavelength coverage to the HE 2148–1247 spectrum, which we refer to as the “cut” spectrum of HE 0338–3945. Thus, an analysis of the cut spectrum permits a more strictly homogeneous analysis as exactly the same spectral lines are employed.

**Table 7.** Comparison of the mean abundances relative to Fe, for normal Population II stars of Paper II in the metallicity range  $-2.6 < [\text{Fe}/\text{H}] < -2.2$ , with those for HE 0338–3945. For each element we report the number of measurements  $N$  available from Paper II, the mean value of [X/Fe], and the standard deviation of [X/Fe] for the sample, the abundance for HE 0338–3945  $[\text{X}/\text{Fe}]_*$ , and the difference  $[\text{X}/\text{Fe}]_* - \langle [\text{X}/\text{Fe}] \rangle$ .

Element	$N$	$\langle [\text{X}/\text{Fe}] \rangle$	$\sigma([\text{X}/\text{Fe}])$	$[\text{X}/\text{Fe}]_*$	$\Delta[\text{X}/\text{Fe}]$
C	58	0.18	0.30	2.13	1.95
Mg	58	0.26	0.08	0.30	0.04
Al	55	-0.86	0.23	-0.88	-0.02
Ca	58	0.27	0.10	0.38	0.11
Sc	58	0.08	0.13	0.53	0.45
Ti	58	0.24	0.12	0.37	0.13
V	18	0.00	0.18	0.19	0.19
Cr	58	-0.19	0.14	-0.12	0.07
Mn	58	-0.47	0.15	-0.49	-0.02
Co	53	0.13	0.10	0.23	0.10
Ni	58	-0.06	0.14	0.01	0.07
Sr	58	-0.10	0.29	0.74	0.84
Y	35	-0.20	0.16	0.83	1.03
Zr	16	0.27	0.18	1.20	0.93
Ba	55	-0.30	0.42	2.41	2.71
La	6	-0.06	0.24	2.28	2.34
Nd	10	0.24	0.52	2.30	2.06
Eu	21	0.39	0.29	1.94	1.55



**Fig. 6.** Difference between abundances for HE 0338–3945 (cut spectrum) and HE 2148–1247. The dashed line shows the mean difference.

For our analysis of HE 2148–1247, we adopted the effective temperature from Cohen et al.,  $T_{\text{eff}} = 6380$  K. The abundance results from our analysis of HE 2148–1247 and the cut spectrum of HE 0338–3945 are compared in Fig. 6. The abundance patterns of the two stars are seen to be very similar. An offset of -0.2 dex between the abundances is seen, but this could easily be due to errors in  $T_{\text{eff}}$  which were not derived consistently. The scatter is consistent with the uncertainties. From this we conclude that the stars have identical abundance patterns for the elements considered, within the errors in the analysis.

**Table 8.** Data for stars of different classes. For  $r$ -I stars and stars not  $n$ -enhanced, see Paper II, Burris et al. (2000), and Fulbright (2000), although data are presented for some outliers of interest from Burris et al. When several investigations exist, either the most reliable source or the mean of recent data has been taken. The mean of abundance ratios [X/Y], Y $\neq$ Fe, was calculated from the mean of the abundances relative to Fe, and not from individual measurements of [X/Y]. (.) indicates an insecure measurement, and such cases are not considered when calculating the means. References: (1) This work, (2) Aoki et al. (2001), (3) Aoki et al. (2002), (4) Barbuy et al. (2005), (5) Paper II, (6) Unpublished preliminary result obtained in Paper II, (7) Burris et al. (2000), (8) Cohen et al. (2003), (9) Hill et al. (2000), (10) Hill et al. (2002) and Plez et al. (2004), (11) Honda et al. (2004a), (12) Ivans et al. (2005), (13) Johnson & Bolte (2002), (14) Johnson & Bolte (2004), (15) Jonsell et al. (2005), (16) Lucatello et al. (2003), (17) Preston & Sneden (2001), (18) Sivarani et al. (2004), (19) Sneden et al. (2003a), (20) Van Eck et al. (2003), (21) Zacs et al. (1998), (22) mean of presented data, (23) mean of (4) and (17).

<i>r + s</i> stars																		
Name	$T_{\text{eff}}$	$\log g$	[Fe/H]	[C/Fe]	[Sr/Fe]	[Y/Fe]	[Zr/Fe]	[Ba/Fe]	[La/Fe]	[Ce/Fe]	[Eu/Fe]	[Pb/Fe]	[Ba/Eu]	[La/Eu]	[Ce/Eu]	[Pb/Ba]	Ref.	
CS 22183-015	5200	2.5	-3.12	2.2:		0.45	0.62	2.09	1.59	1.55	1.39	3.17	0.70	0.20	0.16	1.08	(13)	
CS 22898-027			-2.20	2.08	0.76	0.84	1.20	2.25	2.21	2.13	1.91	2.84	0.34	0.30	0.22	0.59	(22)	
	6250	3.7	-2.25	2.2	0.92	0.73	1.01	2.23	2.13	2.13	1.88	2.84	0.35	0.25	0.25	0.61	(3)	
	6300	4.0	-2.15	1.95	0.60	0.95	1.39	2.27	2.28		1.94		0.33	0.34			(17)	
CS 22948-027			-2.52	2.27	0.90	1.00		1.97	2.32	2.20	1.88	2.72	0.09	0.44	0.32	0.75	(23)	
	4800	1.8	-2.47	2.43	0.90	1.00		2.26	2.32	2.20	1.88	2.72	0.38	0.44	0.32	0.46	(4)	
	4600	0.8	-2.57	2.10				1.67									(17)	
	4800	1.80	-2.46	1.8	0.90	1.00		2.26	2.32	2.20	2.10		0.16	0.22	0.10		(9)	
CS 29497-030			-2.64	2.43	1.09	0.84	1.42	2.25	2.16	2.12	1.72	3.60	0.53	0.44	0.40	1.35	(22)	
	7000	4.1	-2.57	2.47	1.34	0.97	1.40	2.32	2.22	2.10	1.99	3.65	0.33	0.23	0.11	1.33	(12)	
	6650	3.5	-2.70	2.38	0.84	0.71	1.43	2.17	2.10	2.14	1.44	3.55	0.73	0.66	0.70	1.38	(18)	
CS 29497-034	4800	1.8	-2.90	2.63	1.00	1.10		2.03	2.12	1.95	1.80	2.95	0.23	0.32	0.15	0.92	(4)	
	4800	1.80	-2.91	2.3	1.00	1.10		2.03	2.12	1.95	2.25		-0.22	-0.13	-0.30		(9)	
CS 29526-110	6500	3.2	-2.38	2.2	0.88		1.11	2.11	1.69	2.01:	1.73	3.30	0.38	-0.04	0.28	1.19	(3)	
CS 31062-012	6250	4.5	-2.55	2.1	0.30	0.59		1.98	2.02	2.12	1.62	2.40	0.36	0.40	0.50	0.42	(3)	
CS 31062-050			-2.37	2.00	0.91	0.48	0.94	2.55	2.28	2.02	1.82	2.86	0.73	0.46	0.20	0.31	(22)	
	5500	2.70	-2.42	2.00		0.48	0.85	2.80	2.12	2.02	1.79	2.81	1.01	0.33	0.23	0.01	(14)	
	5600	3.0	-2.32	2.00	0.91		1.02	2.30	2.44	2.10:	1.84	2.90	0.46	0.60	0.26	0.60	(3)	
	HE 0024-2523	6625	4.3	-2.71	2.6	0.34	<0.91	<1.22	1.46	1.80		<1.10	3.30	>0.36	>0.70		1.84	(16)
HE 0131-3953	5928	3.83	-2.71	2.45	0.46			2.20	1.94	1.93	1.62		0.58	0.32	0.31		(5)	
HE 0338-3945	6160	4.13	-2.42	2.13	0.74	0.83	1.20	2.41	2.28	2.16	1.94	3.10	0.47	0.34	0.22	0.69	(1)	
	6162	4.09	-2.41	2.07	0.73	0.73		2.41	2.26	2.21	1.89		0.52	0.37	0.32		(5)	
HE 1046-1352	5540	3.0:	-2.6:	2.2:	0.8:	0.5:		2.1:	1.8:	1.8:	1.4:		0.7:	0.4:	0.4:		(6)	
HE 1105+0027	6132	3.45	-2.42	2.00	0.73	0.75		2.45	2.10		1.81		0.64	0.29			(5)	
HE 1405-0822	5400	2.0:	-2.1:	1.7:		0.6:	0.8:	1.8:	1.2:	1.3:	1.1:		0.7:	0.1:	0.2:		(6)	
HE 2148-1247	6380	3.9	-2.28	1.91	0.76	0.83	1.47	2.36	2.38	2.28	1.98	3.12	0.38	0.40	0.30	0.76	(8)	
LP 625-44	5500	2.8	-2.71	2.1	1.15	0.99	1.34	2.74	2.46	2.27	1.97	2.55	0.77	0.49	0.30	-0.19	(2)	
LP 706-7	6000	3.8	-2.74	2.15	0.15	0.25	<1.16	2.01	1.81	1.86	1.40	2.28	0.61	0.41	0.46	0.27	(2)	
<i>s</i> stars																		
Name	$T_{\text{eff}}$	$\log g$	[Fe/H]	[C/Fe]	[Sr/Fe]	[Y/Fe]	[Zr/Fe]	[Ba/Fe]	[La/Fe]	[Ce/Fe]	[Eu/Fe]	[Pb/Fe]	[Ba/Eu]	[La/Eu]	[Ce/Eu]	[Pb/Ba]	Ref.	
CS 22880-074			-1.85	1.41	0.27	0.16	0.73	1.33	1.16	1.22	0.53	1.9	0.80	0.63	0.69	0.57	(22)	
	5850	3.8	-1.93	1.3	0.39	0.16		1.31	1.07	1.22	0.5	1.9	0.81	0.57	0.72	0.59	(3)	
	6050	4.0	-1.76	1.51	0.14	0.6:	0.73	1.34	1.24		0.55		0.79	0.69			(17)	
CS 22881-036	6200	4.0	-2.06	1.96	0.59	1.01		1.93	1.59		1.00		0.93	0.59			(17)	
	5000	2.4	-2.64	2.0	1.4	1.58	1.69	1.71	1.53	1.54	0.79	$\leq 1.6$	0.92	0.74	0.75	$\leq -0.11$	(22)	
CS 30301-015	4900	1.8	-2.67	2.2:	1.4			1.50	1.85		0.8:		0.7:	1.05:			(17)	
	4750	0.8	-2.64	1.6	0.3:	0.29		1.45	0.84	1.16	0.2:	1.7	1.25:	0.64:	0.96:	0.25	(3)	
	HD 196944		-2.33	1.31	0.84	0.57	0.63	1.26	1.13	1.20	0.17	2.00	1.09	0.96	1.03	0.74	(22)	
	5250	1.7	-2.23					1.14									(15)	
HD 196944	5250	1.7	-2.4			0.6			1.00	1.10		2.10					(20)	
	5250	1.8	-2.25	1.2	0.84	0.56	0.66	1.10	0.91	1.01	0.17	1.90	0.93	0.74	0.84	0.80	(3)	
	5250	1.7	-2.45	1.42		0.58		1.56	1.49	1.50							(21)	
HE 0202-2204	5280	1.65	-1.98	1.16	0.57	0.41	0.47	1.41	1.36	1.30	0.49		0.92	0.87	0.81		(5)	
HE 1135+0139	5487	1.80	-2.33	1.19	0.66	0.36	0.46	1.13	0.93	1.17	0.33		0.80	0.60	0.84		(5)	
<i>r-II</i> stars																		
Name	$T_{\text{eff}}$	$\log g$	[Fe/H]	[C/Fe]	[Sr/Fe]	[Y/Fe]	[Zr/Fe]	[Ba/Fe]	[La/Fe]	[Ce/Fe]	[Eu/Fe]	[Pb/Fe]	[Ba/Eu]	[La/Eu]	[Ce/Eu]	[Pb/Ba]	Ref.	
CS 22183-031	5270	2.80	-2.93	0.42	0.10	0.21		0.38			1.16						(11)	
	CS 22892-052	4800	1.50	-3.1	0.88	0.61	0.44	0.82	0.99	1.09	1.02	1.64	<0.9	-0.65	-0.55	-0.62	<-0.09	(19)
CS 29491-069	4884	1.81	-2.95	1.00	0.61	0.45			1.19	1.02		1.54		-0.35	-0.52		(5)	
	5103	2.45	-2.81	0.18	0.07	0.00		0.34				1.06		-0.72			(5)	
CS 29497-004	5013	2.23	-2.81	0.22	0.57	0.66	0.94	1.21	1.21		1.62		-0.41	-0.41			(5)	
CS 31082-001	4825	1.5	-2.92	0.2	0.65	0.43	0.73	1.17	1.13	1.01	1.63	0.40	-0.46	-0.50	-0.62	-0.77	(10)	
	4922	1.90	-2.78	0.22	0.53	0.56	0.89	1.18	1.18	1.06	1.66		-0.48	-0.48	-0.60		(5)	
HE 0430-4901	5296	3.12	-2.72	0.09	-0.01	0.02		0.50					1.16	-0.66				(5)
HE 0432-0923	5131	2.64	-3.19	0.24	0.47	0.51	0.88	0.72				1.25		-0.53			(5)	
HE 1127-1143	5224	2.64	-2.73	0.54	0.24	0.22			0.63			1.08		-0.45			(5)	
HE 1219-0312	5140	2.40	-2.81	-0.08	0.20	0.29	0.65	0.51	0.91		1.41		-0.90	-0.50			(5)	
HE 2224+0143	5198	2.66	-2.58	0.35	0.23	0.13	0.58	0.59	0.65		1.05		-0.46	-0.40			(5)	
HE 2327-5642	5048	2.22	-2.95	0.43	0.31	0.12		0.66	0.67		1.22		-0.56	-0.55			(5)	
Outlier "normal" stars																		
Name	$T_{\text{eff}}$	$\log g$	[Fe/H]	[C/Fe]	[Sr/Fe]	[Y/Fe]	[Zr/Fe]	[Ba/Fe]	[La/Fe]	[Ce/Fe]	[Eu/Fe]	[Pb/Fe]	[Ba/Eu]	[La/Eu]	[Ce/Eu]	[Pb/Ba]	Ref.	
HD 13979	5075	1.90	-2.26		-0.07	-0.63	-0.34	-0.50	-0.10		-0.38		-0.12	0.28			(7)	
HD 25532	5300	1.90	-1.46		0.14	-0.28	0.14	0.17	-0.07		0.10		0.07	-0.17			(7)	
HD 105546	5300	2.50	-1.27		0.45	0.13	0.45	0.42	0.05			0.32		0.10	-0.27		(7)	

#### 5.4. Comparisons with other similar stars

In addition to HE 0338–3945 and HE 2148–1247 there are several other stars that have recently been found to show both enhanced *r*- and *s*-elements. In total 17 such stars are known to us, and these are listed in Table 8.

We have investigated the similarities within the *r*- and *s*-element enhanced group, and the differences relative to other metal-poor stars. The *r*- and *s*-enhanced stars were compared with other stars of  $[Fe/H] < -1.0$  investigated in Paper II, by Burris et al. (2000), and Fulbright (2000), and some other groups listed in Table 8. When several sources were available, data were adopted from the source considered most reliable or the mean of recent data was taken.

The *r*- and *s*-enhanced stars HE 1046–1352 and HE 1405–0822 were analysed in the work of Paper II, but unpublished due to the significant pollution of the spectrum by molecular carbon features making the analysis method uncertain. The presented data for these stars must therefore be regarded as uncertain and preliminary. However, higher quality data of the star HE 1405–0822 is considered in the HERES paper Sivarani et al. (in preparation). It must also be observed that the error bars were quite large for some stars in several investigations, and that errors in stellar parameters may be the reason for stars with seemingly peculiar abundance patterns.

The plot in Fig. 7 shows a clear separation of different kinds of stars. The criteria of the classes in Table 9 in Sect. 6 were defined to achieve this. The stars were classified as *r*-I, *r*-II, *s*-element enhanced, *r* + *s*, i.e. both *r*- and *s*-element enhanced, and/or Pb enhanced. Stars with no significant enhancement, and thus not falling under any of these criteria, are in this paper called “normal” metal-poor stars.

The *r* + *s* stars are rich in both *r*- and *s*-elements, residing in the upper right corner of Fig. 7. All of these stars have  $[Ba/Eu] > 0.0$ . This is also true for the Pb stars with measurements of both Ba and Eu found in the literature. These Pb stars are also then classified as *r* + *s* stars. The *s*-enhanced stars also have  $[Ba/Eu] > 0.0$ , and may be located along the same sequence in the figure as the *r* + *s* stars. The *r*-II and *r*-I stellar classes both have  $[Ba/Eu] < 0.0$ . The *r*-II stars have in general a typically lower content of  $[Eu/Fe]$  than *r* + *s* stars, and in Fig. 7 seem to separate clearly from the *r* + *s* stars. The gap in between the groups might be due to poor statistics; this separation was also noted by Cohen et al. (2003) (however, with some overlap). The normal stars, i.e. stars with no *n*-capture element enhancement, have predominantly  $[Ba/Eu] < 0.0$ , although positive ratios may also be found. The same pattern was also found by Cohen et al. (2003, Fig. 13) and Johnson & Bolte (2004, Fig. 5), although the nomenclature is different in these studies (see Sect. 7).

The plots in Fig. 8 show the ratio of the different heavy *n*-capture *s*-elements (*hs* = Ba, La, Ce) compared to the *r*-element Eu for stars of different classes. As is also seen in Fig. 7 there is a clear separation between one hand *r* + *s* and *s* stars, and on the other hand *r*-II, *r*-I and normal stars. The different classes of stars seem to follow a single well-defined sequence in the [La/Eu] vs. [Ce/Eu] plot, but less so in the other abundance diagrams. [Ba/Eu] shows the greatest variation

**Table 9.** Definition of classes for *n*-capture-rich stars. In this article we define stars not falling under any of these classes as “normal”.


Class	Constraints in [X/Fe]	Other constraints
<i>r</i> -I	$+0.3 \leq [Eu/Fe] \leq +1.0$	$[Ba/Eu] < 0.0$
<i>r</i> -II	$[Eu/Fe] > +1.0$	$[Ba/Eu] < 0.0$
<i>r</i> + <i>s</i>	$[Ba/Fe] > +1.0$ $[Eu/Fe] > +1.0$	$[Ba/Eu] > 0.0$
<i>s</i>	$[Ba/Fe] > +1.0$ $[Eu/Fe] \leq +1.0$	$[Ba/Eu] > 0.0$
Pb	$[Ba/Fe] \geq +1.0$	$[Pb/Ba] \geq +1.0$

amplitude in the diagrams, while La and Ce seem vary less and follow each other. According to Paper II this may well be an observational effect; as seen in Table 5 the abundances of Ba and of the other elements react differently to changes in effective temperature and gravity which is essentially a consequence of the significance of the strong damping wings of the dominating Ba II line  $\lambda 4554$ . Therefore, stellar-parameter errors cancel for the ratios [La/Eu] and [Ce/Eu], but not for [Ba/Eu]. The Pb enhanced *r* + *s* star CS 29526-110 has a rather low abundance of La (Aoki et al. 2002), which makes the ratio [La/Eu] negative. This is astonishing since the other *hs*-elements (Ba, Ce) are considerably enhanced. The authors note that the effective temperature and gravity are rather uncertain for this star.

There are some outliers among the normal stars in the upper panel of Fig. 8. The parameters of these stars may be viewed in Table 8. These stars reside in or quite near the same region as the *r* + *s* stars, but are neither *r*- nor *s*-enhanced. The star BD+17°3248 is rather close to being regarded as an *r* + *s* star, as it has abundances  $[Ba/Fe] = 0.97$ ,  $[Eu/Fe] = 0.96$ , and  $[Ba/Eu] = 0.01$ . However, in Fig. 7 it lies closer to the normal stars. We note that Cowan et al. (2002) find a much lower Ba abundance for this star, which would place it firmly as a *r*-I star.

The histograms in Fig. 9 display the abundance of the heavy *s*-element Ba compared to the light *s*-element (*ls*) Sr for the different stellar classes. In spite of the small number statistics, there are clear separations in mean ratios for different classes. Normal stars have a slightly negative mean [Ba/Sr], although the spread is wide. The mean *r*-I star has slightly more Ba than Sr, and for *r*-II stars Ba dominates. The *s* stars in the diagram are very few, but tend to have even higher [Ba/Sr] than the *r*-II stars. The *r* + *s* stars are, however, the most overabundant in Ba relative to Sr of all groups, with a mean of  $[Ba/Sr] \sim 1.6$ .

We note from Table 8 that there may also be a difference in the metallicity distribution of the stars within these groups. The *r*-II stars are centred on a metallicity of  $[Fe/H] \sim -2.84$  with a scatter on the order of 0.16 dex (see also Paper II). The *r*-I stars are found across the entire range of metallicities investigated in Paper II, i.e.  $-3.4 < [Fe/H] < -1.5$ . The *r* + *s* stars from Table 8 are centred on  $[Fe/H] \sim -2.55$  with a scatter on the order of 0.26 dex. Thus, the *r* + *s* stars seem on average to have a higher metallicity and a wider range of metallicity than the *r*-II stars.



**Fig. 7.** A plot comparing the abundance of the *s*-element Ba to the *r*-element Eu for stars of different classes (see definitions in Table 9). The *r+s* stars, enhanced in both *r*- and *s*-elements, are shown as open circles, and the *r+s* stars also known to be enhanced in Pb are shown as circles with asterisks inside. The stars enriched only in *r*-elements are marked as small (*r*-I) and large (*r*-II) open squares, and stars with only *s*-enhancement are shown as diamonds. Stars not enhanced in either *r*- or *s*-elements, in this article called normal stars, are marked by small filled circles. The diagonal dotted line marks the one-to-one relation. Data are taken from Table 8 and for normal and *r*-I stars from Paper II, Burris et al. (2000), and Fulbright (2000).


## 6. Discussion

In this section we will discuss both the results for HE 0338–3945, and the *r+s* stars in general. We first summarise the abundances of HE 0338–3945 and then briefly discuss the classification of the *r+s* stars, in the light of our survey of stars with *n*-capture enhancement in Sect. 5.4. Finally, we present a detailed discussion of the possible formation scenarios for the *r+s* stars, based on both our results for the abundance pattern of HE 0338–3945 and our survey of *r+s* stars from the literature.

### 6.1. The abundances of HE 0338–3945

Abundances were derived for 33 elements and we estimated upper limits for an additional 6 elements for the star HE 0338–3945. We have confirmed the high content of the *r*-elements Eu, Gd, Dy, Er, and Tm with a mean of  $[r/\text{Fe}] \sim 2.10$  ( $[\text{Eu}/\text{Fe}] \sim 1.94$ ). The upper limits on Tb, Ho, Lu support an overall high *r*-element abundance, although the Ag upper limit is surprisingly low and not consistent with a scaled solar *r*-process distribution as normalised to Eu. This could, however, be an effect of over-ionisation of Ag I. We could detect neither Th nor U, as the spectral lines of these elements are weak and reside in heavily blended regions; only rather high upper limits were obtained.

Although the star is also enhanced in *s*-elements, there is a pronounced difference among the *ls* and the *hs* nuclei. The mean of the *ls* overabundance is  $[\text{Sr}, \text{Y}, \text{Zr}/\text{Fe}] = 0.92$ , and the *hs*  $[\text{Ba}, \text{La}, \text{Ce}/\text{Fe}] = 2.27$ . Obviously, the light *s*-elements do not fit the solar *s*-element distribution of *s*-elements when normalised to Ba. However, they do fit the *r*-element distribution as normalised to Eu. This behaviour has been noted in other stars as well, such as HE 2148–1247 (Cohen et al. 2003).




**Fig. 8.** Plots showing the ratio of the different heavy *n*-capture *s*-elements (Ba, La, Ce) compared to the *r*-element Eu for stars of different classes. Symbols and data as in Fig. 7.

The abundance of Pb is higher than expected from scaled solar *s*-element abundances as normalised to Ba, by about 1 dex. The elements that for the Sun were produced in significant amounts by both the *r*- and *s*-processes (e.g. Nd, Sm and Yb) are not unexpectedly overabundant compared to both the *r*- and the *s*-element solar distributions.

The overabundance of C is confirmed, and we have derived an isotopic ratio of  $^{12}\text{C}/^{13}\text{C} \sim 10$ . Also N and O are overabundant in the star. Na to Ni seem normal for matter contaminated by SNe of type II, the exception being Sc which is significantly enhanced compared to normal Population II stars (but see Sect. 4.1). The abundance of Cu is low relative to iron, and comparable with results for normal stars of similar metallicity (Mishenina et al. 2002; Bihain et al. 2004).

The NLTE and 3D effects discussed in Sect. 4.1 may in many cases be severe. We lack estimates of corrections for most elements, and, moreover, the estimates we have are often quite uncertain. It is thus risky to draw conclusions about



**Fig. 9.** Histograms showing the distribution of different classes of stars when comparing the heavy  $n$ -capture  $s$ -element Ba to the light  $n$ -capture  $s$ -element Sr. Data as in Fig. 7.

nucleosynthesis sites by comparing abundances of different elements. One may hope that, e.g., the net effect of NLTE and 3D could well be of the same order of magnitude for the  $ls$ -element Sr and the  $hs$ -element Ba, but the different character of the lines used may speak against such an optimistic view. Comparisons between different groups of elements, when carried out differentially relative to other stars with similar temperatures and gravities and overall metallicities may, however, be discussed with some confidence, in particular when we have some guidance from theoretical estimates from NLTE and 3D calculations. Unfortunately, 3D corrections for Eu are unavailable at present, and these effects could perhaps be of some importance.

## 6.2. The class $r + s$ stars

With the background of Sect. 5.4 we shall here discuss the classification of  $r + s$  stars. We propose to use  $[Eu/Fe]$  as a measure for the  $r$ -elements, and  $[Ba/Fe]$  for the  $s$ -elements. This choice of classification criteria, depending on single  $n$ -capture element abundances instead of e.g. all  $hs$ -elements, is motivated by simplicity and practicality. It could be of value to base a classification system on many criteria to suppress effects of errors and individual abundance fluctuations, but in practice both Eu and Ba have strong lines which are easy to observe in metal-poor stars and moreover, ambiguities are minimised. Also, this simple classification could be transformed to a purely empirical system, independent of uncertainties in the abundance analysis.

The criteria chosen for the  $r + s$  stars are presented in Table 9. It is natural to set the limit for  $s$ -elements as  $[Ba/Fe] > +1.0$ , see Fig. 7, but the limit for the  $r$ -elements may be debated. For simplicity we have made a choice analogous to that for the  $s$ -elements, and require  $[Eu/Fe] > +1.0$ , although there is no natural separation between the  $r + s$  stars and the  $s$  stars in this respect. However, the distribution of the stars in the  $[Ba/Eu]$  versus  $[Ce/Eu]$  and  $[La/Eu]$  versus  $[Ce/Eu]$  diagrams (see Fig. 8) suggests that there might be a natural separation of these classes of stars. As seen from the plots of Fig. 8 it is also natural to set the criteria  $[Ba/Eu] > 0.0$  for the  $r + s$  stars. Note that we do not require the  $r + s$  stars to be rich in C, although all such stars known to us have  $[C/Fe] \geq +1.7$

The definition of the class  $r + s$  only partially overlaps with the class “ $r/s$  stars” defined by Beers & Christlieb (2005). The  $r/s$  criteria is set to  $0.0 < [Ba/Eu] < +0.5$ , and thus also includes stars enhanced in neither  $r$ - nor  $s$ -elements. The  $r + s$  class, on the other hand, includes stars with  $[Ba/Eu] > +0.5$ .

As seen in Table 9, we use the same criteria for  $r$ -I and  $r$ -II stars as Beers & Christlieb (2005), but this is not the case for  $s$  stars. Stars highly enhanced in  $s$ -elements, only moderately in  $r$ -elements, and with  $[Ba/Eu] > 0.0$  are called  $s$  stars here. This definition differs from the classification of Beers & Christlieb, who define  $s$  stars as having  $[Ba/Fe] > +1.0$  and  $[Ba/Eu] > +0.5$ .

Lead stars are defined as  $[Pb/hs] = [Pb/Ba, La, Ce] > +1.0$  by Van Eck et al. (2003). According to the same principles as discussed above, we propose to use only Ba as the measure for the  $hs$ -elements, and thus advocate the classification given in Table 9. We do not require the Pb stars to also be  $r + s$  stars, only that they are enhanced in Ba and Pb, and much more in the latter element.

## 6.3. Scenarios for the formation of $r + s$ stars

The abundance patterns of stars enriched in both  $r$ - and  $s$ -elements may give important clues to the sites of nucleosynthesis for these elements. Many different scenarios have been proposed, most of them invoking two more or less independent processes, e.g. with one site for the production of the  $r$ -elements and one for the  $s$ -elements. A basic question is whether such scenarios may be retained, in view of the homogeneity of the  $r + s$  group, and its separation from the other classes of Population II stars. Another factor of significance is the frequency of  $r + s$  stars, relative to stars with more normal Population II composition. From Paper II we estimate that on the order of 1% of the Population II stars are  $r + s$  stars, though we note this may be significantly underestimated due to the bias against CH strong stars. Here, these different  $r+s$  scenarios will be discussed.

I. **Radiative levitation:** Peculiar abundance patterns may arise due to radiative levitation of elements with many absorption lines, while heavy elements with few lines may sink in the atmosphere. This is seen in some Ap stars, which display enormous enhancements of the rare-earth elements.

Cohen et al. (2003) discussed this possibility as an explanation for the  $r + s$  stars and dismissed it. The  $r + s$  stars do not show the same abundance patterns as the Ap stars. They presumably are not hot enough to produce any significant radiative levitation in their atmospheres. In particular, they have much deeper convective zones, efficiently mixing and diluting the surface material with the stellar interior. Note also that there are several  $r + s$  stars on the giant branch, where the convective zone contains most of the stellar mass.

II.  **$r$ -rich ISM and self-pollution:** The star may have been formed out of  $r$ -enriched material from e.g. an early supernova, and later self-enriched its surface with  $s$ -elements during He shell flashes in the AGB phase (as discussed by

Hill et al. 2000; Cohen et al. 2003). We note that low-mass metal-poor stars may also be affected by dredge-up at the He flash (Fujimoto et al. 2000). However, both HE 0338–3945 and HE 2148–1247 are located in the turn-off region and presumably have not passed the red-giant phase yet, so in the case of at least these  $r+s$  stars it seems possible to dismiss the hypothesis.

**III. Binary system out of  $r$ -rich ISM and AGB-pollution:** The star could have been a secondary in a binary system, formed out of  $r$ -enriched material, and later polluted with C and  $s$ -elements by mass transfer from its AGB companion (discussed by e.g. Hill et al. 2000; Cohen et al. 2003; Ivans et al. 2005). We note in passing that metal-poor stars may well produce  $s$ -elements even without any Fe seed nuclei, according to Siess et al. (2002).

The mass transfer from a companion AGB star is the scenario generally adopted for the formation of CH stars, known to be single-line spectroscopic binaries (McClure 1983, 1984; and McClure & Woodsworth 1990). It is not clear, however, if all  $r+s$  stars also are CH stars, although all 17  $r+s$  stars in Table 8 have considerably enhanced carbon abundances, as noted in Sect. 6.2. Many CH stars are defined as  $s$  stars; the  $s$  stars are often quite C rich (see Table 8), while the  $r$  stars are generally less C rich. A relatively high fraction of the CH stars seem, however, to be  $r+s$  stars (e.g. Aoki et al. 2002).

There are severe constraints set on the period and orbit for this mechanism to result in a C and  $s$ -enriched matter transfer, as the mass transfer should occur in the AGB phase where the  $s$ -elements are produced and not (only) in the RGB phase (Jorissen & Boffin 1992). The transfer across the binary has to be efficient to reach the levels of contamination seen in  $r+s$  stars. Also the right amounts of elements should be transferred. A glance in Table 8 shows that the carbon enrichment is remarkably constant among the  $r+s$  stars. The low  $^{12}\text{C}/^{13}\text{C}$  ratio of HE 0338–3945 suggests that a rather large fraction of the carbon (which presumably was produced by He burning in the primary) has been CNO-processed; N and O abundances also clearly suggest heavy processing but the dominant C relative to N and O puts constraints on this. There are obviously several steps that have to work, from the initially  $r$ -enriched material from which the star happened to form, via the formation of carbon, the formation of the  $s$ -elements in the AGB inter-shell nucleosynthesis, the CNO processing of C and the pollution of the surface of the companion with sufficient amounts of matter.

One may discuss whether this is a realistic scenario, in view of the many  $r+s$  stars discovered. In general, if a mechanism with two independent steps is in action, one producing the  $r$ -enrichment and another the  $s$ -enrichment, and the two are stochastically independent, one would expect the probability to find a star affected by both mechanisms,  $p(r,s)$  to be the product of the probabilities of finding a star subjected to only one of the two,  $p(r)$  and  $p(s)$ , respectively, i.e.  $p(r,s) = p(r) \cdot p(s)$ . Presently, in the HERES survey there are 8  $r$ -II stars, two  $s$  stars and three  $r+s$  stars, among the 253 stars without strong CH lines analysed in Paper II. The CH-strong rejection criteria may lead to a strong bias, suppressing the number of  $s$  stars, and even  $r+s$  stars. In order for a two-step mechanism to

produce 3  $r+s$  stars, however, a considerable fraction of all Population II stars should show  $s$ -star characteristics. In fact, we find from elementary statistical arguments that the probability of finding 3 or more  $r+s$  stars out of 250 would be only 0.9% if the fraction of  $s$  stars would be 0.1 of all Population II stars. Even if the total fraction of  $s$  stars were 0.3, the chance to find 3  $r+s$  stars would be only 22%. One should also note that in Paper II the  $r+s$  stars can only be detected when they are warm enough such that the CH lines are not so strong as to render the method used uncertain. Only a small fraction of the  $r$ -II stars found in the HERES survey are so warm, which suggests that the ratio of stars with  $r+s$  abundances relative to  $r$ -II stars is considerably higher than suggested by the HERES results. Thus, the probability that an  $r+s$  star is just a star which happened to become  $r$ -enriched, e.g. due to initial inhomogeneities in the ISM or a later nearby SN, and independently of that had an AGB enrichment from a companion, is negligible and should be dismissed. Still another argument against such a two-step scenario is the chemical homogeneity within the  $r+s$  group and with a clear separation from  $r$ -II stars and  $s$  stars. The scenario is also problematic as it is not commonly accepted that the ISM was sufficiently inhomogeneous (e.g. Qian & Wasserburg 2001) to contain such great  $s$ - and  $r$ -element overabundances relative to Fe at  $[\text{Fe}/\text{H}] > -3$ .

A possible modification of this scenario would be that the formation of the binary system is triggered by a supernova which also provides the  $r$ -elements (discussed by Gallino et al. 2005 and Ivans et al. 2005). Vanhala & Cameron (1998) have performed simulations of triggered star formation due to shock waves impacting molecular cloud cores, suggesting that weakly evolved cores may fragment during collapse and form low-mass binaries. Such a scenario is appealing since the  $r$ - and  $s$ -enrichments are no longer stochastically independent and thus the high frequency of  $r+s$  stars might be explained. However, it is presently unclear whether this scenario could be so prevalent as to explain the observed frequency.

Cohen et al. (2003) argue for a  $s$ -enhancement due to an AGB contamination, although they prefer scenario V below. They point out that several of the stars we now define as  $r+s$  stars are binaries (CS 29526-110, HE 2148–1247, HE 0024–2523, LP 625–44), although we cannot confirm this for HE 0338–3945, nor for several other  $r+s$  stars. Preston & Sneden (2001) report on  $r+s$  stars not exhibiting radial velocity variations exceeding  $0.5 \text{ km s}^{-1}$  over an 8 year period. This discussion may also be applied to scenarios IV, V and VI.

**IV. Triple system with SN- and AGB-pollution:** The star could have been a tertiary in a triple system, in which the primary exploded as a  $r$ -element producing supernova, and the other companion evolved into an AGB star dumping  $s$ -rich material on the least massive star.

This scenario is also discussed and dismissed by Cohen et al. (2003) as being not very plausible. The likelihood that a star would survive a close SN explosion, that the secondary would not drift away, and that the secondary could subsequently transfer processed matter onto the star at the right time does not seem very great. If such events do happen, it is not

probable that they would be common enough or sufficiently constrained to create so many  $r + s$  stars with similar abundance patterns, notably with similar relative amounts of C and of  $r$ - and  $s$ -elements.

**V. Binary system with AGB- and 1.5 SN-pollution:** The star could be a secondary in a binary system where the companion first contributed  $s$ -elements as an AGB star, and later exploded as a “Type 1.5” supernova (Zijlstra 2004; Wanajo et al. 2005), producing the  $r$ -elements. The SNe of type 1.5 are proposed to be more common among metal-poor stars than for Population I, due to the strong metallicity dependence of the mass loss during the AGB phase leading to a different initial-final-mass relation for low-metallicity stars. Thus, intermediate-mass metal-poor stars may end up with higher final masses after the AGB phase, and more easily reach the Chandrasekhar mass and explode as a type 1.5 supernova.

**VI. Binary system with AGB- and AIC-pollution:** The star could be a secondary in a binary system, in which the companion first contributed  $s$ -elements as an AGB star and then evolved to a white dwarf. Later, the primary might have undergone an accretion-induced collapse (AIC) in the white dwarf stage, triggered by mass transfer in the other direction across the system. The collapse created a neutrino wind and  $r$ -elements which were transferred across the binary and contaminated the surface of the secondary star (Qian & Wasserburg 2003, Cohen et al. 2003). Cohen et al. also noted that the system might disrupt during the AIC event, which would explain an apparently single  $r + s$  star.

This hypothesis is physically uncertain due to uncertainties in neutrino and neutron-star physics (Qian & Woosley 1996), and it is thus uncertain whether it works at all. It is also unclear how the white dwarf would accrete matter from the secondary star, which in several cases has been found not to be evolved beyond the turn-off phase (HE 0338–3945, HE 2148–1247). One may question how probable is it that the white dwarf is close enough for mass transfer, in view of the mass-loss it has experienced which increases the distance between the components.

**VII. Binary system with only AGB-pollution:** The star might be a secondary in a metal-poor binary system with an AGB companion producing  $n$ -capture elements in a hypothetical high neutron density  $s$ -process. The relatively high flux of neutrons has been proposed to not only produce “normal”  $s$ -elements such as Ba and La, but Eu which otherwise is mainly assumed to be produced in the  $r$ -process with much higher neutron densities.

This hypothesis has been discussed by Aoki et al. (2002), Cohen et al. (2003) and Johnson & Bolte (2004). Aoki et al. and Cohen et al. found some support from the calculations by Goriely & Mowlavi (2000). Cohen et al. found a ratio of Ba/Eu  $\sim 100$  for the star HE 2148–1247 (we similarly find Ba/Eu  $\sim 115$  for HE 0338–3945), and Goriely & Mowlavi predicted a surface abundance ratio of 105 for their most metal-poor ( $Z = 0.001$ ) AGB-star model with dredge up from a long

sequence of He shell flashes. This was, however, the resulting ratio from a relatively mild enhancement (0.48 dex in Ba, 0.07 in Eu) of material with initially solar Ba/Fe and Eu/Fe ratios. The dredge-up material in the model of Goriely & Mowlavi has a Ba/Eu ratio of about 500. Anyhow, Cohen et al. dismissed the scenario since they found the predicted absolute abundances of the produced elements to be quite insufficient to explain the observations. Johnson & Bolte, on the other hand, found evidence in the  $r$ -element ratios (Eu/Tb, Eu/Dy, Th/Eu) which seem difficult to explain with any combination of the normal  $r$ - and  $s$ -processes. They suggested that another form of  $s$ -process is at work, and found some support for this in parametrised calculations made by Malaney (1987). In Malaney’s Table II, yields of the  $s$ -process acting at the unusually high neutron density of  $10^{12} \text{ cm}^{-3}$  are presented, which reduced the Ba/Eu ratios to 150–200. Adopting a neutron exposure of  $\tau_0 = 0.05$  Johnson & Bolte found the predictions of Malaney to agree reasonably well with the observed La/Eu ratio for the  $r + s$  star CS 31062-050, and a generally good overall fit of the abundance pattern of that star from La to Hf. They noted, however, that the observations of the  $^{151}\text{Eu}/^{153}\text{Eu}$  ratio of this star by Aoki et al. (2003) seem to indicate that the neutron density was rather around the more normal value of  $10^8$ .

In order to study the consequences of a high neutron-density  $s$ -process we set up a small program to calculate  $s$ -process yields in the interval from La to Gd, simultaneously solving the rate equations for 88 nuclides in this interval. The  $n$ -capture cross sections were adopted from Bao et al. (2000) with some modifications such as the new data from Best et al. (2001). The  $\beta$ -decay rates and the electron-capture rates were from Takahashi & Yokoi (1987). In stationary-flow calculations we found the abundances of Eu relative to La, Ce, Pr, Nd, Sm and Gd to stay significantly below the observed values and vary rather little when the neutron density increased from  $10^8$  to  $10^{14} \text{ cm}^{-3}$ , with some tendencies to peak around  $10^{12}$ ; only Eu/Sm and Eu/Pr came then close to the observed values. For the time-dependent case we varied neutron exposures and temperatures, but again in this case we could not recover the high observed Eu/ $X$  ratios for  $X = \text{La}, \text{Nd}, \text{Ce}$ , or Gd with any combination of parameters tried. Although these results are preliminary, and a number of nuclear rates are still rather uncertain, we find it less probable that a high-neutron density  $s$ -process would be responsible for the characteristics of the  $r + s$  stars.

**VIII. A relation to blue stragglers?** The fraction of  $r + s$  stars in Table 8 that have  $T_{\text{eff}} > 6000 \text{ K}$  is fairly high, as compared with the fraction of hot  $s$  stars and  $r$ -II stars. This may reflect small-number statistics or selection effects – it is certainly more difficult to detect an  $r$ -II star than a  $r + s$  star at the turn-off point (TOP) due to the fact that the former, at least in our sample, tend to have both lower metallicities and lower [Eu/Fe]. Furthermore, since  $r + s$  stars typically have [C/Fe]  $> 2.0$  (see Table 8), the cooler  $r + s$  might have been rejected from the sample of Paper II due to their strong CH lines. The dominance of detections of  $r + s$  stars at the TOP might also point at some relation between the  $r + s$  stars and the blue stragglers in globular

clusters and in the field. These objects, lying above and blue-ward of the turn-off in the colour-magnitude diagram of clusters, are generally believed to be the result of close-binary evolution with mass transfer, of stellar collisions or of mixing of the stellar interior (for a review, see, e.g. Trimble 1993). Sneden et al. (2003b) recently analysed spectra of six field blue stragglers with  $[\text{Fe}/\text{H}] < -2$ . Three of them were spectroscopic binaries and were found to have enhanced carbon and Sr and Ba. This was interpreted as the result of mass transfer from companions in the AGB stage. One of these stars was remarkably rich in Pb and had in fact a high, though uncertain, Eu abundance – that star is CS 29497-030 and is included as an  $r + s$  star in our Table 8. The idea that sub-giant CH stars may be related to blue stragglers was already put forward by Luck & Bond (1991). However, in order for this scenario to apply to the  $r + s$  stars, and not only as a provider of the  $s$ -elements, an idea which is already included in the scenarios III–VI above, one has to invoke a production mechanism for the  $r$ -elements. One might hope that the alternative origin of blue stragglers as the result of stellar collisions might provide not only overall mixing of CNO and He (e.g. Sills et al. 2005) but possibly also heavy neutron fluxes at head-on collisions, leading to formation of Eu and other  $r$ -elements. It is, however, highly uncertain whether such a speculative mechanism, presumably acting in the centre of dense globular clusters, could contribute enough  $r + s$  stars in the field.

**IX. Origin from a common cloud of  $r+s$  stars:** In view of the similarity in chemical composition of the  $r + s$  stars it is not inconceivable that they were formed out of gas in a common cloud, and even were accreted by the Galaxy at a later stage. To explore this hypothesis we have calculated space velocities for the stars. In the absence of accurate distances for but one star (CS 31062-012), we used photometric distances estimated by reading off the colour–magnitude diagram of the metal-poor globular cluster M 92 (Ruelas-Mayorga & Sánchez 2005) at the appropriate reddening-corrected  $B - V$ . Alternatively, we have compared to theoretical isochrones of Kim et al. (2002) with  $[\text{Fe}/\text{H}] = -2.5$  and an age of 12 Gyr in the  $T_{\text{eff}}-\log g$  diagram and derived the absolute magnitudes from that. For the four stars with well known kinematic data we find space velocities with a spread of approximately  $100 \text{ km s}^{-1}$  in the  $U$ ,  $V$  and  $W$  components, and conclude from this that there is no suggestion of a common space motion.

## 7. Conclusions

In this paper we have analysed a high-quality spectrum of the  $r$ - and  $s$ -element rich star HE 0338–3945, deriving abundances for 33 elements and upper limits for an additional 6 elements. We found the abundances of this star to be very similar to those of HE 2148–1247, which is supported by an homogeneous analysis of the two spectra. In fact the abundance patterns among known  $r + s$  stars in the literature seem to be quite similar considering the differences between analyses. Based on our results for HE 0338–3945 and our survey of other  $r + s$  stars, we have discussed a range of scenarios for the formation of the

$r + s$  stars. Some scenarios have considerable merit, but all have also drawbacks or problems. Scenario I is already dismissed. Scenarios II and III need the ISM to be highly inhomogeneous to produce such high  $r$ -process abundances as detected in these stars. We note that the [Eu/Fe] ratios for the  $r + s$  stars are systematically higher than those of the  $r$ -II stars which might suggest that a process with two independent steps, one of which is also responsible for the  $r$ -II stars is not very probable. However, this is uncertain due to the possibly significant contribution of Eu by the  $s$ -process, and the possible relative effects of convective mixing between the  $r + s$  stars, which are predominantly TOP stars, and the  $r$ -II stars, which are all giants (see also discussion below). Scenario II is also improbable as it requires all  $r + s$  stars to have passed at least the RGB phase. Scenario III presently seems unlikely in view of the frequency of  $r + s$  stars and their apparent chemical homogeneity. The discussed modification of scenario III to include supernova-triggered binary formation seems plausible, though it is presently unclear whether such events can provide the observed frequency of  $r + s$  stars. Scenario IV is improbable as it requires strict stellar orbit constraints to produce the chemical homogeneity of the  $r + s$  class. Scenarios V and VI may be considered, but as they depend heavily on physical processes and parameters that are poorly understood at present, these scenarios at best have to be regarded as uncertain. The observed chemical uniformity is not an obvious consequence of them. Scenario VII still needs a site which can produce the high neutron densities at suitable exposures; the scenario also has to be supported by renewed and more realistic  $s$ -process calculations. It is also not clear how well this scenario would explain the chemical homogeneity. A probable merit of it is, however, that the observed great but varying amounts of Pb may result naturally from the high neutron fluxes. The absence of observed radial velocity variations for several stars (see Preston & Sneden 2001; and Aoki et al. 2002) is a problem, although not very severe as yet, for all scenarios III–VIII. Scenario VIII seems hypothetical and it is also questionable whether it is efficient enough. The available astrometric data, though admittedly meager, does not support Scenario IX.

The majority of the  $r + s$  stars in Table 8 are TOP stars, while all the  $r$ -II stars are giants. One should note the great difference between the TOP stars and the red giants as regards two important aspects: first, the convective zone of the TOP stars is less than 5% of the stellar mass, while it is typically 75% of the mass for the giants. Thus, the dilution of enriched material transferred to the stellar surface of a TOP star is at least a factor of 10 less than for a giant. So, if the enhancements of  $n$ -capture elements are due to mass transfer from a companion, the amount transferred has to be at least one order of magnitude greater for the  $r$ -II stars than for most of the TOP stars. Second, the space volume surveyed in apparent-magnitude limited surveys is at least a factor of 200 greater for the giants than for the  $r + s$  stars. This suggests that the space density of  $r + s$  stars is considerably higher than that of  $r$ -II stars. In spite of their scarcity, it is interesting that no  $r$ -II stars have been found at the TOP, since a much smaller transfer of  $r$ -elements would raise the atmospheric abundances to considerable values. More statistically controlled surveys are desirable.

Future detailed studies, with accurate analysis of all known and suspected stars, are worthwhile. Spectra with high  $S/N$  and high resolution, also in the UV, are important to obtain better limits on Ag, Th and U and to get results for  $r$ -elements that have lines in the UV such as Os and Ir. Attempts to measure Eu isotopic ratios for these stars should be continued. The analysis of certain important elements, such as Eu and Ag, should be scrutinised for NLTE and 3D effects. More detailed calculations of the  $s$ -process at high neutron densities should be carried out. Finally, all known  $r + s$  stars should be monitored in an attempt to disclose their possible binarity.

**Acknowledgements.** We thank Nikolai Piskunov for his invaluable help during the reductions, and Oleg Kochukhov for his kind help with merging the spectra. We are grateful to Kjell Eriksson for computing MARCS models, as well as certain checks of the effective temperature. Nils Ryde and Rurik Wahlin are thanked for providing molecular line lists. We are grateful to Judith Cohen for providing us with her spectrum of HE 2148–1247. Roberto Gallino and Sara Bisterzo are thanked for valuable discussions. We made use of the NASA ADS and VALD databases.

The Uppsala group acknowledges the support of the Swedish Research Council (VR). N.C. acknowledges the support of Deutsche Forschungsgemeinschaft (grant Ch 214/3-1). T.C.B. acknowledges partial support from grants AST 04-06784, and PHY 02-16783, Physics Frontier Centers/JINA: Joint Institute for Nuclear Astrophysics, awarded by the US National Science Foundation.

## References

- Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Aoki, W., Norris, J. E., Ryan, S. G., et al. 2000, *ApJ*, 536, L97
- Aoki, W., Ryan, S. G., Norris, J. E., et al. 2001, *ApJ*, 561, 346
- Aoki, W., Ryan, S. G., Norris, J. E., et al. 2002, *ApJ*, 580, 1149
- Aoki, W., Ryan, S. G., Iwamoto, N., et al. 2003, *ApJ*, 592, L67
- Aoki, W., Honda, S., Beers, T. C., et al. 2005, *ApJ*, 632, 611
- Arlandini, C., Käppeler, F., Wissak, K., et al. 1999, *ApJ*, 525, 886
- Asplund, M. 2004, *Mem. Soc. Astron. It.*, 75, 300
- Asplund, M. 2005, *ARA&A*, 43, 481
- Asplund, M., & García Pérez, A. E. 2001, *A&A*, 372, 601
- Bao, Z. Y., Beer, H., Käppeler, F., et al. 2000, *At. Data Nucl. Data Tables*, 76, 70
- Barbuy, B., Cayrel, C., Spite, M., et al. 1997, *A&A*, 317, L63
- Barbuy, B., Spite, M., Spite, F., et al. 2005, *A&A*, 429, 1031
- Bard, A., Kock, A., & Kock, M. 1991, *A&A*, 248, 315
- Barklem, P. S., Christlieb, N., Beers, T. C., et al. 2005, *A&A*, 439, 129
- Barklem, P. S., Stempels, H. C., Allende Prieto, C., et al. 2002, *A&A*, 385, 951
- Baumüller, D., & Gehren, T. 1997, *A&A*, 325, 1088
- Baumüller, D., Butler, K., & Gehren, T. 1998, *A&A*, 338, 637
- Beers, T. C., & Christlieb, N. 2005, *ARA&A*, 43, 531
- Beers, T. C., Preston, G. W., & Shectman, S. A. 1992, *AJ*, 103, 1987
- Best, J., Stoll, H., Arlandini, C., et al. 2001, *Phys. Rev. C*, 64, 015801
- Biémont, E., & Lowe, R. M. 1993, *A&A*, 273, 665
- Biémont, E., Baudoux, M., Kurucz, R. L., et al. 1991, *A&A*, 249, 539
- Biémont, E., Dutrieux, J.-F., Martin, I., et al. 1998, *J. Phys. B*, 31, 3321
- Biémont, E., Garnir, H. P., Palmeri, P., et al. 2000, *MNRAS*, 312, 116
- Biémont, E., Grevesse, N., Hannaford, P., et al. 1981, *ApJ*, 248, 867
- Biémont, E., Grevesse, N., Hannaford, P., et al. 1989, *ApJ*, 222, 307
- Bihain, G., Israelian, G., Rebolo, R., et al. 2004, *A&A*, 473, 777
- Blackwell, D. E., Booth, A. J., Petford, A. D., et al. 1989, *MNRAS*, 236, 235
- Blackwell, D. E., Menon, S. L., Petford, A. D., et al. 1982a, *MNRAS*, 201, 611
- Blackwell, D. E., Petford, A. D., Shallis, M. J., et al. 1982b, *MNRAS*, 199, 21
- Booth, A. J., Blackwell, D. E., Petford, A. D., et al. 1984, *MNRAS*, 208, 147
- Bridges, J. M., & Kornblith, R. L. 1974, *ApJ*, 192, 793
- Burstein, D., & Heiles, C. 1982, *AJ*, 87, 1165
- Burris, D. L., Pilachowski, C. A., Armandroff, T. E., et al. 2000, *ApJ*, 544, 302
- Busso, M., Gallino, R., & Wasserburg, G. J. 1999, *ARA&A*, 37, 239
- Cardon, B. L., Smith, P. L., Scalo, J. M., et al. 1982, *ApJ*, 260, 395
- Cayrel, R., Depagne, E., Spite, M., et al. 2004, *A&A*, 416, 1117
- Christlieb, N. 2003, *Rev. Mod. Astron.*, 16, 191
- Christlieb, N., Beers, T. C., Barklem, P. S., et al. 2004, *A&A*, 428, 1027
- Cohen, J. G., Christlieb, N., Qian, Y.-Z., et al. 2003, *ApJ*, 588, 1082
- Collet, R., Asplund, M., & Thévenin, F. 2005, *A&A*, 442, 643
- Corliss, C. H., & Bozman, W. R. 1962, *Experimental Transitions Probabilities for Spectral lines of Seventy Elements*, NBS Monographs 32 (Washington: GPO)
- Cowan, J. J., Sneden, C., Burles, S., et al. 2002, *ApJ*, 572, 861
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, 2MASS All Sky Catalog of Point Sources, VizieR Online Data Catalog, 2246
- Den Hartog, E. A., Lawler, J. E., Sneden, C., et al. 2003, *ApJS*, 148, 543
- Fedchak, J. A., Den Hartog, E. A., Lawler, J. E., et al. 2000, *ApJ*, 542, 1109
- Fuhr, J. R., & Wiese, W. L. 1996, *NIST Atomic Transition Probability Tables*, in *Chemical Rubber Company Handbook of Chemistry and Physics*, ed. D. R. Lide (Boca Raton: CRC), 10
- Fuhr, J. R., Martin, G. A., & Wiese, W. L. 1988, *J. Phys. Chem. Ref. Dat.* 17, Suppl. 4
- Fujimoto, M. Y., Ikeda, Y., & Iben, I. 2000, *ApJ*, 529, L25
- Fulbright, J. 2000, *AJ*, 120, 1841
- Gallagher, A. 1967, *Phys. Rev.*, 157, 24
- Gallino, R., Arlandini, C., Busso, M., et al. 1998, *ApJ*, 497, 388
- Gallino, R., Wasserburg, G. J., Busso, M., & Straniero, O. 2005, in *From Lithium to Uranium: Elemental Tracers of Early Cosmic Evolution*, ed. V. Hill, P. Francois, & F. Primas, Proc. IAU Symp., 228, in press
- Gehren, T., Liang, Y. C., Shi, J. R., et al. 2004, *A&A*, 413, 1045
- Goriely, S., & Mowlavi, N. 2000, *A&A*, 362, 599
- Gratton, R. G., & Sneden, C. 1994, *A&A*, 287, 927
- Grevesse, N., Blackwell, D. E., & Petford, A. D. 1989, *A&A*, 208, 157
- Griffin, R., Gustafsson, B., Vieira, T., et al. 1982, *MNRAS*, 198, 637
- Hannaford, P., Lowe, R. M., Grevesse, N., et al. 1982, *ApJ*, 261, 736
- Heise, C., & Kock, M. 1990, *A&A*, 230, 244
- Hill, V., Barbuy, B., Spite, M., et al. 2000, *A&A*, 353, 557
- Hill, V., Plez, B., Cayrel, R., et al. 2002, *A&A*, 387, 560
- Holt, R. A., Scholl, T. J., & Rosner, S. D. 1999, *MNRAS*, 306, 107
- Honda, S., Aoki, W., Kajino, T., et al. 2004a, *ApJ*, 607, 474
- Honda, S., Aoki, W., Ando, H., et al. 2004b, *ApJS*, 152, 113
- Huber, M. C. E., & Sandeman, R. J. 1980, *A&A*, 86, 95
- Irwin, A. W. 1981, *ApJS*, 45, 621
- Ivans, I. I., Sneden, C., Gallino, R., et al. 2005, *ApJ*, 627, L145
- Ivarsson, S., Litzén, U., & Wahlgren, G. M. 2001, *Phys. Scr.*, 64, 455
- Johnson, J. A. 2002, *ApJS*, 139, 219
- Johnson, J. A., & Bolte, M. 2002, *ApJ*, 579, L87

- Johnson, J. A., & Bolte, M. 2004, *ApJ*, 605, 462
- Jonsell, K., Edvardsson, B., Gustafsson, B., et al. 2005, *A&A*, 440, 321
- Jørgensen, U. G., Larsson, M., Iwamae, A., et al. 1996, *A&A*, 315, 204
- Jorissen, A., & Boffin, H. M. J. 1992, in *Binaries as Tracers of Star Formation*, ed. A. Duquennoy, & M. Mayor (Cambridge University Press), 110
- Kim, Y.-C., Demarque, P., Yi, S. K., et al. 2002, *ApJS*, 143, 499
- Kling, R., & Griesmann, U. 2000, *ApJ*, 531, 1173
- Kroll, S., & Kock, M. 1987, *A&AS*, 67, 225
- Korn, A. J., Shi, J., & Gehren, T. 2003, *A&A*, 407, 691
- Kupka, F., Piskunov, N. E., Ryabchikova, T. A., et al. 1999, *A&AS*, 138, 119
- Kurucz, R. L. 1995, CD-ROMs, Cambridge, SAO
- Lawler, J. E., & Dakin, J. T. 1989, *J. Opt. Soc. Am. B*, 6, 1457
- Lawler, J. E., Bonvallet, G., & Sneden, C. 2001a, *ApJ*, 556, 452
- Lawler, J. E., Wickliffe, M. E., Cowley, C. R., et al. 2001b, *ApJS*, 137, 341
- Lawler, J. E., Wickliffe, M. E., Den Hartog, E. A., et al. 2001c, *ApJ*, 563, 1075
- Lawler, J. E., Wyart, J.-F., & Blaise, J. 2001d, *ApJS*, 137, 315
- Lefèvre, P. H., Garnir, H. P., & Biémont, E. 2003, *A&A*, 404, 1153
- Lucatello, S., Gratton, R., Cohen, J. G., et al. 2003, *AJ*, 125, 875
- Lucatello, S., Tsangarides, S., Beers, T. C., et al. 2005, *ApJ*, 625, 825
- Luck, R. E., & Bond, H. E. 1991, *ApJS*, 77, 515
- Malaney, R. A. 1987, *Ap&SS*, 137, 251
- Manning, T. E., Anderson, C. E., & Watson, W. W. 1950, *Phys. Rev.*, 78, 417
- Mårtensson-Pendrill, A.-M., Gough, D. S., & Hannaford, P. 1994, *Phys. Rev. A*, 49, 3351
- Martin, G. A., Fuhr, J. R., & Wiese, W. L. 1988, *J. Phys. Chem. Ref. Dat.* 17, Suppl. 3
- McClure, R. D. 1983, *ApJ*, 268, 264
- McClure, R. D. 1984, *ApJ*, 280, L31
- McClure, R. D., & Woodsworth, A. W. 1990, *ApJ*, 352, 709
- McWilliam, A. 1998, *ApJ*, 115, 1640
- McWilliam, A., Preston, G. W., Sneden, C., et al. 1995, *AJ*, 109, 2757
- Migdalek, J., & Baylis, W. E. 1987, *Can. J. Phys.*, 65, 1612
- Mishenina, T. V., Kovtyukh, V. V., Soubiran, C., et al. 2002, *A&A*, 396, 189
- Moity, J. 1983, *A&AS*, 52, 37
- Musiol, K., & Labuz, S. 1983, *Phys. Scr.* 27, 422
- Nilsson, H., Ivarsson, S., Johansson, S., et al. 2002a, *A&A*, 381, 1090
- Nilsson, H., Zhang, Z., Lundberg, H., et al. 2002b, *A&A*, 382, 368
- Nissen, P. E., Primas, F., Asplund, M., et al. 2002, *A&A*, 390, 235
- Nitz, D. E., Kunau, A. E., Wilson, K. L., et al. 1999, *ApJS*, 122, 557
- O'Brian, T. R., Wickliffe, M. E., Lawler, J. E., et al. 1991, *J. Opt. Soc. Am. B*, 8, 1185
- Palmeri, P., Quinet, P., Wyart, J.-F., et al. 2000, *Phys. Scr.* 61, 323
- Pickering, J. C. 1996, *ApJS*, 107, 811
- Pickering, J. C., Thorne, A. P., & Perez, R. 2001, *ApJS*, 132, 403
- Pinnington, E. H., Berends, R. W., & Lumsden, M. 1995, *J. Phys. B*, 28, 2095
- Pinnington, E. H., Rieger, G., & Kernahan, J. A. 1997, *Phys. Rev. A*, 56, 2421
- Piskunov, N. E., & Valenti, J. A. 2002, *A&A*, 385, 1095
- Plez, B., Hill, V., Cayrel, R., et al. 2004, *A&A*, 428, L9
- Prantzos, H., Hashimoto, M., & Nomoto, K. 1990, *A&A*, 234, 211
- Preston, G. W., & Sneden, C. 2001, *AJ*, 122, 1545
- Qian, Y.-Z., & Woosley 1996, *ApJ*, 471, 331
- Qian, Y.-Z., & Wasserburg, G. J. 2001, *ApJ*, 552, L55
- Qian, Y.-Z., & Wasserburg, G. J. 2003, *ApJ*, 588, 1099
- Quinet, P., Palmeri, P., & Biémont, E., et al. 1999, *MNRAS*, 307, 934
- Ruelas-Mayorga, A., & Sánchez, L. J. 2005, *Rev. Mex.. A&A*, 41, 507
- Ryabchikova, T. A., Hill, G. M., Landstreet, J. D., et al. 1994, *MNRAS*, 267, 697
- Ryan, S. G., Norris, J. E., & Beers, T. C. 1996, *ApJ*, 471, 254
- Ross, J. E., & Aller, L. H. 1972, *Sol. Phys.*, 25, 30
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schnabel, R., Schultz-Johanning, M., & Kock, M. 2004, *A&A*, 414, 1169
- Siess, L., Livio, M., & Lattanzio, J. 2002, *ApJ*, 570, 329
- Sills, A., Adams, T., & Davies, M. B. 2005, *MNRAS*, in press
- Simmerer, J., Sneden, C., Cowan, J. J., et al. 2004, *ApJ*, 617, 1091
- Sivarani, T., Bonifacio, P., Molaro, P., et al. 2004, *A&A*, 413, 1073
- Smith, G. 1981, *A&A*, 103, 351
- Sneden, C. 1973, *ApJ*, 184, 839
- Sneden, C., McWilliam, A., Preston, G. W., et al. 1996, *ApJ*, 467, 819
- Sneden, C., Cowan, J. J., Ivans, I. I., et al. 2000, *ApJ*, 533, L139
- Sneden, C., Cowan, J. J., Lawler, J. E., et al. 2003a, *ApJ*, 591, 936
- Sneden, C., Preston, G. W., & Cowan, J. J. 2003b, *ApJ*, 592, 504
- Takahashi, K., & Yokoi, K. 1987, *At. Data Nucl. Data Tables*, 36, 375
- Trimble, V. 1993, in *Blue Stragglers*, ed. R. Saffer (SanFrancisco: ASP), *ASP Conf. Ser.*, 53, 155
- Valenti, J. A., & Piskunov, N. 1996, *A&AS*, 118, 595
- Van Eck, S., Goriely, S., Jorissen, A., et al. 2001, *Nature*, 412, 793
- Van Eck, S., Goriely, S., Jorissen, A., et al. 2003, *A&A*, 404, 291
- Vanhala, H. A. T., & Cameron, A. G. W. 1998, *ApJ*, 508, 291
- Villemoes, P., Arnesen, A., Heijkenskjold, F., et al. 1993, *J. Phys. B*, 26, 4289
- Wanajo, S., Nomoto, K., Iwamoto, N., et al. 2005, *ApJ*, in press
- Westin, J., Sneden, C., Gustafsson, B., et al. 2000, *ApJ*, 530, 783
- Wickliffe, M. E., Lawler, J. E., & Nave, G. 2000, *J. Quant. Spec. Radiat. Transf.*, 66, 363
- Wiese, W. L., & Martin, G. A. 1980, *Wavelengths and Transition Probabilities for Atoms and Atomic Ions*, NSRDS-NBS No. 68, Part II
- Wisotzki, L., Christlieb, N., Bade, N., et al. 2000, *A&A*, 358, 77
- Zacs, L., Nissen, P. E., & Schuster, W. J. 1998, *A&A*, 337, 216
- Zijlstra, A. 2004, *MNRAS*, 348, L23

# **Online Material**

**Table 2.** Line data and abundances from an analysis of the r- and s-process enhanced star HE0338-3945. Each column gives: **1)** Definition of the line: d=dummy line for defining the wavelength and region, h=hyperfine structure (hfs) line, i=isotopic splitting (isot) line, b=blending line. **2)** The element or molecule producing the line, including isotopic information if necessary. **3)** Ionic stage: 1=neutral, 2=singly ionised. **4)**  $\lambda$ , the central air wavelength in Ångströms. **5)**  $\chi_{\text{lower}}$ , the lower transition level excitation energy. **6)**  $\log g_f$ , the oscillator strength multiplied with the statistical weight, all in logarithm. **7)**  $\log(\Gamma_{\text{vdW}}^{\text{dW}} / N_H)$ , the van der Waals damping constant. If the value is < 0 then it is the log of the FWHM per perturber at 10000 K in the reported units. If the value is > 0 then the data are from the broadening theory of Anstee, Barklem & O'Mara. The broadening cross section ( $\sigma$ ) for a collision speed 1000 m/s, and the velocity parameter ( $\alpha$ ) are packed into a single parameter given by  $\text{INT}(\sigma) + \alpha$ , such that  $\text{INT}(\sigma)$  is the nearest integer to  $\sigma$ . For example, if the value is 207.345, then  $\sigma = 207$  au. and  $\alpha = 0.345$ . The broadening cross section is in atomic units ( $2.80 \times 10^{-21} \text{ m}^2$ ). The velocity parameter is dimensionless. See Anstee & O'Mara (1995; MNRAS 276, 859) for details of how to compute the line width. **8)**  $\log(\Gamma_{10000 \text{ K}}^{\text{Stark}} / N_e)$ , the Stark broadening constant, the log of the FWHM per perturber at 10000K. **9)**  $\log(\Gamma^{\text{rad}})$ , Radiation damping constant, the log of the FWHM. **10)**  $\delta\lambda_{\text{blue}}$ , the range from central wavelength to blue end of spectral window, in milliÅngström. The spectral window employed for the line is  $\lambda + \delta\lambda_{\text{blue}}$  to  $\lambda + \delta\lambda_{\text{red}}$ . **11)**  $\delta\lambda_{\text{red}}$ , the range from central wavelength to red end of spectral window, in milliÅngström. The spectral window employed for the line is  $\lambda + \delta\lambda_{\text{blue}}$  to  $\lambda + \delta\lambda_{\text{red}}$ . **12)** References and notes for the line data.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log g_f$	$\log(\Gamma_{\text{vdW}}^{\text{dW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma_{10000 \text{ K}}^{\text{Stark}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
d	CH	1	3638.6169	0.002	-2.551				-150.	150.	Jørgensen et al. (1996)
d	CH	1	3661.4150	0.051	-2.146				-165.	165.	Jørgensen et al. (1996)
d	CH	1	4310.1299	0.000	-9.999				-100.	2300.	Dummy line. Region 4310.030 - 4312.430
	CH	1	4310.0381	0.096	-3.183						Hill et al. (2002)
	CH	1	4310.0898	0.096	-1.412						Hill et al. (2002)
	CH	1	4310.1099	0.096	-1.474						Hill et al. (2002)
	CH	1	4310.1489	0.431	-3.202						Hill et al. (2002)
	CH	1	4310.1621	0.096	-3.025						Hill et al. (2002)
	CH	1	4310.2031	0.431	-1.443						Hill et al. (2002)
	CH	1	4310.2202	0.431	-1.505						Hill et al. (2002)
	CH	1	4310.2720	0.431	-3.039						Hill et al. (2002)
	CH	1	4310.4038	0.096	-3.183						Hill et al. (2002)
	CH	1	4310.4302	0.408	-1.581						Hill et al. (2002)
	CH	1	4310.4580	0.096	-1.412						Hill et al. (2002)
	CH	1	4310.5039	0.432	-3.202						Hill et al. (2002)
	CH	1	4310.5078	0.408	-2.966						Hill et al. (2002)
	CH	1	4310.5562	0.432	-1.443						Hill et al. (2002)
	CH	1	4310.6792	0.072	-1.550						Hill et al. (2002)
	CH	1	4310.7290	0.408	-1.581						Hill et al. (2002)
	CH	1	4310.7568	0.072	-2.952						Hill et al. (2002)
	CH	1	4310.8071	0.408	-2.966						Hill et al. (2002)
	CH	1	4310.8110	0.408	-3.155						Hill et al. (2002)
	CH	1	4310.8892	0.408	-1.508						Hill et al. (2002)
	CH	1	4310.9370	0.388	-1.677						Hill et al. (2002)
	CH	1	4310.9692	1.107	-2.269						Hill et al. (2002)
	CH	1	4310.9912	0.072	-1.550						Hill et al. (2002)
	CH	1	4311.0449	0.388	-2.879						Hill et al. (2002)
	CH	1	4311.0688	0.072	-2.952						Hill et al. (2002)
	CH	1	4311.0752	0.072	-3.136						Hill et al. (2002)
	CH	1	4311.0752	0.408	-3.155						Hill et al. (2002)
	CH	1	4311.0840	0.987	-4.123						Hill et al. (2002)
	CH	1	4311.1450	0.987	-4.123						Hill et al. (2002)
	CH	1	4311.1528	0.072	-1.477						Hill et al. (2002)
	CH	1	4311.1528	0.388	-1.677						Hill et al. (2002)
	CH	1	4311.1528	0.408	-1.508						Hill et al. (2002)
	CH	1	4311.1958	1.108	-2.229						Hill et al. (2002)
	CH	1	4311.1958	1.108	-3.922						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Starck}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
CH	1	4311.2612	0.388	-2.879							Hill et al. (2002)
CH	1	4311.3281	0.371	-1.805							Hill et al. (2002)
CH	1	4311.3481	0.072	-3.136							Hill et al. (2002)
CH	1	4311.3940	0.388	-3.105							Hill et al. (2002)
CH	1	4311.4238	0.072	-1.477							Hill et al. (2002)
CH	1	4311.4731	0.371	-1.805							Hill et al. (2002)
CH	1	4311.4761	0.371	-2.771							Hill et al. (2002)
CH	1	4311.5020	0.051	-1.646							Hill et al. (2002)
CH	1	4311.5020	0.388	-1.587							Hill et al. (2002)
CH	1	4311.5449	1.108	-2.269							Hill et al. (2002)
CH	1	4311.5469	0.347	-2.386							Hill et al. (2002)
CH	1	4311.5669	0.357	-2.002							Hill et al. (2002)
CH	1	4311.5801	0.388	-3.105							Hill et al. (2002)
CH	1	4311.5918	0.347	-2.386							Hill et al. (2002)
CH	1	4311.6118	0.051	-2.865							Hill et al. (2002)
CH	1	4311.6182	0.662	-2.054							Hill et al. (2002)
CH	1	4311.6182	0.662	-2.054							Hill et al. (2002)
CH	1	4311.6211	0.371	-2.771							Hill et al. (2002)
CH	1	4311.6548	0.357	-2.002							Hill et al. (2002)
CH	1	4311.6880	0.388	-1.587							Hill et al. (2002)
CH	1	4311.7222	1.108	-3.923							Hill et al. (2002)
CH	1	4311.7271	0.051	-1.646							Hill et al. (2002)
CH	1	4311.7290	1.108	-2.229							Hill et al. (2002)
CH	1	4311.7759	0.357	-2.632							Hill et al. (2002)
CH	1	4311.8369	0.051	-2.866							Hill et al. (2002)
CH	1	4311.8608	0.347	-2.432							Hill et al. (2002)
CH	1	4311.8628	0.357	-2.632							Hill et al. (2002)
CH	1	4311.8970	0.371	-3.052							Hill et al. (2002)
CH	1	4311.9058	0.347	-2.432							Hill et al. (2002)
CH	1	4311.9780	0.051	-3.085							Hill et al. (2002)
CH	1	4312.0171	0.371	-3.052							Hill et al. (2002)
CH	1	4312.0449	0.371	-1.688							Hill et al. (2002)
CH	1	4312.0879	0.051	-1.557							Hill et al. (2002)
CH	1	4312.3042	0.033	-1.773							Hill et al. (2002)
CH	1	4312.3042	0.033	-2.759							Hill et al. (2002)
CH	1	4312.3169	0.358	-3.007							Hill et al. (2002)
CH	1	4312.3862	0.358	-3.007							Hill et al. (2002)
CH	1	4312.4561	0.033	-2.759							Hill et al. (2002)
CH	1	4312.5269	0.358	-1.831							Hill et al. (2002)
b	CH	1	4312.5942	0.019	-1.969						Hill et al. (2002)
b	CH	1	4312.5942	0.358	-1.831						Hill et al. (2002)
b	CH	1	4313.6299	0.000	-9.999						Hill et al. (2002)
d	CH	1	4313.6299	0.020	-2.985						Dummy line. Region 4313.430 - 4313.830
b	CH	1	4313.3770	0.020	-2.985						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{He}\alpha}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{He}\alpha}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
	CH	1	4313.4458	0.020	-2.985						Hill et al. (2002)
	CH	1	4313.5898	0.020	-1.800						Hill et al. (2002)
	CH	1	4313.6602	0.020	-1.800						Hill et al. (2002)
b		CH	1	4313.8760	0.009	-3.033					Hill et al. (2002)
b		CH	1	4313.9058	0.009	-3.033					Hill et al. (2002)
d		CH	1	4363.2998	0.000	-9.999					Hill et al. (2002)
b		CH	1	4362.0210	1.247	-4.054					Hill et al. (2002)
b		CH	1	4362.0610	1.247	-1.976					Hill et al. (2002)
b		CH	1	4362.1719	1.247	-1.943					Hill et al. (2002)
b		CH	1	4362.2021	0.777	-1.982					Hill et al. (2002)
b		CH	1	4362.2041	1.247	-4.149					Hill et al. (2002)
b		CH	1	4362.2549	0.777	-3.284					Hill et al. (2002)
	CH	1	4362.5308	0.777	-1.917						Hill et al. (2002)
	CH	1	4362.5488	0.557	-1.749						Hill et al. (2002)
	CH	1	4362.5508	0.557	-3.392						Hill et al. (2002)
	CH	1	4362.6968	0.777	-1.982						Hill et al. (2002)
	CH	1	4362.7490	0.557	-1.705						Hill et al. (2002)
	CH	1	4362.7500	0.777	-3.285						Hill et al. (2002)
	CH	1	4362.9849	0.777	-1.917						Hill et al. (2002)
	CH	1	4363.0859	0.227	-3.380						Hill et al. (2002)
	CH	1	4363.0869	0.227	-1.716						Hill et al. (2002)
	CH	1	4363.1621	1.248	-4.055						Hill et al. (2002)
	CH	1	4363.1860	1.248	-1.976						Hill et al. (2002)
	CH	1	4363.2422	1.248	-1.943						Hill et al. (2002)
	CH	1	4363.2759	1.248	-4.149						Hill et al. (2002)
	CH	1	4363.2910	0.227	-1.673						Hill et al. (2002)
	CH	1	4363.4609	0.558	-1.749						Hill et al. (2002)
	CH	1	4363.4629	0.558	-3.392						Hill et al. (2002)
	CH	1	4363.6050	0.558	-1.705						Hill et al. (2002)
	CH	1	4364.0342	0.228	-3.380						Hill et al. (2002)
	CH	1	4364.0361	0.228	-1.716						Hill et al. (2002)
	CH	1	4364.1138	1.289	-4.093						Hill et al. (2002)
	CH	1	4364.1670	1.289	-1.947						Hill et al. (2002)
	CH	1	4364.1812	0.228	-1.673						Hill et al. (2002)
	CH	1	4364.2529	1.289	-1.916						Hill et al. (2002)
	CH	1	4364.2949	1.289	-4.181						Hill et al. (2002)
d		CH	1	4363.2651	0.000	-9.999					Hill et al. (2002)
b		CH	1	4365.4160	1.290	-4.093					Hill et al. (2002)
b		CH	1	4365.4502	1.290	-1.947					Hill et al. (2002)
b		CH	1	4366.0098	0.987	-3.729					Hill et al. (2002)
		CH	1	4366.2300	0.987	-3.828					Hill et al. (2002)
		CH	1	4366.3120	0.987	-3.828					Hill et al. (2002)
		CH	1	4366.3979	0.597	-3.426					Hill et al. (2002)
		CH	1	4366.4072	0.597	-1.699					Hill et al. (2002)
									2900.	3700.	Dummy line. Region 4366.165 - 4366.965

**Table 2.** Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Stark}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
CH	1	4366.4819	0.268	-3.414							Hill et al. (2002)
CH	1	4366.4971	0.268	-1.666							Hill et al. (2002)
CH	1	4366.5200	1.333	-4.129							Hill et al. (2002)
CH	1	4366.5732	0.597	-1.659							Hill et al. (2002)
CH	1	4366.5879	1.333	-1.920							Hill et al. (2002)
CH	1	4366.6470	1.333	-1.892							Hill et al. (2002)
CH	1	4366.6631	0.268	-1.627							Hill et al. (2002)
CH	1	4366.6992	1.333	-4.211							Hill et al. (2002)
CH	1	4366.8970	0.805	-1.903							Hill et al. (2002)
CH	1	4366.9282	0.987	-3.495							Hill et al. (2002)
CH	1	4366.9302	0.805	-3.326							Hill et al. (2002)
CH	1	4366.9912	0.987	-3.495							Hill et al. (2002)
CH	1	4367.1719	0.805	-1.848							Hill et al. (2002)
Fe	1	3872.5100	0.000	-9.999							Hill et al. (2002)
b	b	CN	1	3870.0049	2.086	-3.592					-2110. -1010. Dummy line. Region 3870.400 - 3871.500
b	b	CN	1	3870.0081	0.962	-3.473					Hill et al. (2002)
b	b	CN	1	3870.0139	0.651	-3.556					Hill et al. (2002)
b	b	CN	1	3870.0181	2.575	-0.038					Hill et al. (2002)
b	b	CN	1	3870.0190	4.153	-3.504					Hill et al. (2002)
b	b	CN	1	3870.0281	2.086	-0.469					Hill et al. (2002)
b	b	CN	1	3870.0459	4.973	-0.109					Hill et al. (2002)
b	b	CN	1	3870.0581	0.651	0.062					Hill et al. (2002)
b	b	CN	1	3870.0730	2.168	-1.086					Hill et al. (2002)
b	b	CN	1	3870.0769	0.962	0.336					Hill et al. (2002)
b	b	CN	1	3870.0801	2.086	-0.486					Hill et al. (2002)
b	b	CN	1	3870.0850	2.168	-1.160					Hill et al. (2002)
b	b	CN	1	3870.0901	2.168	-2.993					Hill et al. (2002)
b	b	12CH	1	3870.0930	0.463	-6.524					Jorgensen et al. (1996)
b	b	CN	1	3870.0950	4.973	-0.142					Hill et al. (2002)
b	b	CN	1	3870.1121	0.651	0.051					Hill et al. (2002)
b	b	12CH	1	3870.1179	0.002	-3.834					Jorgensen et al. (1996)
b	b	CN	1	3870.1311	0.333	-3.079					Hill et al. (2002)
b	b	12CH	1	3870.1440	0.463	-4.228					Jorgensen et al. (1996)
b	b	CN	1	3870.1499	0.962	0.330					Hill et al. (2002)
b	b	CN	1	3870.1509	0.333	-0.289					Hill et al. (2002)
b	b	CN	1	3870.1760	0.333	-0.314					Hill et al. (2002)
b	b	12CH	1	3870.1980	0.019	-7.205					Jorgensen et al. (1996)
b	b	12CH	1	3870.1990	0.463	-4.111					Jorgensen et al. (1996)
b	b	CN	1	3870.2029	3.619	-1.809					Hill et al. (2002)
b	b	CN	1	3870.2119	2.602	-3.832					Hill et al. (2002)
b	b	CN	1	3870.2161	2.840	-4.705					Jorgensen et al. (1996)
b	b	12CH	1	3870.2280	1.860	-4.776					Hill et al. (2002)
b	b	CN	1	3870.2319	0.632	-3.556					Jorgensen et al. (1996)
b	b	12CH	1	3870.2500	0.463	-6.492					Hill et al. (2002)
b	b	CN	1	3870.2700	2.603	-0.020					Jorgensen et al. (1996)
b	b	12CH	1	3870.2729	0.019	-7.205					Jorgensen et al. (1996)
b	b	CN	1	3870.2759	0.632	0.051					Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{HeNe}}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{Sark}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{HeNe}}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
b	CN	1	3870.2900	2.840	-1.501						Hill et al. (2002)
b	12CH	1	3870.2981	0.357	-4.313						Jørgensen et al. (1996)
b	CN	1	3870.3081	2.840	-1.494						Hill et al. (2002)
b	CN	1	3870.3149	3.619	-1.694						Hill et al. (2002)
b	CN	1	3870.3220	3.774	-1.212						Hill et al. (2002)
b	CN	1	3870.3291	0.632	0.040						Hill et al. (2002)
b	CN	1	3870.3320	3.774	-1.287						Hill et al. (2002)
b	CN	1	3870.3401	0.341	-3.079						Hill et al. (2002)
b	CN	1	3870.3501	2.602	-0.029						Hill et al. (2002)
b	CN	1	3870.3621	0.341	-0.267						Hill et al. (2002)
b	12CH	1	3870.3689	0.357	-4.313						Jørgensen et al. (1996)
b	CN	1	3870.3870	0.341	-0.289						Hill et al. (2002)
b	CN	1	3870.4319	0.614	-3.556						Hill et al. (2002)
b	CN	1	3870.4509	2.098	-3.591						Hill et al. (2002)
b	12CH	1	3870.4509	0.019	-7.359						Jørgensen et al. (1996)
b	CN	1	3870.4629	3.177	-1.395						Hill et al. (2002)
b	CN	1	3870.4751	0.614	0.040						Hill et al. (2002)
b	CN	1	3870.4761	2.098	-0.451						Hill et al. (2002)
b	Ca	1	3870.4771	2.521	-1.044	-7.269		-4.03	7.180		VALD/Kupka et al. (1999)
b	CN	1	3870.4819	5.152	0.226						Hill et al. (2002)
b	CN	1	3870.4971	5.536	-0.288						Hill et al. (2002)
b	CN	1	3870.4990	5.152	0.239						Hill et al. (2002)
b	CN	1	3870.5000	5.536	-0.353						Hill et al. (2002)
b	CN	1	3870.5259	4.097	0.334						Hill et al. (2002)
b	CN	1	3870.5259	0.614	0.029						Hill et al. (2002)
b	12CH	1	3870.5259	0.019	-7.359						Jørgensen et al. (1996)
b	CN	1	3870.5291	2.098	-0.469						Hill et al. (2002)
b	CN	1	3870.5300	0.351	-3.079						Hill et al. (2002)
b	CN	1	3870.5349	4.649	-0.517						Hill et al. (2002)
b	CN	1	3870.5491	2.630	-3.829						Hill et al. (2002)
b	CN	1	3870.5530	0.351	-0.245						Hill et al. (2002)
b	12CH	1	3870.5559	1.758	-4.459						Jørgensen et al. (1996)
b	CN	1	3870.5710	3.177	-1.356						Hill et al. (2002)
b	CN	1	3870.5769	4.649	-0.520						Hill et al. (2002)
b	CN	1	3870.5801	0.351	-0.267						Hill et al. (2002)
b	CN	1	3870.5891	2.166	-1.160						Hill et al. (2002)
b	CN	1	3870.5979	4.169	0.227						Hill et al. (2002)
b	CN	1	3870.6001	2.166	-1.249						Hill et al. (2002)
b	CN	1	3870.6030	2.166	-2.896						Hill et al. (2002)
b	CN	1	3870.6069	2.630	-0.011						Hill et al. (2002)
b	CN	1	3870.6121	0.596	-3.556						Hill et al. (2002)
b	CN	1	3870.6499	0.010	-0.600						Hill et al. (2002)
b	CN	1	3870.6541	0.596	0.029						Hill et al. (2002)
b	CN	1	3870.6589	4.169	0.224						Hill et al. (2002)
b	CN	1	3870.6621	0.010	-0.664						Hill et al. (2002)
b	CN	1	3870.6699	0.010	-2.629						Hill et al. (2002)
b	CN	1	3870.6880	2.630	-0.020						Hill et al. (2002)

**Table 2.** Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}/\dot{N}_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}/\dot{N}_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{Trad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
	CN	1	3870.6931	4.097	0.338						Hill et al. (2002)
	CN	1	3870.7009	0.360	-3.255						Hill et al. (2002)
	CN	1	3870.7041	0.596	0.018						Hill et al. (2002)
b	12CH	1	3870.7051	1.654	-5.155						Jørgensen et al. (1996)
	CN	1	3870.7251	0.360	-0.224						Hill et al. (2002)
	CN	1	3870.7300	4.472	-1.632						Hill et al. (2002)
	CN	1	3870.7329	0.932	-3.473						Hill et al. (2002)
	CN	1	3870.7339	4.169	-3.771						Hill et al. (2002)
	CN	1	3870.7529	0.360	-0.245						Hill et al. (2002)
	CN	1	3870.7729	0.578	-3.556						Hill et al. (2002)
	CN	1	3870.8010	0.933	0.330						Hill et al. (2002)
	CN	1	3870.8140	0.578	0.018						Hill et al. (2002)
	CN	1	3870.8311	5.437	0.468						Hill et al. (2002)
	CN	1	3870.8521	0.371	-3.255						Hill et al. (2002)
	CN	1	3870.8530	4.472	-1.833						Hill et al. (2002)
	CN	1	3870.8630	0.578	0.006						Hill et al. (2002)
b	Ce	2	3870.8669	0.684	-0.568						VALD; Kupka et al. (1999)
	CN	1	3870.8730	0.932	0.323						Hill et al. (2002)
	CN	1	3870.8760	0.371	-0.204						Hill et al. (2002)
	CN	1	3870.8831	5.646	0.522						Hill et al. (2002)
	CN	1	3870.8921	2.658	-3.827						Hill et al. (2002)
	CN	1	3870.8960	2.110	-3.590						Hill et al. (2002)
b	12CH	1	3870.8989	0.358	-5.238						Jørgensen et al. (1996)
	CN	1	3870.9060	4.372	-0.515						Hill et al. (2002)
	CN	1	3870.9060	0.371	-0.224						Hill et al. (2002)
	CN	1	3870.9150	0.561	-3.556						Hill et al. (2002)
	CN	1	3870.9221	2.110	-0.435						Hill et al. (2002)
	CN	1	3870.9509	2.658	-0.002						Hill et al. (2002)
	12CH	1	3870.9529	0.358	-5.238						Jørgensen et al. (1996)
b	CN	1	3870.9541	0.561	0.006						Hill et al. (2002)
	CN	1	3871.0020	0.561	-0.006						Hill et al. (2002)
	CN	1	3871.0090	0.381	-0.185						Hill et al. (2002)
	CN	1	3871.0100	4.372	-0.564						Hill et al. (2002)
b	12CH	1	3871.0161	0.358	-4.201						Jørgensen et al. (1996)
	CN	1	3871.0330	2.658	-0.011						Hill et al. (2002)
b	U	1	3871.0349	0.000	0.230						VALD; Kupka et al. (1999)
	CN	1	3871.0371	0.545	-3.556						Hill et al. (2002)
	CN	1	3871.0400	0.381	-0.204						Hill et al. (2002)
b	12CH	1	3871.0701	0.358	-4.201						Jørgensen et al. (1996)
	CN	1	3871.0750	0.545	-0.006						Hill et al. (2002)
	CN	1	3871.0759	5.437	0.466						Hill et al. (2002)
b	12CH	1	3871.0759	0.020	-6.891						Jørgensen et al. (1996)
	CN	1	3871.0950	0.392	-3.255						Hill et al. (2002)
	CN	1	3871.1069	2.164	-1.249						Hill et al. (2002)
	CN	1	3871.1150	2.164	-1.361						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{He}\alpha}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarX}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{He}\alpha}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
<b>b</b>	CN	1	3871.1179	2.164	-2.783						Hill et al. (2002)
	CN	1	3871.1221	0.392	-0.167						Hill et al. (2002)
	CN	1	3871.1221	0.545	-0.019						Hill et al. (2002)
	12CH	1	3871.1331	0.020	-6.891						Jorgensen et al. (1996)
	CN	1	3871.1389	0.529	-3.556						Hill et al. (2002)
	CN	1	3871.1550	0.392	-0.185						Hill et al. (2002)
	CN	1	3871.1770	0.529	-0.019						Hill et al. (2002)
	CN	1	3871.1799	5.119	-0.261						Hill et al. (2002)
	CN	1	3871.1880	0.404	-3.255						Hill et al. (2002)
	CN	1	3871.2151	0.404	-0.150						Hill et al. (2002)
	CN	1	3871.2229	0.529	-0.031						Hill et al. (2002)
	CN	1	3871.2229	0.513	-3.255						Hill et al. (2002)
<b>b</b>	CN	1	3871.2429	2.687	-3.825						Hill et al. (2002)
	CN	1	3871.2490	0.404	-0.167						Hill et al. (2002)
	CN	1	3871.2590	0.513	-0.031						Hill et al. (2002)
	CN	1	3871.2600	0.416	-3.255						Hill et al. (2002)
	CN	1	3871.2649	6.492	-0.354						Hill et al. (2002)
	CN	1	3871.2771	5.119	-0.298						Hill et al. (2002)
	CN	1	3871.2859	0.498	-3.255						Hill et al. (2002)
	CN	1	3871.2891	0.416	-0.133						Hill et al. (2002)
	CN	1	3871.2949	5.963	-0.151						Hill et al. (2002)
	CN	1	3871.3030	2.687	0.007						Hill et al. (2002)
	CN	1	3871.3040	0.513	-0.045						Hill et al. (2002)
	CN	1	3871.3130	0.428	-3.255						Hill et al. (2002)
<b>b</b>	CN	1	3871.3220	0.498	-0.045						Hill et al. (2002)
	CN	1	3871.3240	0.416	-0.150						Hill et al. (2002)
	12CH	1	3871.3269	1.846	-2.075						Jorgensen et al. (1996)
	CN	1	3871.3311	0.483	-3.255						Hill et al. (2002)
	CN	1	3871.3401	2.122	-3.588						Hill et al. (2002)
	CN	1	3871.3430	0.428	-0.117						Hill et al. (2002)
	CN	1	3871.3469	0.441	-3.255						Hill et al. (2002)
	CN	1	3871.3560	0.469	-3.255						Hill et al. (2002)
	CN	1	3871.3601	0.007	-0.664						Hill et al. (2002)
	CN	1	3871.3611	0.455	-3.255						Hill et al. (2002)
	CN	1	3871.3650	0.498	-0.058						Hill et al. (2002)
	CN	1	3871.3650	0.483	-0.058						Hill et al. (2002)
<b>b</b>	CN	1	3871.3669	2.122	-0.419						Hill et al. (2002)
	CN	1	3871.3701	0.007	-0.737						Hill et al. (2002)
	CN	1	3871.3721	1.846	-4.675						Hill et al. (2002)
	CN	1	3871.3730	4.142	0.338						Jorgensen et al. (1996)
	CN	1	3871.3760	0.007	-2.571						Hill et al. (2002)
	CN	1	3871.3779	0.441	-0.102						Hill et al. (2002)
	CN	1	3871.3789	2.854	-5.025						Hill et al. (2002)
	CN	1	3871.3799	0.428	-0.133						Hill et al. (2002)
	CN	1	3871.3840	2.687	-0.002						Hill et al. (2002)
	CN	1	3871.3889	0.469	-0.072						Hill et al. (2002)
	CN	1	3871.3931	0.455	-0.087						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Stark}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	12CH	1	3871.3940	0.096	-1.640						Jørgensen et al. (1996)
b	12CH	1	3871.4021	1.846	-4.754						Jørgensen et al. (1996)
	CN	1	3871.4070	0.483	-0.072						Hill et al. (2002)
	CN	1	3871.4160	0.441	-0.117						Hill et al. (2002)
	CN	1	3871.4241	2.122	-0.435						Hill et al. (2002)
	CN	1	3871.4299	0.469	-0.087						Hill et al. (2002)
	CN	1	3871.4331	0.455	-0.102						Hill et al. (2002)
	CN	1	3871.4370	0.904	-3.473						Hill et al. (2002)
b	12CH	1	3871.4470	1.846	-2.057						Jørgensen et al. (1996)
b	CN	1	3871.4561	2.854	-1.509						Hill et al. (2002)
	CN	1	3871.4741	2.854	-1.501						Hill et al. (2002)
b	CN	1	3871.4851	5.963	-0.113						Hill et al. (2002)
b	CN	1	3871.5039	0.904	0.323						Hill et al. (2002)
b	CN	1	3871.5110	5.168	-0.598						Hill et al. (2002)
b	God	2	3871.5430	0.079	-1.588						VALD:Kupka et al. (1999)
b	CN	1	3871.5500	4.142	0.347						Hill et al. (2002)
b	12CH	1	3871.5530	0.464	-6.524						Jørgensen et al. (1996)
b	CN	1	3871.5540	6.492	-0.434						Hill et al. (2002)
b	12CH	1	3871.5620	2.368	-7.016						Jørgensen et al. (1996)
b	12CH	1	3871.5640	0.464	-4.111						Jørgensen et al. (1996)
b	CN	1	3871.5730	5.168	-0.686						Hill et al. (2002)
b	12CH	1	3871.5730	0.072	-1.707						Jørgensen et al. (1996)
b	CN	1	3871.5759	0.904	0.316						Hill et al. (2002)
b	CN	1	3871.6011	2.715	-3.822						Hill et al. (2002)
b	12CH	1	3871.6040	0.464	-4.228						Jørgensen et al. (1996)
b	12CH	1	3871.6150	0.464	-6.492						Jørgensen et al. (1996)
b	CN	1	3871.6250	2.163	-1.361						Hill et al. (2002)
b	Dy	2	3871.6260	0.928	-0.697						VALD:Kupka et al. (1999)
b	CN	1	3871.6311	2.163	-1.516						Hill et al. (2002)
b	CN	1	3871.6340	2.163	-2.666						Hill et al. (2002)
b	La	2	3871.6350	0.126	-0.127						VALD:Kupka et al. (1999)
b	12CH	1	3871.6399	2.368	-9.811						Jørgensen et al. (1996)
b	CN	1	3871.6621	2.716	0.016						Hill et al. (2002)
b	12CH	1	3871.6660	0.096	-3.117						Jørgensen et al. (1996)
b	12CH	1	3871.6809	0.096	-1.598						Jørgensen et al. (1996)
d	CN	1	3881.8750	0.000	-9.999						675. Dummy line. Region 3882.550 - 3883.400
b	CN	1	3882.0500	3.410	-3.766						Hill et al. (2002)
b	CN	1	3882.0559	0.072	-2.997						Hill et al. (2002)
b	12CH	1	3882.0559	1.978	-7.188						Jørgensen et al. (1996)
b	CN	1	3882.0669	4.530	-0.379						Hill et al. (2002)
b	CN	1	3882.0759	0.072	-0.231						Hill et al. (2002)
b	CN	1	3882.0791	0.364	-3.473						Hill et al. (2002)
b	12CH	1	3882.0859	1.978	-6.586						Jørgensen et al. (1996)
b	CN	1	3882.0901	2.227	-3.414						Hill et al. (2002)
b	CN	1	3882.0969	5.201	-0.198						Hill et al. (2002)
b	CN	1	3882.0979	0.072	-0.257						Hill et al. (2002)
b	CN	1	3882.0991	6.294	-0.040						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{He}\alpha}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{Sark}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{He}\alpha}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
b	CN	1	3882.1050	2.227	-0.647						Hill et al. (2002)
b	CN	1	3882.1230	0.364	0.123						Hill et al. (2002)
b	CN	1	3882.1279	3.410	0.188						Hill et al. (2002)
b	CN	1	3882.1321	2.227	-0.674						Hill et al. (2002)
b	Ti	1	3882.1450	2.017	0.439	-7.606		-4.40	8.150		VALD:Kupka et al. (1999)
b	12CH	1	3882.1450	3.069	-5.690						Jørgensen et al. (1996)
b	12CH	1	3882.1509	1.978	-3.808						Jørgensen et al. (1996)
b	12CH	1	3882.1699	3.069	-8.782						Jørgensen et al. (1996)
b	CN	1	3882.1731	0.364	0.112						Hill et al. (2002)
b	CN	1	3882.1750	2.823	-3.345						Hill et al. (2002)
b	CN	1	3882.1760	2.823	-4.039						Hill et al. (2002)
b	CN	1	3882.1890	5.201	-0.228						Hill et al. (2002)
b	CN	1	3882.2019	3.410	0.180						Hill et al. (2002)
b	12CH	1	3882.2051	2.883	-7.470						Jørgensen et al. (1996)
b	CN	1	3882.2061	6.294	-0.037						Hill et al. (2002)
b	CN	1	3882.2749	0.080	-2.997						Hill et al. (2002)
b	Ti	2	3882.2910	1.116	-1.940	-7.896		-6.67	8.360		VALD:Kupka et al. (1999)
b	CN	1	3882.2949	0.080	-0.207						Hill et al. (2002)
b	CN	1	3882.2959	0.346	-3.473						Hill et al. (2002)
b	CN	1	3882.3191	0.080	-0.231						Hill et al. (2002)
b	CN	1	3882.3379	0.346	0.112						Hill et al. (2002)
b	12CH	1	3882.3450	2.883	-4.580						Jørgensen et al. (1996)
b	12CH	1	3882.3501	3.069	-5.681						Jørgensen et al. (1996)
b	CN	1	3882.3511	2.549	-3.851						Hill et al. (2002)
b	CN	1	3882.3540	3.098	-3.859						Hill et al. (2002)
b	CN	1	3882.3879	0.346	0.101						Hill et al. (2002)
b	CN	1	3882.3970	2.549	-0.120						Hill et al. (2002)
b	CN	1	3882.4121	2.129	-2.256						Jørgensen et al. (1996)
b	12CH	1	3882.4189	2.884	-4.590						Hill et al. (2002)
b	CN	1	3882.4341	3.098	-3.528						Hill et al. (2002)
b	Ce	2	3882.4451	0.322	0.041						VALD:Kupka et al. (1999)
b	CN	1	3882.4729	0.089	-2.997						Hill et al. (2002)
b	12CH	1	3882.4890	2.129	-4.851						Jørgensen et al. (1996)
b	CN	1	3882.4919	0.328	-3.473						Hill et al. (2002)
b	CN	1	3882.4939	2.549	-0.131						Hill et al. (2002)
b	CN	1	3882.4951	0.089	-0.184						Hill et al. (2002)
b	CN	1	3882.4961	4.452	0.231						Hill et al. (2002)
b	Sm	2	3882.5049	0.277	-1.046						VALD:Kupka et al. (1999)
b	CN	1	3882.5200	0.089	-0.207						Hill et al. (2002)
b	12CH	1	3882.5220	1.656	-2.487						Jørgensen et al. (1996)
b	Hf	1	3882.5281	1.943	-0.080						VALD:Kupka et al. (1999)
b	CN	1	3882.5330	0.328	0.101						Hill et al. (2002)
b	12CH	1	3882.5349	1.656	-2.509						Jørgensen et al. (1996)
b	CN	1	3882.5410	4.452	0.232						Hill et al. (2002)
b	12CH	1	3882.5459	2.129	-2.241						Jørgensen et al. (1996)
b	12CH	1	3882.5759	1.656	-4.675						Jørgensen et al. (1996)
b	12CH	1	3882.5779	0.521	-5.008						Jørgensen et al. (1996)

**Table 2.** Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{StarK}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	CN	1	3882.5811	0.328	0.089						Hill et al. (2002)
	CN	1	3882.6240	2.235	-3.413						Hill et al. (2002)
	CN	1	3882.6270	2.822	-3.345						Hill et al. (2002)
	CN	1	3882.6311	2.822	-3.643						Hill et al. (2002)
	CN	1	3882.6389	2.235	-0.623						Hill et al. (2002)
	12CH	1	3882.6389	2.965	-4.170						Jorgensen et al. (1996)
	CN	1	3882.6499	0.098	-2.997						Hill et al. (2002)
	CN	1	3882.6670	0.311	-3.473						Hill et al. (2002)
	CN	1	3882.6680	2.235	-0.647						Hill et al. (2002)
	CN	1	3882.6731	0.098	-0.162						Hill et al. (2002)
	CN	1	3882.6880	5.347	0.173						Hill et al. (2002)
	CN	1	3882.6919	5.347	0.150						Hill et al. (2002)
	CN	1	3882.7000	0.098	-0.184						Hill et al. (2002)
	CN	1	3882.7080	0.311	0.089						Hill et al. (2002)
b	12CH	1	3882.7119	2.964	-4.179						Jorgensen et al. (1996)
	CN	1	3882.7241	4.610	0.394						Hill et al. (2002)
b	12CH	1	3882.7439	2.965	-7.263						Jorgensen et al. (1996)
	CN	1	3882.7520	3.448	-3.764						Hill et al. (2002)
	CN	1	3882.7549	0.311	0.077						Hill et al. (2002)
	CN	1	3882.8069	0.108	-3.173						Hill et al. (2002)
	CN	1	3882.8220	0.294	-3.473						Hill et al. (2002)
	CN	1	3882.8301	3.448	0.196						Hill et al. (2002)
	CN	1	3882.8311	0.108	-0.141						Hill et al. (2002)
	CN	1	3882.8589	0.108	-0.162						Hill et al. (2002)
	CN	1	3882.8589	2.572	-3.849						Hill et al. (2002)
	CN	1	3882.8611	0.294	0.077						Hill et al. (2002)
	CN	1	3882.8630	3.113	-4.447						Hill et al. (2002)
b	Er	2	3882.8860	0.886	-0.214						VALD:Kupka et al. (1999)
b	Ti	1	3882.8911	2.041	0.686						VALD:Kupka et al. (1999)
	CN	1	3882.9031	3.448	0.188						Hill et al. (2002)
	CN	1	3882.9060	2.572	-0.110						Hill et al. (2002)
	CN	1	3882.9070	0.294	0.064						Hill et al. (2002)
	CN	1	3882.9431	0.118	-3.173						Hill et al. (2002)
	CN	1	3882.9460	3.113	-3.859						Hill et al. (2002)
	CN	1	3882.9529	3.000	-1.672						Hill et al. (2002)
	CN	1	3882.9561	0.278	-3.473						Hill et al. (2002)
	CN	1	3882.9680	0.118	-0.122						Hill et al. (2002)
	CN	1	3882.9729	3.000	-1.645						Hill et al. (2002)
	CN	1	3882.9810	4.026	-0.784						Hill et al. (2002)
	CN	1	3882.9839	4.026	-0.733						Hill et al. (2002)
	CN	1	3882.9939	0.278	0.064						Hill et al. (2002)
	CN	1	3882.9971	0.118	-0.141						Hill et al. (2002)
	CN	1	3883.0039	2.572	-0.120						Hill et al. (2002)
	CN	1	3883.0190	4.610	0.398						Hill et al. (2002)
b	12CH	1	3883.0359	2.761	-4.775						Jorgensen et al. (1996)
	CN	1	3883.0391	0.278	0.052						Hill et al. (2002)
	CN	1	3883.0581	0.129	-3.173						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{He}\alpha}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarX}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{He}\alpha}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
b	CN	1	3883.0691	0.262	-3.172						Hill et al. (2002)
	CN	1	3883.0740	3.365	-1.820						Hill et al. (2002)
	CN	1	3883.0840	0.129	-0.103						Hill et al. (2002)
	CN	1	3883.1060	0.262	0.052						Hill et al. (2002)
	12CH	1	3883.1121	2.761	-4.786						Jørgensen et al. (1996)
	CN	1	3883.1150	0.129	-0.122						Hill et al. (2002)
	CN	1	3883.1499	0.262	0.038						Hill et al. (2002)
	CN	1	3883.1531	0.140	-3.173						Hill et al. (2002)
	CN	1	3883.1589	2.243	-3.412						Hill et al. (2002)
	CN	1	3883.1609	0.247	-3.172						Hill et al. (2002)
	CN	1	3883.1750	2.243	-0.599						Hill et al. (2002)
	CN	1	3883.1799	0.140	-0.085						Hill et al. (2002)
	CN	1	3883.1880	3.365	-1.735						Hill et al. (2002)
	CN	1	3883.1970	0.247	0.038						Hill et al. (2002)
	CN	1	3883.2061	2.243	-0.623						Hill et al. (2002)
	CN	1	3883.2119	0.140	-0.103						Hill et al. (2002)
	CN	1	3883.2271	0.152	-3.173						Hill et al. (2002)
	CN	1	3883.2329	0.232	-3.172						Hill et al. (2002)
	CN	1	3883.2400	0.247	0.025						Hill et al. (2002)
b	12CH	1	3883.2419	2.761	-7.781						Jørgensen et al. (1996)
	CN	1	3883.2549	0.152	-0.067						Hill et al. (2002)
	CN	1	3883.2681	0.232	0.025						Hill et al. (2002)
b	12CH	1	3883.2690	0.775	-3.655						Jørgensen et al. (1996)
b	Cr	1	3883.2700	3.887	-0.448						VALD: Kupka et al. (1999)
b	12CH	1	3883.2771	1.803	-7.293						Jørgensen et al. (1996)
b	CN	1	3883.2800	0.164	-3.173						Hill et al. (2002)
b	Fe	1	3883.2800	3.251	-0.604						VALD: Kupka et al. (1999)
b	CN	1	3883.2839	0.217	-3.172						Hill et al. (2002)
b	CN	1	3883.2881	0.152	-0.085						Hill et al. (2002)
b	12CH	1	3883.2959	1.803	-5.432						Jørgensen et al. (1996)
b	CN	1	3883.3091	0.232	0.011						Hill et al. (2002)
b	CN	1	3883.3091	0.164	-0.050						Hill et al. (2002)
b	CN	1	3883.3120	0.177	-3.173						Hill et al. (2002)
b	CN	1	3883.3140	0.203	-3.172						Hill et al. (2002)
b	CN	1	3883.3181	0.217	0.011						Hill et al. (2002)
b	CN	1	3883.3230	0.190	-3.172						Hill et al. (2002)
b	12CH	1	3883.3259	0.775	-6.216						Jørgensen et al. (1996)
b	CN	1	3883.3350	0.190	-0.019						Hill et al. (2002)
b	CN	1	3883.3420	0.177	-0.034						Hill et al. (2002)
b	CN	1	3883.3440	0.164	-0.004						Hill et al. (2002)
b	CN	1	3883.3469	0.164	-0.067						Hill et al. (2002)
b	CN	1	3883.3621	5.945	-0.003						Hill et al. (2002)
b	CN	1	3883.3750	2.595	-3.846						Hill et al. (2002)
b	CN	1	3883.3779	0.177	-0.050						Hill et al. (2002)
b	CN	1	3883.3850	0.203	-0.019						Hill et al. (2002)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{StarK}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{Irad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	CN	1	3883.3921	0.190	-0.034						Hill et al. (2002)
b	12CH	1	3883.4060	0.775	-3.675						Jørgensen et al. (1996)
b	CN	1	3883.4070	4.422	-0.508						Hill et al. (2002)
b	CN	1	3883.4080	5.282	-0.452						Hill et al. (2002)
b	CN	1	3883.4231	2.595	-0.100						Hill et al. (2002)
b	CN	1	3883.4309	4.422	-0.509						Hill et al. (2002)
b	CN	1	3883.4370	5.282	-0.470						Hill et al. (2002)
b	12CH	1	3883.4700	1.483	-7.559						Jørgensen et al. (1996)
b	CN	1	3883.4780	3.485	-3.761						Hill et al. (2002)
b	12CH	1	3883.4800	2.359	-5.453						Jørgensen et al. (1996)
b	CN	1	3883.4890	3.129	-4.447						Hill et al. (2002)
b	CN	1	3883.5210	2.595	-0.110						Hill et al. (2002)
b	12CH	1	3883.5291	2.397	-4.058						Jørgensen et al. (1996)
b	CN	1	3883.5300	5.945	0.000						Hill et al. (2002)
b	Ce	2	3883.5330	0.536	-1.146						VALD:Kupka et al. (1999)
b	CN	1	3883.5581	3.485	0.203						Hill et al. (2002)
b	CN	1	3883.5601	5.712	0.444						Hill et al. (2002)
b	12CH	1	3883.5979	2.397	-7.148						Jørgensen et al. (1996)
b	CN	1	3883.6150	2.823	-3.643						Hill et al. (2002)
b	CN	1	3883.6160	2.823	-3.346						Hill et al. (2002)
b	12CH	1	3883.6250	2.397	-7.148						Jørgensen et al. (1996)
b	CN	1	3883.6289	3.485	0.196						Hill et al. (2002)
b	CN	1	3883.6589	3.830	-3.185						Hill et al. (2002)
b	CN	1	3883.6680	5.258	-0.447						Hill et al. (2002)
b	12CH	1	3883.6951	2.397	-4.047						Jørgensen et al. (1996)
b	CN	1	3883.6960	2.251	-3.412						Hill et al. (2002)
b	O	1	7771.9409	9.146	0.370	-7.469		-5.55	7.520	-400.	VALD:Kupka et al. (1999), loggf:Jørgensen et al. (2005)
b	O	1	7774.1611	9.146	0.240	-7.469		-5.55	7.520	-400.	VALD:Kupka et al. (1999), loggf:Jørgensen et al. (2005)
d	Na	1	5889.9780	0.000	-9.999	381.256		-5.64	7.800	-400.	Dummy line. Region 5889.578 - 5890.378
h	Na	1	5889.9580	0.000	-0.775	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5889.9590	0.000	-0.791	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5889.9590	0.000	-0.678	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5889.9780	0.000	-0.314	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5889.9790	0.000	-0.678	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5889.9790	0.000	-0.777	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5895.9380	0.000	-9.999	381.256		-5.64	7.800	-400.	Dummy line. Region 5895.538 - 5896.118
d	Na	1	5895.9268	0.000	-0.680	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5895.9302	0.000	-1.224	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5895.9468	0.000	-0.777	381.256		-5.64	7.800		Wiese & Martin (1980), hfs:McWilliam et al.(1995)
h	Na	1	5895.9502	0.000	-0.680	381.256		-5.64	7.800	-400.	VALD:Kupka et al. (1999)
h	Na	1	8183.2549	2.102	0.230			-4.56	2.340	-100.	Barklem et al.(2005)
Mg	1	3829.3550	2.707	-0.208	709.301		-4.56	2.340	-300.	300. Barklem et al.(2005)	
Mg	1	4571.0962	0.000	-5.393	-6.970		-3.98	8.720	-300.	230. Barklem et al.(2005)	
Mg	1	4702.9902	4.330	-0.380	2806.269		-3.98	8.720	-300.	230. Barklem et al.(2005)	
Mg	1	5172.7002	2.710	-0.380	729.238		-3.98	7.990	-450.	450. Wiese & Martin (1980)	
Mg	1	5183.6201	2.720	-0.160	729.238		-3.98	7.990	-130.	900. Wiese & Martin (1980)	
Mg	1	5528.4199	4.350	-0.340	-6.979		-4.46	8.720	-300.	300. Wiese & Martin (1980)	

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta/\delta_{\text{std}}$ [mÅ]	$\delta/\delta_{\text{std}}$ [mÅ]	References and notes
Mg	1	5711.0879	4.346	-1.870			8.710	-300.	300.	VALD:Kupka et al.(1999), loggf:Jonsell et al.(2005)	
d	A1	1	3961.5291	0.014	-9.999	655.243	-5.37	8.480	-75.	90.	Dummy line. Region 3961.454 - 3961.619
h	A1	1	3961.5259	0.014	-1.670	655.243	-5.37	8.480			Wiese & Martin(1980), hfs:McWilliam et al.(1995)
h	A1	1	3961.5271	0.014	-1.126	655.243	-5.37	8.480			Wiese & Martin(1980), hfs:McWilliam et al.(1995)
h	A1	1	3961.5291	0.014	-0.762	655.243	-5.37	8.480			Wiese & Martin(1980), hfs:McWilliam et al.(1995)
h	A1	1	3961.5310	0.014	-1.239	655.243	-5.37	8.480			Wiese & Martin(1980), hfs:McWilliam et al.(1995)
h	A1	1	3961.5320	0.014	-1.126	655.243	-5.37	8.480			Wiese & Martin(1980), hfs:McWilliam et al.(1995)
h	A1	1	3961.5339	0.014	-1.223	655.243	-5.37	8.480			Wiese & Martin(1980), hfs:McWilliam et al.(1995)
b	C	1	3961.4031	7.685	-2.140						VALD: Kupka et al. (1999)
b	12CH	1	3961.0620	1.520	-2.828						Jorgensen et al. (1996)
b	12CH	1	3961.1030	1.520	-2.843						Jorgensen et al. (1996)
b	12CH	1	3961.2241	1.290	-2.695						Jorgensen et al. (1996)
b	12CH	1	3961.2620	1.290	-4.658						Jorgensen et al. (1996)
b	12CH	1	3961.3059	1.290	-2.644						Jorgensen et al. (1996)
b	13CH	1	3961.4961	0.230	-2.385						Jorgensen et al. (1996)
b	13CH	1	3961.6289	0.230	-2.353						Jorgensen et al. (1996)
b	13CH	1	3961.7390	1.520	-3.782						Jorgensen et al. (1996)
b	13CH	1	3961.7720	0.230	-4.160						Jorgensen et al. (1996)
b	13CH	1	3961.7891	1.290	-3.649						Jorgensen et al. (1996)
b	13CH	1	3961.7910	1.520	-3.797						Jorgensen et al. (1996)
b	13CH	1	3961.8789	1.290	-3.598						Jorgensen et al. (1996)
b	12CH	1	3961.8931	0.990	-3.305						Jorgensen et al. (1996)
b	12CH	1	3961.9609	0.990	-3.305						Jorgensen et al. (1996)
b	Ca	1	4226.7280	0.000	0.244	372.238	-6.03	8.360	-100.	170.	Wiese & Martin(1980)
b	12CH	1	4226.8110	2.790	-1.112						Jorgensen et al. (1996) predicted 0.8mA
Ca	1	4425.4370	1.879	-0.358	947.274	-5.61	8.025	-215.			
Ca	1	5349.4702	2.710	-0.310	-7.98	-5.89	6.430	-250.			
Ca	1	5588.7598	2.530	0.360	-7.538	-6.07	7.850	-300.			
Ca	1	6102.7300	1.880	-0.790	-7.189	-5.32	7.860	-300.			
Ca	1	6122.2300	1.890	-0.320	-7.189	-5.32	7.860	-300.			
Ca	1	6162.1802	1.900	-0.090	-7.189	-5.32	7.860	-300.			
Ca	1	6169.5630	2.526	-0.570	-7.145	-4.99	7.270	-280.			
Ca	1	6439.0801	2.530	0.390	-7.569	-6.07	7.650	-300.			
Ca	1	7148.1499	2.709	0.208	-7.798	-6.01	7.660	-300.			
Sc	2	4246.8369	0.315	-9.999	-7.910	-6.37	8.164	-65.	220.	Dummy line. Region 4246.772 - 4247.057	
h	Sc	2	4246.8320	0.315	-0.798	-7.910	-6.37	8.164			
h	Sc	2	4246.8359	0.315	-0.386	-7.910	-6.37	8.164			
h	Sc	2	4246.8389	0.315	-0.809	-7.910	-6.37	8.164			
h	Sc	2	4246.8389	0.315	-0.783	-7.910	-6.37	8.164			
h	Sc	2	4246.8408	0.315	-0.982	-7.910	-6.37	8.164			
h	Sc	2	4246.8418	0.315	-1.480	-7.910	-6.37	8.164			
h	Sc	2	4246.8428	0.315	-2.906	-7.910	-6.37	8.164			
h	Sc	2	4246.8428	0.315	-0.956	-7.910	-6.37	8.164			
h	Sc	2	4246.8442	0.315	-1.158	-7.910	-6.37	8.164			
h	Sc	2	4246.8452	0.315	-0.798	-7.910	-6.37	8.164			
h	Sc	2	4246.8462	0.315	-0.982	-7.910	-6.37	8.164			
h	Sc	2	4246.8462	0.315	-0.809	-7.910	-6.37	8.164			

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{vUV}}/\text{N}_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}/\text{N}_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	12CH	1	4246.8062	2.520	-1.501						Jorgensen et al. (1996)
b	12CH	1	4246.9048	2.520	-1.515						Jorgensen et al. (1996)
b	12CH	1	4246.9390	2.519	-1.936						Dummy line. Region 4415.433 - 4415.683
d	Sc	2	4415.5630	0.595	-9.999	-7.920	-6.68	8.253	-130.		
h	Sc	2	4415.5459	0.595	-1.866	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5508	0.595	-1.708	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5552	0.595	-1.719	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5571	0.595	-1.296	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5591	0.595	-1.892	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5591	0.595	-1.694	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5620	0.595	-2.390	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5630	0.595	-3.816	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5649	0.595	-2.068	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5688	0.595	-1.892	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5688	0.595	-1.866	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5698	0.595	-1.719	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	4415.5698	0.595	-1.708	-7.920	-6.68	8.253			Lawler & Dakin (1989), hfs; McWilliam et al.(1995)
h	Sc	2	5031.0200	1.360	-0.400	-7.904	-6.71	8.120	-150.	150.	Lawler & Dakin (1989)
Sc	2	5526.8198	1.770	-0.020	-7.870	-6.64	8.330	-150.	150.	Lawler & Dakin (1989)	
Sc	2	5657.8799	1.510	-0.600	-7.932	-6.60	8.170	-150.	150.	Lawler & Dakin (1989)	
Ti	1	3998.6360	0.048	0.000	305.269	-6.19	7.810	-150.	150.	Grevesse et al. (1989), Blackwell et al. (1982b)	
Ti	1	4533.2490	0.848	0.530	347.242	-5.14	8.083	-220.	300.	Grevesse et al. (1989), Blackwell et al. (1982a)	
b	12CH	1	4533.2021	0.955	-2.364	345.242	-5.31	8.079	-180.	180.	Jorgensen et al. (1996)
Ti	1	4534.7759	0.836	0.340	322.245	-6.17	7.897	-190.	190.	Grevesse et al. (1989), Blackwell et al. (1982a)	
Ti	1	4981.7310	0.848	0.504	322.245	-6.17	7.897	-190.	190.	Martin et al. (1988)	
Ti	1	4991.0649	0.836	0.380	-7.629	-6.17	7.890	-160.	130.	VALD: Kupka et al. (1999)	
b	Ce	2	4991.0142	1.402	-0.701						VALD: Kupka et al. (1999)
Ti	1	5022.8701	0.830	-0.430	-7.633	-6.17	7.890	-200.	200.	Fuhr & Weise (1996)	
Ti	1	5024.8501	0.820	-0.600	-7.635	-6.17	7.880	-200.	200.	Fuhr & Weise (1996)	
Ti	1	5039.9600	0.020	-1.130	-7.720	-6.35	6.880	-200.	200.	Martin et al. (1988)	
Ti	1	5064.6602	0.050	-0.990	-7.719	-6.33	6.850	-200.	200.	Martin et al. (1988)	
Ti	1	5192.9800	0.020	-1.010	-7.727	-6.32	6.820	-200.	200.	Martin et al. (1988)	
Ti	1	5210.3901	0.050	-0.880	-7.724	-6.31	6.810	-200.	200.	Martin et al. (1988)	
Ti	2	4053.8340	1.893	-1.060	-7.908	-6.56	8.380	-70.	110.	VALD: Kupka et al. (1999)	
b	12CH	1	4053.9710	1.248	-2.975						Jorgensen et al. (1996)
b	12CH	1	4053.9031	1.248	-2.915						Jorgensen et al. (1996)
Ti	2	4417.7192	1.165	-1.190	-7.930	-6.67	8.225	-85.	150.	Pickering et al. (2001)	
b	12CH	1	4417.6680	1.324	-2.362						Jorgensen et al. (1996)
b	12CH	1	4417.7969	1.598	-1.984						Jorgensen et al. (1996)
Ti	2	4443.7939	1.080	-0.720	-7.920	-6.51	8.119	-300.	250.	Pickering et al. (2001)	
b	12CH	1	4443.8062	1.581	-2.064						Jorgensen et al. (1996)
b	12CH	1	4443.8579	2.119	-1.272						Jorgensen et al. (1996)
Ti	2	4468.5068	1.131	-0.600	-7.930	-6.72	8.207	-270.	270.	Ryabchikova et al. (1994)	
Ti	2	4501.2729	1.116	-0.770	-7.930	-6.73	8.199	-180.	150.	Pickering et al. (2001)	
b	12CH	1	4501.2280	1.694	-2.167						Jorgensen et al. (1996)
b	12CH	1	4501.3071	3.179	-1.059						Jorgensen et al. (1996)
Ti	2	4563.7612	1.221	-0.690	-7.960	-6.71	8.217	-220.	220.	Pickering et al. (2001)	

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \chi_{\text{osc}}$ [mÅ]	$\delta \chi_{\text{rad}}$ [mÅ]	References and notes
b	Ti	2	4571.9678	1.572	-0.320	-7.890	-6.68	8.367	-200.	150.	Pickering et al. (2001)
b	12CH	1	4571.9331	0.873	-2.206						Jørgensen et al. (1996)
b	12CH	1	4572.0171	2.897	-1.541						Jørgensen et al. (1996)
b	12CH	1	4572.0562	3.273	-1.132						Jørgensen et al. (1996)
b	Ti	2	4589.9580	1.237	-1.620	-7.960	-6.71	8.217	-150.	150.	Ryabchikova et al. (1994)
b	12CH	1	4589.9810	2.519	-1.435						Jørgensen et al. (1996)
b	Ti	2	5185.9102	1.890	-1.490	-7.908	-6.53	8.370	-300.	300.	Martin et al. (1988)
b	Ti	2	5188.6982	1.582	-1.210	-7.948	-6.66	8.230	-160.	40.	Martin et al. (1988)
b	Ti	2	5226.5498	1.570	-1.300	-7.953	-6.71	8.220	-170.	120.	Martin et al. (1988)
b	Ti	2	5336.7900	1.580	-1.700	-7.953	-6.72	8.210	-85.	120.	Martin et al. (1988)
b	Ti	2	5381.0098	1.570	-1.920	-7.956	-6.73	8.200	-200.	200.	Martin et al. (1988)
b	V	1	4111.7700	0.300	0.410	-7.610	-6.19	8.080	-110.	130.	Führ & Weise (1996)
b	V	2	3530.7600	1.070	-0.600	-7.921	-6.58	8.380	-180.	180.	Führ & Weise (1996)
b	V	2	3545.1899	1.100	-0.390	-7.922	-6.57	8.390	-150.	140.	Führ & Weise (1996)
b	V	2	3592.0300	1.100	-0.370	-7.922	-6.61	8.400	-140.	140.	Führ & Weise (1996)
b	Nd	2	3592.0620	1.503	-0.638						VALD:Kupka et al. (1999)
b	V	2	4005.7100	1.820	-0.460	-7.975	-6.64	8.360	-200.	110.	Führ & Weise (1996)
b	12CH	1	4005.7571	1.084	-3.307						Jørgensen et al. (1996)
b	Cr	1	4254.3320	0.000	-0.114	291.267	-6.24	7.410	-220.	200.	Martin et al. (1988)
b	13CH	1	4254.2441	0.522	-2.510						Jørgensen et al. (1996)
b	12CH	1	4254.3149	0.033	-3.645						Jørgensen et al. (1996)
b	Cr	1	5206.0400	0.940	0.020	-7.597	-6.15	7.720	-150.	250.	Martin et al. (1988)
b	Cr	1	5296.7002	0.980	-1.400	-7.621	-6.12	7.720	-300.	300.	Führ & Weise (1996)
b	Cr	1	5345.8101	1.000	-0.980	-7.620	-6.12	7.720	-300.	300.	Martin et al. (1988)
b	Cr	1	5348.3301	1.000	-1.290	-7.620	-6.11	7.720	-300.	300.	Führ & Weise (1996)
b	Cr	1	5409.7998	1.030	-0.720	-7.620	-6.11	7.720	-300.	300.	Martin et al. (1988)
b	Cr	2	4588.2002	4.070	-0.630	-7.963	-6.66	8.410	-175.	150.	Führ & Weise (1996)
b	Cr	2	4848.2300	3.860	-1.140	-7.934	-6.63	8.410	-200.	200.	Führ & Weise (1996)
b	12CH	1	4848.1909	3.384	-1.454						Jørgensen et al. (1996)
d	Mn	1	4030.7529	0.000	-9.999	250.245	-6.28	7.204	-150.	65.	Dummy line. Region 4030.603 - 4030.818
h	Mn	1	4030.7300	0.000	-1.037	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7441	0.000	-1.955	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7461	0.000	-1.177	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7556	0.000	-3.172	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7578	0.000	-1.779	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7593	0.000	-1.335	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7671	0.000	-2.820	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7686	0.000	-1.753	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7698	0.000	-1.519	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7754	0.000	-2.695	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7766	0.000	-1.820	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7776	0.000	-1.741	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7812	0.000	-2.774	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7820	0.000	-1.996	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
h	Mn	1	4030.7825	0.000	-2.026	250.245	-6.28	7.204		Booth et al. (1984), hfs:Lefèvre et al. (2003)	
b	12CH	1	4030.7710	1.293	-3.260						Jørgensen et al. (1996)
b	12CH	1	4030.7910	1.916	-2.283						Jørgensen et al. (1996)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{vUV}}^{1000\text{ K}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{1000\text{ K}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{Trad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	12CH	1	4030.8411	1.916	-2.299			-6.20	8.117		Jørgensen et al. (1996) VALD:Kupka et al. (1999)
b	Fe	1	4030.8899	3.694	-1.067	-7.840					Jørgensen et al. (1996)
b	12CH	1	4030.9221	0.431	-2.098	249.245	-6.28	7.173	-75.	70.	Dummy line, Region 4032.988 - 4033.133 Booth et al. (1984), hfs:Lefèvre et al. (2003)
d	Mn	1	4033.0630	0.000	-9.990	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0437	0.000	-1.200	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0457	0.000	-1.978	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0564	0.000	-1.978	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0583	0.000	-1.463	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0598	0.000	-1.815	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0686	0.000	-1.815	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0701	0.000	-1.794	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0713	0.000	-1.810	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0776	0.000	-1.810	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0789	0.000	-2.241	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0796	0.000	-1.912	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0840	0.000	-1.912	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0847	0.000	-2.940	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0852	0.000	-2.174	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
h	Mn	1	4033.0872	0.000	-2.174	249.245	-6.28	7.173			Booth et al. (1984), hfs:Lefèvre et al. (2003)
b	Ga	1	4032.9761	0.000	-0.620						VALD:Kupka et al. (1999)
b	12CH	1	4032.9890	1.197	-3.003						Jørgensen et al. (1996)
b	13CH	1	4033.0410	0.519	-2.294						Jørgensen et al. (1996)
b	12CH	1	4033.0750	1.491	-2.513						Jørgensen et al. (1996)
b	13CH	1	4033.1160	0.520	-2.292						Jørgensen et al. (1996)
b	12CH	1	4033.1321	1.197	-2.944						Jørgensen et al. (1996)
b	12CH	1	4033.1941	1.491	-2.468						Jørgensen et al. (1996)
b	C	1	4033.2280	7.946	-2.280						VALD:Kupka et al. (1999)
d	Mn	2	3460.3120	1.810	-9.999	-7.988	-6.74	8.430	-130.	160.	Dummy line, Region 3460.182 - 3460.472 Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3030	1.810	-2.071	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3032	1.810	-2.168	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3047	1.810	-2.615	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3049	1.810	-1.906	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3052	1.810	-1.804	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3079	1.810	-2.582	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3081	1.810	-1.820	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3081	1.810	-1.552	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3125	1.810	-2.728	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3125	1.810	-1.830	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3125	1.810	-1.352	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3184	1.810	-3.093	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3184	1.810	-1.184	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3460.3186	1.810	-1.996	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3482.9050	1.833	-9.999	-7.988	-6.74	8.430	-150.	45.	Dummy line, Region 3482.755 - 3482.950 Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3482.8972	1.833	-2.125	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3482.8972	1.833	-2.669	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3482.8987	1.833	-1.937	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)
h	Mn	2	3482.8989	1.833	-3.669	-7.988	-6.74	8.430			Kling & Griesmann (2000) via NBS, hfs:Holt et al (1999)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \chi_{\text{osc}}$ [mÅ]	$\delta \chi_{\text{rad}}$ [mÅ]	References and notes
h	Mn	2	3482.8992	1.833	-2.125	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9011	1.833	-1.907	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9016	1.833	-2.305	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9019	1.833	-1.937	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9048	1.833	-2.050	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9055	1.833	-1.764	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9060	1.833	-1.907	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9106	1.833	-1.406	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3482.9114	1.833	-2.050	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
b	12CH	1	3482.8999	1.198	-2.712						Jorgensen et al. (1996)
d	Mn	2	3488.6760	1.850	-9.999	-7.988	-6.74	8.430	-40.	160. Dummy line. Region 3488.636 - 3488.836	
h	Mn	2	3488.6660	1.850	-1.758	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3488.6663	1.850	-2.126	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3488.6724	1.850	-1.749	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3488.6736	1.850	-2.670	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3488.6736	1.850	-1.758	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3488.6829	1.850	-1.494	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
h	Mn	2	3488.6838	1.850	-1.749	-7.988	-6.74	8.430	Kling & Griesmann (2000) via NBS, hfs; Holt et al (1999)		
Fe	1	3865.5229	1.011	-0.950	282.266	-6.24	8.217	-200.	200. O' Brian et al. (1991)		
Fe	1	3899.7190	0.090	-1.520	-7.770	-6.32	7.150	-220.	220. O' Brian et al. (1991)		
b	12CH	1	3899.6260	1.142	-2.833						Jorgensen et al. (1996)
b	12CH	1	3899.6841	0.777	-3.312						Jorgensen et al. (1996)
Fe	1	3917.1841	0.990	-2.150	-7.695	-6.24	8.220	-160.	60. O' Brian et al. (1991)		
b	Ce	2	3917.2500	0.435	-1.195						VALD; Kupka et al. (1999)
Fe	1	3949.9590	2.180	-1.250	-7.821	-6.22	7.570	-200.	250. O' Brian et al. (1991)		
b	12CH	1	3950.0239	0.639	-3.912						Jorgensen et al. (1996)
Fe	1	4005.2419	1.557	-0.583	328.252	-6.21	8.009	-230.	230. O' Brian et al. (1991)		
Fe	1	4114.4448	2.831	-1.303	302.269	-6.16	8.097	-250.	250. O' Brian et al. (1991)		
Fe	1	4132.9082	2.850	-1.010	-7.659	-6.17	8.100	-200.	200. O' Brian et al. (1991)		
b	12CH	1	4132.8442	1.955	-1.112						Jorgensen et al. (1996)
b	12CH	1	4132.8452	2.052	-1.101						Jorgensen et al. (1996)
Fe	1	4143.8682	1.557	-0.511	317.258	-6.21	8.009	-200.	200. O' Brian et al. (1991)		
b	13CH	1	4143.9160	1.490	-2.113						Jorgensen et al. (1996)
Fe	1	4147.6748	1.490	-2.070	-7.648	-6.20	7.880	-160.	170. O' Brian et al. (1991)		
b	12CH	1	4147.7139	1.610	-2.593						Jorgensen et al. (1996)
Fe	1	4443.1938	2.858	-1.043	224.263	-6.34	7.799	-90.	160. O' Brian et al. (1991)		
b	12CH	1	4443.0059	2.517	-1.515						VALD; Kupka et al. (1999)
b	Zr	2	4443.0078	1.486	-0.330						Jorgensen et al. (1996)
b	12CH	1	4443.1699	2.589	-1.599						Jorgensen et al. (1996)
b	12CH	1	4443.2510	2.818	-1.208						Jorgensen et al. (1996)
Fe	1	4494.5630	2.198	-1.143	416.302	-6.08	8.600	-190.	190. O' Brian et al. (1991)		
b	12CH	1	4494.5190	1.848	-1.606						Jorgensen et al. (1996)
b	12CH	1	4494.5962	1.703	-1.848						Jorgensen et al. (1996)
Fe	1	4528.6138	2.176	-0.887	407.301	-6.08	8.607	-50.	220. O' Brian et al. (1991)		
b	Ce	2	4528.4731	0.864	0.075						VALD; Kupka et al. (1999)
b	12CH	1	4528.6021	2.129	-1.529						Jorgensen et al. (1996)
Fe	1	4602.9409	1.485	-2.208	296.260	-6.20	8.079	-175.	175. O' Brian et al. (1991)		

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma_{\text{vUV}}^{1000}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{1000 K}}^{1000}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{1000})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	Fe	1	4647.4302	2.950	-1.350	-7.685	-6.32	7.300	-170.	170.	Fuhr & Wiese (1996)
b	Fe	1	4872.1372	2.882	-0.567	754.235	-5.49	8.004	-90.	200.	O'Brian et al. (1991)
b	12CH	1	4872.0542	3.221	-1.477						Jorgensen et al. (1996)
b	12CH	1	4872.0952	0.488	-2.864						Jorgensen et al. (1996)
b	12CH	1	4872.2090	1.197	-3.285						Jorgensen et al. (1996)
b	Fe	1	4891.4922	2.852	-0.112	750.237	-5.49	8.009	-230.	230.	O'Brian et al. (1991)
b	12CH	1	4891.3940	1.138	-3.052						Jorgensen et al. (1996)
b	12CH	1	4891.4468	0.729	-3.390						Jorgensen et al. (1996)
b	12CH	1	4891.5332	1.394	-3.209						Jorgensen et al. (1996)
b	Fe	1	4903.3159	2.880	-0.930	-7.259	-5.49	8.000	-190.	190.	O'Brian et al. (1991)
b	12CH	1	4903.3418	3.548	-1.096						Jorgensen et al. (1996)
b	Fe	1	4918.9941	2.845	-0.342	750.237	-5.49	8.009	-350.	350.	O'Brian et al. (1991)
b	12CH	1	4918.9180	1.916	-2.739						Jorgensen et al. (1996)
b	Fe	1	4918.9541	4.154	-0.672	-7.769	-4.53	8.400			VALD:Kupka et al. (1999)
b	12CH	1	4919.0469	0.388	-3.847						Jorgensen et al. (1996)
b	Fe	1	4938.8140	2.875	-1.077	739.237	-5.49	8.009	-140.	200.	O'Brian et al. (1991)
b	12CH	1	4938.7158	0.784	-3.039						Jorgensen et al. (1996)
b	Fe	1	4966.0898	3.330	-0.840	-7.218	-5.30	7.810	-190.	190.	Fuhr & Wiese (1996)
b	Fe	1	4994.1382	0.920	-2.970	-7.744	-6.25	7.180	-190.	190.	Jorgensen et al. (1996)
b	Fe	1	5001.8701	3.880	0.050	-7.273	-5.38	7.970	-200.	200.	Bridges & Komblith (1974)
b	Fe	1	5014.9399	3.940	-0.270	-7.268	-5.54	7.960	-170.	170.	Fuhr & Wiese (1996)
b	Fe	1	5051.6401	0.920	-2.760	-7.746	-6.25	7.190	-200.	200.	O'Brian et al. (1991)
b	Fe	1	5068.7710	2.940	-1.040	-7.265	-5.49	8.010	-200.	200.	O'Brian et al. (1991)
b	Fe	1	5150.8521	0.990	-3.040	-7.742	-6.25	7.180	-70.	160.	O'Brian et al. (1991)
b	Fe	1	5166.2842	0.000	-4.120	-7.826	-6.33	3.540	-180.	180.	O'Brian et al. (1991)
b	Fe	1	5171.6099	1.490	-1.720	-7.687	-6.20	6.330	-210.	220.	O'Brian et al. (1991)
b	Fe	1	5192.3530	3.000	-0.420	-7.266	-5.49	8.010	-150.	150.	O'Brian et al. (1991)
b	Fe	1	5194.9492	1.560	-2.020	-7.680	-6.20	6.290	-200.	170.	O'Brian et al. (1991)
b	Fe	1	5202.3398	2.180	-1.840	-7.603	-6.18	8.230	-200.	200.	Fuhr & Wiese (1996)
b	Fe	1	5202.2510	4.256	-0.639	-7.765	-6.01	8.320			VALD:Kupka et al. (1999)
b	Fe	1	5216.2832	1.610	-2.080	-7.674	-6.20	6.220	-200.	200.	O'Brian et al. (1991)
b	Fe	1	5266.5630	3.000	-0.390	-7.273	-5.49	8.010	-225.	70.	O'Brian et al. (1991)
b	Si	1	5266.6602	5.619	-1.010						VALD:Kupka et al. (1999)
b	12CH	1	5266.7271	1.432	-2.600						Jorgensen et al. (1996)
b	Fe	1	5281.7979	3.040	-0.830	-7.266	-5.49	8.010	-200.	200.	O'Brian et al. (1991)
b	Fe	1	5283.6289	3.240	-0.520	-7.221	-5.45	7.880	-210.	200.	O'Brian et al. (1991)
b	Fe	1	5324.1909	3.211	-1.03	-7.235	-5.50	7.880	-230.	230.	Bard et al. (1991)
b	Fe	1	5328.5420	1.560	-1.850	-7.686	-6.23	6.850	-50.	200.	O'Brian et al. (1991)
b	Cr	1	5328.3770	2.914	0.460	-7.225	-3.97	8.030			VALD:Kupka et al. (1999)
b	Fe	1	5339.9370	3.270	-0.720	-7.221	-5.45	7.870	-200.	200.	O'Brian et al. (1991)
b	Fe	1	5341.0239	1.608	-1.953	-7.680	-6.24	6.830	-200.	200.	VALD:Kupka et al. (1999)
b	Fe	1	5364.8701	4.450	0.230	-7.136	-5.08	8.270	-200.	200.	Fuhr & Wiese (1996)
b	Fe	1	5367.4702	4.410	0.440	-7.153	-5.13	8.270	-220.	220.	Fuhr & Wiese (1996)
b	Fe	1	5369.9741	4.370	0.540	-7.179	-5.06	8.260	-220.	180.	O'Brian et al. (1991)
b	Fe	1	5383.3799	4.310	0.650	-7.219	-5.18	8.250	-230.	230.	O'Brian et al. (1991)
b	Fe	1	5393.1758	3.240	-0.720	-7.235	-5.50	7.870	-200.	60.	Bard et al. (1991) Sneden 2003, add
b	Fe	1	5410.9102	4.470	0.400	-7.132	-5.06	8.210	-200.	200.	Fuhr & Wiese (1996)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{radW}}/N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarK}}/N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta\chi_{\text{osc}}$ [mÅ]	$\delta\chi_{\text{rad}}$ [mÅ]	References and notes
b	Fe	1	5415.2002	4.390	0.640	-7.182	-4.76	8.240	-220.	220.	Führ & Weise (1996)
	Fe	1	5429.7061	0.960	-1.880	-7.755	-6.30	7.160	-300.	300.	O'Brian et al. (1991)
	Fe	1	5429.8271	4.473	-0.527	-7.783	-5.08	8.210			VALD; Kupka et al. (1999)
	Fe	1	5434.5342	1.010	-2.130	-7.749	-6.30	7.150	-230.	230.	O'Brian et al. (1991)
	Fe	1	5445.0400	4.390	0.040	-7.189	-4.58	8.240	-190.	190.	Führ & Weise (1996)
	Fe	1	5446.9238	0.990	-1.910	-7.753	-6.30	7.150	-250.	250.	O'Brian et al. (1991)
	Fe	1	5506.7910	0.990	-2.790	-7.753	-6.30	7.160	-200.	190.	O'Brian et al. (1991)
	Fe	1	5586.7710	3.370	-0.100	-7.221	-5.45	7.880	-240.	240.	Bard et al. (1991)
	Fe	1	5615.6582	3.330	0.050	-7.234	-5.50	7.890	-250.	250.	Bard et al. (1991)
	Fe	1	6136.6240	2.450	-1.410	-7.609	-6.33	7.880	-230.	230.	O'Brian et al. (1991)
	Fe	1	6137.7021	2.590	-1.350	-7.589	-6.11	8.010	-230.	230.	O'Brian et al. (1991)
b	Fe	1	6191.5708	2.430	-1.420	-7.615	-6.33	7.910	-220.	220.	O'Brian et al. (1991)
	Fe	1	6393.6118	2.430	-1.570	-7.622	-6.31	7.970	-220.	220.	O'Brian et al. (1991)
	Fe	1	6400.0088	3.600	-0.290	-7.232	-5.50	7.900	-250.	250.	Bard et al. (1991)
	Fe	1	6430.8560	2.180	-1.950	-7.704	-6.19	8.220	-220.	220.	O'Brian et al. (1991)
	Fe	1	7511.0200	4.178	0.099	-7.626	-5.30	8.220	-240.	240.	VALD; Kupka et al. (1999)
	Fe	1	8387.7725	2.176	-1.493	-7.727	-6.24	7.290	-300.	300.	VALD; Kupka et al. (1999)
	Fe	2	4508.2891	2.856	-2.312	-7.950	-6.67	8.617	-30.	170.	Biemont et al. (1991)
	b	12CH	1	4508.1909	3.278	-1.040					Jørgensen et al. (1996)
	b	12CH	1	4508.3281	3.278	-1.049					Schnabel et al. (2004)
	b	12CH	1	4515.3428	2.844	-2.362	-7.950	-6.60	8.487	-120.	100.
b	Fe	2	4515.3320	0.680	-3.746						Schnabel et al. (2004)
	Fe	1	4515.3398	0.680	-2.666						Jørgensen et al. (1996)
	Fe	2	4520.2241	2.807	-2.550	-7.950	-6.60	8.491	-160.	40.	Möity (1983)
	b	12CH	1	4520.2139	2.060	-1.896					Jørgensen et al. (1996)
	Fe	2	4923.9268	2.891	-1.206	-7.910	-6.58	8.489	-100.	240.	Schnabel et al. (2004)
	Fe	2	5197.5771	3.230	-2.100	-7.824	-6.60	8.480	-200.	200.	VALD; Kupka et al. (1999)
	b	Fe	2	5197.5679	10.398	-2.116	-7.824	-5.84	9.050		VALD; Kupka et al. (1999)
	Fe	2	5234.6299	3.220	-2.270	-7.946	-6.60	8.490	-200.	200.	Heise & Kock (1990), Möity (1983)
	Fe	2	5276.0020	3.200	-1.960	-7.946	-6.60	8.490	-200.	200.	Kroll & Kock (1987), Möity (1983)
	Fe	2	5284.1001	2.890	-3.010	-7.914	-6.60	8.530	-160.	160.	Führ & Wiese (1996)
d	Co	1	3412.3330	0.514	-9.999	-7.666	-6.30	8.030	-80.	160.	Dummy line. Region 3412.253 - 3412.493
	Co	1	3412.3242	0.514	-1.513	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3242	0.514	-1.309	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3250	0.514	-1.776	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3252	0.514	-1.139	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3271	0.514	-0.993	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3328	0.514	-2.408	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3326	0.514	-1.709	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3335	0.514	-0.748	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3357	0.514	-1.361	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
h	Co	1	3412.3374	0.514	-2.253	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3376	0.514	-0.643	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)
	Co	1	3412.3401	0.514	-1.367	-7.666	-6.30	8.030			Cardon et al. (1982), his Pickering (1996)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma_{\text{vUV}}^{10000\text{ K}/N_H})$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{10000 K}/N_e}^{\text{Star}})$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{Irad}})$ [rad.s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
h	Co	1	3412.3425	0.514	-2.253	-7.666	-6.30	8.030			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3412.3455	0.514	-1.440	-7.666	-6.30	8.030			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3412.3486	0.514	-2.350	-7.666	-6.30	8.030			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3412.3516	0.514	-1.651	-7.666	-6.30	8.030			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3412.3557	0.514	-2.554	-7.666	-6.30	8.030			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3412.3635	0.514	-2.952	-7.666	-6.30	8.030			Cardon et al. (1982), hfs Pickering (1996)
d	Co	1	3489.3989	0.923	-9.999	-7.610	-6.27	8.220	-120.	160.	Dummy line. Region 3489.279 - 3489.559
h	Co	1	3489.3804	0.923	-2.434	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3845	0.923	-2.024	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3879	0.923	-1.802	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3899	0.923	-1.320	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3911	0.923	-1.677	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3923	0.923	-1.124	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3938	0.923	-1.626	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3943	0.923	-1.070	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3960	0.923	-1.092	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3960	0.923	-1.656	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3970	0.923	-1.183	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.3977	0.923	-1.371	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.4006	0.923	-0.480	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.4009	0.923	-1.626	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.4019	0.923	-0.621	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.4019	0.923	-1.238	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.4023	0.923	-0.785	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
h	Co	1	3489.4023	0.923	-0.983	-7.610	-6.27	8.220			Cardon et al. (1982), hfs Pickering (1996)
d	Co	1	3894.0730	1.049	-9.990	310.267	-6.27	7.972	-80.	170.	Dummy line. Region 3893.993 - 3894.243
h	Co	1	3894.0576	1.049	-1.298	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0576	1.049	-1.686	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0596	1.049	-1.043	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0632	1.049	-0.845	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0652	1.049	-1.431	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0688	1.049	-1.243	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0688	1.049	-1.716	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0745	1.049	-1.152	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0762	1.049	-0.540	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0764	1.049	-1.686	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0818	1.049	-1.130	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0857	1.049	-1.737	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0913	1.049	-1.184	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.0969	1.049	-1.862	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.1025	1.049	-1.380	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	3894.1101	1.049	-2.084	310.267	-6.27	7.972			Nitz et al. (1999), hfs Pickering (1996)
b	12CH	1	3894.0049	1.068	-3.582						Jorgensen et al. (1996)
d	Co	1	4121.3110	0.922	-9.999	257.242	-6.27	7.167	-45.	170.	Dummy line. Region 4121.266 - 4121.481
h	Co	1	4121.2939	0.922	-0.973	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3008	0.922	-1.078	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3071	0.922	-1.194	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \chi_{\text{osc}}$ [mÅ]	$\delta \chi_{\text{rad}}$ [mÅ]	References and notes
h	Co	1	4121.3130	0.922	-1.323	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3164	0.922	-1.981	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3184	0.922	-1.469	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3208	0.922	-1.770	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3228	0.922	-1.639	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3242	0.922	-1.697	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3267	0.922	-1.843	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3271	0.922	-1.691	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3296	0.922	-1.738	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3301	0.922	-2.106	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3311	0.922	-1.843	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3325	0.922	-2.039	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3359	0.922	-3.282	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3369	0.922	-2.738	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3374	0.922	-2.884	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3379	0.922	-2.583	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3379	0.922	-2.680	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Co	1	4121.3384	0.922	-2.583	257.242	-6.27	7.167			Nitz et al. (1999), hfs Pickering (1996)
h	Ni	1	3519.7581	0.275	-1.407	-7.689	-6.29	7.870	-80.	180.	VALD: Kupka et al. (1999)
Ni	1	3664.0850	0.275	-2.115	-7.710	-6.30	8.010	-130.	130.	VALD: Kupka et al. (1999)	
Ni	1	3775.5649	0.423	-1.408	280.265	-6.30	7.687	-70.	200.	Blackwell et al. (1989)	
Ni	1	3783.5239	0.423	-1.304	227.253	-6.30	7.083	-70.	160.	Huber & Sandeman (1980)	
Ni	1	3807.1379	0.423	-1.220	278.263	-6.30	7.719	-200.	50.	Blackwell et al. (1989)	
Ni	1	3858.2920	0.423	-0.951	274.260	-6.30	7.855	-200.	65.	Blackwell et al. (1989)	
b	12CH	1	3858.3501	0.431	-4.033	-7.777	-6.30	8.010	-130.	Jorgensen et al. (1996)	
Ni	1	5035.3701	3.630	0.290	-7.231	-5.23	8.190	-200.	200.	Fuhr & Weisse (1996)	
Ni	1	5476.9209	1.830	-0.890	-7.781	-6.13	8.200	-85.	130.	Fuhr et al. (1988)	
d	Cu	1	3247.5300	0.000	-9.999				-40.	120.	Dummy line. Region 3247.490 - 3247.65
h	Cu	1	3247.5100	0.000	-1.379						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5120	0.000	-1.028						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5120	0.000	-1.379						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5129	0.000	-1.028						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5129	0.000	-1.777						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5139	0.000	-1.426						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5510	0.000	-0.581						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5520	0.000	-0.932						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5530	0.000	-1.028						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5540	0.000	-1.727						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5540	0.000	-1.379						Fuhr & Weisse (1996), hfs: Kunucz (?)
h	Cu	1	3247.5549	0.000	-2.078						Fuhr & Weisse (1996), hfs: Kunucz (?)
b	Co	1	3247.1489	3.067	-0.824	-7.766	-5.23	8.260			VALD: Kupka et al. (1999)
b	Co	1	3247.1721	1.883	-0.024	-7.830	-6.27	8.370			VALD: Kupka et al. (1999)
b	Fe	2	3247.1750	3.889	-1.220	-7.945	-6.64	8.520			VALD: Kupka et al. (1999)
b	Fe	1	3247.2100	2.608	-1.728	-7.842	-6.32	7.780			VALD: Kupka et al. (1999)
b	Fe	1	3247.2791	2.469	-1.265	-7.756	-5.78	8.000			VALD: Kupka et al. (1999)
b	Fe	2	3247.3889	4.154	-2.150	-7.945	-6.64	8.530			VALD: Kupka et al. (1999)
b	Tm	2	3247.4651	1.087	-0.010						VALD: Kupka et al. (1999)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Stark}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	Nb	2	3247.4709	0.980	-0.280						VALD: Kupka et al. (1999)
b	Fe	1	3248.2041	2.449	-0.673	-7.257		-5.48	8.000		VALD: Kupka et al. (1999)
b	13CH	1	3247.2749	1.000	-2.766						Jorgensen et al. (1996)
b	13CH	1	3247.4041	0.910	-2.181						Jorgensen et al. (1996)
b	12CH	1	3247.6250	0.360	-5.513						Jorgensen et al. (1996)
b	13CH	1	3247.7100	0.700	-1.828						Jorgensen et al. (1996)
b	12CH	1	3247.7410	0.910	-1.227						Jorgensen et al. (1996)
b	12CH	1	3247.7439	0.360	-5.416						Jorgensen et al. (1996)
b	12CH	1	3247.8169	1.000	-1.812						Jorgensen et al. (1996)
b	13CH	1	3247.8750	1.030	-2.355						Jorgensen et al. (1996)
d	Cu	1	3273.9500	0.000	-9.999						-170. Dummy line. Region 3273.78 - 3274.12
h	Cu	1	3273.9250	0.000	-1.375						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9270	0.000	-1.024						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9290	0.000	-2.074						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9299	0.000	-1.723						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9690	0.000	-1.024						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9700	0.000	-1.375						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9719	0.000	-1.024						Fuhr & Weise (1996), hfs: Kurucz (?)
h	Cu	1	3273.9729	0.000	-1.375						Fuhr & Weise (1996), hfs: Kurucz (?)
b	Zr	2	3273.0669	0.164	-0.300						VALD: Kupka et al. (1999)
b	Nd	2	3273.2529	0.205	-1.384						VALD: Kupka et al. (1999)
b	Sm	2	3273.3169	0.434	-1.464						VALD: Kupka et al. (1999)
b	Er	2	3273.3210	0.892	-0.945						VALD: Kupka et al. (1999)
b	Sm	2	3273.4800	0.000	-1.083						VALD: Kupka et al. (1999)
b	Fe	2	3273.4741	9.069	-3.253	-7.815		-5.85	9.080		VALD: Kupka et al. (1999)
b	Fe	2	3273.4900	4.154	-2.320	-7.945		-6.61	8.640		VALD: Kupka et al. (1999)
b	Sc	1	3273.6311	0.021	0.257	-7.479		-6.04	8.480		VALD: Kupka et al. (1999)
b	Hf	2	3273.6489	0.787	-0.960						VALD: Kupka et al. (1999)
b	Co	1	3273.9360	3.067	-0.495	-7.782		-5.15	8.160		VALD: Kupka et al. (1999)
b	Co	1	3274.0061	3.016	-0.736	-7.765		-5.07	8.130		VALD: Kupka et al. (1999)
b	Fe	1	3274.0259	2.588	-2.687	-7.855		-6.32	7.430		VALD: Kupka et al. (1999)
b	Ti	1	3274.0449	1.460	-1.119	-7.708		-4.69	8.200		VALD: Kupka et al. (1999)
b	Gd	2	3274.1860	1.288	-0.138						VALD: Kupka et al. (1999)
b	Fe	1	3274.2310	2.223	-3.548	-7.806		-6.18	7.890		VALD: Kupka et al. (1999)
b	Ti	2	3274.4209	1.165	-1.361	-7.919		-6.66	8.380		VALD: Kupka et al. (1999)
b	12CH	1	3273.6399	0.640	-5.106						Jorgensen et al. (1996)
b	12CH	1	3273.6460	0.640	-5.044						Jorgensen et al. (1996)
b	12CH	1	3274.1470	0.270	-5.536						Jorgensen et al. (1996)
b	12CH	1	3274.2920	0.270	-5.423						Jorgensen et al. (1996)
b	Sr	1	4607.3398	0.000	0.283	404.257		-100.	100.		Migdalek & Baylis (1987)
b	12CH	1	4607.3120	3.467	-0.866						Jorgensen et al. (1996)
b	12CH	1	4607.3159	1.795	-2.574						Jorgensen et al. (1996)
b	12CH	1	4607.3691	1.795	-2.589						Jorgensen et al. (1996)
b	12CH	1	4607.3970	3.303	-1.391						Jorgensen et al. (1996)
b	Sr	2	3464.4600	3.040	0.530						Sneden et al. (2003)
b	12CH	1	3464.3970	0.488	-2.893						Jorgensen et al. (1996)
d	Sr	2	4077.7241	0.000	-9.999	262.208		-150.	140.		Dummy line. Region 4077.574 - 4077.864

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma_{1000 \text{ K}}^{\text{radW}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000 \text{ K}}^{\text{Sark}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{rad}})$ [rad/s]	$\delta_{\text{obs}}$ [mÅ]	$\delta_{\text{pred}}$ [mÅ]	References and notes
i	84Sr	2	4077.7219	0.000	-2.094	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
i	86Sr	2	4077.7231	0.000	-0.848	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
i	87Sr	2	4077.7112	0.000	-2.219	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
h	Sr	2	4077.7112	0.000	-1.779	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
h	Sr	2	4077.7112	0.000	-1.485	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
h	Sr	2	4077.7390	0.000	-1.754	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
h	Sr	2	4077.7390	0.000	-1.779	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
h	Sr	2	4077.7390	0.000	-1.956	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
i	88Sr	2	4077.7241	0.000	0.075	262.208					Pinnington et al. (1995), isot&hfis;Borghs et al. (1983)
i	Sr	2	4161.7900	2.940	-0.600						
d	Sr	2	4215.5400	0.000	-9.999	262.205					
i	84Sr	2	4215.5420	0.000	-2.406	262.205					
i	86Sr	2	4215.5410	0.000	-1.161	262.205					
i	87Sr	2	4215.5239	0.000	-1.791	262.205					
h	Sr	2	4215.5293	0.000	-1.967	262.205					
h	Sr	2	4215.5537	0.000	-2.231	262.205					
h	Sr	2	4215.5591	0.000	-1.791	262.205					
h	Sr	2	4215.5400	0.000	-0.238	262.205					
b	12CH	1	4215.3809	0.458	-4.114						
b	Fe	1	4215.4229	2.990	-1.756	-7.865	-6.31	7.286			
b	Fe	1	4215.4590	2.759	-3.882	-7.857	-6.24	7.888			
b	12CH	1	4215.5068	1.862	-1.796						
b	12CH	1	4215.5132	1.862	-1.716						
Y	2	3242.2700	0.180	0.210							
Y	2	3327.8799	0.410	0.130							
Y	2	3549.0100	0.130	-0.280							
Y	2	3600.7400	0.180	0.280							
Y	2	3611.0400	0.130	0.010							
b	12CH	1	3611.0061	0.034	-4.405						
Y	2	3774.3311	0.130	0.210							
Y	2	3788.6941	0.104	-0.070							
Y	2	3950.3521	0.104	-0.490							
b	12CH	1	3950.4270	1.368	-2.877						
b	Ce	2	3950.4290	0.635	-0.603						
Y	2	4883.6841	1.084	0.070							
b	12CH	1	4883.6948	1.080	-3.173						
Y	2	5087.4302	1.080	-0.170							
Y	2	5205.7300	1.030	-0.340							
Zr	2	3356.0879	0.095	-0.513							
Zr	2	3430.5300	0.470	-0.160							
b	12CH	1	3430.5330	1.055	-3.007						
Zr	2	3438.2300	0.090	0.420							
b	12CH	1	3438.3049	0.709	-3.358						
Zr	2	3457.5601	0.560	-0.530							
b	12CH	1	3457.6589	0.408	-3.055						
Zr	2	3458.9299	0.960	-0.520							
Zr	2	3479.3899	0.710	0.170							

Table 2. Continued.

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{vUV}}^{\text{StarK}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{Irad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	Zr	2	3505.66699	0.160	-0.360				-80.	110.	Biemont et al. (1981) Jorgensen et al. (1996)
b	12CH	1	3505.7251	0.910	-2.998				-210.	150.	Biemont et al. (1981)
b	Zr	2	3551.9600	0.090	-0.310				-110.	140.	Biemont et al. (1981)
b	Zr	2	3573.0801	0.320	-1.040				-90.	140.	Biemont et al. (1981)
b	Zr	2	3766.8201	0.410	-0.810				-140.	130.	Biemont et al. (1981)
b	Zr	2	3998.9700	0.560	-0.670				-80.	VALD:Kupka et al. (1999)	
b	Zr	2	4029.6841	0.713	-0.597				-80.	VALD:Kupka et al. (1999)	
b	12CH	1	4029.6040	1.263	-3.186				-85.	100.	Jorgensen et al. (1996)
b	12CH	1	4029.6379	1.723	-2.337				-200.	200.	Biemont et al. (1981)
b	12CH	1	4029.7000	1.723	-2.353				-100.	80.	Jorgensen et al. (1996)
b	12CH	1	4029.7290	1.138	-3.123				-120.	20.	Jorgensen et al. (1996)
b	Zr	2	4050.3301	0.710	-1.000				-100.	80.	Biemont et al. (1981)
b	Zr	2	4090.5100	0.760	-1.100				-120.	20.	Biemont et al. (1981)
b	12CH	1	4090.4109	1.569	-2.234				-120.	20.	Jorgensen et al. (1996)
b	12CH	1	4090.4580	1.394	-2.983				-120.	20.	Jorgensen et al. (1996)
b	12CH	1	4090.4961	1.569	-2.188				-120.	20.	Jorgensen et al. (1996)
b	Zr	2	4161.2129	0.713	-0.720				-100.	80.	Biemont et al. (1981)
b	12CH	1	4161.1802	2.761	-1.021				-100.	80.	Jorgensen et al. (1996)
b	12CH	1	4161.1841	1.917	-1.420				-100.	80.	Jorgensen et al. (1996)
b	Zr	2	4208.9849	0.713	-0.460				-100.	80.	Biemont et al. (1981)
b	12CH	1	4208.9082	1.986	-2.383				-100.	80.	Jorgensen et al. (1996)
b	12CH	1	4208.9771	1.986	-2.400				-100.	80.	Jorgensen et al. (1996)
d	Fe	2	3281.3040	1.040	-2.990	-7.770	-7.770	-6.22	7.890	-684.	-584. Dummy line. Region 3280.620 - 3280.720
h	Ag	1	3280.6770	0.000	-0.480						Führ & Weise (1996), hfs: Führ & Aller (1972)
h	Ag	1	3280.6780	0.000	-0.460						Führ & Weise (1996), hfs: Führ & Aller (1972)
h	Ag	1	3280.6831	0.000	-0.940						Führ & Weise (1996), hfs: Führ & Aller (1972)
h	Ag	1	3280.6851	0.000	-0.960						Führ & Weise (1996), hfs: Führ & Aller (1972)
b	Ce	2	3280.4810	0.553	-0.673						VALD:Kupka et al. (1999)
b	Zr	2	3280.7351	0.710	-1.100						VALD:Kupka et al. (1999)
b	Fe	1	3280.7471	3.018	-2.528	-7.837	-7.837	-6.32	7.900		VALD:Kupka et al. (1999)
b	Mn	1	3280.7571	2.143	-2.232	-7.781	-7.781	-5.64	6.840		VALD:Kupka et al. (1999)
b	14NH	1	3280.8440	0.104	-1.209						Kurucz CDROM 15 (1993)
b	14NH	1	3280.33770	1.956	-2.982						Kurucz CDROM 15 (1993)
b	14NH	1	3280.4460	2.043	-3.542	-7.541	-7.541	-5.41	7.490		Kurucz CDROM 15 (1993)
b	14NH	1	3280.5459	2.043	-3.600						Kurucz CDROM 15 (1993)
b	14NH	1	3280.5801	1.369	-4.289						Kurucz CDROM 15 (1993)
b	14NH	1	3280.7009	1.711	-3.639						Kurucz CDROM 15 (1993)
b	14NH	1	3280.7151	1.369	-4.240						Kurucz CDROM 15 (1993)
b	14NH	1	3280.7590	1.711	-1.373						Kurucz CDROM 15 (1993)
b	14NH	1	3280.8589	1.612	-1.386						Kurucz CDROM 15 (1993)
b	14NH	1	3280.8740	1.817	-3.646						Kurucz CDROM 15 (1993)
b	14NH	1	3280.8740	1.956	-4.177						Kurucz CDROM 15 (1993)
b	14NH	1	3280.8950	1.817	-6.989						Kurucz CDROM 15 (1993)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{Sark}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}} / N_e)$ [rad/s]	$\delta \chi_{\text{stat}}$ [mÅ]	$\delta \chi_{\text{rad}}$ [mÅ]	References and notes
b	14NH	1	3280.9199	1.956	-2.960						Kurucz CDROM 15 (1993)
b	14NH	1	3280.9290	1.711	-3.656						Kurucz CDROM 15 (1993)
b	14NH	1	3280.9390	1.612	-3.649						Kurucz CDROM 15 (1993)
b	14NH	1	3280.9570	1.711	-1.388						Kurucz CDROM 15 (1993)
b	14NH	1	3280.9570	1.612	-1.416						Kurucz CDROM 15 (1993)
b	14NH	1	3280.9570	1.711	-1.402						Kurucz CDROM 15 (1993)
b	14NH	1	3280.9570	1.612	-1.402						Kurucz CDROM 15 (1993)
b	14NH	1	3281.0000	1.817	-1.360						Kurucz CDROM 15 (1993)
d	Ag	1	3382.8999	0.000	-9.990					-20.	Dummy line. Region 3382.88 - 3382.95
h	Ag	1	3382.8979	0.000	-0.770						Führ & Weise (1996), his; Ross & Aller (1972)
h	Ag	1	3382.8999	0.000	-0.760						Führ & Weise (1996), his; Ross & Aller (1972)
h	Ag	1	3382.9050	0.000	-1.250						Führ & Weise (1996), his; Ross & Aller (1972)
h	Ag	1	3382.9080	0.000	-1.270						VALD:Kupka et al. (1999)
b	Cr	2	3382.6750	2.455	-6.339	-7.930	-6.63	8.370			VALD:Kupka et al. (1999)
b	Cr	2	3382.7839	4.750	-1.929	-7.926	-6.60	8.580			VALD:Kupka et al. (1999)
b	Fe	1	3382.8081	3.018	-2.320	-7.676	-5.27	7.650			VALD:Kupka et al. (1999)
b	Fe	1	3382.7839	4.750	-1.929	-7.842	-6.29	8.340			VALD:Kupka et al. (1999)
b	Co	1	3382.9551	2.080	-1.577	-7.906	-6.56	8.360			VALD:Kupka et al. (1999)
b	Ti	2	3383.5691	1.231	-1.845	-7.921	-6.19	7.850			VALD:Kupka et al. (1999)
b	Ce	2	3383.6760	0.529	-0.344	-7.921	-6.56	8.280			VALD:Kupka et al. (1999)
b	Fe	1	3383.6919	2.198	-1.400	-7.921	-6.19	7.850			VALD:Kupka et al. (1999)
b	Ti	2	3383.7681	0.000	0.142	-7.906	-6.56	8.280			VALD:Kupka et al. (1999)
b	14NH	1	3382.6550	1.401	-1.204						Kurucz CDROM 15 (1993)
b	14NH	1	3382.6831	1.401	-3.544						Kurucz CDROM 15 (1993)
b	14NH	1	3382.6970	1.401	-1.225						Kurucz CDROM 15 (1993)
b	14NH	1	3382.7361	2.039	-3.045						Kurucz CDROM 15 (1993)
b	14NH	1	3382.7539	1.859	-6.602						Kurucz CDROM 15 (1993)
b	14NH	1	3382.8201	2.039	-3.045						Kurucz CDROM 15 (1993)
b	14NH	1	3382.8560	1.401	-3.529						Kurucz CDROM 15 (1993)
b	14NH	1	3382.8689	0.779	-2.309						Kurucz CDROM 15 (1993)
b	14NH	1	3382.8860	0.759	-2.718						Kurucz CDROM 15 (1993)
b	14NH	1	3382.9670	0.779	-2.888						Kurucz CDROM 15 (1993)
b	14NH	1	3382.9951	0.759	-2.875						Kurucz CDROM 15 (1993)
b	14NH	1	3383.0640	1.105	-5.648						Kurucz CDROM 15 (1993)
b	14NH	1	3383.1179	1.179	-4.836						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
b	14NH	1	3383.1311	0.759	-3.322						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
b	14NH	1	3383.1599	1.179	-2.100						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
b	14NH	1	3383.2161	0.426	-3.053						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
b	14NH	1	3383.2410	1.179	-3.083						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
b	14NH	1	3383.2971	0.426	-2.536						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
d	Ba	2	4166.0030	2.720	-9.990					-210.	Dummy line. Region 4165.793 - 4166.203
i	134Ba	2	4166.0030	2.720	-2.039						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
i	135Ba	2	4166.0007	2.720	-2.805						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
i	135Ba	2	4166.0013	2.720	-2.805						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
i	135Ba	2	4166.0013	2.720	-3.203						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
i	135Ba	2	4166.0013	2.720	-2.600						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
i	135Ba	2	4166.0025	2.720	-2.600						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)
i	135Ba	2	4166.0025	2.720	-2.504						Gallagher (1967), isot&hfs lower level only; Villemoes et al. (1993)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{vUV}}^{\text{Star}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000 \text{ K}}^{\text{Star}} / N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{Irad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
i	135Ba	2	4166.0025	2.720	-2.658						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	135Ba	2	4166.0044	2.720	-2.658						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	135Ba	2	4166.0044	2.720	-2.056						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	136Ba	2	4166.0030	2.720	-1.522						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0005	2.720	-2.571						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0011	2.720	-2.571						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0011	2.720	-2.969						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0011	2.720	-2.367						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0025	2.720	-2.367						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0025	2.720	-2.270						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0025	2.720	-2.425						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0045	2.720	-2.425						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4166.0045	2.720	-1.823						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
i	138Ba	2	4166.0029	2.720	-0.563						Gallagher (1967), isot&hfs lower level only: Villemoes et al. (1993)
d	Ba	2	4524.9250	2.512	-9.999						-230. 75. Dummy line. Region 4165.7730 - 4166.0780
i	134Ba	2	4524.9280	2.512	-1.979						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	135Ba	2	4524.9223	2.512	-2.745						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	135Ba	2	4524.9223	2.512	-2.046						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	135Ba	2	4524.9314	2.512	-2.046						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	135Ba	2	4524.9314	2.512	-2.046						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	136Ba	2	4524.9280	2.512	-1.462						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4524.9217	2.512	-2.511						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4524.9217	2.512	-1.812						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4524.9318	2.512	-1.812						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	137Ba	2	4524.9318	2.512	-1.812						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
i	138Ba	2	4524.9280	2.512	-0.504						VALD: Kupka et al. (1999), isot&hfs lower level only: Villemoes et al. (1993)
b	12CH	1	4524.8711	1.862	-1.796						Jorgensen et al. (1996)
b	12CH	1	4524.8770	1.862	-1.803						Jorgensen et al. (1996)
d	Ba	2	4553.9878	0.000	-9.999			303.222			-825. 820. Dummy line. Region 4553.1630 - 4554.8080
i	134Ba	2	4554.0000	0.000	-1.449			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	135Ba	2	4553.9690	0.000	-1.816			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	135Ba	2	4553.9700	0.000	-1.816			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	135Ba	2	4553.9710	0.000	-2.214			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	135Ba	2	4554.0170	0.000	-1.369			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	135Ba	2	4554.0200	0.000	-1.816			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	135Ba	2	4554.0210	0.000	-2.514			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	136Ba	2	4554.0000	0.000	-0.932			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	137Ba	2	4553.9650	0.000	-1.583			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	137Ba	2	4553.9670	0.000	-1.583			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	137Ba	2	4553.9680	0.000	-1.981			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	137Ba	2	4554.0200	0.000	-1.135			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	137Ba	2	4554.0220	0.000	-1.583			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	138Ba	2	4554.0230	0.000	-2.281			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
i	138Ba	2	4554.0000	0.000	0.026			303.222			Gallagher (1967), isot&hfs: McWilliam (1998)
b	12CH	1	4554.0791	3.056	-1.171						Jorgensen et al. (1996)
b	12CH	1	4554.1001	1.853	-2.642						Jorgensen et al. (1996)
d	L <sub>a</sub>	2	3949.1001	0.400	-9.990						-180. 200. Dummy line. Region 3948.92 - 3949.30

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{Flower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{radW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{StarK}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{std}}$ [mÅ]	$\delta \lambda_{\text{std}}$ [mÅ]	References and notes
h	La	2	3949.0359	0.400	-1.337						Lawler et al. (2001), hfs; the same
h	La	2	3949.0369	0.400	-1.191						Lawler et al. (2001), hfs; the same
h	La	2	3949.0425	0.400	-0.995						Lawler et al. (2001), hfs; the same
h	La	2	3949.0442	0.400	-1.008						Lawler et al. (2001), hfs; the same
h	La	2	3949.0452	0.400	-1.668						Lawler et al. (2001), hfs; the same
h	La	2	3949.0540	0.400	-0.761						Lawler et al. (2001), hfs; the same
h	La	2	3949.0564	0.400	-0.886						Lawler et al. (2001), hfs; the same
h	La	2	3949.0579	0.400	-1.559						Lawler et al. (2001), hfs; the same
h	La	2	3949.0703	0.400	-0.576						Lawler et al. (2001), hfs; the same
h	La	2	3949.0732	0.400	-0.825						Lawler et al. (2001), hfs; the same
h	La	2	3949.0757	0.400	-1.580						Lawler et al. (2001), hfs; the same
h	La	2	3949.0918	0.400	-0.420						Lawler et al. (2001), hfs; the same
h	La	2	3949.0952	0.400	-0.821						Lawler et al. (2001), hfs; the same
h	La	2	3949.0981	0.400	-1.690						Lawler et al. (2001), hfs; the same
h	La	2	3949.1179	0.400	-0.284						Lawler et al. (2001), hfs; the same
h	La	2	3949.1223	0.400	-0.887						Lawler et al. (2001), hfs; the same
h	La	2	3949.1257	0.400	-1.902						Lawler et al. (2001), hfs; the same
h	La	2	3949.1494	0.400	-0.163						Lawler et al. (2001), hfs; the same
h	La	2	3949.1543	0.400	-1.092						Lawler et al. (2001), hfs; the same
h	La	2	3949.1584	0.400	-2.305						Lawler et al. (2001), hfs; the same
d	La	2	3988.5100	0.403	-9.999						Dummy line. Region 3988.310 - 3988.710
h	La	2	3988.4402	0.403	-1.362						Lawler et al. (2001), hfs; the same
h	La	2	3988.4409	0.403	-1.839						Lawler et al. (2001), hfs; the same
h	La	2	3988.4473	0.403	-1.140						Lawler et al. (2001), hfs; the same
d	La	2	3988.4485	0.403	-9.999						Dummy - a zero component kept for bookkeeping
h	La	2	3988.4492	0.403	-1.362						Lawler et al. (2001), hfs; the same
h	La	2	3988.4595	0.403	-1.041						Lawler et al. (2001), hfs; the same
h	La	2	3988.4612	0.403	-1.985						Lawler et al. (2001), hfs; the same
h	La	2	3988.4626	0.403	-1.140						Lawler et al. (2001), hfs; the same
h	La	2	3988.4768	0.403	-1.015						Lawler et al. (2001), hfs; the same
h	La	2	3988.4792	0.403	-1.355						Lawler et al. (2001), hfs; the same
h	La	2	3988.4810	0.403	-1.041						Lawler et al. (2001), hfs; the same
h	La	2	3988.4993	0.403	-1.068						Lawler et al. (2001), hfs; the same
h	La	2	3988.5022	0.403	-0.969						Lawler et al. (2001), hfs; the same
h	La	2	3988.5046	0.403	-1.015						Lawler et al. (2001), hfs; the same
h	La	2	3988.5271	0.403	-1.263						Lawler et al. (2001), hfs; the same
h	La	2	3988.5305	0.403	-0.683						Lawler et al. (2001), hfs; the same
h	La	2	3988.5334	0.403	-1.068						Lawler et al. (2001), hfs; the same
h	La	2	3988.5640	0.403	-0.455						Lawler et al. (2001), hfs; the same
h	La	2	3988.5674	0.403	-1.263						Lawler et al. (2001), hfs; the same
b	12CH	1	3988.5901	0.911	-3.407						Jorgensen et al. (1996)
d	La	2	4086.7090	0.000	-9.999						Dummy line. Region 4086.529 - 4086.889
h	La	2	4086.6960	0.000	-1.266						Lawler et al. (2001), hfs; the same
h	La	2	4086.6997	0.000	-1.108						Lawler et al. (2001), hfs; the same
h	La	2	4086.7029	0.000	-1.119						Lawler et al. (2001), hfs; the same
h	La	2	4086.7061	0.000	-1.292						Lawler et al. (2001), hfs; the same
h	La	2	4086.7083	0.000	-0.696						Lawler et al. (2001), hfs; the same

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Stark}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
h	La	2	4086.7097	0.000	-1.094						Lawler et al. (2001), hfs; the same
h	La	2	4086.7107	0.000	-1.790						Lawler et al. (2001), hfs; the same
h	La	2	4086.7114	0.000	-3.216						Lawler et al. (2001), hfs; the same
h	La	2	4086.7122	0.000	-1.468						Lawler et al. (2001), hfs; the same
h	La	2	4086.7178	0.000	-1.292						Lawler et al. (2001), hfs; the same
h	La	2	4086.7192	0.000	-1.119						Lawler et al. (2001), hfs; the same
h	La	2	4086.7207	0.000	-1.108						Lawler et al. (2001), hfs; the same
h	La	2	4086.7217	0.000	-1.266						Lawler et al. (2001), hfs; the same
d	La	2	4196.5439	0.321	-9.999						Lawler et al. (2001), hfs; the same
h	La	2	4196.5391	0.321	-0.926						Lawler et al. (2001), hfs; the same
h	La	2	4196.5391	0.321	-1.496						Lawler et al. (2001), hfs; the same
h	La	2	4196.5474	0.321	-1.496						Lawler et al. (2001), hfs; the same
h	La	2	4196.5474	0.321	-1.324						Lawler et al. (2001), hfs; the same
h	La	2	4196.5479	0.321	-1.338						Lawler et al. (2001), hfs; the same
h	La	2	4196.5542	0.321	-1.338						Lawler et al. (2001), hfs; the same
h	La	2	4196.5542	0.321	-2.020						Lawler et al. (2001), hfs; the same
h	La	2	4196.5547	0.321	-1.349						Lawler et al. (2001), hfs; the same
h	La	2	4196.5596	0.321	-1.349						Lawler et al. (2001), hfs; the same
h	La	2	4196.5596	0.321	-3.446						Lawler et al. (2001), hfs; the same
h	La	2	4196.5601	0.321	-1.522						Lawler et al. (2001), hfs; the same
h	La	2	4196.5630	0.321	-1.522						Lawler et al. (2001), hfs; the same
h	La	2	4196.5635	0.321	-1.698						Lawler et al. (2001), hfs; the same
b	Ce	2	4196.3330	0.417	-0.377						Lawler et al. (2001), hfs; the same
b	12CH	1	4196.2358	0.522	-4.120						Jorgensen et al. (1996)
d	La	2	4526.1099	0.770	-9.990						Dummy line. Region 4525.96 - 4526.26
h	La	2	4526.0952	0.770	-1.192						Lawler et al. (2001), hfs; the same
h	La	2	4526.0996	0.770	-1.932						Lawler et al. (2001), hfs; the same
h	La	2	4526.1030	0.770	-2.932						Lawler et al. (2001), hfs; the same
h	La	2	4526.1055	0.770	-1.363						Lawler et al. (2001), hfs; the same
h	La	2	4526.1094	0.770	-1.749						Lawler et al. (2001), hfs; the same
h	La	2	4526.1123	0.770	-2.514						Lawler et al. (2001), hfs; the same
h	La	2	4526.1143	0.770	-1.570						Lawler et al. (2001), hfs; the same
h	La	2	4526.1172	0.770	-1.716						Lawler et al. (2001), hfs; the same
h	La	2	4526.1196	0.770	-2.280						Lawler et al. (2001), hfs; the same
h	La	2	4526.1211	0.770	-1.841						Lawler et al. (2001), hfs; the same
h	La	2	4526.1235	0.770	-1.775						Lawler et al. (2001), hfs; the same
h	La	2	4526.1250	0.770	-2.134						Lawler et al. (2001), hfs; the same but only lower state
h	La	2	4526.1260	0.770	-2.259						Lawler et al. (2001), hfs; the same but only lower state
h	La	2	4526.1274	0.770	-1.958						Lawler et al. (2001), hfs; the same but only lower state
h	La	2	4526.1284	0.770	-2.037						Lawler et al. (2001), hfs; the same but only lower state
b	12CH	1	4526.1929	2.119	-2.607						Jorgensen et al. (1996)
d	La	2	4920.9800	0.130	-9.999						Dummy line. Region 4920.89 - 4921.18
h	La	2	4920.9692	0.130	-2.261						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9692	0.130	-2.407						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9702	0.130	-2.738						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9702	0.130	-2.078						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9702	0.130	-2.065						Lawler et al. (2001a), hfs; the same but only lower state

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{Sark}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta_{\lambda_{\text{line}}}$ [mÅ]	$\delta_{\lambda_{\text{rad}}}$ [mÅ]	References and notes
h	La	2	4920.9722	0.130	-2.629						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9722	0.130	-1.956						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9722	0.130	-1.831						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9751	0.130	-2.650						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9751	0.130	-1.895						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9751	0.130	-1.646						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9785	0.130	-2.760						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9785	0.130	-1.891						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9785	0.130	-1.490						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9834	0.130	-2.972						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9834	0.130	-1.957						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9834	0.130	-1.354						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9888	0.130	-3.375						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9888	0.130	-2.162						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4920.9888	0.130	-1.233						Lawler et al. (2001a), hfs; the same but only lower state
d	La	2	4921.7798	0.240	-9.999						Dummy line. Region 4921.58 - 4921.98
h	La	2	4921.7778	0.240	-3.601						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7778	0.240	-2.220						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7778	0.240	-1.139						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7788	0.240	-3.207						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7788	0.240	-3.601						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7788	0.240	-2.005						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7788	0.240	-1.233						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7798	0.240	-3.010						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7798	0.240	-1.927						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7798	0.240	-1.334						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7808	0.240	-2.923						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7808	0.240	-1.915						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7808	0.240	-1.445						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7812	0.240	-2.939						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7812	0.240	-1.955						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7812	0.240	-1.566						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7817	0.240	-3.123						Lawler et al. (2001a), hfs; the same but only lower state
h	La	2	4921.7817	0.240	-2.006						Lawler et al. (2001a), hfs; the same but only lower state
b	La	2	4921.7817	0.240	-2.053						Jorgensen et al. (1996)
h	La	2	4921.7817	0.240	-1.700						Dummy line. Region 5114.46 - 5114.71
h	La	2	5114.5601	0.230	-9.999						Lawler et al. (2001), hfs; the same
h	La	2	5114.5127	0.230	-1.624						Lawler et al. (2001), hfs; the same
h	La	2	5114.5293	0.230	-1.820						Lawler et al. (2001), hfs; the same
h	La	2	5114.5566	0.230	-1.820						Lawler et al. (2001), hfs; the same
h	La	2	5114.5732	0.230	-3.005						Lawler et al. (2001), hfs; the same
h	La	2	5114.5864	0.230	-1.824						Lawler et al. (2001), hfs; the same
h	La	2	5114.6084	0.230	-1.824						Lawler et al. (2001), hfs; the same
h	La	2	5114.6216	0.230	-2.079						Lawler et al. (2001), hfs; the same
Ce	2	3655.8401	0.320	-0.020	-140.	175.					

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{varW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{StarK}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	Ce	2	3940.3301	0.320	-0.270				-150.	150.	Palmeri et al.(2000)
b	12CH	1	3940.3711	1.720	-2.597						Jorgensen et al.(1996)
b	12CH	1	3940.3789	0.072	-3.846				-150.	150.	Jorgensen et al.(1996)
b	Ce	2	3978.6460	0.536	-0.082						VALD:Kupka et al.(1999)
b	Ce	2	3992.3799	0.450	-0.170				-100.	145.	Palmeri et al.(2000)
b	Ce	2	4083.2219	0.701	0.240				-150.	150.	Sneden et al.(1996)
b	12CH	1	4083.1389	1.509	-2.775						Jorgensen et al.(1996)
b	12CH	1	4083.2549	1.860	-1.565						Jorgensen et al.(1996)
b	13CH	1	4083.2571	0.431	-2.954						Jorgensen et al.(1996)
b	Ce	2	4115.3701	0.920	0.100				-120.	120.	Palmeri et al.(2000)
b	12CH	1	4115.3652	1.703	-2.101						Jorgensen et al.(1996)
b	Ce	2	4118.1401	0.700	0.190				-150.	120.	Palmeri et al.(2000)
b	Ce	2	4120.8271	0.320	-0.240				-40.	170.	Sneden et al.(1996)
b	Ce	2	4145.0000	0.700	0.130				-120.	120.	Palmeri et al.(2000)
b	Ce	2	4165.6001	0.910	0.530				-80.	120.	Palmeri et al.(2000)
b	13CH	1	4165.5122	0.991	-2.212						Jorgensen et al.(1996)
b	13CH	1	4165.5552	0.991	-2.196						Jorgensen et al.(1996)
b	12CH	1	4165.5688	2.234	-1.554						Jorgensen et al.(1996)
b	13CH	1	4165.5791	0.639	-2.941						Jorgensen et al.(1996)
b	12CH	1	4165.6348	2.234	-1.570				-150.	150.	Palmeri et al.(2000)
b	Ce	2	4539.7402	0.330	-0.020				-110.	90.	Graffton & Sneden(1994)
b	Ce	2	4562.3589	0.478	0.330				-90.	150.	Palmeri et al.(2000)
b	Ce	2	4572.2798	0.680	0.290				-90.	150.	Palmeri et al.(2000)
b	12CH	1	4572.2739	2.897	-1.528						Jorgensen et al.(1996)
b	Ce	2	4628.1611	0.516	0.260				-150.	150.	Palmeri et al.(2000)
b	Ce	2	5274.2300	1.040	0.150				-150.	150.	Palmeri et al.(2000)
b	Pr	2	3964.8201	0.050	0.120				-120.	120.	Ivarsson et al.(2001)
b	12CH	1	3964.4709	1.523	-2.843						Jorgensen et al.(1996)
b	Ce	2	3964.4961	0.322	-0.582						VALD:Kupka et al.(1999)
b	Fe	1	3964.5149	2.845	-1.592						VALD:Kupka et al.(1999)
b	Fe	2	3964.5791	2.778	-3.926						VALD:Kupka et al.(1999)
b	Ru	1	3964.8340	0.000	-1.900						VALD:Kupka et al.(1999)
b	Eu	2	3964.8970	1.279	-1.395						VALD:Kupka et al.(1999)
b	Hf	2	3964.9409	2.211	-0.440						VALD:Kupka et al.(1999)
b	Co	1	3964.9961	1.049	-3.130				-7.657	-6.27	8.110
b	Mg	1	3965.1931	4.346	-3.410						VALD:Kupka et al.(1999)
b	C	1	3965.2151	7.946	-3.040						VALD:Kupka et al.(1999)
b	12CH	1	3964.5371	0.770	-4.050						Jorgensen et al.(1996)
b	13CH	1	3964.6489	1.330	-3.601						Jorgensen et al.(1996)
b	13CH	1	3964.6689	1.140	-3.627						Jorgensen et al.(1996)
b	13CH	1	3964.6880	1.140	-3.569						Jorgensen et al.(1996)
b	13CH	1	3964.7241	1.330	-3.550						Jorgensen et al.(1996)
b	12CH	1	3965.1440	0.980	-4.051						Jorgensen et al.(1996)
b	12CH	1	3965.1660	0.980	-4.051						Jorgensen et al.(1996)
b	12CH	1	3965.3269	0.980	-3.260						Jorgensen et al.(1996)
b	12CH	1	3965.3491	0.980	-3.260						Jorgensen et al.(1996)
Pr	2	5219.0498	0.800	-0.050					-100.	100.	Ivarsson et al.(2001)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{Sark}} / N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \chi_{\text{bic}}$ [mÅ]	$\delta \chi_{\text{rad}}$ [mÅ]	References and notes
b	Pr	2	5220.1099	0.800	0.300			-120.	120.	Ivarsson et al. (2001)	
b	Pr	2	5259.7300	0.630	0.110			-120.	120.	Ivarsson et al. (2001)	
b	12CH	1	5259.6890	1.289	-2.981			-180.	180.	Jørgensen et al. (1996)	
b	Nd	2	3780.3999	0.470	-0.350			-150.	150.	Den Hartog (2003)	
b	Nd	2	3784.2500	0.380	0.150			-120.	100.	Den Hartog (2003)	
b	Nd	2	3900.2300	0.470	0.100			-75.	75.	Den Hartog (2003)	
b	Nd	2	3979.4690	0.205	-0.330			-95.	95.	Jørgensen et al. (1996)	
b	12CH	1	3979.5330	1.015	-3.138			-120.	70.	Den Hartog (2003)	
b	Nd	2	3991.7410	0.000	-0.260			-100.	90.	Den Hartog (2003)	
b	Nd	2	4004.0100	0.060	-0.570			-130.	130.	Jørgensen et al. (1996)	
b	12CH	1	4004.0840	1.034	-3.508			-150.	75.	Den Hartog (2003)	
b	Nd	2	4013.2200	0.180	-1.100			-100.	90.	Jørgensen et al. (1996)	
b	Nd	2	4018.8230	0.064	-0.850			-100.	95.	Den Hartog (2003)	
b	12CH	1	4018.7610	0.341	-4.161			-130.	95.	Den Hartog (2003)	
b	Nd	2	4021.3269	0.321	-0.100			-130.	110.	Den Hartog (2003)	
b	Nd	2	4023.0900	0.560	0.040			-90.	110.	Jørgensen et al. (1996)	
b	12CH	1	4022.9629	1.108	-3.275			-90.	100.	Den Hartog (2003)	
b	Nd	2	4051.1499	0.380	-0.300			-130.	135.	Jørgensen et al. (1996)	
b	12CH	1	4051.1040	1.875	-2.305			-90.	110.	Den Hartog (2003)	
b	Nd	2	4051.1799	1.875	-2.321			-130.	135.	Jørgensen et al. (1996)	
b	12CH	1	4061.0850	0.471	0.550			-90.	110.	Den Hartog (2003)	
b	Nd	2	4069.2649	0.064	-0.570			-130.	135.	Jørgensen et al. (1996)	
b	12CH	1	4069.2920	1.455	-2.386			-90.	110.	Den Hartog (2003)	
b	Nd	2	4109.4482	0.321	0.350			-130.	135.	Jørgensen et al. (1996)	
b	12CH	1	4109.3691	0.597	-3.795			-90.	110.	Den Hartog (2003)	
b	12CH	1	4109.4351	0.597	-3.840			-130.	135.	Jørgensen et al. (1996)	
b	12CH	1	4109.4570	0.597	-1.683			-90.	110.	Den Hartog (2003)	
b	Nd	2	4232.3740	0.064	-0.470			-130.	135.	Jørgensen et al. (1996)	
b	12CH	1	4232.5142	0.072	-3.870			-90.	110.	Den Hartog (2003)	
b	12CH	1	4232.5459	2.879	-1.103			-130.	135.	Jørgensen et al. (1996)	
b	Ce	2	4232.5610	0.721	-0.743			-130.	135.	VALD: Kupka et al. (1999)	
b	Nd	2	4446.3838	0.205	-0.350			-110.	60.	Den Hartog (2003)	
b	12CH	1	4446.2920	1.598	-1.970			-110.	60.	Den Hartog (2003)	
b	12CH	1	4446.4600	1.598	-1.900			-110.	60.	Den Hartog (2003)	
b	Nd	2	4462.9790	0.559	0.040			-135.	135.	Den Hartog (2003)	
b	12CH	1	4462.8892	1.596	-1.677			-135.	135.	Jørgensen et al. (1996)	
b	12CH	1	4462.8911	1.596	-1.689			-135.	135.	Jørgensen et al. (1996)	
b	12CH	1	4462.9351	2.037	-1.380			-135.	135.	Jørgensen et al. (1996)	
b	Nd	2	4542.6001	0.740	-0.280			-135.	135.	Den Hartog (2003)	
b	Nd	2	4579.3198	0.740	-0.480			-135.	135.	Den Hartog (2003)	
b	12CH	1	4597.1001	1.520	-2.709			-135.	135.	Jørgensen et al. (1996)	
b	Nd	2	4597.1089	3.137	-1.148			-135.	135.	Jørgensen et al. (1996)	
b	Nd	2	4645.7700	0.560	-0.760			-180.	180.	Den Hartog (2003)	
b	12CH	1	4645.7310	2.131	-2.560			-180.	180.	Jørgensen et al. (1996)	

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Star}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
	Nd	2	4706.5400	0.000	-0.710				-180.	180.	Den Hartog (2003)
	Nd	2	4959.1201	0.060	-0.800				-80.	200.	Den Hartog (2003)
	Nd	2	5092.7900	0.380	-0.610				-200.	200.	Den Hartog (2003)
	Nd	2	5130.5898	1.300	0.450				-100.	200.	Den Hartog (2003)
	Nd	2	5212.3501	0.200	-0.960				-100.	110.	Den Hartog (2003)
	Nd	2	5234.2100	0.550	-0.510				-70.	65.	Den Hartog (2003)
	Nd	2	5249.6001	0.980	0.200				-140.	110.	Den Hartog (2003)
	Nd	2	5250.8101	0.750	-0.720				-200.	200.	Den Hartog (2003)
	Nd	2	5255.5098	0.200	-0.670				-200.	200.	Den Hartog (2003)
	Nd	2	5293.1699	0.820	0.100				-90.	80.	Den Hartog (2003)
	Nd	2	5311.4800	0.990	-0.420				-200.	200.	Den Hartog (2003)
	Nd	2	5319.8198	0.550	-0.140				-100.	80.	Den Hartog (2003)
	Sm	2	3760.7029	0.190	-0.418				-80.	100.	Biemont et al. (1989)
	Sm	2	4318.9351	0.277	-0.268				-150.	150.	Biemont et al. (1989)
	Sm	2	4519.6348	0.544	-0.434				-150.	150.	Biemont et al. (1989)
	Sm	2	4642.2319	0.378	-0.516				-110.	85.	Biemont et al. (1989)
d	Eu	2	3819.6699	0.000	-9.999				-100.	100.	Dummy line. Region 3819.57 - 3819.77
i	151Eu	2	3819.5857	0.000	-0.941						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6028	0.000	-0.832						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6045	0.000	-1.610						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6265	0.000	-0.722						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6289	0.000	-1.419						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6306	0.000	-2.827						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6565	0.000	-0.617						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6602	0.000	-1.366						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6626	0.000	-2.681						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6931	0.000	-0.519						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.6980	0.000	-1.407						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.7014	0.000	-2.768						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.7356	0.000	-0.426						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.7417	0.000	-1.598						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3819.7466	0.000	-3.096						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6304	0.000	-0.903						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6387	0.000	-1.572						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6394	0.000	-0.794						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6504	0.000	-2.789						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6511	0.000	-1.381						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6663	0.000	-1.327						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6812	0.000	-0.481						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6836	0.000	-1.369						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6846	0.000	-2.730						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.6980	0.000	-0.388						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.7029	0.000	-1.560						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3819.7053	0.000	-3.058						Lawler et al. (2001b), hfs r-process isotopic ratios: the same

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \chi_{\text{osc}}$ [mÅ]	$\delta \chi_{\text{rad}}$ [mÅ]	References and notes
b	Fe	1	3819.4929	3.396	-1.300	-7.200	-5.16	8.020			Lawler et al. (2001b), hfs r-process isotopic ratios: the same
b	Nd	2	3819.7019	0.380	-0.900						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
b	Zr	2	3819.7920	1.208	-1.100						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
d	Eu	2	3907.1101	0.207	-9.999						Dummy line. Region 3907.065 - 3907.205
i	151Eu	2	3907.0608	0.207	-0.695						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.0947	0.207	-0.863						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1077	0.207	-1.506						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1228	0.207	-1.063						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1331	0.207	-1.341						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1450	0.207	-1.314						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1460	0.207	-2.603						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1526	0.207	-1.330						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1616	0.207	-1.679						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1628	0.207	-2.239						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1663	0.207	-1.417						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1738	0.207	-2.093						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1743	0.207	-1.582						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	3907.1790	0.207	-2.126						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.0879	0.207	-0.656						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1040	0.207	-0.825						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1086	0.207	-1.468						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1162	0.207	-1.024						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1208	0.207	-1.303						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1252	0.207	-1.276						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1257	0.207	-2.565						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1294	0.207	-1.292						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1321	0.207	-1.641						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1343	0.207	-2.200						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1348	0.207	-1.378						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1377	0.207	-1.544						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1389	0.207	-2.054						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	3907.1404	0.207	-2.088						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
b	Ce	2	3906.9180	0.521	-0.649						Jorgensen et al. (1996)
b	Fe	1	3906.9619	3.283	-1.497						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
b	12CH	1	3907.2280	0.784	-3.291						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
b	Ce	2	3907.2881	1.107	0.515						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
d	Eu	2	4129.7202	0.000	-9.999						Dummy line. Region 4129.640 - 4129.970
i	151Eu	2	4129.6230	0.000	-1.833						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6265	0.000	-1.356						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6401	0.000	-1.637						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6450	0.000	-1.298						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6484	0.000	-1.833						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6650	0.000	-1.578						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6709	0.000	-1.168						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6758	0.000	-1.637						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.6982	0.000	-1.614						Lawler et al. (2001b), hfs r-process isotopic ratios: the same

**Table 2.** Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{vddW}}^{1000\text{ K}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{1000 K}}^{1000\text{ K}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{1000\text{ K}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
i	151Eu	2	4129.7046	0.000	-1.017						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.7100	0.000	-1.578						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.7393	0.000	-1.801						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.7461	0.000	-0.866						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.7529	0.000	-1.614						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.7964	0.000	-0.722						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	151Eu	2	4129.8032	0.000	-1.801						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6777	0.000	-1.795						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6807	0.000	-1.318						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6841	0.000	-1.598						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6875	0.000	-1.260						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6904	0.000	-1.795						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6943	0.000	-1.539						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.6978	0.000	-1.129						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7012	0.000	-1.598						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7095	0.000	-1.576						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7119	0.000	-0.979						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7153	0.000	-1.539						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7300	0.000	-1.763						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7310	0.000	-0.828						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7334	0.000	-1.576						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7549	0.000	-0.683						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
i	153Eu	2	4129.7563	0.000	-1.763						Lawler et al. (2001b), hfs r-process isotopic ratios: the same
b	12CH	1	4129.8848	1.761	-1.123						Jorgensen et al. (1996)
Gd	2	3424.5901	0.350	-0.170					-70.	70.	Sneden et al. (2003)
Gd	2	3439.2300	0.380	0.150					-70.	70.	Sneden et al. (2003)
b	12CH	1	3439.1699	0.371	-3.387						Jorgensen et al. (1996)
b	12CH	1	3439.2820	0.357	-3.725						Jorgensen et al. (1996)
Gd	2	3454.8999	0.030	-0.480					-70.	70.	Sneden et al. (2003)
b	12CH	1	3454.9839	1.208	-3.298						Jorgensen et al. (1996)
Gd	2	3481.8000	0.490	0.230					-70.	70.	Sneden et al. (2003)
Gd	2	3844.5801	0.140	-0.400					-80.	80.	Sneden et al. (2003)
b	12CH	1	3844.6541	0.838	-3.440						Jorgensen et al. (1996)
b	12CH	1	3844.6721	0.784	-3.501						Jorgensen et al. (1996)
Gd	2	4085.5701	0.730	0.070					-70.	70.	Sneden et al. (2003)
Gd	2	4191.0801	0.430	-0.570					-70.	70.	Sneden et al. (2003)
Gd	2	4215.0200	0.430	-0.580					-70.	70.	Sneden et al. (2003)
Ni	1	3510.3320	0.212	-0.670					-1238.	-1138.	Dummy line. Region 3809,094 - 3809,194
b	Ti	2	3509.8350	1.892	-1.360						VALD:Kupka et al. (1999)
b	Tb	2	3509.1440	0.000	0.700						Lawler et al. (2001c)
b	Zr	2	3507.6431	0.959	-1.314						VALD:Kupka et al. (1999)
b	Ni	1	3507.6880	0.165	-2.510						VALD:Kupka et al. (1999)
b	Co	2	3507.7800	2.779	-1.502						VALD:Kupka et al. (1999)
b	Ce	2	3507.9409	0.175	-0.754						VALD:Kupka et al. (1999)
b	Fe	2	3508.2019	1.724	-4.210						VALD:Kupka et al. (1999)
b	Er	2	3508.3789	1.402	-0.005						VALD:Kupka et al. (1999)
b	Fe	1	3508.4741	2.990	-0.882						VALD:Kupka et al. (1999)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{He}^+}$ [eV]	$\log gf$	$\log(\Gamma_{\text{rad}}^{\text{vwN}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{Sark}} / N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{Trad}})$ [rad/s]	$\delta \lambda_{\text{Sark}}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
b	Fe	1	3508.5181	2.559	-1.573	-7.843	-6.32	7.600			VALD; Kupka et al. (1999)
b	Ce	2	3508.5000	0.459	-0.841						VALD; Kupka et al. (1999)
b	Fe	1	3508.6101	2.865	-2.307	-7.696	-4.56	7.950			VALD; Kupka et al. (1999)
b	Ce	2	3508.7041	1.533	0.079						VALD; Kupka et al. (1999)
b	Fe	1	3509.1189	2.832	-2.120	-7.213	-5.63	8.270			VALD; Kupka et al. (1999)
b	Zr	1	3509.3159	0.071	-0.110						VALD; Kupka et al. (1999)
b	Fe	1	3509.7251	2.885	-2.154	-7.686	-4.59	7.850			VALD; Kupka et al. (1999)
b	Co	1	3509.8401	0.582	-0.320	-7.670	-6.30	7.930			VALD; Kupka et al. (1999)
b	Fe	1	3509.8621	2.223	-2.070	-7.679	-6.20	8.310			VALD; Kupka et al. (1999)
b	La	2	3509.9919	0.126	-1.705						VALD; Kupka et al. (1999)
b	C	1	3510.1321	7.488	-2.650						VALD; Kupka et al. (1999)
b	Nb	2	3510.2710	1.990	0.120						VALD; Kupka et al. (1999)
b	Co	1	3510.4141	0.101	-1.250	-7.779	-6.37	7.390			VALD; Kupka et al. (1999)
b	Fe	1	3510.4390	2.484	-1.461	-7.596	-6.06	8.300			VALD; Kupka et al. (1999)
b	Zr	2	3510.4500	0.559	-0.884						VALD; Kupka et al. (1999)
b	Dy	2	3407.8000	0.000	0.180		-100.	160.			Wickliffe et al. (2000)
b	Dy	2	3434.3701	0.000	-0.450		-160.	160.			Wickliffe et al. (2000)
b	12CH	1	3434.4309	0.872	-2.998						Jorgensen et al. (1996)
b	Dy	2	3445.5701	0.000	-0.150		-200.	200.			Wickliffe et al. (2000)
b	12CH	1	3445.5569	0.670	-3.429						Jorgensen et al. (1996)
b	Dy	2	3449.8899	0.540	-0.750		-150.	150.			Wickliffe et al. (2000)
b	Dy	2	3454.3201	0.100	-0.140		-75.	150.			Wickliffe et al. (2000)
b	12CH	1	3454.3169	1.138	-3.251						Jorgensen et al. (1996)
b	12CH	1	3454.3879	0.709	-3.017						Jorgensen et al. (1996)
b	Dy	2	3460.9700	0.000	-0.070						Wickliffe et al. (2000)
b	Dy	2	3506.8101	0.100	-0.600						Wickliffe et al. (2000)
b	Dy	2	3531.7100	0.000	0.770						Wickliffe et al. (2000)
b	Mn	1	3531.8269	2.282	-0.097	-7.751	-5.17	8.290			VALD; Kupka et al. (1999)
b	Dy	2	3538.5200	0.000	-0.020						Wickliffe et al. (2000)
b	Dy	2	3550.2200	0.590	0.270						Wickliffe et al. (2000)
b	Dy	2	3563.1499	0.100	-0.360						Wickliffe et al. (2000)
b	Dy	2	3694.8101	0.100	-0.110						Wickliffe et al. (2000)
b	Dy	2	3788.4399	0.100	-0.570						Wickliffe et al. (2000)
b	Dy	2	3944.6809	0.000	0.100						Wickliffe et al. (2000)
b	Dy	2	3996.6899	0.590	-0.260						Wickliffe et al. (2000)
b	Dy	2	4077.9600	0.100	-0.040						Wickliffe et al. (2000)
b	12CH	1	4077.9290	0.639	-1.998						Jorgensen et al. (1996)
d	Ho	2	4152.5801	0.080	-9.990						Dummy line. Region 4152.53 - 4152.63
h	Ho	2	4152.4429	0.080	-1.685						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.4448	0.080	-3.047						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.5195	0.080	-3.047						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.5210	0.080	-1.765						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.5210	0.080	-2.822						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.5889	0.080	-2.733						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.5903	0.080	-2.822						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.5903	0.080	-1.846						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.6489	0.080	-2.710						Sneden et al. (2003), hfs; the same

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Stark}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
h	Ho	2	4152.6509	0.080	-1.926						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.6519	0.080	-2.733						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7012	0.080	-2.741						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7031	0.080	-2.002						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7051	0.080	-2.710						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7456	0.080	-2.836						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7480	0.080	-2.072						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7505	0.080	-2.741						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7832	0.080	-3.060						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7856	0.080	-2.127						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.7881	0.080	-2.836						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.8159	0.080	-2.157						Sneden et al. (2003), hfs; the same
h	Ho	2	4152.8184	0.080	-3.060						Sneden et al. (2003), hfs; the same
b	Sc	1	4152.3359	1.969	0.448	-7.582					VALD:Kupka et al. (1999)
b	Dy	2	4152.4780	1.169	-0.727						VALD:Kupka et al. (1999)
b	Nb	1	4152.5742	0.086	-0.030						VALD:Kupka et al. (1999)
b	Zr	1	4152.6450	1.366	-0.250						VALD:Kupka et al. (1999)
b	Fe	1	4152.7612	3.430	-3.249	-7.653					VALD:Kupka et al. (1999)
b	La	2	4152.7729	1.754	-0.343						VALD:Kupka et al. (1999)
b	Ce	2	4152.9258	1.676	-0.175						VALD:Kupka et al. (1999)
b	13CH	1	4152.1812	2.450	-3.165						Jorgensen et al. (1996)
b	12CH	1	4152.6289	2.780	-2.650						Jorgensen et al. (1996)
b	12CH	1	4152.6860	2.780	-2.661						Jorgensen et al. (1996)
b	12CH	1	4152.9111	0.520	-4.503						Jorgensen et al. (1996)
Er	2	3559.8999	0.000	-0.740							-90.
Er	2	3729.5200	0.000	-0.500							-80.
Er	2	3938.6299	0.000	-0.520							-100.
b	12CH	1	3938.5310	0.557	-4.071						Kurnucz (1995)
b	12CH	1	3938.6970	0.557	-3.998						Jorgensen et al. (1996)
Tm	2	3462.2000	0.000	0.030							Jorgensen et al. (1996)
b	12CH	1	3462.1951	0.777	-2.835						Jorgensen et al. (1996)
b	12CH	1	3462.3010	0.987	-3.362						Jorgensen et al. (1996)
b	12CH	1	3462.3101	0.458	-2.938						Jorgensen et al. (1996)
b	Fe	1	3462.3521	2.198	-2.110	-7.804					VALD:Kupka et al. (1999)
Tm	2	3700.2600	0.030	-0.380							Kurnucz (1995)
Tm	2	3701.3601	0.000	-0.540							-150.
Tm	2	3761.9099	0.000	-0.430							150.
Tm	2	3289.3701	0.000	-9.999							-180.
d	Yb	2	3289.3701	0.000	-9.999						180.
i	000Yb	2	3289.3647	0.000	-1.495						-300.
i	000Yb	2	3289.3701	0.000	-0.639						Dummy line. Region 3289.07 - 3289.62
i	000Yb	2	3289.3743	0.000	-0.477						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994),isot=Yb170,172,174,176
i	000Yb	2	3289.3782	0.000	-0.876						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994),isot=Yb170,172,174,176
i	171Yb	2	3289.3364	0.000	-1.427						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994),isot=Yb170,172,174,176
i	171Yb	2	3289.3757	0.000	-1.029						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994)
i	171Yb	2	3289.3821	0.000	-1.728						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994)
i	173Yb	2	3289.3486	0.000	-2.107						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994)
i	173Yb	2	3289.3555	0.000	-1.563						Sneden et al.(2003),isot&hfs;Martensson-Pendril etal.(1994)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{He}^+}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{StarK}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{He}^+}$ [mÅ]	$\delta \lambda_{\text{rad}}$ [mÅ]	References and notes
i	173Yb	2	3289.3647	0.000	-1.199						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3289.3821	0.000	-1.676						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3289.3867	0.000	-1.563						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3289.3936	0.000	-1.660						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
b	Fe	2	3289.3501	3.814	-1.620	-7.944	-6.66	8.479			Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
b	V	2	3289.3899	1.096	-0.931	-7.922	-6.58	8.362			Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
b	Fe	1	3289.4299	2.832	-2.720	-7.787	-6.05	8.173			Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
d	Yb	2	3694.1899	0.000	-9.999				-100.		Dummy line. Region 3694.09 - 3694.39
i	000Yb	2	3694.1826	0.000	-1.815						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994), isot=Yb170,172,174,176
i	000Yb	2	3694.1899	0.000	-0.959						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994), isot=Yb170,172,174,176
i	000Yb	2	3694.1958	0.000	-0.797						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994), isot=Yb170,172,174,176
i	000Yb	2	3694.2014	0.000	-1.196						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	171Yb	2	3694.1396	0.000	-1.747						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	171Yb	2	3694.1973	0.000	-1.446						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	171Yb	2	3694.2068	0.000	-1.747						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3694.1680	0.000	-1.582						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3694.1758	0.000	-1.679						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3694.2158	0.000	-2.126						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
i	173Yb	2	3694.2239	0.000	-1.582						Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
b	Fe	1	3694.0061	3.038	0.078	-7.200	5.49	8.200			Snedden et al.(2003), isot&hfs; Martensson-Pendrill et al. (1994)
b	Fe	1	3554.9250	2.832	0.538	-7.744	-6.44	8.270	-555.	-455.	Dummy line. Region 3554.370 - 3554.470
h	Lu	2	3554.3982	2.150	-0.486						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.3987	2.150	-1.056						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4104	2.150	-1.056						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4109	2.150	-0.884						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4121	2.150	-0.898						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4263	2.150	-0.898						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4275	2.150	-1.580						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4287	2.150	-0.909						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4424	2.150	-0.909						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4436	2.150	-3.006						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4448	2.150	-1.082						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4561	2.150	-1.082						Snedden et al.(2003), hfs; the same
h	Lu	2	3554.4573	2.150	-1.258						VALD:Kupka et al.(1999)
h	Zr	2	3554.0791	1.184	-0.682						VALD:Kupka et al.(1999)
b	Fe	1	3554.1179	0.958	-2.206	-7.660	-6.24	7.990			VALD:Kupka et al.(1999)
b	Sm	2	3554.1609	0.333	-1.117						VALD:Kupka et al.(1999)
b	Er	2	3554.3000	0.886	-0.723						VALD:Kupka et al.(1999)
b	Cr	1	3554.3889	3.111	-1.054	-7.99	-5.10	8.230			VALD:Kupka et al.(1999)
b	Fe	1	3554.4409	2.940	-2.491	-7.653	-5.83	7.840			VALD:Kupka et al.(1999)
b	Fe	1	3554.5010	2.882	-1.029	-7.765	-5.15	7.930			VALD:Kupka et al.(1999)
b	Ce	2	3554.6360	1.177	-0.073						Jørgensen et al. (1996)
b	Ce	2	3555.0010	0.320	-0.338						Jørgensen et al. (1996)
b	12CH	1	3554.3279	1.520	-4.270						Jørgensen et al. (1996)
b	12CH	1	3554.4939	0.340	-5.573						Jørgensen et al. (1996)
b	12CH	1	3554.4961	0.340	-5.573						Jørgensen et al. (1996)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma_{\text{vUV}}^{10000 \text{ K}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{10000 K}}^{1000 \text{ K}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{1000 \text{ K}}/N_e)$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	12CH	1	3554.6140	1.320	-4.732						Jørgensen et al. (1996)
b	12CH	1	3554.7090	0.340	-5.014						Jørgensen et al. (1996)
b	12CH	1	3554.7109	0.340	-5.014						Jørgensen et al. (1996)
b	12CH	1	3554.7900	1.440	-4.877						Jørgensen et al. (1996)
b	Hf	2	3399.7930	0.000	-0.490						VALD: Kupka et al. (1999)
b	Fe	1	3399.3330	2.198	-0.622	-7.667					VALD: Kupka et al. (1999)
b	Zr	2	3399.3391	0.319	-0.704						VALD: Kupka et al. (1999)
b	Dy	2	3399.3560	1.571	-0.159						VALD: Kupka et al. (1999)
b	Fe	1	3399.3860	3.047	-2.513	-7.811					VALD: Kupka et al. (1999)
b	Gd	2	3399.4021	0.382	-0.863						VALD: Kupka et al. (1999)
b	Co	1	3399.4309	2.080	-1.504	-7.842					VALD: Kupka et al. (1999)
b	Cr	2	3399.5129	4.497	-1.270	-8.002					VALD: Kupka et al. (1999)
b	Cr	2	3399.5510	9.684	-3.469	-7.580					VALD: Kupka et al. (1999)
b	Nb	2	3399.7051	1.818	-0.370						VALD: Kupka et al. (1999)
b	Sm	2	3399.8411	0.659	-1.065						VALD: Kupka et al. (1999)
b	Tm	2	3399.9490	1.111	-0.096						VALD: Kupka et al. (1999)
b	Gd	2	3399.9871	0.354	-0.460						VALD: Kupka et al. (1999)
b	Cr	2	3400.1060	4.297	-2.400	-7.926					VALD: Kupka et al. (1999)
b	Mn	2	3400.1140	4.936	-1.146	-7.950					VALD: Kupka et al. (1999)
b	Ce	2	3400.2500	0.933	-0.604						VALD: Kupka et al. (1999)
b	12CH	1	3399.4121	0.950	-4.775						Jørgensen et al. (1996)
b	12CH	1	3399.5090	0.950	-4.701						Jørgensen et al. (1996)
b	12CH	1	3399.5100	1.520	-2.907						Jørgensen et al. (1996)
b	12CH	1	3399.7219	1.520	-2.875						Jørgensen et al. (1996)
b	13CH	1	3399.9690	1.520	-3.861						Jørgensen et al. (1996)
b	12CH	1	3400.1399	0.730	-5.450						Jørgensen et al. (1996)
b	12CH	1	3400.2629	0.730	-5.385						Jørgensen et al. (1996)
d	Pb	1	3683.4600	0.970	-9.999						110. Dummy line, Region 3683.31 - 3683.57
h	Pb	1	3683.4795	0.970	-2.342						Biemont et al. (2000), isot&zhfs: Manning et al. (1950)
h	Pb	1	3683.4700	0.970	-1.239						Biemont et al. (2000), isot&zhfs: Manning et al. (1950)
h	Pb	1	3683.4812	0.970	-1.683						Biemont et al. (2000), isot&zhfs: Manning et al. (1950)
h	Pb	1	3683.4424	0.970	-1.382						Biemont et al. (2000), isot&zhfs: Manning et al. (1950)
h	Pb	1	3683.4600	0.970	-0.754						Biemont et al. (2000), isot&zhfs: Manning et al. (1950)
d	Pb	1	4057.8150	1.320	-9.999						150. Dummy line, Region 4057.755 - 4057.965
h	Pb	1	4057.8262	1.320	-2.012						Biemont et al. (2000), iso&zhfs: Manning et al. (1950)
h	Pb	1	4057.8150	1.320	-1.019						Biemont et al. (2000), iso&zhfs: Manning et al. (1950)
h	Pb	1	4057.8308	1.320	-1.463						Biemont et al. (2000), iso&zhfs: Manning et al. (1950)
h	Pb	1	4057.8100	1.320	-2.162						Biemont et al. (2000), iso&zhfs: Manning et al. (1950)
h	Pb	1	4057.7583	1.320	-1.208						Biemont et al. (2000), iso&zhfs: Manning et al. (1950)
h	Pb	1	4057.8018	1.320	-0.534						Biemont et al. (2000), iso&zhfs: Manning et al. (1950)
b	12CH	1	4057.7600	2.438	-1.936						Jørgensen et al. (1996)
b	12CH	1	4057.8560	0.371	-2.272						Jørgensen et al. (1996)
b	12CH	1	4057.8621	0.371	-3.164						Jørgensen et al. (1996)
Th	2	4019.1289	0.000	-0.228							50. Nilsson et al. (2002b)
b	Fe	1	4019.0500	2.608	-2.780	-7.720					VALD: Kupka et al. (1999)
b	Ce	2	4019.0569	1.014	-0.213						VALD: Kupka et al. (1999)
b	Ni	1	4019.0581	1.935	-3.174	-7.870					VALD: Kupka et al. (1999)

Table 2. Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\lambda_{\text{lower}}$ [eV]	$\log g f$	$\log(\Gamma^{\text{radW}} / N_H)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{Sark}} / N_e)$ [rad.cm <sup>-3</sup> /s]	$\log(\Gamma^{\text{rad}})$ [rad/s]	$\delta \lambda_{\text{blue}}$ [mÅ]	$\delta \lambda_{\text{red}}$ [mÅ]	References and notes
b	Co	1	4019.2891	0.582	-3.232	-7.860	-6.30	7.030			VALD; Kupka et al. (1999)
b	Co	1	4019.2991	0.629	-3.769	-7.860	-6.30	7.350			VALD; Kupka et al. (1999)
b	13CH	1	4018.8899	1.334	-3.389						Jorgensen et al. (1996)
b	13CH	1	4018.9690	0.463	-2.345						Jorgensen et al. (1996)
b	13CH	1	4019.0471	0.463	-2.337						Jorgensen et al. (1996)
b	12CH	1	4019.3911	1.219	-3.262						Jorgensen et al. (1996)
d	Fe	1	3859.9109	0.000	-9.999				-390.	-290.	Dummy line. Region 3859.521 - 3859.621
b	U	2	3859.5710	0.036	-0.067						Nilsson et al. (2002a)
b	Fe	1	3859.2129	2.404	-0.749	-7.720	-6.33	7.310			VALD; Kupka et al. (1999)
b	Nd	2	3859.4180	0.205	-1.308						VALD; Kupka et al. (1999)
b	Cr	1	3859.5879	3.395	-0.017	-7.700	-5.06	8.230			VALD; Kupka et al. (1999)
b	Fe	1	3859.9109	0.000	-0.710	-7.780	-6.32	7.190			VALD; Kupka et al. (1999)
b	CN	1	3859.3330	3.009	-4.898						Hill et al. (2002)
b	CN	1	3859.3740	3.009	-1.189						Hill et al. (2002)
b	CN	1	3859.3760	3.081	-3.777						Hill et al. (2002)
b	CN	1	3859.3811	0.558	-3.160						Hill et al. (2002)
b	CN	1	3859.3979	0.558	-0.447						Hill et al. (2002)
b	CN	1	3859.3979	1.018	-3.635						Hill et al. (2002)
b	CN	1	3859.4150	0.558	-0.476						Hill et al. (2002)
b	CN	1	3859.4480	1.018	0.041						Hill et al. (2002)
b	CN	1	3859.4541	3.081	0.163						Hill et al. (2002)
b	CN	1	3859.4590	1.894	-3.587						Hill et al. (2002)
b	CN	1	3859.4661	3.081	0.155						Hill et al. (2002)
b	CN	1	3859.4719	3.009	-1.170						Hill et al. (2002)
b	CN	1	3859.4871	1.894	-0.417						Hill et al. (2002)
b	CN	1	3859.5020	1.018	0.032						Hill et al. (2002)
b	CN	1	3859.5310	1.894	-0.433						Hill et al. (2002)
b	CN	1	3859.5750	3.118	-3.774						Hill et al. (2002)
b	CN	1	3859.5869	1.939	-2.791						Hill et al. (2002)
b	CN	1	3859.5911	1.939	-1.258						Hill et al. (2002)
b	CN	1	3859.5979	1.939	-1.370						Hill et al. (2002)
b	CN	1	3859.6130	4.838	-0.252						Hill et al. (2002)
b	CN	1	3859.6499	0.089	-0.162						Hill et al. (2002)
b	CN	1	3859.6541	3.118	0.170						Hill et al. (2002)
b	CN	1	3859.6609	3.118	0.163						Hill et al. (2002)
b	CN	1	3859.6669	0.566	-3.159						Hill et al. (2002)
b	CN	1	3859.6631	2.312	-0.443						Hill et al. (2002)
b	CN	1	3859.6721	4.838	-0.256						Hill et al. (2002)
b	CN	1	3859.6721	0.997	-3.635						Hill et al. (2002)
b	CN	1	3859.7090	2.312	-0.460						Hill et al. (2002)
b	CN	1	3859.7209	0.997	0.032						Hill et al. (2002)
b	CN	1	3859.7310	4.632	-0.700						Hill et al. (2002)
b	CN	1	3859.7310	2.312	-3.581						Hill et al. (2002)

**Table 2.** Continued.

Def	Elem/mol	Ion	$\lambda$ [Å]	$\chi_{\text{lower}}$ [eV]	$\log gf$	$\log(\Gamma_{1000\text{ K}}^{\text{vdW}}/N_H)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{1000\text{ K}}^{\text{Stark}}/N_e)$ [rad.cm <sup>3</sup> /s]	$\log(\Gamma_{\text{rad}}^{\text{rad}})$ [rad/s]	$\delta\lambda_{\text{blue}}$ [mÅ]	$\delta\lambda_{\text{red}}$ [mÅ]	References and notes
b	CN	1	3859.7510	2.721	-4.559						Hill et al. (2002)
b	CN	1	3859.7739	0.997	0.023						Hill et al. (2002)
b	CN	1	3859.7810	1.206	-3.556						Hill et al. (2002)
b	CN	1	3859.7871	3.155	-3.772						Hill et al. (2002)
b	12CH	1	3859.1960	0.051	-5.595						Jørgensen et al. (1996)
b	12CH	1	3859.2261	1.962	-4.042						Jørgensen et al. (1996)
b	12CH	1	3859.2700	0.072	-5.589						Jørgensen et al. (1996)
b	12CH	1	3859.3479	0.051	-5.595						Jørgensen et al. (1996)
b	12CH	1	3859.5210	0.072	-5.589						Jørgensen et al. (1996)
b	12CH	1	3859.6479	0.072	-5.438						Jørgensen et al. (1996)
b	12CH	1	3859.8650	0.072	-5.438						Jørgensen et al. (1996)
b	12CH	1	3859.8850	0.096	-5.434						Jørgensen et al. (1996)
b	12CH	1	3859.9309	1.365	-3.103						Jørgensen et al. (1996)
b	12CH	1	3859.9641	1.365	-3.119						Jørgensen et al. (1996)

## Appendix A: Line list

The complete line list is presented in Table 2, which is available only electronically. Here we briefly comment on line selection issues and data sources for each element, particularly for the oscillator strengths, hfs and isotopic splitting. Other data such as wavelengths, excitation potentials were collected from various sources, often the Vienna Atomic Line Database (VALD, Kupka et al. 1999), the NIST compilations or Sneden et al. (2003a).

**Carbon:** The abundance of carbon was derived from the  $A-X$  bands at 4310–4313 Å (0, 0) and 4362–4367 Å (1, 1), along with two  $B^2\Sigma-X^2\Pi$  (1, 0) CH lines at 3638 Å and 3661 Å. The band regions used in this programme are the same as those used by Paper II. The two single lines were selected from the list of Jørgensen et al. (1996) as they are unblended, of moderate strength, and have reasonable  $S/N$  in the observed spectrum. The C abundance was derived from  $^{12}\text{CH}$  features assuming an isotopic ratio of  $^{12}\text{C}/^{13}\text{C} = \infty$  as in Paper II.

**Nitrogen:** The nitrogen abundance was derived using the  $B\Sigma-X\Sigma$  bands of CN at 3872 Å (1, 1) and 3882 Å (0, 0). The line data was extracted from a list made available by Plez (1998, private communication) and is described by Hill et al. (2002).

**Oxygen:** The data for the oxygen lines were compiled from VALD, except for the log  $gf$  values which were astrophysical values taken from Jonsell et al. (2005). The differences from the NIST log  $gf$  values in VALD were less than 0.02 dex.

**Sodium:** For two of the three Na I lines the  $f$ -values of Wiese & Martin (1980) and hfs from McWilliam et al. (1995) were used. The  $f$ -value for the 8183 Å line was taken from VALD. No hfs data were available for this line, and as it is weak this should not be important.

**Magnesium:** A total of 7 lines of Mg I were used. Data for the lines at 3829, 4571, and 4703 Å were taken from Paper II. Due to contamination by CH, Balmer lines, and some residual reduction artefacts not all lines from Paper II could be used. Oscillator strengths for the lines at 5173, 5184, and 5528 Å originate from Wiese & Martin (1980). For the 5711 Å line the astrophysical log  $gf$ -value from Jonsell et al. (2005) was adopted.

**Aluminium:** The Al I resonance line at 3961 Å was used for the analysis. The hfs was taken from McWilliam et al. (1995).

**Calcium:** The  $f$ -values for the 10 Ca I lines were taken from Wiese & Martin (1980), Smith (1981), Jonsell et al. (2005), and VALD.

**Scandium:** The abundance of scandium was determined from 5 Sc II lines, using  $f$ -values from Lawler & Dakin (1989). Hfs for the lines 4247 and 4415 Å was taken from McWilliam et al. (1995). The remaining lines are very weak and hfs was not considered.

**Titanium:** A total of 11 lines of Ti I and 13 of Ti II were analysed. The oscillator strengths were gathered from various sources: Blackwell et al. (1982a), Blackwell et al. (1982b), Fuhr & Wiese (1996), Grevesse et al. (1989), Martin et al. (1988), Pickering et al. (2001), Ryabchikova et al. (1994) and VALD.

**Vanadium:** The vanadium abundance is based on one V I and 4 V II lines. The  $f$ -value data were taken from Fuhr & Wiese (1996).

**Chromium:** The  $f$ -values for the 6 Cr I and 2 Cr II lines were taken from Fuhr & Wiese (1996) and Martin et al. (1988).

**Manganese:** For the two lines of Mn I we adopted the log  $gf$  data from Booth et al. (1984), and hfs was computed using the data compiled in Lefèvre et al. (2003). For the three Mn II lines we used data from Kling & Griesmann (2000), with hfs from Holt et al. (1999).

**Iron:** A total of 61 Fe I lines and 8 Fe II lines were analysed. The neutral line oscillator strengths are taken from O'Brian et al. (1991), Fuhr & Wiese (1996), Bard et al. (1991), Bridges & Kornblith (1974), and VALD. For the Fe II lines  $f$ -values were adopted from Biémont et al. (1991), Fuhr & Wiese (1996), Heise & Kock (1990), Kroll & Kock (1987), Moity (1983), Schnabel et al. (2004), and VALD.

**Cobalt:** The cobalt abundance was derived from 4 Co I lines using  $f$ -values from Cardon et al. (1982) and Nitz et al. (1999), and hfs data from Pickering (1996).

**Nickel:** The 8 lines of Ni I have oscillator strengths from Blackwell et al. (1989), Fuhr et al. (1988), Fuhr & Wiese (1996), Huber & Sandeman (1980), and VALD.

**Copper:** We analysed two lines of Cu I, both in the UV. The log  $gf$  data were taken from Fuhr & Wiese (1996), and hfs data from Kurucz (1995).

**Strontium:** One Sr I line and 4 Sr II lines were analysed. Hyperfine and isotopic splitting are accounted for in the 4077 and 4215 Å lines (see discussion in Paper II). We did not find hyperfine structure data for the remaining lines; however, as they are weak, this should not be important. The  $f$ -values were gathered from Migdalek & Baylis (1987), Pinnington et al. (1995), and Sneden et al. (2003a).

**Yttrium:** The  $f$ -values for the 11 lines of Y II are from Hannaford et al. (1982).

**Zirconium:** All but two of the 16 lines of Zr II were analysed with  $f$ -values from Biémont et al. (1981). The remaining lines at 3356 and 4019 Å have  $f$ -values from VALD.

**Silver:** The upper limit to the silver abundance was derived from two Ag I lines. The oscillator strengths were taken from Fuhr & Wiese (1996). The hfs data was taken from Ross & Aller (1972).

**Barium:** The barium abundance is derived using three Ba II lines. The log  $gf$  data came from Gallagher (1967) and VALD. The isotopic splitting and hfs were taken from Villemoes et al. (1993) for the two weak lines, noting that data for only the lower levels are available. For the 4554 Å line, hfs and isotopic splitting from McWilliam (1998) was adopted. In all cases, the solar isotopic composition from Anders & Grevesse (1989) was assumed. We note that if a pure  $r$ -process isotopic composition is assumed we find an abundance only 0.03 dex lower.

**Lanthanum:** The 8 lines of La II were analysed employing oscillator strengths and hfs data from Lawler et al. (2001a). For the lines at 4921 and 4922 Å only the hfs of the lower levels was considered, as no data were available for the upper levels.

**Cerium:** Abundances were derived from 15 lines of Ce II. The  $f$ -values were gathered from various sources, Palmeri et al. (2000), Sneden et al. (1996), Gratton & Sneden (1994), and VALD.

**Praseodymium:** The log  $gf$  values for the 4 lines of Pr II were taken from Ivarsson et al. (2001).

**Neodymium:** A total of 34 Nd II lines were analysed. The oscillator strengths were taken from Den Hartog et al. (2003).

**Samarium:** The 4 lines of Sm II were analysed using  $f$ -values from Biémont et al. (1989).

**Europium:** The abundance of europium is based on 3 Eu II lines, with  $f$ -values, hfs and isotopic splitting taken from Lawler et al. (2001c). Solar  $r$ -process isotopic fractions were assumed.

**Gadolinium:** The  $f$ -values for the 8 Gd II lines were collected from Sneden et al. (2003a); the paper of Cowan et al. (2002) should be consulted for details.

**Terbium:** The  $f$ -value for the single line of Tb II was taken from Lawler et al. (2001d).

**Dysprosium:** The  $f$ -values for 15 of the 16 Dy II lines analysed were taken from Wickliffe et al. (2000). For the line at 3944 Å the  $f$ -value is from Biémont & Lowe (1993).

**Holmium:** The single Ho II line at 4152 Å has atomic data including hfs taken from Sneden et al. (2003a).

**Erbium:** The  $f$ -values for the 3 Er II lines were taken from Musiol & Labuz (1983) and Kurucz (1995).

**Thulium:** The 4 Tm II lines have oscillator strengths from Kurucz (1995). According to Sneden et al. (1996) these are rescaled laboratory data from Corliss & Bozman (1962).

**Ytterbium:** The  $f$ -values used for our two Yb II lines are from Sneden et al. (2003a). They have renormalised theoretical oscillator strengths from Biémont et al. (1998) using lifetimes from Pinnington et al. (1997). The isotopic and hyperfine splitting data are taken from Mårtensson-Pendrel et al. (1994), and the solar isotopic ratios from Anders & Grevesse (1989) have been assumed.

**Lutetium:** For the single Lu II line we adopted the log  $gf$  and hfs data from Sneden et al. (2003a). The oscillator strength is from Quinet et al. (1999), renormalised using lifetimes from Fedchak et al. (2000).

**Hafnium:** We have one line of Hf II with the log  $gf$  taken from VALD.

**Lead:** The abundance is derived from two Pb I lines. The oscillator strengths were taken from Biémont et al. (2000) and the hfs and isotopic splitting data from Manning et al. (1950). Solar isotopic ratios from Anders & Grevesse (1989) have been assumed.

**Thorium:** An upper limit for the thorium abundance is derived from the Th II line at 4019.129 Å employing the oscillator strength from Nilsson et al. (2002b).

**Uranium:** The  $f$ -value for the uranium line is taken from Nilsson et al. (2002a).