



**HAL**  
open science

## Regulation of the generation of reactive oxygen species during photosynthetic electron transport

Anja Krieger-Liszkay, Ginga Shimakawa

► **To cite this version:**

Anja Krieger-Liszkay, Ginga Shimakawa. Regulation of the generation of reactive oxygen species during photosynthetic electron transport. *Biochemical Society Transactions*, 2022, 50 (2), pp.1025-1034. 10.1042/BST20211246 . hal-03744901

**HAL Id: hal-03744901**

**<https://hal.science/hal-03744901>**

Submitted on 3 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Regulation of the generation of reactive oxygen species during photosynthetic electron transport**

2 Anja Krieger-Liszkay<sup>1,\*</sup> and Ginga Shimakawa<sup>2</sup>

3 <sup>1</sup>Université Paris-Saclay, Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, 91198 Gif-sur-  
4 Yvette cedex, France

5 <sup>2</sup>Department of Bioscience, School of Biological and Environmental Sciences, Kwansei-Gakuin  
6 University, 2-1 Gakuen, Sanda, Hyogo 669-1337, Japan

7 \*Corresponding author: phone: +33 1 6908 1803; E-mail: anja.krieger-liszkay@cea.fr

8

## 9 **Abstract:**

10 Light capture by chlorophylls and photosynthetic electron transport carry the risk of the generation of  
11 reactive oxygen species (ROS) including singlet oxygen, superoxide anion radicals and hydrogen  
12 peroxide. Rapid changes in light intensity, electron fluxes and accumulation of strong oxidants and  
13 reductants increase ROS production. Superoxide is mainly generated at the level of photosystem I  
14 while photosystem II is the main source of singlet oxygen. ROS can induce oxidative damage of the  
15 photosynthetic apparatus, however, ROS are also important to tune processes inside the chloroplast  
16 and participate in retrograde signalling regulating the expression of genes involved in acclimation  
17 responses. Under most physiological conditions light harvesting and photosynthetic electron transport  
18 are regulated to keep the level of ROS at a non-destructive level. Photosystem II is most prone to  
19 photoinhibition but can be quickly repaired while photosystem I is protected in most cases. The size of  
20 the transmembrane proton gradient is central for the onset of mechanisms that protect against  
21 photoinhibition. The proton gradient allows dissipation of excess energy as heat in the antenna  
22 systems and it regulates electron transport. pH-dependent slowing down of electron donation to  
23 photosystem I protects it against ROS generation and damage. Cyclic electron transfer and  
24 photoreduction of oxygen contribute to the size of the proton gradient. The yield of singlet oxygen  
25 production in photosystem II is regulated by changes in the midpoint potential of its primary quinone  
26 acceptor. In addition, numerous antioxidants inside the photosystems, the antenna and the thylakoid  
27 membrane quench or scavenge ROS.

28

## 29 **Introduction**

30 Living in an oxygen-rich atmosphere carries the risk of oxidative damage of lipids, proteins and  
31 nucleic acids. Beside oxidative damage, reactive oxygen species (ROS) are also important as signaling  
32 molecules triggering the expression of responsive genes, and they play a crucial role in developmental

33 processes, in plant – microbe interactions and the defense against pathogen attack [1,2]. Oxygen, a  
34 triplet in its ground state, is a strong oxidizing agent, however its reactivity is limited by spin pairing  
35 and reorientation. By contrast, its singlet form,  $^1\text{O}_2$ , and the three intermediate redox states  
36 (superoxide, peroxide and hydroxyl radical) between  $\text{O}_2$  and its fully reduced form  $\text{H}_2\text{O}$  are much more  
37 reactive with the highest reactivity for  $^1\text{O}_2$  and the hydroxyl radical ( $^{\bullet}\text{OH}$ ). ROS are generated by  
38 specialized enzymes like for example NADPH oxidases to fight pathogen attack [3], and ROS generation  
39 occurs as side reactions in electron transport chains like the respiratory chain [4] and in oxygen  
40 photosynthetic organisms in the photosynthetic electron transport chain. Furthermore,  
41 photosynthetic organisms have their own special oxidative stress burden like light-induced generation  
42 of excited states of the chlorophylls that can lead to  $^1\text{O}_2$  generation.

43

#### 44 **Superoxide generation by photosystem I**

45         Superoxide anion radicals ( $\text{O}_2^{\bullet-}$ ) are generated by the reduction of  $\text{O}_2$  at the three main  
46 complexes of the photosynthetic electron transport chain (Fig. 1), photosystem II (PSII) [5], the  
47 cytochrome  $b_6/f$  complex (cyt $b_6/f$ ) [6], photosystem I (PSI) and at the plastoquinone (PQ) pool [7]. The  
48 acceptor side of PSI is the most important site of  $\text{O}_2^{\bullet-}$  generation in the so-called Mehler reaction (Fig.  
49 2). The Mehler reaction is per se not deleterious for PSI when it occurs at the level of the terminal  
50 acceptors  $F_A F_B$  since  $\text{O}_2^{\bullet-}$  generated at the stroma site of PSI is detoxified efficiently.  $\text{O}_2^{\bullet-}$  is first reduced  
51 to  $\text{H}_2\text{O}_2$  by superoxide dismutase (SOD) or spontaneously and  $\text{H}_2\text{O}_2$  is decomposed by ascorbate  
52 peroxidase, peroxiredoxins, and glutathione peroxidase [8]. However, the situation is different when  
53  $\text{O}_2^{\bullet-}$  is generated inside the PSI reaction center. Since the midpoint redox potential of the  $\text{O}_2/\text{O}_2^{\bullet-}$  pair  
54 in thylakoid membranes is assumed to be much more negative (around -550 mV) [9], both,  
55 phylloquinones and [4Fe-4S] clusters are suggested as thermodynamically plausible sites of  $\text{O}_2$   
56 reduction within PSI. Recently, Kozuleva et al. (2021) reported for isolated PSI that  $\text{O}_2$  photoreduction  
57 is observed in the absence of the [4Fe-4S] clusters and that it is enhanced with the longer lifetime of  
58 phylloquinone, suggesting that the A-branch phylloquinone is the site of  $\text{O}_2$  reduction in isolated PSI  
59 (Fig. 2) [10]. Because of low membrane permeability of  $\text{O}_2^{\bullet-}$  ( $2.1 \times 10^{-6} \text{ cm s}^{-1}$ ) [11],  $\text{O}_2^{\bullet-}$  generated within  
60 the protein complex or at the interface protein complex/thylakoid membrane is assumed to oxidatively  
61 attack PSI before it diffuses out of the membrane and is disproportionated enzymatically by SOD.  
62 Alternatively,  $\text{O}_2^{\bullet-}$  is non-enzymatically disproportionated to  $\text{H}_2\text{O}_2$  that reacts with the reduced [4Fe-  
63 4S] centers generating  $^{\bullet}\text{OH}$  when electron transport is strongly limited at the acceptor side of PSI  
64 [12,13]. In this way,  $^{\bullet}\text{OH}$ , being more reactive than  $\text{O}_2^{\bullet-}$ , can be generated in PSI.

65

## 66 Singlet oxygen production in photosystem II

67 When excited chlorophyll in its triplet state ( $^3\text{Chl}$ ) reacts with molecular oxygen ( $^3\text{O}_2$ ) the very  
68 reactive  $^1\text{O}_2$  is generated. In the Chl containing antenna systems,  $^3\text{Chl}$  may be generated from a singlet  
69 excited chlorophyll ( $^1\text{Chl}$ ). However, in an intact antenna system, little  $^1\text{O}_2$  is generated because  
70 carotenoids, including xanthophylls, are in close contact with Chl [14]. Carotenoids quench efficiently  
71  $^3\text{Chl}$  and protect thereby against  $^1\text{O}_2$  production. In the reaction centre of PSII,  $^3\text{Chl}$  is generated by  
72 charge recombination of the primary radical pair ( $\text{P680}^+ \text{Ph}^-$ ), with pheophytin (Ph) being the primary  
73 electron acceptor and P680 the primary chlorophyll electron donor (Fig. 2). When light absorption  
74 exceeds the capacity of photosynthetic electron transport, the quinone acceptors of PSII ( $\text{Q}_\text{A}$ ,  $\text{Q}_\text{B}$  and  
75 the PQ pool) become reduced and the probability of  $^1\text{O}_2$  generation increases. The midpoint potential  
76 of  $\text{Q}_\text{A}$  determines whether charge recombination proceeds via the dangerous route of repopulation of  
77 the primary radical pair ( $\text{P680}^+ \text{Phe}^-$ ) generating with a certain probability  $^3\text{P680}$  and subsequently  $^1\text{O}_2$   
78 or via the safe charge recombination directly into the ground state of P680 without producing  $^1\text{O}_2$  (for  
79 details see [15-17]). The yield of  $^1\text{O}_2$  formation is lowered when  $\text{Q}_\text{A}$  adopts the high potential form by  
80 shifting the midpoint redox potential of  $\text{Q}_\text{A}$  to a more positive value. PSII with high potential  $\text{Q}_\text{A}$  has  
81 been found in PSII with an inactive water-splitting complex when either the obligatory cofactor calcium  
82 was lacking or when the  $\text{Mn}_4\text{CaO}_5$  cluster had been removed [18,19]. Recently, using Fourier  
83 Transformed Infrared (FTIR) spectroscopy instead of chlorophyll fluorescence as a measure for the  
84 reduction state of  $\text{Q}_\text{A}$ , more negative potentials for the  $\text{Em} (\text{Q}_\text{A}/\text{Q}_\text{A}^-)$  have been reported, and hardly  
85 any change in the  $\text{Em} (\text{Q}_\text{A}/\text{Q}_\text{A}^-)$  upon Mn depletion has been observed [20]. However, it has been shown  
86 previously that PSII with an inactivated water-splitting complex generates much less  $^1\text{O}_2$  despite charge  
87 separation similar to active PSII [21,22], supporting the  $\text{Em} (\text{Q}_\text{A}/\text{Q}_\text{A}^-)$  values determined by chlorophyll  
88 fluorescence. Further work is needed to show whether chlorophyll fluorescence or FTIR spectroscopy  
89 measure the reduction state of  $\text{Q}_\text{A}$  more reliably. The  $\text{Em} (\text{Q}_\text{A}/\text{Q}_\text{A}^-)$  is also shifted towards a more  
90 positive value in the absence of the ligand bicarbonate at the non-heme iron at the acceptor side of  
91 PSII. This shift, although much smaller than the one found upon inactivation of the water-splitting  
92 complex, decreases also significantly  $^1\text{O}_2$  production [15]. A cyanobacterial assembly state of PSII has  
93 been recently characterized [23]. Binding of the assembly factor Psb28 induces a large conformational  
94 change at the PSII acceptor side, which distort the binding pocket of  $\text{Q}_\text{B}$  and replace the bicarbonate  
95 ligand of non-heme iron with glutamate. The yield of  $^1\text{O}_2$  generation is lowered in this assembly state  
96 showing that the replacement of bicarbonate by glutamate protects the nascent PSII.

97 PSII centres with high potential  $\text{Q}_\text{A}$  have been observed under different physiological  
98 conditions *in vivo*; (i) in green algae prior to photoactivation, i.e., the light-dependent assembly of the  
99  $\text{Mn}_4\text{CaO}_5$  cluster [17], and (ii) in plant leaves of under high light [24] and drought [25]. Small carboxylic

100 acids generated in the photorespiratory pathway may also modify the midpoint potential of  $Q_A$  by  
101 replacing bicarbonate as has been shown for the *A. thaliana* photorespiration mutant *hpr1-1* that  
102 accumulates glycolate [26]. Such regulation may become physiologically relevant when the  $CO_2$   
103 concentration is temporarily low and the glycolate concentration is high inside the chloroplast. A  
104 slowdown of forward electron transport at the acceptor side of PSII has been reported to occur under  
105 moderate drought stress, a condition favouring photorespiration compared to  $CO_2$  assimilation [25].  
106 This observation may reflect replacement of bicarbonate by glycolate in a wild-type plant. Reversible  
107 inactivation of PSII by glycolate thus represents a way how a photorespiratory metabolite can protect  
108 plants from  $^1O_2$  in high light, at elevated temperature and drought conditions when photorespiration  
109 is more important.

110

### 111 **Photoinhibition in photosystem I and II**

112 ROS generated in photosystems in the light can provoke oxidative damage of the protein  
113 complexes leading to an inactivation of the photosystems, a process known as photoinhibition.  
114 Photoinhibition of PSII is easily observed in intact cells and leaves, when the damage exceeds the repair  
115 of damaged PSII. The molecular mechanism of PSII photoinhibition is discussed controversially in the  
116 literature.  $^1O_2$  is a plausible cause for the damage of the D1 protein [27], but also  $O_2^{\bullet-}$  and  $^{\bullet}OH$   
117 generated at the acceptor side of PSII [28]. These radicals have been shown to oxidise specific amino  
118 acid residues of the D1 protein [29], while the specific amino acids attacked by  $^1O_2$  have still to be  
119 identified. In addition, photoinhibition can also take place at the donor side of PSII inhibiting the  
120  $Mn_4CaO_5$  cluster, thereby limiting electron donation to  $TyrZ^+$  and  $P680^+$  [30]. Upon illumination with  
121 UV light, Mn in the  $Mn_4CaO_5$  cluster is reduced and the cluster falls apart, resulting in donor-side-  
122 induced photoinhibition [31]. Damaged D1 protein is rapidly replaced by newly synthesized D1 protein  
123 in the so-called PSII repair cycle [32]. Not only active PSII but also the repair of damaged PSII is sensitive  
124 to ROS because the translation factor EF-Tu is easily inhibited by oxidation of a cysteine residue [33].  
125 Inactivation of PSII suppresses the electron transport to PSI, protecting it by alleviating the  $O_2^{\bullet-}$   
126 generation in PSI. Therefore, photoinhibition of PSII is often regarded as a predetermined breaking  
127 point to protect the rest of the photosynthetic apparatus against photo-induced oxidative damage.

128 As described above,  $O_2^{\bullet-}$  and  $^{\bullet}OH$  are the candidates to cause the inactivation of PSI. Based on  
129 electron spin resonance spectroscopy in damaged PSI, [4Fe-4S] clusters are the most plausible  
130 damaged sites by ROS [34,35]. Damage of the acceptor side of PSI is often observed as a decrease in  
131 the total amount of P700 (the reaction centre chlorophyll in PSI) that is kept oxidized in saturating  
132 light. Additionally, the inactivation of PSI is almost linearly correlated with the decrease in the rate of  
133 photosynthetic  $CO_2$  assimilation in leaves [36]. Because damaged PSI takes several days or weeks to be

134 replaced [37,38], PSI photoinhibition can be a lethal event for photosynthetic organisms. It should be  
135 noted that except for extreme environmental conditions, including artificial fluctuating light [39] and  
136 chilling stress [40], and for specific mutants [41,42], PSI photoinhibition is hardly observed *in vivo*,  
137 which suggests that  $O_2^{\bullet-}$  is efficiently scavenged before PSI is damaged or that  $O_2^{\bullet-}$  generation is very  
138 low.

139

#### 140 **Scavenging $O_2^{\bullet-}$ and $^1O_2$ by antioxidants**

141 Prenylipids like tocochromanols (tocopherols, especially  $\alpha$ -Toc, tocotrienols and  
142 plastochochromanol-8), plastoquinone-9 (PQ/PQH<sub>2</sub>) and carotenoids (carotenes and xanthophylls) are  
143 lipid-soluble antioxidants in the chloroplasts, which can quench or scavenge ROS [43].  $\alpha$ -Toc and PQH<sub>2</sub>  
144 efficiently scavenge  $O_2^{\bullet-}$  and lipid radicals, and, like carotenoids, also quench and scavenge  $^1O_2$ . The  
145 level of the antioxidants  $\alpha$ -Toc and PQH<sub>2</sub> increases under environmental conditions that promote  
146 oxidative stress. Carotenoid levels and their composition also change in response to changes in  
147 environmental conditions [44]. Carotenoids are essential quenchers of  $^3Chl$  and  $^1O_2$  in the antenna  
148 systems of the photosystems and  $\beta$ -carotene can scavenge  $^1O_2$  inside the PSII reaction centre while its  
149 position is too far to quench directly the triplet state  $^3P680$ .  $\beta$ -carotene protects also PSI against  $^1O_2$   
150 generation as has been shown in isolated PSI complexes from a mutant with a lower  $\beta$ -carotene  
151 content [45]. It has also been recently proposed that  $\alpha$ -Toc is localized near Ph<sub>D1</sub> and the nonheme  
152 iron, protecting amino acids of the D1 protein against oxidative modifications [46]. In a recent study,  
153 the antioxidant activities of  $\alpha$ -Toc, PQ/PQH<sub>2</sub>, zeaxanthin in excess light stress were analysed in  
154 *Arabidopsis thaliana* wild-type and in mutants affected in the individual antioxidants. It was shown  
155 that  $^1O_2$  is first quenched by  $\alpha$ -Toc. When it is consumed, zeaxanthin and PQ take this role [47].  
156 Cyanobacteria contain a special photoactive soluble carotenoid protein, the orange carotenoid protein  
157 that is able to quench efficiently  $^1O_2$  in addition to its role in thermal dissipation of excess energy by  
158 the cyanobacterial antenna, the phycobilisome [48,49]. The water-soluble chlorophyll protein WSCP  
159 has also  $^1O_2$  quenching properties, although it does not contain a carotenoid [50]. The physiological  
160 role of WSCP is far from being understood, and it is only present in some species of the Brassicaceae,  
161 Chenopodiaceae, Amaranthaceae and Polygonaceae.

162

#### 163 **Regulatory mechanisms to suppress ROS generation**

164 Photosynthetic organisms employ a large number of regulatory mechanisms for electron  
165 transport and light utilisation to suppress ROS generation in response to short-term environmental  
166 changes in addition to the ROS scavenging systems described above. When light is sufficient,  
167 photosynthetic electron transport is usually limited by the CO<sub>2</sub> content inside the leaves. This is the

168 case at high light intensities when the plant is under optimal conditions. In abiotic stress conditions  
169 like drought, the stomata are closed and the CO<sub>2</sub> concentration in the cells and chloroplasts drops,  
170 thereby limiting CO<sub>2</sub> assimilation already at light intensities that are non-saturating in the absence of  
171 such stress. In low CO<sub>2</sub>, photorespiration is crucial to keep the electron transport chain oxidized  
172 thereby protecting against photoinhibition [51]. When photosynthetic organisms are exposed to  
173 excess light and the electron transport chain is highly reduced, the potential risk of photosynthetic  
174 electron transport to generate ROS increases unless regulatory mechanisms start to function. Such  
175 regulation mechanisms comprise i) slowing down electron transport at the acceptor side of PSII and at  
176 the *cytb<sub>6</sub>f* complex, ii) dissipating excess energy as heat, and iii) additional electron transport routes  
177 via specialized proteins or directly to oxygen.

178 The most important control over photosynthetic electron transport is exerted by the proton  
179 motive force, composed of the proton gradient ( $\Delta\text{pH}$ ) and the electrical gradient ( $\Delta\Psi$ ), which is created  
180 during electron transport in thylakoid membranes. Protons are liberated at the donor side of PSII  
181 during water oxidation and they are pumped across the thylakoid membrane during the electron  
182 transport in the *cytb<sub>6</sub>f* complex. The  $\Delta\text{pH}$  provides the driving force for ATP synthesis by the CF<sub>0</sub>CF<sub>1</sub>-  
183 ATP synthase. The pH inside the thylakoid lumen is important for the protection of both PSI and PSII  
184 against ROS damage by the so-called photosynthetic control and by initiating quenching mechanisms  
185 in the antenna that allow dissipation of excess energy as heat (qE, the pH-dependent component of  
186 non-photochemical quenching). When the  $\Delta\text{pH}$  becomes larger than needed for ATP synthesis,  
187 photosynthetic control sets in. Photosynthetic control is defined as the pH-dependent decrease of the  
188 activity of the *cyt b<sub>6</sub>f* complex [52-54]. The consequence is a limitation of electron donation to PSI,  
189 leading to an accumulation of P700<sup>+</sup> that alleviates O<sub>2</sub><sup>•-</sup> production, resulting in the protection of PSI  
190 against ROS damage [39]. The lumen pH in Arabidopsis leaves under ambient CO<sub>2</sub> was estimated to  
191 range from approximately pH 7.5 to 6.5 under weak and saturating light, respectively [55]. When net  
192 ATP synthesis is zero, the pH in the lumen can decrease as low as pH 5.2 [53], a pH at which the water-  
193 splitting activity starts to be slowed down [56]. Below pH 5.5, Ca<sup>2+</sup>, an obligatory co-factor of the water-  
194 splitting complex is reversibly removed, evoking a shift of Q<sub>A</sub> to the high potential form [19,57] and  
195 protecting PSII against <sup>1</sup>O<sub>2</sub> generation (see above).

196 Moderate low pH values in the lumen allow, beside photosynthetic control, regulation at the  
197 antenna level via qE. In photosynthetic eukaryotes, qE is triggered by  $\Delta\text{pH}$  and commonly requires  
198 three components: (1) de-epoxidation of xanthophylls, (2) conformational change in the major LHCII  
199 and (3) unique proteins associated with LHCII, including PsbS in land plants, LHCSR in green algae and  
200 basal land plants, and LhcX in diatoms [58-60]. Although the detailed mechanism of qE is still  
201 controversial and different among organisms,  $\Delta\text{pH}$  is known to function in activating the de-epoxidase

202 and/or the protonation of these unique proteins. Therefore, the capability of qE to dissipate excess  
203 energy in the PSII antenna system and to prevent  $^1\text{O}_2$  generation in the PSII reaction centre depends  
204 on the size of  $\Delta\text{pH}$ . In Arabidopsis, the pH value for zeaxanthin accumulation and PsbS protonation was  
205 estimated to be about 6.8 [55].

206         Alternative electron transport pathways like the Mehler reaction and cyclic electron flow help  
207 to protect the photosynthetic apparatus against damage by ROS. They do not lead to the production  
208 of NADPH but theoretically contribute to the  $\Delta\text{pH}$  and permit the synthesis of extra ATP. Cyclic electron  
209 transport around PSI is defined as the electron transport from ferredoxin to PQ. There are two  
210 pathways proposed for cyclic electron flow: one that is insensitive to the inhibitor Antimycin A and  
211 involving the chloroplast NAD(P)H dehydrogenase (NDH) complex using ferredoxin instead of NAD(P)H  
212 as electron donor; and the other Antimycin-A-sensitive one that involves the Cyt *b<sub>6</sub>f* complex and the  
213 elusive proton gradient regulation complex PGR5/PGRL1. However the exact pathway of the second  
214 one is still a matter of debate [61]. The pathway via the NDH complex, a homologue of the complex I  
215 of the mitochondrial respiratory chain, is probably pumping extra protons into the thylakoid lumen  
216 [62] to support chlororespiration and contributes to the photosynthetic control. It should be noted  
217 that the *in vivo* activity of cyclic electron flow has not yet been clarified because of the technical  
218 difficulty to evaluate it separately from other electron transport pathways. In green algae and  
219 cyanobacteria,  $\text{O}_2$  photoreduction to water, catalyzed by the flavodiiron enzymes, represents a  
220 significant electron transport activity (the optimal rate is almost comparable to the photosynthetic  $\text{O}_2$   
221 evolution rate) and contributes to the  $\Delta\text{pH}$ , alleviating ROS damage in PSI [63-66]. The electron flux *via*  
222 flavodiiron proteins is smaller but its physiological roles are still conserved in basal land plants, and  
223 finally lost in angiosperms in the evolutionary history [67,68]. The *in vivo* electron flux *via* the Mehler  
224 reaction (i.e., the non-enzymatic reduction of  $\text{O}_2$  to  $\text{O}_2^{\cdot-}$  in PSI) is smaller than that *via* flavodiiron  
225 proteins. Nevertheless, even in the absence of the flavodiiron proteins, the reduction of  $\text{O}_2$  to  $\text{O}_2^{\cdot-}$   
226 contributes to the  $\Delta\text{pH}$ , and it is enhanced under certain physiological conditions like for example in  
227 plants grown under short photoperiod [69].

228         Alternative electron transport routes are in competition with each other, and their activity is  
229 controlled by the redox state of the chloroplast stroma *via* the thioredoxin system. It has been recently  
230 shown that cyclic electron transport is controlled by thioredoxin *m* and by the NADPH-dependent  
231 thioredoxin system of the chloroplast NTRC [70,71]. In this context it is interesting to mention that  
232 thioredoxin *m* forms a complex with PGRL1 [72,73] that itself contains redox active cysteine residues  
233 [74]. The formation of a complex between reduced thioredoxin *m* and PGRL1 may inhibit cyclic electron  
234 flow by preventing the supercomplex formation required for cyclic electron flow. Possibly,  $\text{O}_2$



235 photoreduction at PSI is activated by the thioredoxin system in a competitive manner with cyclic  
236 electron flow.

237 In addition to cyclic electron flow and O<sub>2</sub> photoreduction at PSI, O<sub>2</sub> can also be reduced by the  
238 plastid terminal oxidase (PTOX) that uses PQH<sub>2</sub> as electron donor [75]. The electron flux *via* PTOX is  
239 usually much lower than that of CO<sub>2</sub> assimilation *in vivo*, but it is up-regulated in response to abiotic  
240 stress at the expression and redox levels of the protein [76,77]. Its membrane association depends on  
241 the ΔpH [78]. Thereby the enzyme may have only access to its substrate in the presence of a high ΔpH.  
242 Such a regulation mechanisms avoids the competition of PTOX with linear electron flow under  
243 conditions that are favorable for CO<sub>2</sub> assimilation. PTOX is possibly functioning as a safety valve at the  
244 acceptor side of PSII to regulate ROS generation in both PSI and PSII.

245

#### 246 **Conclusions and further research directions**

247 Much work has been done to identify the sites of ROS production in the different  
248 photosynthetic complexes. However, most work on the exact sites of ROS production has been  
249 performed using isolated PSII and PSI complexes. It is likely that ROS production and its control differ  
250 depending on the production sites and on the organization of supercomplexes. For example,  
251 differences in ROS production are likely to occur in the different assembly states of the photosystems,  
252 as has been shown for <sup>1</sup>O<sub>2</sub> in cyanobacterial PSII assembly state [23]. A detailed study on ROS  
253 formation of different photosystem complex compositions would be of interest. During the repair cycle  
254 of PSII, it is also likely that the yield of ROS is different in monomeric PSII compared to dimeric PSII and  
255 upon the removal of the cofactors from damaged monomeric PSII. Furthermore, the yield of ROS  
256 production and the stability of the ROS may depend on the site of the thylakoid membrane where ROS  
257 generation takes place. It has been shown recently that the site of <sup>1</sup>O<sub>2</sub> production is important for <sup>1</sup>O<sub>2</sub>  
258 signalling through flu/executer pathway [79,80].

259 A feedback reaction of <sup>1</sup>O<sub>2</sub> generation in PSII by photorespiratory metabolites is a new concept  
260 that links photosynthetic electron transport with primary metabolism. Further research is needed to  
261 show whether under photorespiratory conditions like drought or increased temperature the glycolate  
262 concentration inside the chloroplast can get high enough to exchange with the bicarbonate at the  
263 acceptor side of PSII in wild-type plants.

264 Beside the exact sites of ROS production, further work is required to understand the regulation  
265 of alternative electron transport pathways that lower the yield of ROS production. First work has been  
266 done to identify sites of thioredoxin-dependent regulation of cyclic flow. However, redox regulation of  
267 the Mehler reaction and the flavodiiron proteins remains an open question. The whole complexity of  
268 the interplay of the different components of the thioredoxin system, like the individual thioredoxins,

269 NTRC, peroxiredoxins and other players like H<sub>2</sub>O<sub>2</sub> has also still to be unravelled. Redox regulation of  
270 PTOX has just been described very recently.

271 The relationship of controlling ROS production in photosynthetic electron transport with ROS  
272 cell signalling should be further studied. Super-resolution microscopy techniques using specific  
273 fluorescent dyes or the expression of ROS-sensitive fluorescent proteins may shine in future new light  
274 on the distribution of ROS production inside a single chloroplast, while standard confocal microscopy  
275 may allow differences in ROS production amongst individual chloroplasts. A combination of these  
276 techniques with single cell transcriptomic techniques may help to further understand ROS signalling  
277 networks.

278

279

## 280 **Perspectives**

281 ● ROS generated in photosynthetic electron transport have important roles in signalling and specific  
282 reactions, but their level has to be controlled since to high ROS levels provoke oxidative damage.  
283 The detailed understanding of the mechanism of ROS production and regulation mechanisms  
284 limiting ROS formation is a prerequisite to understand acclimation to abiotic stress.

285

286 ● The molecular mechanism of ROS generation at the photosystems is mainly understood on the  
287 basis of experiments with isolated thylakoid membranes and photosystems. Several players of  
288 regulatory mechanisms have been identified at the molecular level which control ROS production,  
289 but their importance *in vivo* and their interactions in responses to environmental fluctuations are  
290 far from being understood.

291

292 ● Continuous technical improvement of specific spin traps, ROS specific fluorescence dyes,  
293 expression of fluorescent protein-based biosensors and super-resolution microscopy will allow in  
294 future quantitative analyses of the production rates and sites of the individual ROS species *in vivo*  
295 with spatial information and unprecedented sensitivity. Future quantitative analyses of the  
296 interactions or competitions between the different regulation mechanisms of photosynthetic  
297 electron transport will allow a comprehensive understanding of ROS levels inside the chloroplast.

298

299

300

301

## 302 **Conflict of interests**

303 The authors declare no conflict of interest.

304

### 305 **Author contributions**

306 A.K.L. and G.S. wrote the article.

307

### 308 **Acknowledgements**

309 We would like to thank Umama Hani and Lucas Leverne, both Université Paris -Saclay, I2BC, for critical  
310 reading of the manuscript. This work was supported by the Labex Saclay Plant Sciences-SPS (ANR-17-  
311 EUR-0007).

312

### 313 **References**

- 314 1 K.-J. Dietz, I. Turkan and A. Krieger-Liszkay (2016) Redox- and reactive oxygen species-dependent  
315 signaling into and out of the photosynthesizing chloroplast. *Plant Physiol.* **171**, 1541-1550  
316 <http://www.plantphysiol.org/content/171/3/1541.abstract>
- 317
- 318 2 D. Camejo, Á. Guzmán-Cedeño and A. Moreno (2016) Reactive oxygen species, essential molecules,  
319 during plant–pathogen interactions. *Plant Physiol. Biochem.* **103**, 10-23  
320 <https://www.sciencedirect.com/science/article/pii/S0981942816300602>
- 321
- 322 3 S. Pfeilmeier, G.C. Petti, M. Bortfeld-Miller, B. Daniel, C.M. Field, S. Sunagawa, et al. (2021) The  
323 plant NADPH oxidase RBOHD is required for microbiota homeostasis in leaves. *Nature*  
324 *Microbiology* **6**, 852-864 <https://doi.org/10.1038/s41564-021-00929-5>
- 325
- 326 4 E.L. Robb, A.R. Hall, T.A. Prime, S. Eaton, M. Szibor, C. Viscomi, et al. (2018) Control of mitochondrial  
327 superoxide production by reverse electron transport at complex I. *J. Biol. Chem.* **293**, 9869-  
328 9879 <https://doi.org/10.1074/jbc.RA118.003647>
- 329
- 330 5 P. Pospíšil (2012) Molecular mechanisms of production and scavenging of reactive oxygen species  
331 by photosystem II. *Biochim. Biophys. Acta Bioenerg.* **1817**, 218-231  
332 <https://www.sciencedirect.com/science/article/pii/S0005272811001344>
- 333
- 334 6 D. Baniulis, S.S. Hasan, J.T. Stofleth and W.A. Cramer (2013) Mechanism of enhanced superoxide  
335 production in the cytochrome *b<sub>6</sub>f* complex of oxygenic photosynthesis. *Biochemistry* **52**, 8975-  
336 8983 <https://doi.org/10.1021/bi4013534>
- 337
- 338 7 M.M. Mubarakshina and B.N. Ivanov (2010) The production and scavenging of reactive oxygen  
339 species in the plastoquinone pool of chloroplast thylakoid membranes. *Physiol. Plant.* **140**,  
340 103-110 <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1399-3054.2010.01391.x>
- 341
- 342 8 N. Smirnov and D. Arnaud (2019) Hydrogen peroxide metabolism and functions in plants. *New*  
343 *Phytol.* **221**, 1197-1214 <https://nph.onlinelibrary.wiley.com/doi/abs/10.1111/nph.15488>

- 344  
345 9 P. Wardman (1990) Bioreductive activation of quinones: Redox properties and thiol reactivity. *Free*  
346 *Radical Res. Commun.* **8**, 219-229 <https://doi.org/10.3109/10715769009053355>
- 347  
348 10 M. Kozuleva, A. Petrova, Y. Milrad, A. Semenov, B. Ivanov, K.E. Redding, et al. (2021) Phylloquinone  
349 is the principal Mehler reaction site within photosystem I in high light. *Plant Physiol.* **186**, 1848-  
350 1858 <https://doi.org/10.1093/plphys/kiab221>
- 351  
352 11 M.-A. Takahashi and K. Asada (1983) Superoxide anion permeability of phospholipid membranes  
353 and chloroplast thylakoids. *Arch. Biochem. Biophys.* **226**, 558-566  
354 <https://www.sciencedirect.com/science/article/pii/0003986183903259>
- 355  
356 12 B.L. Upham and L.S. Jahnke (1986) Photooxidative reactions in chloroplast thylakoids. Evidence  
357 for a Fenton-type reaction promoted by superoxide or ascorbate. *Photosynth. Res.* **8**, 235-247  
358 <https://doi.org/10.1007/BF00037131>
- 359  
360 13 K. Sonoike (2011) Photoinhibition of photosystem I. *Physiol. Plant.* **142**, 56-64  
361 <http://www.ncbi.nlm.nih.gov/pubmed/21128947>
- 362  
363 14 H. Hashimoto, C. Uragami and R.J. Cogdell, Carotenoids and photosynthesis, in: C. Stange (Ed.)  
364 Carotenoids in Nature: Biosynthesis, Regulation and Function, Springer International  
365 Publishing, Cham, 2016, pp. 111-139 [https://doi.org/10.1007/978-3-319-39126-7\\_4](https://doi.org/10.1007/978-3-319-39126-7_4)
- 366  
367 15 K. Brinkert, S. De Causmaecker, A. Krieger-Liszkay, A. Fantuzzi and A.W. Rutherford (2016)  
368 Bicarbonate-induced redox tuning in Photosystem II for regulation and protection. *Proc. Natl.*  
369 *Acad. Sci. U.S.A.* **113**, 12144-12149 <http://www.pnas.org/content/113/43/12144.abstract>
- 370  
371 16 A.W. Rutherford and A. Krieger-Liszkay (2001) Herbicide-induced oxidative stress in photosystem  
372 II. *Trends Biochem. Sci.* **26**, 648-653  
373 <http://www.sciencedirect.com/science/article/pii/S0968000401019533>
- 374  
375 17 G.N. Johnson, A.W. Rutherford and A. Krieger (1995) A change in the midpoint potential of the  
376 quinone Q<sub>A</sub> in photosystem II associated with photoactivation of oxygen evolution. *Biochim.*  
377 *Biophys. Acta Bioenerg.* **1229**, 202-207  
378 <https://www.sciencedirect.com/science/article/pii/0005272895000032>
- 379  
380 18 A. Krieger, A.W. Rutherford and G.N. Johnson (1995) On the determination of redox midpoint  
381 potential of the primary quinone electron acceptor, Q<sub>A</sub>, in Photosystem II. *Biochim. Biophys.*  
382 *Acta Bioenerg.* **1229**, 193-201  
383 <https://www.sciencedirect.com/science/article/pii/000527289500002Z>
- 384  
385 19 A. Krieger and E. Weis (1993) The role of calcium in the pH-dependent control of Photosystem II.  
386 *Photosynth. Res.* **37**, 117-130 <http://dx.doi.org/10.1007/BF02187470>
- 387

- 388 20 Y. Kato and T. Noguchi (2021) Effects of stromal and luminal side perturbations on the redox  
389 potential of the primary quinone electron acceptor Q<sub>A</sub> in photosystem II. *Biochemistry* **60**,  
390 3697-3706 <https://doi.org/10.1021/acs.biochem.1c00624>
- 391
- 392 21 A. Krieger, A.W. Rutherford, I. Vass and É. Hideg (1998) Relationship between Activity, D1 Loss,  
393 and Mn Binding in Photoinhibition of Photosystem II. *Biochemistry* **37**, 16262-16269  
394 <https://doi.org/10.1021/bi981243v>
- 395
- 396 22 N. Keren, I. Ohad, A.W. Rutherford, F. Drepper and A. Krieger-Liszkay (2000) Inhibition of  
397 photosystem II activity by saturating single turnover flashes in calcium-depleted and active  
398 photosystem II. *Photosynth. Res.* **63**, 209-216 <http://dx.doi.org/10.1023/A:1006435530817>
- 399
- 400 23 J. Zabret, S. Bohn, S.K. Schuller, O. Arnolds, M. Möller, J. Meier-Credo, et al. (2021) Structural  
401 insights into photosystem II assembly. *Nat. Plants* **7**, 524-538 [https://doi.org/10.1038/s41477-](https://doi.org/10.1038/s41477-021-00895-0)  
402 [021-00895-0](https://doi.org/10.1038/s41477-021-00895-0)
- 403
- 404 24 A. Krieger-Liszkay, C. Fufezan and A. Trebst (2008) Singlet oxygen production in photosystem II  
405 and related protection mechanism. *Photosynth. Res.* **98**, 551-564  
406 <https://doi.org/10.1007/s11120-008-9349-3>
- 407
- 408 25 L. Leverne and A. Krieger-Liszkay (2021) Moderate drought stress stabilizes the primary quinone  
409 acceptor QA and the secondary quinone acceptor QB in photosystem II. *Physiol. Plant.* **171**,  
410 260-267 <https://onlinelibrary.wiley.com/doi/abs/10.1111/ppl.13286>
- 411
- 412 26 M. Messant, S. Timm, A. Fantuzzi, W. Weckwerth, H. Bauwe, A.W. Rutherford, et al. (2018)  
413 Glycolate induces redox tuning of photosystem II in vivo: Study of a photorespiration mutant.  
414 *Plant Physiol.* **177**, 1277-1285 <https://doi.org/10.1104/pp.18.00341>
- 415
- 416 27 I. Vass (2011) Role of charge recombination processes in photodamage and photoprotection of  
417 the photosystem II complex. *Physiol. Plant.* **142**, 6-16  
418 <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1399-3054.2011.01454.x>
- 419
- 420 28 P. Pospíšil (2016) Production of reactive oxygen species by photosystem II as a response to light  
421 and temperature stress. *Front. Plant Sci.* **7**,  
422 <https://www.frontiersin.org/article/10.3389/fpls.2016.01950>
- 423
- 424 29 R. Kale, A.E. Hebert, L.K. Frankel, L. Sallans, T.M. Bricker and P. Pospíšil (2017) Amino acid oxidation  
425 of the D1 and D2 proteins by oxygen radicals during photoinhibition of Photosystem II. *Proc.*  
426 *Natl. Acad. Sci. U.S.A.* **114**, 2988-2993 <http://www.pnas.org/content/114/11/2988.abstract>
- 427
- 428 30 A. Zavafer, M.H. Cheah, W. Hillier, W.S. Chow and S. Takahashi (2015) Photodamage to the oxygen  
429 evolving complex of photosystem II by visible light. *Sci. Rep.* **5**, 16363  
430 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4642293/>

431

- 432 31 I. Vass, E. Turcsányi, E. Touloupakis, D. Ghanotakis and V. Petrouleas (2002) The mechanism of  
433 UV-A radiation-induced inhibition of photosystem II electron transport studied by EPR and  
434 chlorophyll fluorescence. *Biochemistry* **41**, 10200-10208 <https://doi.org/10.1021/bi020272+>  
435
- 436 32 S. Järvi, M. Suorsa and E.-M. Aro (2015) Photosystem II repair in plant chloroplasts — Regulation,  
437 assisting proteins and shared components with photosystem II biogenesis. *Biochim. Biophys.*  
438 *Acta Bioenerg.* **1847**, 900-909  
439 <http://www.sciencedirect.com/science/article/pii/S0005272815000171>
- 440
- 441 33 H. Jimbo, R. Yutthanasirikul, T. Nagano, T. Hisabori, Y. Hihara and Y. Nishiyama (2018) Oxidation  
442 of translation factor EF-Tu inhibits the repair of photosystem II. *Plant Physiol.* **176**, 2691-2699  
443 <http://www.plantphysiol.org/content/plantphysiol/176/4/2691.full.pdf>
- 444
- 445 34 K. Sonoike, I. Terashima, M. Iwaki and S. Itoh (1995) Destruction of photosystem I iron-sulfur  
446 centers in leaves of *Cucumis sativus* L. by weak illumination at chilling temperatures. *FEBS Lett.*  
447 **362**, 235-238 <http://www.ncbi.nlm.nih.gov/pubmed/7720879>
- 448
- 449 35 A. Tiwari, F. Mamedov, M. Grieco, M. Suorsa, A. Jajoo, S. Styring, et al. (2016) Photodamage of  
450 iron-sulphur clusters in photosystem I induces non-photochemical energy dissipation. *Nat.*  
451 *Plants* **2**, 16035 <http://dx.doi.org/10.1038/nplants.2016.35>
- 452
- 453 36 G. Shimakawa and C. Miyake (2019) What quantity of photosystem I is optimum for safe  
454 photosynthesis? *Plant Physiol.* **179**, 1479-1485  
455 <http://www.plantphysiol.org/content/plantphysiol/179/4/1479.full.pdf>
- 456
- 457 37 H. Kudoh and K. Sonoike (2002) Irreversible damage to photosystem I by chilling in the light: cause  
458 of the degradation of chlorophyll after returning to normal growth temperature. *Planta* **215**,  
459 541-548 <http://www.ncbi.nlm.nih.gov/pubmed/12172835>
- 460
- 461 38 M. Zivcak, M. Brestic, K. Kunderlikova, O. Sytar and S.I. Allakhverdiev (2015) Repetitive light pulse-  
462 induced photoinhibition of photosystem I severely affects CO<sub>2</sub> assimilation and  
463 photoprotection in wheat leaves. *Photosynth. Res.* **126**, 449-463  
464 <http://www.ncbi.nlm.nih.gov/pubmed/25829027>
- 465
- 466 39 T. Sejima, D. Takagi, H. Fukayama, A. Makino and C. Miyake (2014) Repetitive short-pulse light  
467 mainly inactivates photosystem I in sunflower leaves. *Plant Cell Physiol.* **55**, 1184-1193  
468 <http://www.ncbi.nlm.nih.gov/pubmed/24793753>
- 469
- 470 40 I. Terashima, S. Funayama and K. Sonoike (1994) The site of photoinhibition in leaves of *Cucumis*  
471 *sativus* L. at low temperatures is photosystem I, not photosystem II. *Planta* **193**, 300-306  
472 <http://dx.doi.org/10.1007/BF00192544>
- 473
- 474 41 M. Suorsa, S. Jarvi, M. Grieco, M. Nurmi, M. Pietrzykowska, M. Rantala, et al. (2012) PROTON  
475 GRADIENT REGULATION5 is essential for proper acclimation of Arabidopsis photosystem I to


- 476 naturally and artificially fluctuating light conditions. *Plant Cell* **24**, 2934-2948  
477 <http://www.ncbi.nlm.nih.gov/pubmed/22822205>
- 478
- 479 42 Y. Munekage, M. Hojo, J. Meurer, T. Endo, M. Tasaka and T. Shikanai (2002) PGR5 is involved in  
480 cyclic electron flow around photosystem i and is essential for photoprotection in Arabidopsis.  
481 *Cell* **110**, 361-371 <http://www.sciencedirect.com/science/article/pii/S009286740200867X>
- 482
- 483 43 B. Nowicka, A. Trela-Makowej, D. Latowski, K. Strzalka and R. Szymańska (2021) Antioxidant and  
484 signaling role of plastid-derived isoprenoid quinones and chromanols. *Int. J. Mol. Sci.* **22**, 2950  
485 <https://www.mdpi.com/1422-0067/22/6/2950>
- 486
- 487 44 B. Demmig-Adams, A.M. Gilmore and W.W.A. Iii (1996) In vivo functions of carotenoids in higher  
488 plants. *FASEB J.* **10**, 403-412  
489 <https://faseb.onlinelibrary.wiley.com/doi/abs/10.1096/fasebj.10.4.8647339>
- 490
- 491 45 S. Cazzaniga, Z. Li, K.K. Niyogi, R. Bassi and L. Dall'Osto (2012) The Arabidopsis *szl1* mutant reveals  
492 a critical role of  $\beta$ -carotene in photosystem i photoprotection. *Plant Physiol.* **159**, 1745-1758  
493 <http://www.plantphysiol.org/content/159/4/1745.abstract>
- 494
- 495 46 A. Kumar, A. Prasad, M. Sedlářová, R. Kale, L.K. Frankel, L. Sallans, et al. (2021) Tocopherol controls  
496 D1 amino acid oxidation by oxygen radicals in Photosystem II. *Proc. Natl. Acad. Sci. U.S.A.* **118**,  
497 e2019246118 <https://www.pnas.org/content/pnas/118/4/e2019246118.full.pdf>
- 498
- 499 47 A. Kumar, A. Prasad, M. Sedlářová, B. Ksas, M. Havaux and P. Pospíšil (2020) Interplay between  
500 antioxidants in response to photooxidative stress in Arabidopsis. *Free Radical Biol. Med.* **160**,  
501 894-907 <https://www.sciencedirect.com/science/article/pii/S089158492031234X>
- 502
- 503 48 A. Sedoud, R. López-Igual, A. ur Rehman, A. Wilson, F. Perreau, C. Boulay, et al. (2014) The  
504 cyanobacterial photoactive orange carotenoid protein is an excellent singlet oxygen quencher.  
505 *Plant Cell* **26**, 1781-1791 <http://www.plantcell.org/content/26/4/1781.abstract>
- 506
- 507 49 F. Muzzopappa and D. Kirilovsky (2020) Changing color for photoprotection: The orange  
508 carotenoid protein. *Trends Plant Sci.* **25**, 92-104  
509 <https://www.sciencedirect.com/science/article/pii/S1360138519302559>
- 510
- 511 50 K. Schmidt, C. Fufezan, A. Krieger-Liszkay, H. Satoh and H. Paulsen (2003) Recombinant water-  
512 soluble chlorophyll protein from *Brassica oleracea* var. Botrys binds various chlorophyll  
513 derivatives. *Biochemistry* **42**, 7427-7433 <https://doi.org/10.1021/bi034207r>
- 514
- 515 51 S. Takahashi, H. Bauwe and M. Badger (2007) Impairment of the photorespiratory pathway  
516 accelerates photoinhibition of photosystem II by suppression of repair but not acceleration of  
517 damage processes in Arabidopsis. *Plant Physiol.* **144**, 487-494  
518 <http://www.ncbi.nlm.nih.gov/pubmed/17400706>
- 519

- 520 52 C.H. Foyer, J. Neukermans, G. Queval, G. Noctor and J. Harbinson (2012) Photosynthetic control  
521 of electron transport and the regulation of gene expression. *J. Exp. Bot.* **63**, 1637-1661  
522 <https://dx.doi.org/10.1093/jxb/ers013>
- 523
- 524 53 A.N. Tikhonov (2013) pH-Dependent regulation of electron transport and ATP synthesis in  
525 chloroplasts. *Photosynth. Res.* **116**, 511-534 <http://dx.doi.org/10.1007/s11120-013-9845-y>
- 526
- 527 54 M.A. Schöttler, S.Z. Tóth, A. Boulouis and S. Kahlau (2015) Photosynthetic complex stoichiometry  
528 dynamics in higher plants: biogenesis, function, and turnover of ATP synthase and the  
529 cytochrome *b<sub>6</sub>f* complex. *J. Exp. Bot.* **66**, 2373-2400 <http://dx.doi.org/10.1093/jxb/eru495>
- 530
- 531 55 K. Takizawa, J.A. Cruz, A. Kanazawa and D.M. Kramer (2007) The thylakoid proton motive force in  
532 vivo. Quantitative, non-invasive probes, energetics, and regulatory consequences of light-  
533 induced pmf. *Biochim. Biophys. Acta Bioenerg.* **1767**, 1233-1244  
534 <http://www.sciencedirect.com/science/article/pii/S0005272807001557>
- 535
- 536 56 E. Schlodder and B. Meyer (1987) pH dependence of oxygen evolution and reduction kinetics of  
537 photooxidized chlorophyll *a<sub>II</sub>* (P-680) in Photosystem II particles from *Synechococcus* sp.  
538 *Biochim. Biophys. Acta Bioenerg.* **890**, 23-31  
539 <https://www.sciencedirect.com/science/article/pii/0005272887900648>
- 540
- 541 57 A. Krieger, E. Weis and S. Demeter (1993) Low-pH-induced Ca<sup>2+</sup> ion release in the water-splitting  
542 system is accompanied by a shift in the midpoint redox potential of the primary quinone  
543 acceptor Q<sub>A</sub>. *Biochim. Biophys. Acta Bioenerg.* **1144**, 411-418  
544 <http://www.sciencedirect.com/science/article/pii/0005272893901283>
- 545
- 546 58 N.C.M. Magdaong and R.E. Blankenship (2018) Photoprotective, excited-state quenching  
547 mechanisms in diverse photosynthetic organisms. *J. Biol. Chem.* **293**, 5018-5025  
548 <https://doi.org/10.1074/jbc.TM117.000233>
- 549
- 550 59 A.V. Ruban and S. Wilson (2020) The mechanism of non-photochemical quenching in plants:  
551 Localization and driving forces. *Plant Cell Physiol.* **62**, 1063-1072  
552 <https://doi.org/10.1093/pcp/pcaa155>
- 553
- 554 60 J.M. Buck, P.G. Kroth and B. Lepetit (2021) Identification of sequence motifs in Lhcx proteins that  
555 confer qE-based photoprotection in the diatom *Phaeodactylum tricornutum*. *Plant J.* **108**,  
556 1721-1734 <https://onlinelibrary.wiley.com/doi/abs/10.1111/tpj.15539>
- 557
- 558 61 W.J. Nawrocki, B. Bailleul, P. Cardol, F. Rappaport, F.A. Wollman and P. Joliot (2019) Maximal  
559 cyclic electron flow rate is independent of PGRL1 in *Chlamydomonas*. *Biochim. Biophys. Acta*  
560 *Bioenerg.* **1860**, 425-432  
561 <http://www.sciencedirect.com/science/article/pii/S0005272818302524>
- 562




- 563 62 J.M. Schuller, J.A. Birrell, H. Tanaka, T. Konuma, H. Wulfhorst, N. Cox, et al. (2019) Structural  
564 adaptations of photosynthetic complex I enable ferredoxin-dependent electron transfer.  
565 *Science* **363**, 257-260 <http://science.sciencemag.org/content/sci/363/6424/257.full.pdf>
- 566
- 567 63 Y. Helman, D. Tchernov, L. Reinhold, M. Shibata, T. Ogawa, R. Schwarz, et al. (2003) Genes  
568 encoding A-type flavoproteins are essential for photoreduction of O<sub>2</sub> in cyanobacteria. *Curr.*  
569 *Biol.* **13**, 230-235 <http://www.sciencedirect.com/science/article/pii/S0960982203000460>
- 570
- 571 64 A. Santana-Sanchez, D. Solymosi, H. Mustila, L. Bersanini, E.-M. Aro and Y. Allahverdiyeva (2019)  
572 Flavodiiron proteins 1-to-4 function in versatile combinations in O<sub>2</sub> photoreduction in  
573 cyanobacteria. *eLife* **8**, e45766 <https://doi.org/10.7554/eLife.45766>
- 574
- 575 65 F. Chaux, A. Burlacot, M. Mekhalfi, P. Auroy, S. Blangy, P. Richaud, et al. (2017) Flavodiiron proteins  
576 promote fast and transient O<sub>2</sub> photoreduction in *Chlamydomonas*. *Plant Physiol.* **174**, 1825-  
577 1836 <https://doi.org/10.1104/pp.17.00421>
- 578
- 579 66 P. Sétif, G. Shimakawa, A. Krieger-Liszkay and C. Miyake (2020) Identification of the electron donor  
580 to flavodiiron proteins in *Synechocystis* sp. PCC 6803 by *in vivo* spectroscopy. *Biochim. Biophys.*  
581 *Acta Bioenerg.* **1861**, 148256  
582 <http://www.sciencedirect.com/science/article/pii/S0005272820301067>
- 583
- 584 67 A. Alboresi, M. Storti, L. Cendron and T. Morosinotto (2019) Role and regulation of class-C  
585 flavodiiron proteins in photosynthetic organisms. *Biochem. J.* **476**, 2487-2498  
586 <https://doi.org/10.1042/BCJ20180648>
- 587
- 588 68 G. Shimakawa, H. Hanawa, S. Wada, G.T. Hanke, Y. Matsuda and C. Miyake (2021) Physiological  
589 roles of flavodiiron proteins and photorespiration in the liverwort *Marchantia polymorpha*.  
590 *Front. Plant Sci.* **12**, <https://www.frontiersin.org/article/10.3389/fpls.2021.668805>
- 591
- 592 69 L. Michelet and A. Krieger-Liszkay (2012) Reactive oxygen intermediates produced by  
593 photosynthetic electron transport are enhanced in short-day grown plants. *Biochim. Biophys.*  
594 *Acta Bioenerg.* **1817**, 1306-1313  
595 <http://www.sciencedirect.com/science/article/pii/S0005272811002866>
- 596
- 597 70 A. Courteille, S. Vesa, R. Sanz-Barrio, A.-C. Cazalé, N. Becuwe-Linka, I. Farran, et al. (2012)  
598 Thioredoxin *m*<sub>4</sub> controls photosynthetic alternative electron pathways in Arabidopsis *Plant*  
599 *Physiol.* **161**, 508-520 <https://doi.org/10.1104/pp.112.207019>
- 600
- 601 71 L. Nikkanen, J. Toivola, A. Trotta, M.G. Diaz, M. Tikkanen, E.-M. Aro, et al. (2018) Regulation of  
602 cyclic electron flow by chloroplast NADPH-dependent thioredoxin system. *Plant Direct* **2**,  
603 e00093 <https://onlinelibrary.wiley.com/doi/abs/10.1002/pld3.93>
- 604
- 605 72 Y. Okegawa and K. Motohashi (2020) M-type thioredoxins regulate the PGR5/PGRL1-dependent  
606 pathway by forming a disulfide-linked complex with PGRL1. *Plant Cell* **32**, 3866-3883  
607 <http://www.plantcell.org/content/plantcell/32/12/3866.full.pdf>

- 608  
609 73 B.-C. Wolf, T. Isaacson, V. Tiwari, I. Dangoor, S. Mufkadi and A. Danon (2020) Redox regulation of  
610 PGRL1 at the onset of low light intensity. *Plant J.* **103**, 715-725  
611 <https://onlinelibrary.wiley.com/doi/abs/10.1111/tpj.14764>
- 612  
613 74 A.P. Hertle, T. Blunder, T. Wunder, P. Pesaresi, M. Pribil, U. Armbruster, et al. (2013) PGRL1 is the  
614 elusive ferredoxin-plastoquinone reductase in photosynthetic cyclic electron flow. *Mol. Cell*  
615 **49**, 511-523 <http://www.sciencedirect.com/science/article/pii/S1097276512009896>
- 616  
617 75 Q. Yu, K. Feilke, A. Krieger-Liszkay and P. Beyer (2014) Functional and molecular characterization  
618 of plastid terminal oxidase from rice (*Oryza sativa*). *Biochim. Biophys. Acta Bioenerg.* **1837**,  
619 1284-1292 <http://www.sciencedirect.com/science/article/pii/S0005272814001236>
- 620  
621 76 G.N. Johnson and P. Stepien (2016) Plastid terminal oxidase as a route to improving plant stress  
622 tolerance: Known knowns and known unknowns. *Plant Cell Physiol.* **57**, 1387-1396  
623 <http://dx.doi.org/10.1093/pcp/pcw042>
- 624  
625 77 I. Rog, A.K. Chaturvedi, V. Tiwari and A. Danon (2021) Low light-regulated intramolecular disulfide  
626 fine-tunes the role of PTOX in Arabidopsis. *Plant J.* **109**, 585-597  
627 <https://doi.org/10.1111/tpj.15579>
- 628  
629 78 S. Bolte, E. Marcon, M. Jaunario, L. Moyet, M. Paternostre, M. Kuntz, et al. (2020) Dynamics of  
630 the localization of the plastid terminal oxidase inside the chloroplast. *J. Exp. Bot.* **71**, 2661-  
631 2669 <https://doi.org/10.1093/jxb/eraa074>
- 632  
633 79 V. Dogra and C. Kim (2020) Singlet oxygen metabolism: From genesis to signaling. *Front. Plant Sci.*  
634 **10**, <https://www.frontiersin.org/article/10.3389/fpls.2019.01640>
- 635  
636 80 L. Wang, D. Leister, L. Guan, Y. Zheng, K. Schneider, M. Lehmann, et al. (2020) The *Arabidopsis*  
637 SAFEGUARD1 suppresses singlet oxygen-induced stress responses by protecting grana  
638 margins. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 6918-6927  
639 <https://www.pnas.org/content/pnas/117/12/6918.full.pdf>
- 640  
641



**Figure 1. Photosynthetic electron transport in thylakoid membranes.**

The reaction centre chlorophyll dimers P680 and P700, respectively in photosystem (PS) II and I, are photo-oxidised by light, resulting in photosynthetic electron transport from H<sub>2</sub>O to NADP<sup>+</sup> through the interchain components, including plastoquinone (PQ), cytochrome *b<sub>6</sub>f* complex (*cyt b<sub>6</sub>f*) and plastocyanin (PC). In the acceptor side of PSI, ferredoxin-NADP<sup>+</sup> oxidoreductase (FNR) donates electrons from ferredoxin (Fd) to NADP<sup>+</sup>. The different concentration of H<sup>+</sup> across thylakoid membranes is the motive force for ATP production by ATP synthase (ATPase). Singlet oxygen (<sup>1</sup>O<sub>2</sub>) can be mainly generated at the light-harvesting complex (LHCII) and the reaction centre of PSII. Superoxide anion radical (O<sub>2</sub><sup>•-</sup>) can be mainly generated at the acceptor side of PSI, and also in PSII and *cyt b<sub>6</sub>f*.



**Figure 2. Generation of reactive oxygen species in the reaction centres of photosystems.**

Whereas in photosystem (PS) II, the primary electron transport reaction occurs in the side of D1 protein, in PSI two branches function for the electron transport in PsaA and PsaB proteins. Singlet oxygen (<sup>1</sup>O<sub>2</sub>) is generated by the triplet state of P680 produced by the charge recombination of P680<sup>+</sup> and Ph<sub>D1</sub><sup>-</sup>. Superoxide anion radical (O<sub>2</sub><sup>•-</sup>) is generated by the Mehler reaction in the A-branch phylloquinone (A<sub>1A</sub>). Abbreviations are as follows: Chl, chlorophyll; Ph, pheophytin; Q, quinone; PhQ, phylloquinone; F, [4Fe-4S] cluster.