



HAL
open science

Comments on “A Simple, Rapid, Comparative Evaluation of Multiple Products for Decontamination of Actinide-contaminated Rat Skin Ex Vivo “

Celine Bouvier Capely, Guillaume Phan

► **To cite this version:**

Celine Bouvier Capely, Guillaume Phan. Comments on “A Simple, Rapid, Comparative Evaluation of Multiple Products for Decontamination of Actinide-contaminated Rat Skin Ex Vivo “. Health Physics, 2022, 123 (2), pp.NC. 10.1097/HP.0000000000001575 . hal-03745211

HAL Id: hal-03745211

<https://hal.science/hal-03745211v1>

Submitted on 3 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Céline Bouvier-Capely

celine.bouvier@irsn.fr

Guillaume Phan

guillaume.phan@irsn.fr

Institut de Radioprotection et Sécurité Nucléaire, PSE-SANTE/SESANE/LRSI

31 avenue de la Division Leclerc

92260 Fontenay-aux-Roses, France

Title: Comments on “A Simple, Rapid, Comparative Evaluation of Multiple Products for Decontamination of Actinide-contaminated Rat Skin *Ex Vivo*”

Dear Editors,

This letter is regarding the paper written by Griffiths et al. (Griffiths et al., 2022). We read with interest this article describing a simple and rapid test to compare the efficiency of various products to remove cutaneous contamination by actinides with a rat skin *ex vivo* model.

We would like to draw your attention to some experimental biases in the work of Dr Griffiths and her co-authors that may mislead the interpretations of the published results.

According to our analysis of this paper, we have 3 main remarks.

Our first and main remark relates on the difference in the modalities of treatments application on the skin after the contamination step:

Number of treatment applications: According to the materials and methods section and Table 1, liquid products (water, DTPA 5 mM, or prediluted Trait Rouge) were applied three times on the contaminated area (3×0.5 mL) whilst other products (calixarene nanoemulsion, osmogel, hyaluronic acid or wound dressings) were applied once and rinsed three times by water wash (3×0.5 mL H₂O). Thus, the performances of these liquid products could be overestimated and attributed to an undue greater number of applications compared to other products (gels and wound dressings).

Volume/amount of treatment: various volumes of treatments have been used in this study without explanation about these differences (0.5 mL for liquid treatments, 0.1 g for calixarene nanoemulsion, 0.3 g for gel products (osmogel, hyaluronic acid and algoplaque paste), 1 g for French green clay).

Treatment application on dry or wetted skin: It was not clearly stated in the Figures 2 and 3 whether the different products were applied on wetted skin or dry skin. Skin wetting seems to be crucial condition for gel or low fluidity products (such as calixarene nanoemulsion, osmogel, hyaluronic acid or dressings) as mentioned by the authors themselves, and probably less so for aqueous products (water, DTPA, prediluted Trait Rouge). According to the results mentioned in the discussion section and focused on the comparison of the efficacy between Trait Rouge and calixarene nanoemulsion treatments and cited hereafter:

“Similarly, for Pu 17.3 ± 3.1 % remained in skin after Trait Rouge treatment and 15.6 ± 2.1 % (dry skin) or 2.8 ± 1.2 % (wetted skin) after calixarene nanoemulsion”,

it seems that the results of Figure 2 refer to application of calixarene nanoemulsion on dry skin, despite the protocol described by the authors in Material and Methods section: “the manufacturer’s instructions (Cevibra) were followed: the contaminated skin area was wetted with 100 μ L water and the calixarene nanoemulsion (100 mg) was added”.

Consequently, we believe that the differences in application conditions (different volumes and amounts of each product, number of applications, wetted or dry skin) makes the comparison of the published results dubious. This clearly limits the interpretation of these results. Since the applications

differ from fluid products (water, DTPA, Trait Rouge) to less fluid products (Osmogel, Calixarene, wound dressings), the statistical analyses and comparison presented in Table 2 are irrelevant. The authors should have used other tests such as non-parametric or ANOVA tests, or paired Student t-test for products of the same category (fluid or less fluid).

Secondly, the proposed *ex vivo* model of the Griffiths' study may be simple and rapid but does not comply with the Association Française de Normalisation (AFNOR), which recommends the use of Franz diffusion cells and porcine skin which is a better surrogate for human skin in order to assess the performance of "nuclear, radiological, biological, chemical" decontamination products on intact skin (AFNOR 2016). Yet, the authors have already used such protocols in their previous studies (Tazart et al. 2018) but this was not mentioned in the discussion. Franz diffusion cells are relevant for the assessment of the potential diffusion of contaminants through the skin when the actinides are deposited for a long time such as 2h.

Our last remark concerns the citation of the reference Grives et al 2017 (Grivès et al. 2017) at page 9. The authors have referenced the wrong paper and have misinterpreted the results. In the referenced study, contaminations were performed *in vivo* on wounds model in the rats and not on intact skin as in the present study. The authors probably confused it with the paper Grives et al 2015 (Grives et al. 2015).

To conclude, the authors should reinterpret the results of the experiments in which the products were applied in the same conditions and clearly state that the application conditions are not completely comparable. Otherwise, the authors could perform supplementary tests, ideally on Franz diffusion cells for fair comparison purpose (for example Trait Rouge and calixarene nanoemulsion applied both not diluted and in the same amount/volume on dry skin or wetted skin, or dilute 1:1 both products Trait Rouge and calixarene nanoemulsion before their application).

We thank you for taking into consideration our comments and suggestions.

Sincerely,

Dr Céline Bouvier-Capely and Dr Guillaume Phan,

Fontenay-aux-Roses, January 24th 2022

The authors declare to have developed the calixarene nanoemulsion with financial support of Direction Générale de l'Armement (DGA), Centre National de la Recherche Scientifique (CNRS) and Institut de Radioprotection et Sûreté Nucléaire (IRSN). This product has been patented and is now commercialized by Cevibra Laboratory.

References

- AFNOR. Security and protection of citizens: Cbrn-e-protocols for performance assessment of technologies intended for undamaged skin decontamination. NF X32-122; 2016.
- Griffiths N M, Devilliers K, Laroche P, Van der Meer A. A Simple, Rapid, Comparative Evaluation of Multiple Products for Decontamination of Actinide-contaminated Rat Skin *Ex Vivo* 122(3): 371-382; 2022.
- Grivès S, Phan G, Bouvier-Capely C, Suhard D, Rebière F, Agarande M, Fattal E. Compared *in vivo* efficiency of nanoemulsions unloaded and loaded with calixarene and soapy water in the treatment of superficial wounds contaminated by uranium. *Chemico-Biological Interactions* 267: 33-39; 2017.

- Grivès S, Phan G, Morat G, Suhard D, Rebiere F, Fattal E. Ex vivo uranium decontamination efficiency on wounded skin and in vitro skin toxicity of a calixarene-loaded nanoemulsion. *Journal of Pharmaceutical Sciences* 104: 2008-2017; 2015.
- Tazart A, Bolzinger MA, Lamart S, Coudert S, Angulo JF, Jandard V, Briançon S, Griffiths NM. Actinide-contaminated skin: Comparing decontamination efficacy of water, cleansing gels, and DTPA gels. *Health Physics* 115: 12-20; 2018.