



**HAL**  
open science

## **TCDD aggravates the formation of the atherosclerotic plaque in ApoE KO mice with a sexual dimorphic pattern**

Laetitia Bey, Xavier Coumoul, Min Ji Kim

► **To cite this version:**

Laetitia Bey, Xavier Coumoul, Min Ji Kim. TCDD aggravates the formation of the atherosclerotic plaque in ApoE KO mice with a sexual dimorphic pattern. *Biochimie*, 2022, 195, pp.54-58. 10.1016/j.biochi.2022.01.012 . hal-03745304

**HAL Id: hal-03745304**

**<https://hal.science/hal-03745304v1>**

Submitted on 5 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **TCDD aggravates the formation of the atherosclerotic plaque in ApoE KO mice with a**  
2 **sexual dimorphic pattern**

3

4 **Laetitia Bey<sup>1</sup>, Xavier Coumoul<sup>1</sup>, Min Ji Kim<sup>2</sup>**

5

6 **Institutions :**

7 <sup>1</sup> Université de Paris, INSERM UMR-S1124, Paris, France

8 <sup>2</sup> Université Sorbonne Paris Nord, Bobigny, INSERM UMR-S1124, Paris, France

9 **Correspondance: Min Ji Kim** email: [min-ji.kim@inserm.fr](mailto:min-ji.kim@inserm.fr)

10

11 **Keywords:** atherosclerosis, AhR, TCDD, sexual dimorphism

12

13 **Abbreviations**

14 AhR: Aryl hydrocarbon receptor

15 ApoE KO: Apolipoprotein E Knock-Out

16 bHLH/PAS: basic Helix Loop Helix/Per ARNT Sim

17 CD68: *Cluster* of Differentiation 68

18 ICAM1: InterCellular Adhesion Molecule 1

19 MCP1: Monocyte Chemoattractant Protein 1

20 SEM: standard error of the mean

21  $\alpha$ -SMA:  $\alpha$ -smooth muscle actin

22 TCDD: 2,3,7,8-Tetrachlorodibenzo-*para*-dioxin

23 VCAM1: Vascular Cell Adhesion Molecule 1

24 VSMCs: vascular smooth muscle cells

25

26

27 **Abstract:**

28 Aryl hydrocarbon receptor (AhR) ligands are recognized as aggravating factors in  
29 cardiovascular diseases but little is known about the role of the AhR in atherosclerosis  
30 considering the effects of age and gender. We exposed male and female ApoE knock-out  
31 mice, a model to study the pathogenesis of atherosclerosis, to a potent AhR ligand, 2,3,7,8-  
32 tetrachlorodibenzo-*p*-dioxin (TCDD) by an intraperitoneal injection of 1 µg/kg/week for 8  
33 weeks. Atherosclerotic lesions, histological parameters and critical atherosclerotic markers in  
34 aorta were analysed. TCDD increased atherogenic lesions in 35-week old female mice,  
35 leading to a switch of vascular smooth muscle cells (VSMCs) from a contractile to a pro-  
36 atherogenic phenotype and increased expression for VCAM1. AhR activation accelerates the  
37 formation of atherosclerotic plaques with sex and age differences due to the phenotypical  
38 switch of VSMCs.

39

## 40 **1. Introduction**

41 The AhR is a bHLH/PAS transcriptional factor and a xenobiotic receptor. It is of interest in  
42 the fields of toxicology because of its exogenous agonists such as 2,3,7,8-tetrachlorodibenzo-  
43 *p*-dioxin (TCDD), a potent carcinogen. In addition to the environmental chemicals, dietary  
44 and endogenous AhR ligands have been identified including tryptophan metabolites and  
45 flavonoids [1]. After ligand binding, AhR transactivates the expression of xenobiotic  
46 metabolizing enzymes and many other genes involved in cell proliferation and inflammatory  
47 responses. Endogenous ligands are inferred to be involved in the regulation of cardiac  
48 functions since vascular abnormalities are among the most common phenotypes observed in  
49 AhR knock-out (KO) mice [2–4]. Epidemiological studies suggest the role of AhR in  
50 dyslipidemia, hypertension, carotid atherosclerosis and myocardial ischemia [5,6] and mouse  
51 studies have reported a role for AhR in the increased expression of inflammatory genes in the  
52 aorta [7,8]. One of the mechanisms by which AhR could play a role in atherosclerosis is the  
53 modulation of inflammatory cytokines, observed in multiple models [9–12].

54 Atherosclerosis results from the accumulation of lipids in the arterial wall with the  
55 occurrence of atheromatous plaques which disrupt the permeability of the vessel wall leading  
56 to infiltration of low-density lipoproteins whose oxidized forms are specifically captured by  
57 infiltrated macrophages. This last process is stimulated by the expression of biomarkers  
58 including VCAM1 and ICAM1, two adhesion proteins which attract monocytes which then  
59 differentiate into macrophages. Macrophages and vascular smooth muscle cells (VSMCs)  
60 differentiate into CD68-expressing foam cells following the internalization of oxidized LDLs  
61 [13,14]. Normally present in the medial layer of blood vessels, VSMCs switch from a  
62 contractile phenotype to a migratory/secretory phenotype with an increased extracellular  
63 matrix production. In ApoE KO mice, the progression of atherosclerosis is associated with a  
64 decreased expression of  $\alpha$ -SMA and an increased expression of cell adhesion molecules  
65 (ICAM1 and VCAM1) on VSMCs [15].

66 Although the role of AhR has been studied in the early stages of atherosclerotic development  
67 [7], few studies have focussed on its role in the progression of the pathology. Here, we  
68 studied the effects of chronic TCDD treatment on the progression of atherosclerosis in ApoE  
69 KO mice that are in advanced phases of atherosclerosis. We tested the hypothesis that chronic  
70 stimulation of the AhR aggravates the development of atherosclerosis and assessed the  
71 influence of sex and age as they are important biological variables in atherosclerosis [16–20].

## 72 2. Materials and Methods

### 73 2.1. Animals

74 ApoE KO mice purchased from Charles River were bred in our facility on a 12-hour light-  
75 dark cycle and a standard chow diet (temperature 22°C) in accordance with EU directive  
76 2010/63/EU for animal experiments. 20- and 27-week-old male and female ApoE mice (n =  
77 10 per group) were included in the study. They received an intraperitoneal (ip) injection of  
78 corn-oil or 1 µg/kg of TCDD (LGC Standards) in corn-oil / week for 8 weeks (table 1). **This**  
79 **dose was chosen as the highest dose with no observed effect in a previous study [21].** They  
80 were sacrificed one week after the last injection by intraperitoneal injection of ketamine-HCl  
81 100 mg/kg and xylazine 20 mg/kg.

	Number of mice	
	Corn-oil	TCDD
28-week old females	5	5
35-week old females	5	5
28-week old males	4	6
35-week old males	6	4

82 **Table 1:** Composition of the groups and number of ApoE KO mice studied

83

### 84 2.2. Quantification of atherosclerotic plaques (Oil red O staining)

85 The sizes of atherosclerotic lesions were measured in oil red O-stained frozen transverse  
86 sections of aortas. The root of the aorta was dissected under a microscope and frozen in OCT  
87 embedding medium at -80°C. Five sections of 10 µm (at 200, 400, 600 and 800 µm from the  
88 cusps) were harvested. The sections were counterstained with hematoxylin. The oil red-O  
89 stained slides were observed with AxioObserver Z1 (Zeiss) at a magnification of 5 and  
90 analysed by Leica Qwin software (Leica) to quantify the surface area of the lesion and of the  
91 vessel. The density of the lesion was calculated by dividing the surface area of the lesion by  
92 the surface area of the vessel.

93

### 94 2.3. Immunohistochemistry

95 Ethanol-fixed cryosections of aorta were blocked in 5% BSA and 0,1 % fish gelatin and then  
96 processed for immunostaining with CD68 (rat, Serotec), aSMA (rabbit, Abcam) and VCAM1

97 (goat, R&D) antibodies in 1X PBS – 1% BSA and the anti-rat alexa fluor 647, anti-rabbit  
98 rhodamine and anti-goat FITC secondary antibodies (Jackson ImmunoResearch). The nuclei  
99 were visualized with Hoechst (Molecular Probes).

100

## 101 **2.4. Statistical analysis**

102 The statistical analyses were performed using GraphPad Prism. All data were expressed as  
103 the mean +/- SEM. The control and TCDD-treated groups were compared using either  
104 Kruskal Wallis followed by Mann-Whitney test or directly by Mann-Whitney test. A p-value  
105 of < 0.05 was considered statistically significant.


106

## 107 **3. Results**

### 108 **3.1. Exposure to TCDD aggravates the development of atherogenic lesions specifically in** 109 **ApoE KO female mice**

110 Twenty- and 27-week-old ApoE KO animals were injected with 1 µg of TCDD/kg/week for  
111 8 weeks and sacrificed one week after the last injection. At the end of the study, they were  
112 respectively 28 and 35 weeks old.

113 The atherosclerotic plaque surface was quantified to determine the lesion density, obtained by  
114 dividing the surface area of the lesion by the surface area of the vessel (Figure 1A-D). This  
115 ratio avoids the overestimation of the surface area of lesions and facilitates the comparison  
116 between animals. Both of them were increased by TCDD specifically in 35-week-old ApoE  
117 KO female mice (Figure 1B, E-H), whereas they were not affected in corresponding male  
118 mice. The stage of atherosclerosis development also seems be a differentiating factor: in 28-  
119 week old mice, TCDD had no effect on the lesion density (Figure 1A, C).




120  
 121 **Figure 1.** Lesion density in the aortas of 28- and 35 week-old ApoE KO female (A-B) and  
 122 male (C-D) mice. It is calculated as the ratio, lesion area / aorta area x100, and is expressed  
 123 as mean  $\pm$  SEM. \* :  $p < 0.05$ . Oil red-O staining of aortas from 35 week-old female mice  
 124 exposed or not to TCDD. Two representative aortas are displayed per group (control mice: E-  
 125 F and TCDD-exposed mice: G-H).

126

127 **3.2. Exposure to TCDD leads to a migratory phenotype of VSMCs in the atherosclerotic**  
 128 **plaque of ApoE KO female mice.**

129 We focused on 35 week-old female mice to investigate the effect of TCDD exposure in the  
 130 aorta on the expression of markers: 1) CD68, a marker of foam cells, inflammatory cells and  
 131 of SMCs with a migratory phenotype, 2) VCAM1, expressed on activated endothelial cells  
 132 and on VSMCs from the atherosclerotic plaque, 3)  $\alpha$ -SMA, a marker of VSMCs with a  
 133 contractile phenotype, which normally decreases in the atherosclerotic plaque. The increase in  
 134 the number of CD68+ cells was not significant in the TCDD treated group (Figure 2A-E).  
 135 However, we observed that  $\alpha$ -SMA+ cells were significantly decreased (Figure 2A-D, 2F)  
 136 while a significant increase of VCAM1+ cells was found (2A-D, 2G). The increased  
 137 development of atherosclerotic lesions observed in TCDD-treated mice can be explained by  
 138 the switch of VSMCs from a contractile phenotype to a migratory and secretory phenotype  
 139 which promotes atherosclerotic plaque development along with expression of pro-  
 140 atherosclerotic markers such as VCAM1.



141

142 **Figure 2: Immunostaining and percentage of CD68, α-SMA, VCAM1+ cells in the**  
 143 **aortic root of 35 weeks-old control or TCDD-exposed female mice.** Aortas from 35 week-  
 144 old control or TCDD-exposed female mice were stained for CD68 (pink), α-SMA (red) and  
 145 VCAM1 (green) and the nuclei were visualized with Hoechst. Two representative aorta are  
 146 displayed per group (control: A-B or TCDD-exposed mice: C-D). The number of positive  
 147 cells was normalised by the total cell number and represented as the mean ± SEM for each  
 148 group (\* :  $p < 0.05$ , E-G).

149

#### 150 4. Discussion

151 In this study, we aim to better understand the role of the AhR in the development of  
 152 atherosclerosis. In fact, AhR has been suggested to be involved in the growth of  
 153 atherosclerotic plaques in ApoE-KO mice and the expression of the AhR gene is increased in  
 154 the atherosclerotic mouse aorta and in human carotid artery disease tissue [9]. In the early



155 steps of atherosclerotic development, AhR activation has been shown to aggravate lesions  
156 through an inflammatory response [7]. This is coherent with the many studies which directly  
157 link the AhR pathway to the regulation of pro-inflammatory cytokines [9–12].

158 In this manuscript, we describe for the first time, a gender difference in effects of TCDD on  
159 the progression of atherosclerotic plaques. We exposed 20 and 27 week-old female and male  
160 mice to 8-weeks of ip injections of TCDD, a AhR ligand. Increased lesion densities were  
161 present in the older group of ApoE female mice exposed to TCDD as compared to the same  
162 aged male mice.

163 We then focused on the phenotypic switching of VSMCs. VSMCs play a crucial remodeling  
164 role in the vessel wall potentially recruiting monocyte/macrophages into the plaque [14].  
165 AhR activation leads to a significantly decreased expression of  $\alpha$ -SMA due to a decreased  
166 number of contractile VSMCs and an increased expression of VCAM1. Our data are in  
167 agreement with those of Guo et al. [22] who reported that TCDD promotes proliferation and  
168 migration of human VSMCs with a decreased expression of  $\alpha$ -SMA.

169 The influence of gender and age in the the development of atherosclerosis has been  
170 documented. The atheroprotective effects of estrogen have been reported in ApoE KO mice  
171 on high cholesterol diets [18] but other studies showed larger surface lesions in female ApoE  
172 KO mice as compared to male mice [17,19,20]. **It also can be interesting to explore if the  
173 sexual dimorphism is due to a differential response to AhR activation in aortic cells between  
174 males and females as it was already observed in the liver of C57BL/6 mice [23].**

175 Our results suggest that the development of AhR antagonists may be useful therapeutically,  
176 with age and gender as important parameters to be considered. The plasticity of the AhR  
177 which binds a variety of ligands (aromatic hydrocarbons, tryptophan metabolites, indoles  
178 from the diet...) will be also a key factor [1]. Interestingly, these ligands are reported to have  
179 differential or even opposite effects on Th17 differentiation [24,25]. The competition  
180 between the different types of ligands for the AhR could influence the development of the  
181 atherogenic plaques over time and it might be interesting to consider, as a preventive  
182 pharmacological approach, antagonizing the effect of aromatic hydrocarbons.

183 **Author contributions**

184 Conception of the work: XC, MJK; planning of experiments, collection & analysis of data:  
185 LB, MJK; writing of manuscript: XC, MJK. All authors have read and approved the final  
186 manuscript.

187 **Funding:** This work was supported by INSERM; Université de Paris; Université Sorbonne  
188 Paris Nord.

189 **Acknowledgements:** We thank Dr L.Aggerbeck for critical reading of the manuscript.

190 **Conflicts of Interest:** The authors declare no conflict of interest.

191

192 **References**

- 193 [1] Q. Ma, Influence of light on aryl hydrocarbon receptor signaling and consequences in  
 194 drug metabolism, physiology and disease, *Expert Opin. Drug Metab. Toxicol.* 7 (2011)  
 195 1267–1293. <https://doi.org/10.1517/17425255.2011.614947>.
- 196 [2] A. Vasquez, N. Atallah-Yunes, F.C. Smith, X. You, S.E. Chase, A.E. Silverstone, K.L.  
 197 Vikstrom, A role for the aryl hydrocarbon receptor in cardiac physiology and function  
 198 as demonstrated by AhR knockout mice, *Cardiovasc. Toxicol.* 3 (2003) 153–163.  
 199 <https://doi.org/10.1385/ct:3:2:153>.
- 200 [3] A.K. Lund, M.B. Goens, N.L. Kanagy, M.K. Walker, Cardiac hypertrophy in aryl  
 201 hydrocarbon receptor null mice is correlated with elevated angiotensin II, endothelin-1,  
 202 and mean arterial blood pressure, *Toxicol. Appl. Pharmacol.* 193 (2003) 177–187.  
 203 <https://doi.org/10.1016/j.taap.2003.08.008>.
- 204 [4] V.S. Carreira, Y. Fan, H. Kurita, Q. Wang, C.-I. Ko, M. Naticchioni, M. Jiang, S. Koch,  
 205 X. Zhang, J. Biesiada, M. Medvedovic, Y. Xia, J. Rubinstein, A. Puga, Disruption of Ah  
 206 Receptor Signaling during Mouse Development Leads to Abnormal Cardiac Structure  
 207 and Function in the Adult, *PloS One.* 10 (2015) e0142440.  
 208 <https://doi.org/10.1371/journal.pone.0142440>.
- 209 [5] O. Humblet, L. Birnbaum, E. Rimm, M.A. Mittleman, R. Hauser, Dioxins and  
 210 cardiovascular disease mortality, *Environ. Health Perspect.* 116 (2008) 1443–1448.  
 211 <https://doi.org/10.1289/ehp.11579>.
- 212 [6] P.M. Lind, B. van Bavel, S. Salihovic, L. Lind, Circulating levels of persistent organic  
 213 pollutants (POPs) and carotid atherosclerosis in the elderly, *Environ. Health Perspect.*  
 214 120 (2012) 38–43. <https://doi.org/10.1289/ehp.1103563>.
- 215 [7] D. Wu, N. Nishimura, V. Kuo, O. Fiehn, S. Shahbaz, L. Van Winkle, F. Matsumura,  
 216 C.F.A. Vogel, Activation of aryl hydrocarbon receptor induces vascular inflammation  
 217 and promotes atherosclerosis in apolipoprotein E<sup>-/-</sup> mice, *Arterioscler. Thromb. Vasc.*  
 218 *Biol.* 31 (2011) 1260–1267. <https://doi.org/10.1161/ATVBAHA.110.220202>.
- 219 [8] J.S. Kerley-Hamilton, H.W. Trask, C.J.A. Ridley, E. Dufour, C. Lesseur, C.S.  
 220 Ringelberg, K.L. Moodie, S.L. Shipman, M. Korc, J. Gui, N.W. Shworak, C.R.  
 221 Tomlinson, Inherent and benzo[a]pyrene-induced differential aryl hydrocarbon receptor  
 222 signaling greatly affects life span, atherosclerosis, cardiac gene expression, and body  
 223 and heart growth in mice, *Toxicol. Sci. Off. J. Soc. Toxicol.* 126 (2012) 391–404.  
 224 <https://doi.org/10.1093/toxsci/kfs002>.
- 225 [9] J.B. Kim, M. Pjanic, T. Nguyen, C.L. Miller, D. Iyer, B. Liu, T. Wang, O. Sazonova, I.  
 226 Carcamo-Orive, L.P. Matic, L. Maegdefessel, U. Hedin, T. Quertermous, TCF21 and  
 227 the environmental sensor aryl-hydrocarbon receptor cooperate to activate a pro-  
 228 inflammatory gene expression program in coronary artery smooth muscle cells, *PLoS*  
 229 *Genet.* 13 (2017) e1006750. <https://doi.org/10.1371/journal.pgen.1006750>.
- 230 [10] M.J. Kim, V. Pelloux, E. Guyot, J. Tordjman, L.-C. Bui, A. Chevallier, C. Forest, C.  
 231 Benelli, K. Clément, R. Barouki, Inflammatory pathway genes belong to major targets  
 232 of persistent organic pollutants in adipose cells, *Environ. Health Perspect.* 120 (2012)  
 233 508–514. <https://doi.org/10.1289/ehp.1104282>.
- 234 [11] C.F.A. Vogel, E. Sciallo, P. Wong, P. Kuzmicky, N. Kado, F. Matsumura, Induction of  
 235 proinflammatory cytokines and C-reactive protein in human macrophage cell line U937  
 236 exposed to air pollution particulates, *Environ. Health Perspect.* 113 (2005) 1536–1541.  
 237 <https://doi.org/10.1289/ehp.8094>.
- 238 [12] C. Wang, M.C. Petriello, B. Zhu, B. Hennig, PCB 126 induces monocyte/macrophage  
 239 polarization and inflammation through AhR and NF-κB pathways, *Toxicol. Appl.*  
 240 *Pharmacol.* 367 (2019) 71–81. <https://doi.org/10.1016/j.taap.2019.02.006>.

- 241 [13] S. Allahverdian, A.C. Chehroudi, B.M. McManus, T. Abraham, G.A. Francis,  
242 Contribution of intimal smooth muscle cells to cholesterol accumulation and  
243 macrophage-like cells in human atherosclerosis, *Circulation*. 129 (2014) 1551–1559.  
244 <https://doi.org/10.1161/CIRCULATIONAHA.113.005015>.
- 245 [14] D. Hu, C. Yin, S. Luo, A.J.R. Habenicht, S.K. Mohanta, *Vascular Smooth Muscle Cells*  
246 *Contribute to Atherosclerosis Immunity*, *Front. Immunol.* 10 (2019) 1101.  
247 <https://doi.org/10.3389/fimmu.2019.01101>.
- 248 [15] M. Braun, P. Pietsch, K. Schrör, G. Baumann, S.B. Felix, Cellular adhesion molecules  
249 on vascular smooth muscle cells, *Cardiovasc. Res.* 41 (1999) 395–401.  
250 [https://doi.org/10.1016/s0008-6363\(98\)00302-2](https://doi.org/10.1016/s0008-6363(98)00302-2).
- 251 [16] J.C. Surra, N. Guillén, J.M. Arbonés-Mainar, C. Barranquero, M.A. Navarro, C. Arnal,  
252 I. Orman, J.C. Segovia, J. Osada, Sex as a profound modifier of atherosclerotic lesion  
253 development in apolipoprotein E-deficient mice with different genetic backgrounds, *J.*  
254 *Atheroscler. Thromb.* 17 (2010) 712–721. <https://doi.org/10.5551/jat.3541>.
- 255 [17] G. Caligiuri, A. Nicoletti, X. Zhou, I. Törnberg, G.K. Hansson, Effects of sex and age  
256 on atherosclerosis and autoimmunity in apoE-deficient mice, *Atherosclerosis*. 145  
257 (1999) 301–308. [https://doi.org/10.1016/s0021-9150\(99\)00081-7](https://doi.org/10.1016/s0021-9150(99)00081-7).
- 258 [18] T. Chiba, M. Ikeda, K. Umegaki, T. Tomita, Estrogen-dependent activation of neutral  
259 cholesterol ester hydrolase underlying gender difference of atherogenesis in apoE-/-  
260 mice, *Atherosclerosis*. 219 (2011) 545–551.  
261 <https://doi.org/10.1016/j.atherosclerosis.2011.08.051>.
- 262 [19] D.D. Smith, X. Tan, O. Tawfik, G. Milne, D.J. Stechschulte, K.N. Dileepan, Increased  
263 aortic atherosclerotic plaque development in female apolipoprotein E-null mice is  
264 associated with elevated thromboxane A2 and decreased prostacyclin production, *J.*  
265 *Physiol. Pharmacol. Off. J. Pol. Physiol. Soc.* 61 (2010) 309–316.
- 266 [20] J.J. Man, J.A. Beckman, I.Z. Jaffe, Sex as a Biological Variable in Atherosclerosis, *Circ.*  
267 *Res.* 126 (2020) 1297–1319. <https://doi.org/10.1161/CIRCRESAHA.120.315930>.
- 268 [21] C. Duval, F. Teixeira-Clerc, A.F. Leblanc, S. Touch, C. Emond, M. Guerre-Millo, S.  
269 Lotersztajn, R. Barouki, M. Aggerbeck, X. Coumoul, Chronic Exposure to Low Doses  
270 of Dioxin Promotes Liver Fibrosis Development in the C57BL/6J Diet-Induced Obesity  
271 Mouse Model, *Environ. Health Perspect.* 125 (2017) 428–436.  
272 <https://doi.org/10.1289/EHP316>.
- 273 [22] S. Guo, R. Zhang, Q. Liu, Q. Wan, Y. Wang, Y. Yu, G. Liu, Y. Shen, Y. Yu, J. Zhang,  
274 2,3,7,8-Tetrachlorodibenzo-p-dioxin promotes injury-induced vascular neointima  
275 formation in mice, *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 33 (2019) 10207–  
276 10217. <https://doi.org/10.1096/fj.201900546R>.
- 277 [23] R. Nault, K.A. Fader, J.R. Harkema, T. Zacharewski, Loss of liver-specific and sexually  
278 dimorphic gene expression by aryl hydrocarbon receptor activation in C57BL/6 mice,  
279 *PloS One*. 12 (2017) e0184842. <https://doi.org/10.1371/journal.pone.0184842>.
- 280 [24] N.P. Singh, U.P. Singh, M. Rouse, J. Zhang, S. Chatterjee, P.S. Nagarkatti, M.  
281 Nagarkatti, Dietary Indoles Suppress Delayed-Type Hypersensitivity by Inducing a  
282 Switch from Proinflammatory Th17 Cells to Anti-Inflammatory Regulatory T Cells  
283 through Regulation of MicroRNA, *J. Immunol. Baltim. Md 1950*. 196 (2016) 1108–  
284 1122. <https://doi.org/10.4049/jimmunol.1501727>.
- 285 [25] M. Veldhoen, K. Hirota, A.M. Westendorf, J. Buer, L. Dumoutier, J.-C. Renauld, B.  
286 Stockinger, The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to  
287 environmental toxins, *Nature*. 453 (2008) 106–109.  
288 <https://doi.org/10.1038/nature06881>.
- 289