



HAL
open science

PrPC signalling in neurons: From basics to clinical challenges

Théo Hirsch, Julia Hernandez-Rapp, Séverine Martin-Lannerée, Jean-Marie Launay, Sophie Mouillet-Richard

► **To cite this version:**

Théo Hirsch, Julia Hernandez-Rapp, Séverine Martin-Lannerée, Jean-Marie Launay, Sophie Mouillet-Richard. PrPC signalling in neurons: From basics to clinical challenges. *Biochimie*, 2014, 104, pp.2-11. 10.1016/j.biochi.2014.06.009 . hal-03745692

HAL Id: hal-03745692

<https://hal.science/hal-03745692>

Submitted on 4 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manuscript Number: BIOCHI-D-14-00198R1

Title: PrPC signalling in neurons: from basics to clinical challenges

Article Type: Review Article

Section/Category: Regular issue

Keywords: Prion protein, cell signalling, neurons, A β , prion diseases, Alzheimer's disease

Corresponding Author: Dr. Sophie Mouillet-Richard,

Corresponding Author's Institution:

First Author: Théo Z Hirsch

Order of Authors: Théo Z Hirsch; Julia Hernandez-Rapp; Séverine Martin-Lannerée; Jean-Marie Launay; Sophie Mouillet-Richard

Manuscript Region of Origin: FRANCE

Abstract: The cellular prion protein PrPC was identified over twenty-five years ago as the normal counterpart of the scrapie prion protein PrPSc, itself the main if not the sole component of the infectious agent at the root of Transmissible Spongiform Encephalopathies (TSEs). PrPC is a ubiquitous cell surface protein, abundantly expressed in neurons, which constitute the targets of PrPSc-mediated toxicity. Converging evidence have highlighted that neuronal, GPI-anchored PrPC is absolutely required for prion-induced neuropathogenesis, which warrants investigating into the normal function exerted by PrPC in a neuronal context. It is now well-established that PrPC can serve as a cell signalling molecule, able to mobilize transduction cascades in response to interactions with partners. This function endows PrPC with the capacity to participate in multiple neuronal processes, ranging from survival to synaptic plasticity. A diverse array of data have allowed to shed light on how this function is corrupted by PrPSc. Recently, amyloid A β oligomers, whose accumulation is associated with Alzheimer's disease (AD), were shown to similarly instigate toxic events by deviating PrPC-mediated signalling. Here, we provide an overview of the various signal transduction cascades ascribed to PrPC in neurons, summarize how their subversion by PrPSc or A β oligomers contributes to TSE or AD neuropathogenesis and discuss the ensuing clinical implications.

Opposed Reviewers:

Dear Dr Forest,

Thank you for your letter of June 3rd 2014 regarding our review entitled « **PrP^C signalling in neurons: from basics to clinical challenges** ». I am pleased to provide a revised version of our review, which takes into account the reviewers' comments. I hope this review is now suitable for publication in Biochimie.

Yours sincerely,

Dr S. Mouillet-Richard
INSERM UMR-S1124, Université Paris Descartes,
E-mail: sophie.mouillet-richard@parisdescartes.fr

Detailed answer to the reviewers:

We thank the reviewers for their positive comments.

As pointed by reviewer 3, we have corrected some errors within the text.

We have taken into account the minor modifications suggested by reviewer 4:

1. *In fig. 2, the interaction with α7nAChR could be addressed in the scheme, since it is an important regulator of STII-PrP signal transduction complex Beraldo et al., J Biol Chem. 2010 Nov 19;285(47):36542-50.*

The interaction of PrP^C with the alpha7nAChR is now presented in Fig.2

2. *Provide reference for the statement "This signal, which can be revealed in response to antibody-mediated ligation of PrPC, also occurs upon binding of PrPC to its endogenous ligands STII (Line 13 - 18, page 8)".*

The corresponding reference is now inserted at the end of the sentence.

3. *The subtitle of section 5 does not seem to be appropriate since the section mainly addresses the transmembrane proteins responsible for transducing PrPC signaling into intracellular compartments. Temporal alterations of PrPC signaling cascades were not discussed.*

The subtitle of section 5 has been changed into "PrP^C signalling partners at the membrane".

Author agreement form for

PrP^c signalling in neurons: from basics to clinical challenges

by Théo Z. Hirsch, Julia Hernandez-Rapp, Séverine Martin-Lannerée, Jean-Marie Launay and Sophie Mouillet-Richard

Material submitted is original, all authors are in agreement to have the article published

1
2
3
4 **PrP^C signalling in neurons: from basics to clinical challenges**
5
6

7 **Théo Z. Hirsch^{1,2}, Julia Hernandez-Rapp^{1,2,3}, Séverine Martin-Lannerée^{1,2}, Jean-Marie**
8 **Launay^{4,5} and Sophie Mouillet-Richard^{1,2,*}**
9

10
11 **Abstract**
12

13
14 The cellular prion protein PrP^C was identified over twenty-five years ago as the normal
15 counterpart of the scrapie prion protein PrP^{Sc}, itself the main if not the sole component of
16 the infectious agent at the root of Transmissible Spongiform Encephalopathies (TSEs). PrP^C
17 is a ubiquitous cell surface protein, abundantly expressed in neurons, which constitute the
18 targets of PrP^{Sc}-mediated toxicity. Converging evidence have highlighted that neuronal, GPI-
19 anchored PrP^C is absolutely required for prion-induced neuropathogenesis, which warrants
20 investigating into the normal function exerted by PrP^C in a neuronal context. It is now well-
21 established that PrP^C can serve as a cell signalling molecule, able to mobilize transduction
22 cascades in response to interactions with partners. This function endows PrP^C with the
23 capacity to participate in multiple neuronal processes, ranging from survival to synaptic
24 plasticity. A diverse array of data have allowed to shed light on how this function is
25 corrupted by PrP^{Sc}. Recently, amyloid A β oligomers, whose accumulation is associated with
26 Alzheimer's disease (AD), were shown to similarly instigate toxic events by deviating PrP^C-
27 mediated signalling. Here, we provide an overview of the various signal transduction
28 cascades ascribed to PrP^C in neurons, summarize how their subversion by PrP^{Sc} or A β
29 oligomers contributes to TSE or AD neuropathogenesis and discuss the ensuing clinical
30 implications.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Highlights Hirsch et al.

- Prion diseases are caused by the conversion of PrP^C into its scrapie PrP^{Sc} isoform.
- PrP^C can act as a cell surface receptor able to instruct signalling events.
- PrP^{Sc} exerts its toxicity by subverting PrP^C-dependent signalling cascades in neurons.
- PrP^C may transduce toxic signals after binding to A β oligomers.
- Targeting PrP^C-associated pathways may alleviate prion and Alzheimer's diseases.

PrP^C signalling in neurons: from basics to clinical challenges

Théo Z. Hirsch^{1,2}, Julia Hernandez-Rapp^{1,2,3}, Séverine Martin-Lannerée^{1,2}, Jean-Marie Launay^{4,5} and Sophie Mouillet-Richard^{1,2,*}

¹ INSERM UMR-S1124, 75006 Paris France.

² Université Paris Descartes, Sorbonne Paris Cité, UMR-S1124, 75006 Paris France.

³ Université Paris Sud 11, ED419 Biosigne, 91400 Orsay, France

⁴ AP-HP Service de Biochimie, Fondation FondaMental, INSERM U942 Hôpital Lariboisière, 75010 Paris, France

⁵ Pharma Research Department, F. Hoffmann-La-Roche Ltd., CH-4070 Basel, Switzerland.

* Correspondence

Dr. Sophie Mouillet-Richard

INSERM U-1124,

Université Paris Descartes

45, rue des Saints Pères

75006 Paris, France

Tel.: +33 1 42 86 42 40

email: sophie.mouillet-richard@parisdescartes.fr

Abstract

1
2 The cellular prion protein PrP^C was identified over twenty-five years ago as the normal
3
4 counterpart of the scrapie prion protein PrP^{Sc}, itself the main if not the sole component
5
6 of the infectious agent at the root of Transmissible Spongiform Encephalopathies (TSEs).
7
8 PrP^C is a ubiquitous cell surface protein, abundantly expressed in neurons, which
9
10 constitute the targets of PrP^{Sc}-mediated toxicity. Converging evidence have highlighted
11
12 that neuronal, GPI-anchored PrP^C is absolutely required for prion-induced
13
14 neuropathogenesis, which warrants investigating into the normal function exerted by
15
16 PrP^C in a neuronal context. It is now well-established that PrP^C can serve as a cell
17
18 signalling molecule, able to mobilize transduction cascades in response to interactions
19
20 with partners. This function endows PrP^C with the capacity to participate in multiple
21
22 neuronal processes, ranging from survival to synaptic plasticity. A diverse array of data
23
24 have allowed to shed light on how this function is corrupted by PrP^{Sc}. Recently, amyloid
25
26 A β oligomers, whose accumulation is associated with Alzheimer's disease (AD), were
27
28 shown to similarly instigate toxic events by deviating PrP^C-mediated signalling. Here, we
29
30 provide an overview of the various signal transduction cascades ascribed to PrP^C in
31
32 neurons, summarize how their subversion by PrP^{Sc} or A β oligomers contributes to TSE
33
34 or AD neuropathogenesis and discuss the ensuing clinical implications.
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 **Keywords:** Prion protein, cell signalling, neurons, A β , prion diseases, Alzheimer's
50
51 disease
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abbreviations: AD, Alzheimer's disease; APP, amyloid precursor protein; cPLA2, cytosolic phospholipase A2; GPI, glycosylphosphatidylinositol; GPCR, G-protein coupled receptor; GSK3 β , glycogen synthase kinase 3 β ; Ln γ 1, laminin γ 1 chain; LRP-1, low-density lipoprotein-related protein 1; LTD, long-term depression; LTP, long-term potentiation; mTOR, mammalian target of rapamycin; NCAM, neural cell adhesion molecule; PDK1, phosphoinositide-dependent kinase 1; PI3K, phosphoinositide 3 kinase; PKA, protein kinase A; PLC, phospholipase C; PrP^C, cellular prion protein; PrP^{Sc}, scrapie prion protein; PSD, postsynaptic density; ROS, reactive oxygen species; STI-1, stress-inducible protein 1; TACE, TNF α -converting enzyme; TSE, Transmissible Spongiform Encephalopathies; UPR: unfolded protein response;

1. Introduction

1
2
3 The cellular prion protein PrP^C has received considerable attention due to its central
4
5 role in the development of Transmissible Spongiform Encephalopathies (TSEs). These
6
7 fatal neurodegenerative disorders arise from the autocatalytic conversion of PrP^C into
8
9 its aggregate-prone conformational variant PrP^{Sc}, standing for scrapie isoform of the
10
11 prion protein [1]. PrP^{Sc} actually constitutes the only identified component of the
12
13 infectious TSE agent, which propagates by acting as a seed to promote the template-
14
15 directed misfolding of endogenous PrP^C [1]. Accordingly, PrP-null mice fail to develop
16
17 TSE upon prion inoculation [2]. PrP^C is a normal host-encoded protein of about 230
18
19 residues, present in all cell types and most abundantly found in neurons [3]. PrP^C
20
21 possesses a glycosylphosphatidylinositol (GPI) anchor that sustains its attachment to
22
23 the plasma membrane and can be hydrolyzed by endogenous phospholipases to yield
24
25 secreted PrP^C (see Fig. 1) [4, 5]. Its main features include the presence of octapeptide
26
27 repeats in the unstructured N-terminal region, involved in the binding of divalent
28
29 cations such as copper [6] or zinc [7], as well as a disulfide bond encompassing two
30
31 asparagine residues that can be potentially N-glycosylated [8]. PrP^C can undergo several
32
33 proteolytic cleavages: (i) in the octarepeat region (beta-cleavage), (ii) at position
34
35 111/112 (alpha-cleavage) or (iii) in the extreme C-terminus to generate an anchorless
36
37 isoform (reviewed in [5, 9], [10]). The so-called alpha cleavage, giving rise to the N1 N-
38
39 terminal and the C1 C-terminal fragments, has been most extensively studied as it
40
41 occurs physiologically, notably in the brain [11] and because it precludes the conversion
42
43 into pathological PrP^{Sc} [12]. Thus, while often referred to as a unique entity, PrP^C
44
45 actually encompasses a great diversity of isoforms, whose distribution may vary
46
47 according to the cell type and whose respective functions await further investigation [3].
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Like many GPI-anchored protein, PrP^C segregates in sphingolipid-rich membrane
2 microdomains known as "lipid-rafts" (reviewed in [13]). These specialized subcellular
3 compartments are enriched in receptors, second messengers and effectors and are
4 considered as "signal transduction centres" [14]. Over the past fifteen years, evidence
5 has accumulated that PrP^C can act as a cell surface receptor, co-receptor or ligand, able
6 to recruit downstream signal transduction pathways. Such function is thought to be
7 driven by the interaction of PrP^C with promiscuous partners, within multimolecular
8 complexes [3]. Among the vast array of PrP^C interactors and effectors in different cell
9 types and tissues, special interest is devoted to those that may play a role in prion-
10 induced neurotoxicity. Indeed, it is now well-acknowledged that pathogenic prions exert
11 their deleterious action by subverting the normal function of PrP^C in the brain [15].
12 Their toxicity is fully dependent on the expression of PrP^C in neurons [16, 17] and
13 requires that it is GPI-anchored [18]. Deciphering how PrP^{Sc} deviates PrP^C-dependent
14 signalling cascades is attracting all the more attention since this pathogenic mechanism
15 appears to be also at play in Alzheimer's disease (AD), with amyloid A β oligomers
16 hijacking PrP^C-dependent pathways to instigate toxic events (reviewed in [19]).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 In the present review, we summarize the current knowledge of PrP^C-dependent
53 signalling in neurons, its pathophysiological implications and highlight some therapeutic
54 prospects to alleviate TSE and AD-related pathogenesis.
55
56
57
58
59
60
61
62
63
64
65

2. How is PrP^C signalling initiated?

A current view is that the mobilization of PrP^C-mediated cell signalling events occurs in response to the association of this protein with various partners at the plasma membrane [3], a typical scheme for GPI-anchored proteins [14]. Indeed, in neurons as in

1 other cell types, the majority of PrP^C molecules are GPI-anchored, which drives their
2 subcellular localization in lipid rafts [13]. Two transmembrane variants of PrP^C have
3 further been described that are associated with neurodegeneration and whose
4 physiological relevance remains elusive [20]. Beyond rafts, PrP^C is also found in
5 postsynaptic densities (PSDs) [21], which may associate with postsynaptic membrane
6 rafts [22]. The special composition of these compartments allows the spatial segregation
7 of receptors and signal transduction effectors and favours their effective interaction
8 [23]. Classically, the mobilization of signalling cascades by GPI-anchored proteins is
9 induced by their clustering in response to ligand binding [23]. Such clustering can be
10 triggered by antibody-mediated ligation, an experimental paradigm that allowed our
11 laboratory to unveil various PrP^C-dependent cascades in neuronal cells [24-27]. Data
12 from the Tazelt laboratory further established that the recruitment of neuroprotective
13 signals by PrP^C necessitates its dimerization via its hydrophobic domain (residues 113-
14 133) [24]. Interestingly, PrP^C dimerization appears to enhance its alpha-cleavage and
15 production of the N1 fragment [25] as well as its shedding [26]. PrP^C cleavage may
16 additionally be influenced by the binding of copper [10], which also impacts on PrP^C
17 trafficking and signalling activity [6, 27]. Shed PrP^C may act in cis or in trans to
18 homodimerize [28], interact with the neural cell adhesion molecule NCAM [29] or
19 putatively other yet-to-be-identified cell surface molecules [30, 31]. Besides,
20 physiologically relevant interactions of PrP^C with several extracellular molecules, acting
21 as ligands, have been documented and are summarized in (Fig. 1). These include the
22 matrix components vitronectin and laminin or the extracellular chaperone stress-
23 inducible protein 1 (STI-1), whose binding to PrP^C causes reciprocal conformational
24 changes [32]. The repertoire of PrP^C interactors also includes the amyloid precursor
25 protein APP [33, 34] and its aggregation-prone cleavage product A β [19, 35] (see Fig. 1),
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 whose unbalanced production lies at the root of AD, according to the amyloid cascade
2 hypothesis [36]. While it is unclear whether the PrP^C-APP interaction instructs
3 downstream cell signalling events or whether the binding of PrP^C to A β may occur under
4 physiological conditions, it is now well-established that A β oligomers bind to PrP^C in AD
5 brain [37] and subvert PrP^C-dependent signalling cascades to affect various neuronal
6 processes (reviewed in [19]) (see below).
7
8
9
10
11
12
13

14 The multiplicity of PrP^C interactors on the one hand and isoforms on the other hand
15 raises the issue as to which PrP^C entities interact with a given partner and whether
16 several ligands may compete for binding to a given PrP^C entity, as recently suggested for
17 STI-1 and A β oligomers [38].
18
19
20
21
22
23
24
25
26
27

28 **3. PrP^C signalling in neuronal cells: a plethora of effectors**

29 The ability of PrP^C to trigger cell signalling events, initially suspected from its location in
30 lipid rafts (reviewed in [13]) and the reported ability of GPI-anchored proteins to
31 activate src kinases [39], has been examined using multiple experimental paradigms. A
32 well-characterized proximal effector of PrP^C is Fyn, a tyrosine kinase of the src family
33 that we first identified as a neuronal-restricted target of PrP^C using the inducible
34 neuronal 1C11 cell line [40], and that associates with PrP^C to form a signalling complex
35 [41]. In 1C11-derived differentiated cells, the PrP^C-Fyn coupling is relayed by the
36 membrane protein caveolin [40], now recognized as an important regulator of neuronal
37 intracellular signalling [42]. The PrP^C-dependent activation of Fyn was subsequently
38 shown to be involved in NCAM-dependent neurite outgrowth in primary hippocampal
39 neurons [29]. Apart from caveolin and NCAM, the natures of the PrP^C partners that may
40 contribute to Fyn activation in a physiological context remain to be identified. From a
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

pathological point of view, the PrP^C-Fyn platform has received renewed interest in the last few years as it is hijacked both by PrP^{Sc} [43] and by A β oligomers [19, 44] to transduce their toxicity (see below).

In a recent study, we documented that neuronal PrP^C has the capacity to recruit another member of the src kinase family, the Lyn kinase [27], previously found to be present in the plasma microenvironment of PrP^C in primary cerebellar neurons [45]. This signal, which can be revealed in response to antibody-mediated ligation of PrP^C, also occurs upon binding of PrP^C to its endogenous ligand STI-1 [27]. Noteworthy, like Fyn, the activation of Lyn driven by PrP^C is transduced via caveolin [27], further supporting a central role for caveolin in PrP^C-dependent signalling.

Other reported intracellular effectors of PrP^C signalling include the phosphoinositide 3 kinase (PI3K) [27, 46], Akt [27, 47], protein kinase A (PKA) [48-50], the reactive oxygen species (ROS) generating enzyme NADPH oxidase [51], the MAP kinases ERK1/2 [49-52], the glycogen synthase kinase GSK3 β [27], the TNF α -converting enzyme (TACE) metalloprotease [53], the mammalian target of rapamycin mTOR [47], as well as several transcription factors: CREB, Egr-1 and c-fos [54]. Links with sphingolipid [55] and calcium signalling [56-59] have also been depicted. The diverse array of PrP^C signalling effectors is summarized in Table 1.

Some of these effectors have been shown to be under the control of Fyn (e.g. NADPH oxidase) or Lyn (e.g. GSK3 β). Some other pathways, such as those featuring PKA, remain to be delineated. More generally, an important challenge for the next few years will be to translate the repertoire of PrP^C targets into transduction cascades in relation with the initial triggering signal (e.g. STI-1 or laminin), cell type and biological response, as outlined in Fig. 2. Altogether, achieving this goal appears as a prerequisite to ultimately

1 reach a global view of the various PrP^C-dependent signalling platforms and pathways,
2 their potential crosstalks, as well as their integrated cellular outcome.
3
4
5
6

7 **4. An intimate connection with neurotransmitter receptors**

8
9
10 Another important class of lipid raft residents in neurons is neurotransmitter receptors
11 [60]. A number of studies have unveiled functional interactions between PrP^C and
12 diverse ionotropic or G-protein coupled neurotransmitter receptors, which are
13 summarized in Table 2. Focusing on serotonergic receptors, our laboratory reported on
14 the PrP^C-dependent modulation of the signalling activity of 5-HT_{1B}, 5-HT_{2A} and 5-HT_{2B}
15 receptors, all coupled to G-proteins, in serotonergic 1C11^{5-HT} neuronal cells [61].
16 Interestingly, we observed both positive and negative actions of PrP^C, depending on the
17 G-protein and pathway considered, and that the modulatory effect of PrP^C depends on
18 caveolin [61]. Very recently, we delineated an additional link between PrP^C and the 5-
19 HT_{1B} receptor, by showing that the PrP^C-dependent inactivation of GSK3 β interferes
20 with the subcellular localization and functional activity of this receptor and thereby
21 potentiates neurotransmitter release [27]. Metabotropic glutamate receptors constitute
22 another class of G-protein coupled neurotransmitter receptors (GPCR) that has captured
23 much attention in relation to PrP^C function. PrP^C has the capacity to interact with
24 mGlu1AR, mGlu1BR as well as mGlu5R [56], which all belong to the Group I of
25 metabotropic glutamate receptors and are coupled to Gq-mediated phospholipase C
26 (PLC) activation [62]. Further, the recruitment of the PrP^C-mGluR signalling complex by
27 the laminin γ 1 chain (Ln γ 1) was shown to trigger calcium release, ERK1/2 activation
28 and to sustain neurite outgrowth [56]. In a pathological context, mGlu5R receptors have
29 been reported to relay the activation of Fyn [63] and to mediate the facilitation of long
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 term depression (LTD) [64] in response to the binding of A β oligomers to PrP^C (see
2 below).

3
4
5 The capacity of PrP^C to interact with neurotransmitter receptors is not restricted to
6
7 GPCRs. Indeed, it may form a complex with the ionotropic nicotinic acetylcholine
8
9 receptor α 7nAChR, whose mobilization in the hippocampus by the STI-1 ligand
10
11 promotes intracellular calcium release [57]. Finally, multiple links with NMDA receptors
12
13 have been depicted, in relation with the critical involvement of these receptors in long-
14
15 term potentiation and synaptic plasticity [65]. For instance, PrP^C was reported to exert a
16
17 negative regulation on GluN2D-containing NMDA receptors, thereby providing a
18
19 neuroprotective mechanism against glutamate excitotoxicity [66]. It also appears to
20
21 sustain the internalization of GluN2B-containing NMDA receptors via Fyn-mediated
22
23 phosphorylation in response to ethanol exposure [67]. Of note, several lines of evidence
24
25 indicate that the PrP^C-dependent modulation of NMDA receptors is lost in the presence
26
27 of toxic oligomers, including A β , leading to reduced receptor desensitization [44, 68, 69].
28
29
30 Altogether, these observations substantiate an important role for PrP^C in the regulation
31
32 of diverse neurotransmitter receptors, which may account for the subtle behavioural
33
34 and / or cognitive deficits reported in PrP-null mice [70]. Refining our current
35
36 knowledge of the interplay between PrP^C and neurotransmitter receptors has obvious
37
38 implications for unravelling the global impact of prions or A β oligomers on neuronal
39
40 activity and homeostasis, as will be discussed below.
41
42
43
44
45
46
47
48
49
50
51
52

53 **5. PrP^C signalling partners at the membrane**

54
55
56 Lipid rafts typically allow a tight spatial and temporal control of cell signalling events
57
58 [71]. As for PrP^C-dependent signalling cascades, multiple regulatory mechanisms have
59
60
61
62
63
64
65

1 been highlighted. First, the interaction of PrP^C with caveolin [40, 41], likely sustains the
2 compartmentation of PrP^C with appropriate signalling partners, e.g. src kinases or
3 neurotransmitter receptors [42]. Adding further complexity to the picture, lipid rafts
4 may vary in their composition [72]. Such raft heterogeneity may account for the spatial
5 restriction of the PrP^C-caveolin-Lyn complex to neuronal cell bodies, to the opposite of
6 the PrP^C-caveolin-Fyn complex, which is found in neuronal processes [27]. Whether
7 each platform involves distinct subsets of PrP^C molecules remains to be investigated.
8 The nature of potential transmembrane partners contributing to the spatial control of
9 PrP^C-signalling complexes also deserves attention. Both NCAM [29] and mGlu5R [63]
10 have been reported to transduce PrP^C-dependent Fyn activation. Recently, we
11 highlighted the contribution of the transmembrane low-density receptor protein 1 (LRP-
12 1) to the recruitment of effectors downstream from the PrP^C-caveolin-Lyn platform [27].
13 LRP-1 was previously identified as a PrP^C partner that regulates its internalization [73],
14 and whose interaction with PrP^C is enhanced by copper [74]. Among its multiple ligands,
15 LRP-1 is known to interact with A β , and recent data indicate that LRP-1 is mandatory for
16 the PrP^C-dependent toxicity of A β [75]. Beyond favouring the interaction with LRP-1,
17 the binding of copper to PrP^C was also shown to enhance its endocytosis together with
18 Glypican-1 [76], a GPI-anchored heparan sulfate proteoglycan that mediates the
19 association of PrP^C with lipid rafts and, in a pathological context, favours the formation
20 of PrP^{Sc} [77]. While it may act as a cofactor for prion conversion, whether Glypican-1
21 modulates interactions of PrP^C with partners and downstream cell signalling events
22 remains to be investigated. The same question applies to the 37kDa/67kDa laminin
23 receptor, reported to promote the internalization of PrP^C [78] and to sustain PrP^{Sc}
24 propagation in prion-infected cells [79]. Whatever the partners involved, PrP^C
25 endocytosis appears to be mandatory for the recruitment of some effectors, such as the
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 activation of ERK1/2 in response to STI-1 [52]. Finally, as will be emphasized below,
2 alterations in the dynamics of PrP^C interactions with its partners, such as its increased
3 cell-surface localization upon binding to A β oligomers [80], may disrupt the strength
4 and duration of PrP^C-related signalling and instigate toxic cascades.
5
6
7
8
9

10 11 12 **6. Translating PrP^C-mediated signalling pathways in neurons into a biological** 13 **response** 14 15 16

17 Overall, through the various signalling cascades brought to light in a neuronal context,
18 PrP^C appears to be connected with nearly all aspects of neuronal physiology. Its
19 contribution to neurite outgrowth may be sustained by its interaction with NCAM and
20 downstream Fyn activation [29], vitronectin [81], Ln γ 1 [82, 83] or STI-1 [50] and may
21 also involve an interplay with integrins [41, 84]. Of note, a synergistic action of Ln γ 1
22 and STI-1 in triggering PrP^C-dependent calcium signalling and neuritogenesis has been
23 described [59]. Beraldo et al. also highlighted the requirement of mGlu1R to transduce
24 the PrP^C-Ln γ 1-mediated induction of process outgrowth [56]. In vivo, the PrP^C-Ln γ 1
25 interaction was shown to modulate memory consolidation, with a suggested implication
26 of PKA and ERK1/2 [85]. This property is shared by the PrP^C-STI-1 duo [86], which also
27 sustains protein synthesis through the mobilization of mTOR [47]. Interestingly, mTOR
28 also emerges as an important player in synaptic plasticity [87], like other PrP^C targets
29 such as ERK1/2, CREB, or egr-1 [88]. Another possible signalling intermediate in these
30 processes is the GSK3 β kinase, which we recently identified as a downstream target of
31 the PrP^C-STI-1 coupling [27] and has been shown both to play a pivotal role in synaptic
32 plasticity and long-term potentiation (LTP) [89] and to regulate mTOR activity [90].
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

may suspect that inhibition of this kinase instigated by the interaction of PrP^C with STI-1 also contributes to the well-established neuroprotective action of these two molecules [49, 92-94].

Multiple links between PrP^C and neurotransmitter-associated functions have further been uncovered. For instance, through a ROS-TACE-TNF α pathway, PrP^C promotes bioamine degradation [53], an observation which has to be brought together with the contribution of bioamine turnover to memory consolidation [95, 96]. Our recent work further established that the negative regulation of the serotonergic 5-HT_{1B} receptor activity downstream from the PrP^C-Lyn-GSK3 β cascade potentiates neurotransmitter release [27]. Adding to this view is the interplay between PrP^C and diverse neurotransmitter receptors, as summarized above. Finally, we may note that the multifaceted contribution of PrP^C to neuronal homeostasis and activity is in keeping with the wide array of alterations depicted in the context of prion infection [97-99] or after binding of A β oligomers to PrP^C (reviewed in [19]).

7. Causes and consequences of the subversion of PrP^C signalling in neurons

That the deviation of PrP^C-dependent-signalling may contribute to neurodegeneration is now well established in the case of prion infection and also in Alzheimer's disease. Prions fail to induce any pathogenesis when endogenous PrP^C is selectively knocked-out in neurons [17] or is exclusively anchorless [18]. These pioneering studies have set the stage for investigating how PrP^{Sc} may usurp PrP^C normal signalling function to exert its toxic action. For instance, pathological PrP conformers were shown to induce a switch from anti-apoptotic to pro-apoptotic signalling of PrP^C [24]. PrP^{Sc} aggregation was further reported to trigger the clustering of its GPI anchors at high densities and thereby

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

cause alterations in membrane composition, pathological activation of cytosolic phospholipase A2 (cPLA2) and synaptic damage [100]. In a recent work, we documented that pathogenic prions hijack the PrP^C-Fyn platform to constitutively recruit downstream effectors, including NADPH oxidase, and cause impairment in the clearance of amyloid A β [43] (Fig. 3). The former observation recalls the central role established for the NADPH oxidase subunit NOX2 in prion-induced neuronal damage [101, 102]. Altogether, these data support the notion that the conversion of PrP^C into PrP^{Sc} may promote the constitutive recruitment of proximal downstream effectors in lipid rafts. Overactivation of PrP^C downstream targets may similarly account for the development of a spontaneous neurodegenerative illness in transgenic mice overexpressing PrP^C by 5 to 10 fold [103]. Besides, the accumulation of PrP^C in lipid raft during aging, proposed to explain the occurrence of sporadic prion diseases [104], might also amplify the activation of downstream cascades. Importantly, the pathological deviation of PrP^C-mediated signalling may be further self-sustained by various subsequent alterations such as aberrant distribution of caveolin-1 [105], impaired cholesterol metabolism [106], decreased protective alpha-cleavage [107] or reduced clearance of A β [43, 108].

The pathological implications of the subversion of PrP^C function are not restricted to prion infection since PrP^C can convey neurotoxic signals after binding to various types of beta-sheet rich conformers, including A β [68]. Regarding the AD connection, converging data gathered in the last 5 years demonstrate that PrP^C can act as a cell-surface receptor for oligomeric A β (reviewed in [19]). Multiple deleterious outcomes of the interaction of A β with PrP^C have been depicted, including synaptic dysfunction and dendritic spine retraction [44], neuronal death [109], inhibition of LTP [35], exacerbation of LTD [64], axonal degeneration [110], as well as induction of autophagy [111]. From a mechanistic

1 point of view, A β oligomers bind to PrP^C at different sites within the N-terminal region
2 (see Fig. 1) and recruit downstream effectors (Fig. 3). Among these, Fyn has captured
3 the most attention since this kinase has long been known to be mandatory for A β -
4 mediated toxicity [112, 113]. In recent years, several outstanding observations have
5 been reported, including the central role of Fyn in promoting phosphorylation of the
6 GluN2B subunit of NMDA receptors and dendritic spine loss [44], as well as tau
7 hyperphosphorylation [114] after engagement of A β oligomers with PrP^C. mGlu5Rs
8 were further shown to act as co-receptors for PrP^C-bound A β oligomers to promote Fyn
9 activation [63]. Whether this property is shared by other metabotropic or ionotropic
10 receptors deserves to be investigated, in view of the broad implication of
11 neurotransmitter receptors in AD pathology [115]. Adding to the above data, PrP^C was
12 also shown to relay the A β -dependent activation of cPLA2 [116], caspase 3 [109],
13 ERK1/2 [80], NADPH oxidase [117] and phosphoinositide-dependent kinase-1 (PDK1),
14 itself leading to decreased TACE activity [107]. Further studies are needed to explore the
15 relationship between these diverse events and Fyn activation. Finally, from a
16 therapeutic perspective, it is worth noting that some studies have addressed the
17 relevance of observations gained in vitro or in mice and have for instance established
18 that A β oligomers and PrP^C interact in the brain of AD patients but not healthy controls
19 [37]. The overactivation of PDK1 found in AD mouse models to be dependent on PrP^C
20 expression was corroborated in AD patients as well [107]. Overall, while PrP^C certainly
21 is not the sole relay of A β -mediated toxicity [118], these cumulative data provide several
22 promising avenues for therapeutic intervention in prion and Alzheimer's diseases,
23 which are discussed below.

8. Therapeutic prospects

1
2
3 At present, TSEs remain fatal diseases with no effective treatment despite numerous
4
5 trials (see [119] for example). This also holds true for AD, which represents a major
6
7 health burden with an estimated 25 million people affected worldwide. The
8
9 development of effective therapies for prion and Alzheimer's diseases faces multiple
10
11 challenges: patients are diagnosed at a stage when brain damage has already occurred,
12
13 drugs need to cross the blood-brain-barrier, and last but not least, the normal functions
14
15 of the targeted proteins need to be preserved. Beyond strategies aiming at limiting the
16
17 production and / or accumulation of PrP^{Sc}, which will not be addressed here, one major
18
19 axis followed for the treatment of TSEs is that of immunotherapy (reviewed in [120]).
20
21 Some passive immunotherapy studies have notably demonstrated a substantial benefit
22
23 for a panel of anti-PrP^C antibodies in delaying onset of disease, although whether this
24
25 strategy can still prove useful when treatment is initiated after the appearance of clinical
26
27 symptoms has yet to be demonstrated (reviewed in [120]). More recently, a subset of
28
29 the same antibodies has been tested for their ability to counteract the interaction of A β
30
31 oligomers with PrP^C. In vitro, the binding of A β oligomers to PrP^C was successfully
32
33 antagonized using 6D11 [35, 44, 109, 114], ICSM35 [121] or Sha31 [122] anti-PrP^C
34
35 antibodies. The 6D11 antibody was further shown to protect against cognitive deficits in
36
37 a mouse model of AD [123]. Finally, brain-derived A β -mediated impairment of LTP
38
39 could be efficiently prevented following immunotargeting of PrP^C using ICSM35 or
40
41 ICSM18 antibodies or the D13 antibody fragment (Fab) [124].
42
43
44
45
46
47
48
49
50
51

52
53 To the opposite of these studies, some antibodies targeting PrP^C exhibit neurotoxic
54
55 potential. Beyond some conflicting results [125, 126], an extensive study from the
56
57 Aguzzi laboratory has allowed to map the epitopes of a panel of toxicity-inducing anti-
58
59 PrP^C antibodies [102]. These neurotoxic antibodies were shown to systematically target
60
61
62
63
64
65

1 the globular domain (residues 124-230) - although some antibodies recognizing this
2 region remained innocuous - and to promote neurodegeneration via calpain and NOX2
3 activation [102], in a manner similar to infectious prions [101]. Since NADPH oxidase is
4 a downstream effector of PrP^C signalling [51] and is overactivated in prion infection
5 [43], NOX2, as suggested above [101, 102], and putatively other NADPH oxidase
6 subunits, may prove valuable targets to combat TSEs. The indispensable role of NOX2 in
7 host defence would, however, need to be taken into account, should its inhibition be
8 further considered for therapeutic purposes [127].
9

10 Further in relation with the above-mentioned subversion of PrP^C-mediated pathways by
11 PrP^{Sc} or A β , the potential benefit of targeting Fyn with the small molecule inhibitor
12 saracatinib is now being evaluated within a clinical trial focusing on AD [128]. In view of
13 the central role exerted by Fyn in the recruitment of pathogenic cascades by prions [43],
14 whether the same strategy may apply to TSEs seems worth investigating. Likewise,
15 PDK1 overactivation was recently identified as another common feature of TSEs and AD,
16 whose inhibition allows alleviating both diseases [107]. Of note, although further work
17 is needed to develop inhibitory molecules devoid of intrinsic toxicity, treatment was
18 shown to be effective while starting at a relatively late stage [107]. Besides, the
19 observation that PDK1 overactivation in AD can be antagonized through silencing of
20 PrP^C [107] points to an interesting link between PrP^C signalling and PDK1, which
21 deserves further investigation.
22

23 The group of Mallucci adopted a different approach based on targeting a generic
24 pathway sustaining neurodegeneration. A commonality between TSEs, AD and
25 Parkinson's disease is the impairment in the unfolded protein response (UPR) [129].
26 One effector of this pathway is the eIF2 α elongation factor, whose phosphorylation halts
27 protein synthesis. Following the demonstration that translational repression by
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 phosphorylated eIF2 α mediates prion neurodegeneration [130], Moreno et al.
2 documented that inhibition of the UPR using an orally-administered pharmacological
3 compound could abrogate the development of clinical disease in prion-infected mice,
4 even when starting treatment after the emergence of behavioural signs [131]. Whether
5 the advantage of this treatment is comparable to that afforded by others such as PDK1
6 inhibition is unclear at present. In this respect, the development of standardized
7 methods using defined mouse models and prion strains would help launch comparative
8 studies to assess the relative benefit of various therapeutic approaches since the onset
9 and progression of prion pathogenesis is highly synchronous in inoculated mice,
10 contrary to AD mouse models. Notwithstanding, it is worthy to note that the
11 suppression of eIF2 α phosphorylation was similarly shown to be beneficial in a mouse
12 model of AD [132], suggesting that targeting the UPR may represent a powerful strategy
13 to combat various neurodegenerative disorders [129]. Because transient
14 phosphorylation of eIF2 α is involved in proteostasis, one important challenge for the
15 next years would be to develop UPR-targeting therapeutic approaches that preserve the
16 beneficial function of eIF2 α , as recently outlined [129]. Some progress towards this
17 goal may be achieved through further delineating the link between PrP^C signalling and
18 eIF2 α , which has been but barely addressed until now [47].

19 Finally, an emerging axis focuses on the protective properties of the N1 PrP^C fragment.
20 Multiple studies have ascribed the neuroprotective role of PrP^C to the N-terminal
21 domain and have suggested that this action is, at least partly, supported by its soluble N1
22 fragment [92, 133]. Recently, building upon the mapping of the binding sites of A β
23 oligomers to the N-terminal region of PrP^C (see Fig. 1), several groups have documented
24 that soluble N1 is endowed with the capacity to trap A β oligomers and thereby protect

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

against A β -induced neurotoxicity in various paradigms [25, 134-136]. The challenge now is to translate these observations into therapeutic strategies [137].

Conclusion

The recent array of convergent data supporting the view that the subversion of PrP^C-dependent signalling cascades is a pathogenic trigger in both prion and Alzheimer's diseases have sparked renewed interest on the physiological neuronal function exerted by PrP^C. While considerable progress has been achieved in deciphering the signalling events instructed by PrP^C, the challenge now is to reach a global view of the pathways orchestrated by PrP^C according to the ligands at the play and the neuronal subtype considered, as well as their potential interconnections. Besides, further increasing our knowledge of PrP^C-dependent signalling cascades may allow the identification of additional effectors, a subset of which may represent potential targets according to their status in disease. Overall, multiple strategies are now being exploited to counteract the toxic cascades arising from the deviation of PrP^C-mediated signalling and we may hope some will reach success in the near future.

Acknowledgements

We apologize to colleagues in the field whose work we were unable to cite owing to space limitations. We acknowledge financial support from the ANR, INSERM, as well as the Region Ile de France (DIM-Stem Pôle).

The authors declare no conflict of interest.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1:
Downstream targets of PrP^C-mediated signalling in a neuronal context (IEG: immediate early genes; SMase: sphingomyelinase)

Targets	Cell type / tissue	Physiological response	Reference
Src kinases			
<i>Fyn</i>	1C11-derived bioaminergic neuronal cells		[40]
	Primary hippocampal neurons	Neurite outgrowth	[29]
	PC12 neuronal cells	Neurite outgrowth	[41]
<i>Lyn</i>	1C11-derived serotonergic neuronal cells	Potentialiation of neurotransmitter release	[27]
	Raphe	Potentialiation of neurotransmitter release	[27]
PI3K/Akt			
PKA	N2a neuroblastoma cells	Cell survival	[46]
	Primary hippocampal neurons	Protein synthesis	[47]
	1C11-derived serotonergic neuronal cells	Potentialiation of neurotransmitter release	[27]
PKA	Rat retina	Neuroprotection	[49]
	Primary hippocampal neurons	Neuroprotection	[50]
	Hippocampal slices	Synaptic plasticity	[48]
NADPH oxidase			
ERK1/2	1C11-derived bioaminergic neuronal cells		[51]
	Rat retina	Neuroprotection	[49]
	SN56 cells		[52]
	Primary hippocampal neurons	Neurite outgrowth	[50]
	1C11-derived bioaminergic neuronal cells		[51]
GSK3β	Primary hippocampal neurons	Neurite outgrowth	[56]
	1C11-derived serotonergic neuronal cells	Potentialiation of neurotransmitter release	[27]
	Raphe	Potentialiation of neurotransmitter release	[27]
TACE			
mTOR	1C11-derived bioaminergic neuronal cells	Neurotransmitter catabolism	[53]
	Primary hippocampal neurons	Protein synthesis	[47]
CREB			
SMase	1C11-derived bioaminergic neuronal cells	Transcription of IEG	[54]
	Hippocampal cell lines, mouse brain		[55]
Ca²⁺	Hippocampal neurons		[58]
	Primary hippocampal neurons	Neuroprotection, neuronal survival	[57]
	Primary hippocampal neurons	Neurite outgrowth	[56]
	Dorsal root ganglia neurons	Axonal growth	[59]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 2:
Functional interactions between PrP^C-mediated signalling with neurotransmitter receptors in physiological or pathological contexts

Receptor	Cell type / tissue	Link with PrP ^C signalling	References
Serotonergic			
<i>5-HT1BR, 5-HT2AR, 5-HT2BR</i>	1C11-derived serotonergic neuronal cells	Modulation of couplings	[40]
<i>5-HT1BR</i>	1C11-derived serotonergic neuronal cells	Relocalization of receptor, blockade of activity	[27]
Glutamatergic metabotropic			
<i>mGlu1R, mGluR5</i>	Primary hippocampal neurons	LNγ1-induced neurite outgrowth	[56]
<i>mGlu5R</i>	Cortical neurons	Aβ-induced alteration of neuronal processes	[63]
<i>mGlu5R</i>	Hippocampus	Aβ-induced facilitation of LTD	[64]
Glutamatergic ionotropic			
<i>GluN2DR</i>	Hippocampal slices	Negative regulation	[66]
<i>GluN2BR</i>	Hippocampal slices	Internalization	[67]
<i>NMDAR</i>	SH-SY5Y cells	Aβ-induced neurotoxicity	[68]
<i>GluN2BR</i>	Cortical neurons	Aβ-induced Internalization	[44]
<i>NMDAR</i>	Hippocampal neurons	Aβ-induced neurotoxicity	[69]
Cholinergic ionotropic			
<i>α7nAChR</i>	Hippocampal neurons	Neuroprotection, neuronal survival	[57]

Legends to figures

Figure 1: Secondary structure of mouse PrP^C. The alpha-cleavage (green) occurs at position 111/112 and generates the N1 and C1 fragments. The beta-cleavage (red) occurs in the vicinity of octapeptide repeats, which bind copper ions, and generates the N2 and C2 fragments. PrP^C can also be shed from the plasma membrane through the action of ADAM10. Two main binding sites have been mapped for A β to the very N-terminus (AA 23–50) and near the alpha-cleavage site at amino acids 92–110. The protective factor STI-1 and the matrix protein laminin bind to PrP^C at amino acids 113–128 and 173–182, respectively. The alpha helices (α) and beta sheets (β) are indicated.

Figure 2: Summary of the signalling cascades mobilized by PrP^C in response to the binding of STI-1 or laminin, as well as their biological outcomes. Responses to PrP^C-dependent signalling include neuroprotection [49] / neuronal survival [57], protein synthesis [47], potentiation of neurotransmitter release [27], axonal growth [59] and neurite outgrowth [56].

Figure 3: Schematic representation of the corruption of PrP^C-mediated signalling in prion and Alzheimer's diseases. PrP^{Sc} accumulation and A β oligomers promote the recruitment of various effectors of PrP^C, including Fyn, NADPH oxidase and the MAP kinases ERK1/2. In the context of prion infection, it leads to a strong reduction in MMP-9 mRNAs and activity, which drives impairment in A β clearance, causing A β accumulation [43]. In Alzheimer's disease, the A β -PrP^C-Fyn cascade promotes Tau hyperphosphorylation [114] as well as impairment of the NMDA receptors surface localization [44]. PrP^C is also involved in the deleterious A β -dependent overactivation of PDK1 [107].

References

- 1 [1] A. Aguzzi, A.M. Calella, Prions: protein aggregation and infectious diseases,
2 *Physiol Rev.* 89 (2009) 1105-1152.
- 3 [2] H. Bueler, A. Aguzzi, A. Sailer, R.A. Greiner, P. Autenried, M. Aguet, C. Weissmann,
4 Mice devoid of PrP are resistant to scrapie, *Cell* 73 (1993) 1339-1347.
- 5 [3] R. Linden, V.R. Martins, M.A. Prado, M. Cammarota, I. Izquierdo, R.R. Brentani,
6 Physiology of the prion protein, *Physiol Rev.* 88 (2008) 673-728.
- 7 [4] E.T. Parkin, N.T. Watt, A.J. Turner, N.M. Hooper, Dual mechanisms for shedding of
8 the cellular prion protein, *J Biol Chem* 279 (2004) 11170-11178.
- 9 [5] B. Puig, H. Altmeppen, M. Glatzel, The GPI-anchoring of PrP: Implications in
10 sorting and pathogenesis, *Prion* 8 (2014).
- 11 [6] C.L. Haigh, S.Y. Marom, S.J. Collins, Copper, constitutive endoproteolytic
12 processing of the prion protein and cell signalling, *Front Biosci.* 15 (2010) 1086-1104.
- 13 [7] N.T. Watt, N.M. Hooper, The prion protein and neuronal zinc homeostasis, *Trends*
14 *Biochem Sci* 28 (2003) 406-410.
- 15 [8] M. Ermonval, S. Mouillet-Richard, P. Codogno, O. Kellermann, J. Botti, Evolving
16 views in prion glycosylation: functional and pathological implications, *Biochimie* 85
17 (2003) 33-45.
- 18 [9] F. Checler, B. Vincent, Alzheimer's and prion diseases: distinct pathologies,
19 common proteolytic denominators, *Trends Neurosci.* 25 (2002) 616-620.
- 20 [10] A.J. McDonald, J.P. Dibble, E.G. Evans, G.L. Millhauser, A new paradigm for
21 enzymatic control of alpha-cleavage and beta-cleavage of the prion protein, *J Biol Chem*
22 289 (2014) 803-813.
- 23 [11] I. Laffont-Proust, B.A. Faucheux, R. Hassig, V. Sazdovitch, S. Simon, J. Grassi, J.J.
24 Hauw, K.L. Moya, S. Haik, The N-terminal cleavage of cellular prion protein in the human
25 brain, *FEBS Lett.* 579 (2005) 6333-6337.
- 26 [12] L. Westergard, J.A. Turnbaugh, D.A. Harris, A naturally occurring C-terminal
27 fragment of the prion protein (PrP) delays disease and acts as a dominant-negative
28 inhibitor of PrP^{Sc} formation, *J Biol Chem* 286 (2011) 44234-44242.
- 29 [13] V. Lewis, N.M. Hooper, The role of lipid rafts in prion protein biology, *Front*
30 *Biosci.* 16 (2011) 151-168.
- 31 [14] R.G. Parton, M.A. del Pozo, Caveolae as plasma membrane sensors, protectors and
32 organizers, *Nat Rev Mol Cell Biol* 14 (2013) 98-112.
- 33 [15] D.A. Harris, H.L. True, New insights into prion structure and toxicity, *Neuron.* 50
34 (2006) 353-357.
- 35 [16] S. Brandner, A. Raeber, A. Sailer, T. Blattler, M. Fischer, C. Weissmann, A. Aguzzi,
36 Normal host prion protein (PrP^C) is required for scrapie spread within the central
37 nervous system, *Proc Natl Acad Sci U S A* 93 (1996) 13148-13151.
- 38 [17] G. Mallucci, A. Dickinson, J. Linehan, P.C. Klohn, S. Brandner, J. Collinge, Depleting
39 neuronal PrP in prion infection prevents disease and reverses spongiosis, *Science* 302
40 (2003) 871-874.
- 41 [18] B. Chesebro, M. Trifilo, R. Race, K. Meade-White, C. Teng, R. LaCasse, L. Raymond,
42 C. Favara, G. Baron, S. Priola, B. Caughey, E. Masliah, M. Oldstone, Anchorless prion
43 protein results in infectious amyloid disease without clinical scrapie, *Science* 308 (2005)
44 1435-1439.
- 45 [19] J. Lauren, Cellular prion protein as a therapeutic target in Alzheimer's disease, *J*
46 *Alzheimers Dis* 38 (2014) 227-244.

- 1 [20] E. Biasini, J.A. Turnbaugh, U. Unterberger, D.A. Harris, Prion protein at the
2 crossroads of physiology and disease, *Trends Neurosci* 35 (2011) 92-103.
- 3 [21] M.O. Collins, H. Husi, L. Yu, J.M. Brandon, C.N. Anderson, W.P. Blackstock, J.S.
4 Choudhary, S.G. Grant, Molecular characterization and comparison of the components
5 and multiprotein complexes in the postsynaptic proteome, *J Neurochem* 97 Suppl 1
6 (2006) 16-23.
- 7 [22] T. Suzuki, J. Zhang, S. Miyazawa, Q. Liu, M.R. Farzan, W.D. Yao, Association of
8 membrane rafts and postsynaptic density: proteomics, biochemical, and ultrastructural
9 analyses, *J Neurochem* 119 (2011) 64-77.
- 10 [23] D. Lingwood, K. Simons, Lipid rafts as a membrane-organizing principle, *Science*
11 327 (2010) 46-50.
- 12 [24] A.S. Rambold, V. Muller, U. Ron, N. Ben-Tal, K.F. Winklhofer, J. Tatzelt, Stress-
13 protective signalling of prion protein is corrupted by scrapie prions, *EMBO J.* 27 (2008)
14 1974-1984.
- 15 [25] M. Beland, J. Motard, A. Barbarin, X. Roucou, PrP(C) homodimerization stimulates
16 the production of PrPC cleaved fragments PrPN1 and PrPC1, *J Neurosci* 32 (2012)
17 13255-13263.
- 18 [26] M. Beland, X. Roucou, Taking advantage of physiological proteolytic processing of
19 the prion protein for a therapeutic perspective in prion and Alzheimer diseases, *Prion* 8
20 (2013).
- 21 [27] J. Hernandez-Rapp, S. Martin-Lanneree, T.Z. Hirsch, E. Pradines, A. Alleaume-
22 Butaux, B. Schneider, A. Baudry, J.M. Launay, S. Mouillet-Richard, A PrPC-caveolin-Lyn
23 complex negatively controls neuronal GSK3 β and serotonin 1B receptor, *Sci Rep* (in
24 press).
- 25 [28] C. Hundt, S. Weiss, PrP dimerization in the yeast two-hybrid system, *International*
26 *symposium on the characterization and diagnosis of prion diseases in animals and man*
27 (1999).
- 28 [29] A. Santuccione, V. Sytnyk, I. Leshchyn'ska, M. Schachner, Prion protein recruits its
29 neuronal receptor NCAM to lipid rafts to activate p59fyn and to enhance neurite
30 outgrowth, *J Cell Biol* 169 (2005) 341-354.
- 31 [30] S. Chen, A. Mange, L. Dong, S. Lehmann, M. Schachner, Prion protein as trans-
32 interacting partner for neurons is involved in neurite outgrowth and neuronal survival,
33 *Mol Cell Neurosci* 22 (2003) 227-233.
- 34 [31] J. Kanaani, S.B. Prusiner, J. Diacovo, S. Baekkeskov, G. Legname, Recombinant
35 prion protein induces rapid polarization and development of synapses in embryonic rat
36 hippocampal neurons in vitro, *J Neurochem* 95 (2005) 1373-1386.
- 37 [32] S.A. Romano, Y. Cordeiro, L.M. Lima, M.H. Lopes, J.L. Silva, D. Foguel, R. Linden,
38 Reciprocal remodeling upon binding of the prion protein to its signaling partner
39 hop/STI1, *FASEB J* 23 (2009) 4308-4316.
- 40 [33] D.M. Kaiser, M. Acharya, P.L. Leighton, H. Wang, N. Daude, S. Wohlgemuth, B. Shi,
41 W.T. Allison, Amyloid beta precursor protein and prion protein have a conserved
42 interaction affecting cell adhesion and CNS development, *PLoS One* 7 (2012) e51305.
- 43 [34] G. Schmitt-Ulms, K. Hansen, J. Liu, C. Cowdrey, J. Yang, S.J. DeArmond, F.E. Cohen,
44 S.B. Prusiner, M.A. Baldwin, Time-controlled transcardiac perfusion cross-linking for the
45 study of protein interactions in complex tissues, *Nat Biotechnol.* 22 (2004) 724-731.
- 46 [35] J. Lauren, D.A. Gimbel, H.B. Nygaard, J.W. Gilbert, S.M. Strittmatter, Cellular prion
47 protein mediates impairment of synaptic plasticity by amyloid-beta oligomers, *Nature.*
48 457 (2009) 1128-1132.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [36] C. Haass, D.J. Selkoe, Soluble protein oligomers in neurodegeneration: lessons
2 from the Alzheimer's amyloid beta-peptide, *Nat Rev Mol Cell Biol* 8 (2007) 101-112.
- 3 [37] F. Dohler, D. Sepulveda-Falla, S. Krasemann, H. Altmepfen, H. Schluter, D.
4 Hildebrand, I. Zerr, J. Matschke, M. Glatzel, High molecular mass assemblies of amyloid-
5 beta oligomers bind prion protein in patients with Alzheimer's disease, *Brain* 137
6 (2014) 873-886.
- 7 [38] V.G. Ostapchenko, F.H. Beraldo, A.H. Mohammad, Y.F. Xie, P.H. Hirata, A.C.
8 Magalhaes, G. Lamour, H. Li, A. Maciejewski, J.C. Belrose, B.L. Teixeira, M. Fahnstock,
9 S.T. Ferreira, N.R. Cashman, G.N. Hajj, M.F. Jackson, W.Y. Choy, J.F. Macdonald, V.R.
10 Martins, V.F. Prado, M.A. Prado, The prion protein ligand, stress-inducible
11 phosphoprotein 1, regulates amyloid-beta oligomer toxicity, *J Neurosci* 33 (2013)
12 16552-16564.
- 13 [39] I. Stefanova, V. Horejsi, I.J. Ansotegui, W. Knapp, H. Stockinger, GPI-anchored cell-
14 surface molecules complexed to protein tyrosine kinases, *Science* 254 (1991) 1016-
15 1019.
- 16 [40] S. Mouillet-Richard, M. Ermonval, C. Chebassier, J.L. Laplanche, S. Lehmann, J.M.
17 Launay, O. Kellermann, Signal transduction through prion protein, *Science* 289 (2000)
18 1925-1928.
- 19 [41] B. Pantera, C. Bini, P. Cirri, P. Paoli, G. Camici, G. Manao, A. Caselli, PrPc activation
20 induces neurite outgrowth and differentiation in PC12 cells: role for caveolin-1 in the
21 signal transduction pathway, *J Neurochem* 110 (2009) 194-207.
- 22 [42] C.M. Stern, P.G. Mermelstein, Caveolin regulation of neuronal intracellular
23 signaling, *Cell Mol Life Sci* 67 (2010) 3785-3795.
- 24 [43] E. Pradines, J. Hernandez-Rapp, A. Villa-Diaz, C. Dakowski, H. Ardila-Osorio, S.
25 Haik, B. Schneider, J.M. Launay, O. Kellermann, J.M. Torres, S. Mouillet-Richard,
26 Pathogenic prions deviate PrP(C) signaling in neuronal cells and impair A-beta
27 clearance, *Cell Death Dis* 4 (2013) e456.
- 28 [44] J.W. Um, H.B. Nygaard, J.K. Heiss, M.A. Kostylev, M. Stagi, A. Vortmeyer, T.
29 Wisniewski, E.C. Gunther, S.M. Strittmatter, Alzheimer amyloid-beta oligomer bound to
30 postsynaptic prion protein activates Fyn to impair neurons, *Nat Neurosci* 15 (2012)
31 1227-1235.
- 32 [45] N. Loberto, S. Prioni, A. Bettiga, V. Chigorno, A. Prinetti, S. Sonnino, The
33 membrane environment of endogenous cellular prion protein in primary rat cerebellar
34 neurons, *J Neurochem* 95 (2005) 771-783.
- 35 [46] N. Vassallo, J. Herms, C. Behrens, B. Krebs, K. Saeki, T. Onodera, O. Windl, H.A.
36 Kretschmar, Activation of phosphatidylinositol 3-kinase by cellular prion protein and
37 its role in cell survival, *Biochem Biophys Res Commun.* 332 (2005) 75-82.
- 38 [47] M. Roffe, F.H. Beraldo, R. Bester, M. Nunziante, C. Bach, G. Mancini, S. Gilch, I.
39 Vorberg, B.A. Castilho, V.R. Martins, G.N. Hajj, Prion protein interaction with stress-
40 inducible protein 1 enhances neuronal protein synthesis via mTOR, *Proc Natl Acad Sci U*
41 *S A* 107 (2010) 13147-13152.
- 42 [48] M.D. Caiati, V.F. Safiulina, G. Fattorini, S. Sivakumaran, G. Legname, E. Cherubini,
43 PrPC controls via protein kinase A the direction of synaptic plasticity in the immature
44 hippocampus, *J Neurosci* 33 (2013) 2973-2983.
- 45 [49] L.B. Chiarini, A.R. Freitas, S.M. Zanata, R.R. Brentani, V.R. Martins, R. Linden,
46 Cellular prion protein transduces neuroprotective signals, *Embo J.* 21 (2002) 3317-
47 3326.
- 48 [50] M.H. Lopes, G.N. Hajj, A.G. Muras, G.L. Mancini, R.M. Castro, K.C. Ribeiro, R.R.
49 Brentani, R. Linden, V.R. Martins, Interaction of cellular prion and stress-inducible
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 protein 1 promotes neuritogenesis and neuroprotection by distinct signaling pathways, *J*
2 *Neurosci* 25 (2005) 11330-11339.

3 [51] B. Schneider, V. Mutel, M. Pietri, M. Ermonval, S. Mouillet-Richard, O. Kellermann,
4 NADPH oxidase and extracellular regulated kinases 1/2 are targets of prion protein
5 signaling in neuronal and nonneuronal cells, *Proc Natl Acad Sci U S A* 100 (2003) 13326-
6 13331.

7 [52] F.A. Caetano, M.H. Lopes, G.N. Hajj, C.F. Machado, C. Pinto Arantes, A.C. Magalhaes,
8 P. Vieira Mde, T.A. Americo, A.R. Massensini, S.A. Priola, I. Vorberg, M.V. Gomez, R.
9 Linden, V.F. Prado, V.R. Martins, M.A. Prado, Endocytosis of prion protein is required for
10 ERK1/2 signaling induced by stress-inducible protein 1, *J Neurosci*. 28 (2008) 6691-
11 6702.

12 [53] E. Pradines, D. Loubet, S. Mouillet-Richard, P. Manivet, J.M. Launay, O. Kellermann,
13 B. Schneider, Cellular prion protein coupling to TACE-dependent TNF-alpha shedding
14 controls neurotransmitter catabolism in neuronal cells, *J Neurochem*. 110 (2009) 912-
15 923.

16 [54] E. Pradines, D. Loubet, B. Schneider, J.M. Launay, O. Kellermann, S. Mouillet-
17 Richard, CREB-dependent gene regulation by prion protein: impact on MMP-9 and beta-
18 dystroglycan, *Cell Signal*. 20 (2008) 2050-2058.

19 [55] R. Schmalzbauer, S. Eigenbrod, S. Winoto-Morbach, W. Xiang, S. Schutze, U.
20 Bertsch, H.A. Kretzschmar, Evidence for an association of prion protein and
21 sphingolipid-mediated signaling, *J Neurochem*. 106 (2008) 1459-1470.

22 [56] F.H. Beraldo, C.P. Arantes, T.G. Santos, C.F. Machado, M. Roffe, G.N. Hajj, K.S. Lee,
23 A.C. Magalhaes, F.A. Caetano, G.L. Mancini, M.H. Lopes, T.A. Americo, M.H. Magdesian, S.S.
24 Ferguson, R. Linden, M.A. Prado, V.R. Martins, Metabotropic glutamate receptors
25 transduce signals for neurite outgrowth after binding of the prion protein to laminin
26 gamma1 chain, *FASEB J* 25 (2011) 265-279.

27 [57] F.H. Beraldo, C.P. Arantes, T.G. Santos, N.G. Queiroz, K. Young, R.J. Rylett, R.P.
28 Markus, M.A. Prado, V.R. Martins, Role of alpha7 nicotinic acetylcholine receptor in
29 calcium signaling induced by prion protein interaction with stress-inducible protein 1, *J*
30 *Biol Chem* 285 (2010) 36542-36550.

31 [58] M. Fuhrmann, T. Bittner, G. Mitteregger, N. Haider, S. Moosmang, H. Kretzschmar,
32 J. Herms, Loss of the cellular prion protein affects the Ca²⁺ homeostasis in hippocampal
33 CA1 neurons, *J Neurochem*. 98 (2006) 1876-1885.

34 [59] T.G. Santos, F.H. Beraldo, G.N. Hajj, M.H. Lopes, M. Roffe, F.C. Lupinacci, V.G.
35 Ostapchenko, V.F. Prado, M.A. Prado, V.R. Martins, Laminin-gamma1 chain and stress
36 inducible protein 1 synergistically mediate PrPC-dependent axonal growth via Ca²⁺
37 mobilization in dorsal root ganglia neurons, *J Neurochem* 124 (2013) 210-223.

38 [60] J.A. Allen, R.A. Halverson-Tamboli, M.M. Rasenick, Lipid raft microdomains and
39 neurotransmitter signalling, *Nat Rev Neurosci*. 8 (2007) 128-140.

40 [61] S. Mouillet-Richard, M. Pietri, B. Schneider, C. Vidal, V. Mutel, J.M. Launay, O.
41 Kellermann, Modulation of serotonergic receptor signaling and cross-talk by prion
42 protein, *J Biol Chem* 280 (2005) 4592-4601.

43 [62] S.S. Willard, S. Koochekpour, Glutamate, glutamate receptors, and downstream
44 signaling pathways, *Int J Biol Sci* 9 (2013) 948-959.

45 [63] J.W. Um, A.C. Kaufman, M. Kostylev, J.K. Heiss, M. Stagi, H. Takahashi, M.E. Kerrisk,
46 A. Vortmeyer, T. Wisniewski, A.J. Koleske, E.C. Gunther, H.B. Nygaard, S.M. Strittmatter,
47 Metabotropic glutamate receptor 5 is a coreceptor for Alzheimer abeta oligomer bound
48 to cellular prion protein, *Neuron* 79 (2013) 887-902.

- 1 [64] N.W. Hu, A.J. Nicoll, D. Zhang, A.J. Mably, T. O'Malley, S.A. Purro, C. Terry, J.
2 Collinge, D.M. Walsh, M.J. Rowan, mGlu5 receptors and cellular prion protein mediate
3 amyloid-beta-facilitated synaptic long-term depression in vivo, *Nat Commun* 5 (2014)
4 3374.
- 5 [65] G.L. Collingridge, J.T. Isaac, Y.T. Wang, Receptor trafficking and synaptic plasticity,
6 *Nat Rev Neurosci* 5 (2004) 952-962.
- 7 [66] H. Khosravani, Y. Zhang, S. Tsutsui, S. Hameed, C. Altier, J. Hamid, L. Chen, M.
8 Villemaire, Z. Ali, F.R. Jirik, G.W. Zamponi, Prion protein attenuates excitotoxicity by
9 inhibiting NMDA receptors, *J Cell Biol* 181 (2008) 551-565.
- 10 [67] A. Petit-Paitel, B. Menard, A. Guyon, V. Beringue, J.L. Nahon, N. Zsurgger, J. Chabry,
11 Prion protein is a key determinant of alcohol sensitivity through the modulation of N-
12 methyl-D-aspartate receptor (NMDAR) activity, *PLoS One* 7 (2012) e34691.
- 13 [68] U.K. Resenberger, A. Harmeier, A.C. Woerner, J.L. Goodman, V. Muller, R. Krishnan,
14 R.M. Vabulas, H.A. Kretzschmar, S. Lindquist, F.U. Hartl, G. Multhaup, K.F. Winklhofer, J.
15 Tatzelt, The cellular prion protein mediates neurotoxic signalling of beta-sheet-rich
16 conformers independent of prion replication, *EMBO J* 30 (2011) 2057-2070.
- 17 [69] H. You, S. Tsutsui, S. Hameed, T.J. Kannanayakal, L. Chen, P. Xia, J.D. Engbers, S.A.
18 Lipton, P.K. Stys, G.W. Zamponi, Abeta neurotoxicity depends on interactions between
19 copper ions, prion protein, and N-methyl-D-aspartate receptors, *Proc Natl Acad Sci U S A*
20 109 (2012) 1737-1742.
- 21 [70] M. Schmitz, C. Greis, P. Ottis, C.J. Silva, W.J. Schulz-Schaeffer, A. Wrede, K. Koppe,
22 B. Onisko, J.R. Requena, N. Govindarajan, C. Korth, A. Fischer, I. Zerr, Loss of Prion
23 Protein Leads to Age-Dependent Behavioral Abnormalities and Changes in Cytoskeletal
24 Protein Expression, *Mol Neurobiol* (2014).
- 25 [71] T. Golub, S. Wacha, P. Caroni, Spatial and temporal control of signaling through
26 lipid rafts, *Curr Opin Neurobiol.* 14 (2004) 542-550.
- 27 [72] S. Mishra, P.G. Joshi, Lipid raft heterogeneity: an enigma, *J Neurochem* 103 Suppl
28 1 (2007) 135-142.
- 29 [73] D.R. Taylor, N.M. Hooper, The low-density lipoprotein receptor-related protein 1
30 (LRP1) mediates the endocytosis of the cellular prion protein, *Biochem J.* 402 (2007) 17-
31 23.
- 32 [74] I. Singh, A.P. Sagare, M. Coma, D. Perlmutter, R. Gelein, R.D. Bell, R.J. Deane, E.
33 Zhong, M. Parisi, J. Ciszewski, R.T. Kasper, R. Deane, Low levels of copper disrupt brain
34 amyloid-beta homeostasis by altering its production and clearance, *Proc Natl Acad Sci U*
35 *S A* (2013).
- 36 [75] J.V. Rushworth, H.H. Griffiths, N.T. Watt, N.M. Hooper, Prion protein-mediated
37 toxicity of amyloid-beta oligomers requires lipid rafts and the transmembrane LRP1, *J*
38 *Biol Chem* 288 (2013) 8935-8951.
- 39 [76] F. Cheng, J. Lindqvist, C.L. Haigh, D.R. Brown, K. Mani, Copper-dependent co-
40 internalization of the prion protein and glypican-1, *J Neurochem.* 98 (2006) 1445-1457.
- 41 [77] D.R. Taylor, I.J. Whitehouse, N.M. Hooper, Glypican-1 mediates both prion protein
42 lipid raft association and disease isoform formation, *PLoS Pathog.* 5 (2009) e1000666.
- 43 [78] S. Gauczynski, J.M. Peyrin, S. Haik, C. Leucht, C. Hundt, R. Rieger, S. Krasemann, J.P.
44 Deslys, D. Dormont, C.I. Lasmezas, S. Weiss, The 37-kDa/67-kDa laminin receptor acts as
45 the cell-surface receptor for the cellular prion protein, *EMBO J* 20 (2001) 5863-5875.
- 46 [79] C. Leucht, S. Simoneau, C. Rey, K. Vana, R. Rieger, C.I. Lasmezas, S. Weiss, The 37
47 kDa/67 kDa laminin receptor is required for PrP(Sc) propagation in scrapie-infected
48 neuronal cells, *EMBO Rep* 4 (2003) 290-295.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [80] F.A. Caetano, F.H. Beraldo, G.N. Hajj, A.L. Guimaraes, S. Jurgensen, A.P.
2 Wasilewska-Sampaio, P.H. Hirata, I. Souza, C.F. Machado, D.Y. Wong, F.G. De Felice, S.T.
3 Ferreira, V.F. Prado, R.J. Rylett, V.R. Martins, M.A. Prado, Amyloid-beta oligomers
4 increase the localization of prion protein at the cell surface, *J Neurochem* 117 (2011)
5 538-553.
- 6 [81] G.N. Hajj, M.H. Lopes, A.F. Mercadante, S.S. Veiga, R.B. da Silveira, T.G. Santos, K.C.
7 Ribeiro, M.A. Juliano, S.G. Jacchieri, S.M. Zanata, V.R. Martins, Cellular prion protein
8 interaction with vitronectin supports axonal growth and is compensated by integrins, *J*
9 *Cell Sci.* 120 (2007) 1915-1926.
- 10 [82] E. Graner, A.F. Mercadante, S.M. Zanata, O.V. Forlenza, A.L. Cabral, S.S. Veiga, M.A.
11 Juliano, R. Roesler, R. Walz, A. Minetti, I. Izquierdo, V.R. Martins, R.R. Brentani, Cellular
12 prion protein binds laminin and mediates neuritogenesis, *Brain Res Mol Brain Res* 76
13 (2000) 85-92.
- 14 [83] C.F. Machado, F.H. Beraldo, T.G. Santos, D. Bourgeon, M.C. Landemberger, M.
15 Roffe, V.R. Martins, Disease-associated mutations in the prion protein impair laminin-
16 induced process outgrowth and survival, *J Biol Chem* 287 (2012) 43777-43788.
- 17 [84] D. Loubet, C. Dakowski, M. Pietri, E. Pradines, S. Bernard, J. Callebert, H. Ardila-
18 Osorio, S. Mouillet-Richard, J.M. Launay, O. Kellermann, B. Schneider, Neuritogenesis: the
19 prion protein controls beta1 integrin signaling activity, *FASEB J* 26 (2012) 678-690.
- 20 [85] A.S. Coitinho, A.R. Freitas, M.H. Lopes, G.N. Hajj, R. Roesler, R. Walz, J.I. Rossato, M.
21 Cammarota, I. Izquierdo, V.R. Martins, R.R. Brentani, The interaction between prion
22 protein and laminin modulates memory consolidation, *Eur J Neurosci.* 24 (2006) 3255-
23 3264.
- 24 [86] A.S. Coitinho, M.H. Lopes, G.N. Hajj, J.I. Rossato, A.R. Freitas, C.C. Castro, M.
25 Cammarota, R.R. Brentani, I. Izquierdo, V.R. Martins, Short-term memory formation and
26 long-term memory consolidation are enhanced by cellular prion association to stress-
27 inducible protein 1, *Neurobiol Dis.* 26 (2007) 282-290.
- 28 [87] M. Costa-Mattioli, L.M. Monteggia, mTOR complexes in neurodevelopmental and
29 neuropsychiatric disorders, *Nat Neurosci* 16 (2013) 1537-1543.
- 30 [88] G.M. Thomas, R.L. Huganir, MAPK cascade signalling and synaptic plasticity, *Nat*
31 *Rev Neurosci.* 5 (2004) 173-183.
- 32 [89] C.A. Bradley, S. Peineau, C. Taghibiglou, C.S. Nicolas, D.J. Whitcomb, Z.A.
33 Bortolotto, B.K. Kaang, K. Cho, Y.T. Wang, G.L. Collingridge, A pivotal role of GSK-3 in
34 synaptic plasticity, *Front Mol Neurosci* 5 (2012) 13.
- 35 [90] K. Inoki, H. Ouyang, T. Zhu, C. Lindvall, Y. Wang, X. Zhang, Q. Yang, C. Bennett, Y.
36 Harada, K. Stankunas, C.Y. Wang, X. He, O.A. MacDougald, M. You, B.O. Williams, K.L.
37 Guan, TSC2 integrates Wnt and energy signals via a coordinated phosphorylation by
38 AMPK and GSK3 to regulate cell growth, *Cell* 126 (2006) 955-968.
- 39 [91] O. Kaidanovich-Beilin, J.R. Woodgett, GSK-3: Functional Insights from Cell Biology
40 and Animal Models, *Front Mol Neurosci* 4 (2011) 40.
- 41 [92] U.K. Resenberger, K.F. Winklhofer, J. Tatzelt, Neuroprotective and neurotoxic
42 signaling by the prion protein, *Top Curr Chem* 305 (2011) 101-119.
- 43 [93] X. Roucou, A.C. Leblanc, Cellular prion protein neuroprotective function:
44 implications in prion diseases, *J Mol Med* 83 (2005) 3-11. .
- 45 [94] S.M. Zanata, M.H. Lopes, A.F. Mercadante, G.N. Hajj, L.B. Chiarini, R. Nomizo, A.R.
46 Freitas, A.L. Cabral, K.S. Lee, M.A. Juliano, E. de Oliveira, S.G. Jachieri, A. Burlingame, L.
47 Huang, R. Linden, R.R. Brentani, V.R. Martins, Stress-inducible protein 1 is a cell surface
48 ligand for cellular prion that triggers neuroprotection, *EMBO J* 21 (2002) 3307-3316.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [95] S. Haider, S. Khaliq, D.J. Haleem, Enhanced serotonergic neurotransmission in the
2 hippocampus following tryptophan administration improves learning acquisition and
3 memory consolidation in rats, *Pharmacol Rep.* 59 (2007) 53-57.
- 4 [96] K. Kobayashi, T. Kobayashi, Genetic evidence for noradrenergic control of long-
5 term memory consolidation, *Brain Dev.* 23 (2001) S16-23.
- 6 [97] G.G. Kovacs, H. Budka, Prion diseases: from protein to cell pathology, *Am J Pathol.*
7 172 (2008) 555-565.
- 8 [98] J.M. Ledoux, Effects on the serotonergic system in sub-acute transmissible
9 spongiform encephalopathies: current data, hypotheses, suggestions for
10 experimentation, *Med Hypotheses* 64 (2005) 910-918.
- 11 [99] C. Vidal, C. Herzog, A.M. Haeberle, C. Bombarde, M.C. Miquel, J. Carimalo, J.M.
12 Launay, S. Mouillet-Richard, C. Lasmezas, D. Dormont, O. Kellermann, Y. Bailly, Early
13 dysfunction of central 5-HT system in a murine model of bovine spongiform
14 encephalopathy, *Neuroscience.* 160 (2009) 731-743.
- 15 [100] C. Bate, A. Williams, Neurodegeneration induced by clustering of sialylated
16 glycosylphosphatidylinositols of prion proteins, *J Biol Chem* 287 (2012) 7935-7944.
- 17 [101] J. Falsig, T. Sonati, U.S. Herrmann, D. Saban, B. Li, K. Arroyo, B. Ballmer, P.P.
18 Liberski, A. Aguzzi, Prion pathogenesis is faithfully reproduced in cerebellar organotypic
19 slice cultures, *PLoS Pathog* 8 (2012) e1002985.
- 20 [102] T. Sonati, R.R. Reimann, J. Falsig, P.K. Baral, T. O'Connor, S. Hornemann, S.
21 Yaganoglu, B. Li, U.S. Herrmann, B. Wieland, M. Swayampakula, M.H. Rahman, D. Das, N.
22 Kav, R. Riek, P.P. Liberski, M.N. James, A. Aguzzi, The toxicity of anti-prion antibodies is
23 mediated by the flexible tail of the prion protein, *Nature* 501 (2013) 102-106.
- 24 [103] R. Chiesa, P. Piccardo, E. Biasini, B. Ghetti, D.A. Harris, Aggregated, wild-type prion
25 protein causes neurological dysfunction and synaptic abnormalities, *J Neurosci.* 28
26 (2008) 13258-13267.
- 27 [104] F. Agostini, C.G. Dotti, A. Perez-Canamas, M.D. Ledesma, F. Benetti, G. Legname,
28 Prion protein accumulation in lipid rafts of mouse aging brain, *PLoS One* 8 (2013)
29 e74244.
- 30 [105] M. Russelakis-Carneiro, C. Hetz, K. Maundrell, C. Soto, Prion replication alters the
31 distribution of synaptophysin and caveolin 1 in neuronal lipid rafts, *Am J Pathol.* 165
32 (2004) 1839-1848.
- 33 [106] H.L. Cui, B. Guo, B. Scicluna, B.M. Coleman, V.A. Lawson, L. Ellett, P.J. Meikle, M.
34 Bukrinsky, N. Mukhamedova, D. Sviridov, A.F. Hill, Prion infection impairs cholesterol
35 metabolism in neuronal cells, *J Biol Chem* 289 (2014) 789-802.
- 36 [107] M. Pietri, C. Dakowski, S. Hannaoui, A. Alleaume-Butaux, J. Hernandez-Rapp, A.
37 Ragagnin, S. Mouillet-Richard, S. Haik, Y. Bailly, J.M. Peyrin, J.M. Launay, O. Kellermann, B.
38 Schneider, PDK1 decreases TACE-mediated alpha-secretase activity and promotes
39 disease progression in prion and Alzheimer's diseases, *Nat Med* 19 (2013) 1124-1131.
- 40 [108] J. Hernandez-Rapp, S. Martin-Lannere, T.Z. Hirsch, J.M. Launay, S. Mouillet-
41 Richard, Hijacking PrP(c)-dependent signal transduction: when prions impair Aβ
42 clearance, *Front Aging Neurosci* 6 (2014) 25.
- 43 [109] W. Kudo, H.P. Lee, W.Q. Zou, X. Wang, G. Perry, X. Zhu, M.A. Smith, R.B. Petersen,
44 H.G. Lee, Cellular prion protein is essential for oligomeric amyloid-beta-induced
45 neuronal cell death, *Hum Mol Genet* 21 (2012) 1138-1144.
- 46 [110] D.A. Gimbel, H.B. Nygaard, E.E. Coffey, E.C. Gunther, J. Lauren, Z.A. Gimbel, S.M.
47 Strittmatter, Memory impairment in transgenic Alzheimer mice requires cellular prion
48 protein, *J Neurosci* 30 (2010) 6367-6374.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [111] J. Nah, J.O. Pyo, S. Jung, S.M. Yoo, T.I. Kam, J. Chang, J. Han, A.A.S. Soo, T. Onodera,
2 Y.K. Jung, BECN1/Beclin 1 is recruited into lipid rafts by prion to activate autophagy in
3 response to amyloid beta 42, *Autophagy* 9 (2013) 2009-2021.
- 4 [112] R. Williamson, T. Scales, B.R. Clark, G. Gibb, C.H. Reynolds, S. Kellie, I.N. Bird, I.M.
5 Varndell, P.W. Sheppard, I. Everall, B.H. Anderton, Rapid tyrosine phosphorylation of
6 neuronal proteins including tau and focal adhesion kinase in response to amyloid-beta
7 peptide exposure: involvement of Src family protein kinases, *J Neurosci.* 22 (2002) 10-
8 20.
- 9 [113] R. Williamson, A. Usardi, D.P. Hanger, B.H. Anderton, Membrane-bound beta-
10 amyloid oligomers are recruited into lipid rafts by a fyn-dependent mechanism, *Faseb J.*
11 22 (2008) 1552-1559.
- 12 [114] M. Larson, M.A. Sherman, F. Amar, M. Nuvolone, J.A. Schneider, D.A. Bennett, A.
13 Aguzzi, S.E. Lesne, The complex PrP(c)-Fyn couples human oligomeric Abeta with
14 pathological tau changes in Alzheimer's disease, *J Neurosci* 32 (2012) 16857-16871a.
- 15 [115] A. Thathiah, B. De Strooper, The role of G protein-coupled receptors in the
16 pathology of Alzheimer's disease, *Nat Rev Neurosci* 12 (2011) 73-87.
- 17 [116] C. Bate, A. Williams, Amyloid-beta-induced synapse damage is mediated via
18 cross-linkage of cellular prion proteins, *J Biol Chem* 286 (2011) 37955-37963.
- 19 [117] K.P. Walsh, L.S. Minamide, S.J. Kane, A.E. Shaw, D.R. Brown, B. Pulford, M.D. Zabel,
20 J.D. Lambeth, T.B. Kuhn, J.R. Bamburg, Amyloid-beta and Proinflammatory Cytokines
21 Utilize a Prion Protein-Dependent Pathway to Activate NADPH Oxidase and Induce
22 Cofilin-Actin Rods in Hippocampal Neurons, *PLoS One* 9 (2014) e95995.
- 23 [118] I. Benilova, B. De Strooper, Neuroscience. Promiscuous Alzheimer's amyloid: yet
24 another partner, *Science* 341 (2013) 1354-1355.
- 25 [119] S. Haik, G. Marcon, A. Mallet, M. Tettamanti, A. Welaratne, G. Giaccone, S. Azimi, V.
26 Pietrini, J.R. Fabreguettes, D. Imperiale, P. Cesaro, C. Buffa, C. Aucan, U. Lucca, L. Peckeu,
27 S. Suardi, C. Tranchant, I. Zerr, C. Houillier, V. Redaelli, H. Vespignani, A. Campanella, F.
28 Sellal, A. Krasnianski, D. Seilhean, U. Heinemann, F. Sedel, M. Canovi, M. Gobbi, G. Di Fede,
29 J.L. Laplanche, M. Pocchiari, M. Salmona, G. Forloni, J.P. Brandel, F. Tagliavini,
30 Doxycycline in Creutzfeldt-Jakob disease: a phase 2, randomised, double-blind, placebo-
31 controlled trial, *Lancet Neurol* 13 (2014) 150-158.
- 32 [120] Y. Roettger, Y. Du, M. Bacher, I. Zerr, R. Dodel, J.P. Bach, Immunotherapy in prion
33 disease, *Nat Rev Neurol* 9 (2013) 98-105.
- 34 [121] D.B. Freir, A.J. Nicoll, I. Klyubin, S. Panico, J.M. Mc Donald, E. Risse, E.A. Asante,
35 M.A. Farrow, R.B. Sessions, H.R. Saibil, A.R. Clarke, M.J. Rowan, D.M. Walsh, J. Collinge,
36 Interaction between prion protein and toxic amyloid beta assemblies can be
37 therapeutically targeted at multiple sites, *Nat Commun* 2 (2011) 336.
- 38 [122] K. Alier, L. Ma, J. Yang, D. Westaway, J.H. Jhamandas, Abeta inhibition of ionic
39 conductance in mouse basal forebrain neurons is dependent upon the cellular prion
40 protein PrPC, *J Neurosci* 31 (2011) 16292-16297.
- 41 [123] E. Chung, Y. Ji, Y. Sun, R.J. Kascsak, R.B. Kascsak, P.D. Mehta, S.M. Strittmatter, T.
42 Wisniewski, Anti-PrPC monoclonal antibody infusion as a novel treatment for cognitive
43 deficits in an Alzheimer's disease model mouse, *BMC Neurosci* 11 (2010) 130.
- 44 [124] A.E. Barry, I. Klyubin, J.M. Mc Donald, A.J. Mably, M.A. Farrell, M. Scott, D.M. Walsh,
45 M.J. Rowan, Alzheimer's disease brain-derived amyloid-beta-mediated inhibition of LTP
46 in vivo is prevented by immunotargeting cellular prion protein, *J Neurosci* 31 (2011)
47 7259-7263.
- 48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 [125] P.C. Klohn, M. Farmer, J.M. Linehan, C. O'Malley, M. Fernandez de Marco, W.
2 Taylor, M. Farrow, A. Khalili-Shirazi, S. Brandner, J. Collinge, PrP antibodies do not
3 trigger mouse hippocampal neuron apoptosis, *Science* 335 (2012) 52.
- 4 [126] L. Solfrosi, J.R. Criado, D.B. McGavern, S. Wirz, M. Sanchez-Alavez, S. Sugama, L.A.
5 DeGiorgio, B.T. Volpe, E. Wiseman, G. Abalos, E. Masliah, D. Gilden, M.B. Oldstone, B.
6 Conti, R.A. Williamson, Cross-linking cellular prion protein triggers neuronal apoptosis
7 in vivo, *Science* 303 (2004) 1514-1516.
- 8 [127] H.M. Gao, H. Zhou, J.S. Hong, NADPH oxidases: novel therapeutic targets for
9 neurodegenerative diseases, *Trends Pharmacol Sci* 33 (2012) 295-303.
- 10 [128] H.B. Nygaard, C.H. van Dycck, S.M. Strittmatter, Fyn kinase inhibition as a novel
11 therapy for Alzheimer's disease, *Alzheimers Res Ther* 6 (2014) 8.
- 12 [129] M. Halliday, G.R. Mallucci, Targeting the unfolded protein response in
13 neurodegeneration: A new approach to therapy, *Neuropharmacology* 76 Pt A (2014)
14 169-174.
- 15 [130] J.A. Moreno, H. Radford, D. Peretti, J.R. Steinert, N. Verity, M.G. Martin, M. Halliday,
16 J. Morgan, D. Dinsdale, C.A. Ortori, D.A. Barrett, P. Tsaytler, A. Bertolotti, A.E. Willis, M.
17 Bushell, G.R. Mallucci, Sustained translational repression by eIF2alpha-P mediates prion
18 neurodegeneration, *Nature* 485 (2012) 507-511.
- 19 [131] J.A. Moreno, M. Halliday, C. Molloy, H. Radford, N. Verity, J.M. Axten, C.A. Ortori,
20 A.E. Willis, P.M. Fischer, D.A. Barrett, G.R. Mallucci, Oral treatment targeting the unfolded
21 protein response prevents neurodegeneration and clinical disease in prion-infected
22 mice, *Sci Transl Med* 5 (2013) 206ra138.
- 23 [132] T. Ma, M.A. Trinh, A.J. Wexler, C. Bourbon, E. Gatti, P. Pierre, D.R. Cavener, E.
24 Klann, Suppression of eIF2alpha kinases alleviates Alzheimer's disease-related plasticity
25 and memory deficits, *Nat Neurosci* 16 (2013) 1299-1305.
- 26 [133] M.V. Guillot-Sestier, F. Checler, alpha-Secretase-derived cleavage of cellular prion
27 yields biologically active catabolites with distinct functions, *Neurodegener Dis* 10 (2012)
28 294-297.
- 29 [134] B.R. Fluharty, E. Biasini, M. Stravalaci, A. Sclip, L. Diomede, C. Balducci, P. La
30 Vitola, M. Messa, L. Colombo, G. Forloni, T. Borsello, M. Gobbi, D.A. Harris, An N-terminal
31 fragment of the prion protein binds to amyloid-beta oligomers and inhibits their
32 neurotoxicity in vivo, *J Biol Chem* 288 (2013) 7857-7866.
- 33 [135] M.V. Guillot-Sestier, C. Sunyach, S.T. Ferreira, M.P. Marzolo, C. Bauer, A. Thevenet,
34 F. Checler, alpha-Secretase-derived fragment of cellular prion, N1, protects against
35 monomeric and oligomeric amyloid beta (Abeta)-associated cell death, *J Biol Chem* 287
36 (2012) 5021-5032.
- 37 [136] K. Nieznanski, J.K. Choi, S. Chen, K. Surewicz, W.K. Surewicz, Soluble prion protein
38 inhibits amyloid-beta (Abeta) fibrillization and toxicity, *J Biol Chem* 287 (2012) 33104-
39 33108.
- 40 [137] M. Beland, X. Roucou, Taking advantage of physiological proteolytic processing of
41 the prion protein for a therapeutic perspective in prion and Alzheimer diseases, *Prion* 8
42 (2014).
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1
[Click here to download high resolution image](#)




Figure 1

Figure 2
[Click here to download high resolution image](#)




Figure 2

Figure 3
[Click here to download high resolution image](#)




Figure 3