



**HAL**  
open science

## Asymptotic-Möbius maps

Georg Alexander Gruetzner

► **To cite this version:**

| Georg Alexander Gruetzner. Asymptotic-Möbius maps. 2022. hal-03746707v4

**HAL Id: hal-03746707**

**<https://hal.science/hal-03746707v4>**

Preprint submitted on 16 Dec 2022 (v4), last revised 21 Feb 2023 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Asymptotic-Möbius maps.

Georg Grützner

## Abstract

Roughly speaking, a map between metric spaces is an asymptotic-Möbius map if it almost preserves cross-ratios of points being a large distance apart from each other. We show that under such maps, some large-scale notions of dimension increase: asymptotic dimension for finitely generated nilpotent groups, telescopic dimension for  $CAT(0)$  spaces.

## 1 Introduction

This paper introduces and investigates a class of maps that mimics quasi-Möbius maps in the context of geometric group theory.

In [13] Pierre Pansu introduces a notion of large-scale conformal maps that mimics the infinitesimal behavior of conformal maps. Briefly, large scale conformal maps map families of disjoint balls to families of weakly overlapping quasi-balls. It is a very flexible notion, that includes for example coarse embeddings. This flexibility makes the asymptotic behavior of such maps however less predictable.

In this paper, we introduce a more restrictive and semantically simpler notion that mimics quasi-Möbius behaviors. Roughly speaking, a map between metric spaces is an asymptotic-Möbius map if it almost preserves cross-ratios of points being a large distance apart from each other. In consequence, the results presented in this paper are much stronger than in [13].

### 1.1 Examples

We postpone the precise definition of asymptotic-Möbius ( $AM$ ) maps until later. Sources of examples of  $AM$ -maps are:

1. Quasi-isometric embeddings,
2. Sublinear-Bilipschitz-equivalences (i.e. maps that induce Lipschitz-Equivalences on asymptotic cones [5]),
3. Snowflaking (i.e. replacing a metric by a power of it),
4. Assouad maps from doubling metric spaces to  $\mathbb{R}^N$ .

For instance, every nilpotent Lie group or finitely generated group can be  $AM$ -embedded in Euclidean space of sufficiently high dimension [1]. We shall furthermore encounter infinite dimensional examples.

## 1.2 Results

Our main result is that under  $AM$ -maps some large-scale notion of dimension increases. The relevant notion depends on the class of groups considered.

**Theorem 3.** *Let  $G$  and  $G'$  be finitely generated nilpotent groups and  $f : G \rightarrow G'$  an  $AM$ -map, then  $\text{asdim}(G) \leq \text{asdim}(G')$ . Furthermore, if  $\text{asdim}(G) = \text{asdim}(G')$ , then the asymptotic cones of  $G$  and  $G'$  are isomorphic. Conversely, given nilpotent groups with isomorphic asymptotic cones, there exists an asymptotic-Möbius map between them.*

In the world of  $\text{CAT}(0)$ -spaces, the analogous theorem takes the following form.

**Theorem 4.** *Let  $X$  and  $Y$  be  $\text{CAT}(0)$  spaces. If there exists an  $AM$ -map  $f : X \rightarrow Y$ , then the telescopic dimension increases e.g.  $\text{tele-dim}(X) \leq \text{tele-dim}(Y)$ .*

## 2 Asymptotic-Möbius maps

### 2.1 Quasi-Möbius maps

Let  $X$  be a metric space, and  $x, y, z, w \in X$  a quadruple of distinct points. Their *cross-ratio* is

$$[x, y, z, w] = \frac{d(x, z)d(y, w)}{d(x, w)d(y, z)}.$$

If  $X$  is unbounded, then the cross-ratio extends to the one-point-compactification of  $X$  [14]. A map  $f : X \rightarrow Y$  is *quasi-Möbius* if there exists a homeomorphism  $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that for all quadruples of distinct points  $x, y, z, w \in X$ ,

$$[f(x), f(y), f(z), f(w)] \leq \eta([x, y, z, w]).$$

If  $f$  is a homeomorphism, then  $f$  is quasi-Möbius iff its inverse is quasi-Möbius.

Examples of quasi-Möbius maps are

1. the stereographic projection  $\mathbb{R}^n \rightarrow S^n$ ,
2. the Cayley transforms, the complex, quaternionic and octonionic analoga of the stereographic projection,
3. inversions  $x \mapsto \frac{x}{|x|^2}$  in Banach spaces.

### 2.2 Asymptotic-Möbius maps

Given  $u : \mathbb{R}_+ \rightarrow \mathbb{R}$ , we say that the points  $x$  and  $y$  are separated in  $X$  by  $u$  seen from a fixed origin  $o$ , if  $d(x, y) > u(d(x, o) + d(o, y))$ . For simplicity we often write  $x, y > u$  and  $|x| = d(o, x)$  for the distance to the origin.

**Definition 1.** A function  $u : \mathbb{R}_+ \rightarrow \mathbb{R}$  is admissible if

1. it is non-decreasing,
2.  $u$  grows sublinearly i.e.  $\limsup_{r \rightarrow \infty} \frac{u(r)}{r} = 0$ .

**Definition 2.** A map  $f : X \rightarrow Y$  is an asymptotic-Möbius map or AM-map, if there exists an admissible function  $u$  and a homeomorphism  $\eta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that for every  $x, y, z, w$  in  $X$  such that all 6 distances between  $x, y, z, w > u$ , then

$$[f(x), f(y), f(z), f(w)] < \eta([x, y, z, w]).$$

**Definition 3.** A metric space  $X$  is asymptotically chained, if for every origin  $o \in X$  there exists an admissible function  $v$ , such that for all  $x, y \in X$  there exists a chain  $x_1 = x, \dots, x_{k+1} = y$  that satisfies

$$\max_{i \in 1 \dots k} \{d(x_i, x_{i+1})\} < v(|x| + |y|).$$

Our main technical step will be the following theorem, whose proof, after some preliminary definition, will occupy Section 2.

**Theorem 1.** Let  $f$  be an AM-map from an asymptotically chained space  $X$  to a space  $Y$  that preserves diverging and bounded sequences, then  $f$  induces a continuous, injective, quasi-symmetric map  $g$  between some asymptotic cones of  $X$  and  $Y$ .

## 2.3 Ultralimits and asymptotic-cones

**Definition 4.** A (non-principal) ultrafilter  $\omega$  over  $\mathbb{N}$  is a set of subsets of  $\mathbb{N}$  satisfying the following conditions:

1. If  $A, B \in \omega$  then  $A \cap B \in \omega$ .
2. If  $A \in \omega, A \subset B \subset \mathbb{N}$ , then  $B \in \omega$ .
3. For every  $A \subset \mathbb{N}$  either  $A \in \omega$  or  $\mathbb{N} \setminus A \in \omega$ .
4. No finite subset of  $\mathbb{N}$  is in  $\omega$ .

Equivalently,  $\omega$  is a finitely additive measure on  $\mathbb{N}$  such that each subset has measure either 0 or 1 and every finite subset has measure 0.

If a statement  $A(n)$  holds for all  $n \in B$  with  $B$  belonging to an ultra filter  $\omega$ , we say that  $A(n)$  holds  $\omega$ -almost surely.

**Definition 5.** Let  $\omega$  be a (non-principle) ultrafilter over  $\mathbb{N}$ . An  $\omega$ -limit of a sequence of points  $\{x_n\}$  in a topological space  $X$ , is a point  $x$  in  $X$  such that for every neighborhood  $U$  of  $x$  the relation  $x_n \in U$  holds  $\omega$ -almost surely.

If  $X$  is a Hausdorff-space, then the  $\omega$ -limit of a sequence is unique. We denote this point by  $\lim_{\omega} x_n$ .

**Definition 6.** The ultrapower of a set  $X$  w.r.t. an ultrafilter  $\omega$ , denoted by  $X^{\omega}$ , consists of equivalence classes of sequences  $\{x_n\}$ ,  $x_n \in X$ , where two sequences  $\{x_n\}$  and  $\{y_n\}$  are identified if and only if  $x_n = y_n$   $\omega$ -almost surely.

We adopt the notation  $\{x_n\}_\omega$  for the equivalence class of sequences.

Let  $(X, d)$  be a metric space,  $\omega$  an ultrafilter over  $\mathbb{N}$  and  $o$  an origin in  $X$ . Furthermore, there is given a sequence of numbers  $\{d_n\}$ , with  $\lim_\omega d_n = \infty$ . Then there exists a pseudo-metric on the ultrapower  $X^\omega$  by setting the distance between  $x = \{x_n\}_\omega$  and  $y = \{y_n\}_\omega$  to

$$d_\omega(x, y) = \lim_\omega \frac{d(x_n, y_n)}{d_n}.$$

Let  $X_o^\omega$  be the set of equivalence classes of elements  $x = \{x_n\}_\omega$  in  $X^\omega$  that satisfy  $\lim_\omega \frac{d(x_n, o)}{d_n} < \infty$  and two elements  $x = \{x_n\}_\omega$  and  $y = \{y_n\}_\omega$  of  $X^\omega$  are identified if and only if  $d_\omega(x, y) = 0$ . The metric space  $(X_o^\omega, d_\omega)$  is called an *asymptotic cone* of  $(X, d)$ .

We adopt the notation  $x_\omega$  for points in the asymptotic cone  $(X_o^\omega, d_\omega)$  of  $(X, d)$ .

**Lemma 7.** *If  $x_\omega \neq y_\omega$ , then for all admissible functions  $u$  and all sequences  $\{x_n\}$ ,  $\{y_n\}$  representing  $x_\omega$  and  $y_\omega$  respectively,*

$$d(x_n, y_n) > u(|x_n| + |y_n|), \quad \omega - a.s.$$

**Proof:** If there exists an admissible function  $u$  and sequences  $\{x_n\}$ ,  $\{y_n\}$  such that

$$d(x_n, y_n) \leq u(|x_n| + |y_n|), \quad \omega - a.s.$$

then  $\omega - a.s.$ ,

$$\frac{d(x_n, y_n)}{d_n} \leq \frac{u(|x_n| + |y_n|)}{d_n}.$$

Thus

$$\lim_\omega \frac{d(x_n, y_n)}{d_n} = 0.$$

■

**Lemma 8.** *Let  $X$  be an asymptotically chained metric space. Let  $\{x_n\}$  and  $\{y_n\}$  be two sequences in  $X$  representing the same point  $x_\omega$  different from the origin  $o_\omega$ , and such that  $x_n \neq y_n$   $\omega - a.s.$  Then for every admissible function  $u$ , either*

$$d(x_n, y_n) > u, \quad \omega - a.s.$$

or there exists a sequence  $\{w_n\}$  representing the same point s.t.

$$d(x_n, w_n) > u \quad \omega - a.s.$$

and

$$d(y_n, w_n) > u \quad \omega - a.s.$$

**Proof:** Suppose  $d(x_n, y_n) \leq u$   $\omega - a.s.$  and by assumption,  $|x_n|, |y_n| \rightarrow +\infty$ .

Since  $X$  is asymptotically chained, there exists an admissible function  $v$  and a chain  $x_1^n = o, \dots, x_{k+1}^n = y_n$  such that

$$\max_{i \in \{1, \dots, k\}} \{d(x_i^n, x_{i+1}^n)\} < v(|y_n|).$$

Assume that  $n$  is large enough such that  $2u(4|y_n|) < |y_n|$ , then at least one point  $x_{i_n}^n$  in the chain satisfies

$$2u(4|y_n|) < d(y_n, x_{i_n}^n) \leq 2u(4|y_n|) + v(|y_n|).$$

Define  $w_n = x_{i_n}^n$ .

For  $n$  large enough,

$$\begin{aligned} 2d(x_n, y_n) &\leq 2u(|x_n| + |y_n|) \\ &\leq |x_n| + |y_n| \\ &\leq 2|y_n| + d(x_n, y_n). \end{aligned} \tag{1}$$

Therefore,

$$d(x_n, y_n) \leq u(4|y_n|).$$

and hence

$$u(4|y_n|) < d(x_n, w_n) \leq 3u(4|y_n|) + v(|y_n|). \tag{2}$$

Pick  $N$  s.t.  $\forall n \geq N$

$$3u(4|y_n|) + v(|y_n|) \leq 2|y_n|.$$

Then

$$\begin{aligned} u(|x_n| + |w_n|) &\leq u(|x_n| + |y_n| + d(y_n, w_n)) \\ &\leq u(2|y_n| + d(x_n, y_n) + 2u(4|y_n|) + v(|y_n|)) \\ &\leq u(2|y_n| + 3u(4|y_n|) + v(|y_n|)) \\ &\leq u(4|y_n|) \\ &< d(x_n, w_n). \end{aligned} \tag{3}$$

That means  $x_n, w_n > u$  and

$$\frac{d(x_n, w_n)}{d_n} \leq \frac{3u(4|y_n|) + v(|y_n|)}{d_n} \rightarrow 0.$$

Similarly,

$$\begin{aligned} u(|y_n| + |w_n|) &\leq u(2|y_n| + d(y_n, w_n)) \\ &\leq u(2|y_n| + 2u(4|y_n|) + v(|y_n|)) \\ &\leq u(4|y_n|) \\ &< d(y_n, w_n). \end{aligned} \tag{4}$$

■

## 2.4 Proof of Theorem 1.

**Proof:** Let  $u$  be an admissible function as given by the definition of  $f$ . Fix an origin  $o \in X$ . Let  $B_n$  be the ball of radius  $n$  around  $o$ . For every  $n$ , pick a point  $z_n$  in  $B_n$  such that

$$d(f(o), f(z_n)) \tag{5}$$

is 'maximal'. More precisely, let  $\epsilon > 0$  and pick  $z_n$  such that

$$d(f(o), f(z_n)) > \sup_{x \in B_n} d(f(o), f(x)) - \epsilon. \quad (6)$$

The supremum on the right-hand-side exists. If not we could construct a bounded sequence in  $B_n$  that is mapped by  $f$  to an unbounded sequence. This contradicts the assumptions.

Furthermore, the four point condition

$$[f(x), f(w), f(y), f(z)] \leq \eta([x, w, y, z]),$$

reduces to the three point condition

$$\frac{d(f(x), f(y))}{d(f(x), f(z))} \leq \eta\left(\frac{d(x, y)}{d(x, z)}\right),$$

by taking  $w$  to a point at  $\infty$ .

In the following we often shorten the notation  $f(x)$  to  $x'$ .

Define

$$d_n = d(o, z_n) \quad (7)$$

and

$$d'_n = d(o', z'_n). \quad (8)$$

By construction and assumption on  $f$ , the sequence  $d'_n$  diverges. Hence, also the sequence  $d_n$  must diverge by assumption on  $f$ .

(1) We show that if  $\frac{d(o, x_n)}{d_n}$  is bounded, then  $\frac{d(o', f(x_n))}{d'_n}$  is bounded as well.

If  $x_\omega = o_\omega$ , then  $d(o, x_n) \leq d_n$  for  $n$  large, therefore  $d(o', x'_n) \leq d'_n$ .

If  $x_\omega \neq z_\omega$ ,  $x_\omega \neq o_\omega$ , then by lemma 7  $\{x_n\}, \{z_n\}, \{o\}$  are separated by  $u$   $\omega$ -a.s. and the Möbius-condition gives

$$\frac{d(o', x'_n)}{d'_n} = \frac{d(o', x'_n)}{d(o', z'_n)} < \eta\left(\frac{d(o, x_n)}{d(o, z_n)}\right) = \eta\left(\frac{d(o, x_n)}{d_n}\right) \quad \omega - a.s.$$

Thus  $\frac{d(o', x'_n)}{d'_n}$  is bounded.

If  $x_\omega = z_\omega$ , by lemma 8, there exists  $\{w_n\}$  representing the same point s.t.  $\{w_n\}, \{x_n\}$  are  $u$  separated and  $\{w_n\}, \{z_n\}$  are  $u$  separated. In particular,  $\{w_n\}, \{x_n\}, \{o\}$  and  $\{w_n\}, \{z_n\}, \{o\}$  are  $u$  separated. Using the Möbius-condition again,

$$\begin{aligned} \frac{d(o', x'_n)}{d'_n} &= \frac{d(o', x'_n)}{d(o', z'_n)} \\ &= \frac{d(o', x'_n)}{d(o', w'_n)} \frac{d(o', w'_n)}{d(o', z'_n)} \\ &< \eta\left(\frac{d(o, x_n)}{d(o, w_n)}\right) \eta\left(\frac{d(o, w_n)}{d(o, z_n)}\right) \\ &= \eta\left(\frac{d(o, x_n)}{d_n} \frac{d_n}{d(o, w_n)}\right) \eta\left(\frac{d(o, w_n)}{d_n}\right) \\ &\rightarrow \eta(1)^2 \end{aligned} \quad (9)$$

Thus  $\frac{d(o', x'_n)}{d'_n}$  is bounded.

(2) The map  $\{x_n\} \rightarrow \{f(x_n)\}$  is well-defined from  $X_\omega(d_n)$  to  $Y_\omega(d'_n)$ .

If  $\{x_n\}$  and  $\{y_n\}$  are two sequences representing the same point different from the origin  $o_\omega$ , then by lemma 8 and lemma 7, either  $\{x_n\}, \{y_n\}, \{o\}$  are separated by  $u$ , or there exists a sequence  $\{w_n\}$  representing the same point s.t.  $\{x_n\}, \{w_n\}, \{o\}$  and  $\{y_n\}, \{w_n\}, \{o\}$  are separated by  $u$ .

In the first case,

$$\begin{aligned} \frac{d(x'_n, y'_n)}{d'_n} &= \frac{d(o', x'_n)}{d'_n} \frac{d(x'_n, y'_n)}{d(x'_n, o')} \\ &\leq \frac{d(o', x'_n)}{d'_n} \eta\left(\frac{d(x_n, y_n)}{d(x_n, o)}\right) \\ &= \frac{d(o', x'_n)}{d'_n} \eta\left(\frac{d(x_n, y_n)}{d_n} \frac{d_n}{d(x_n, o)}\right) \\ &\rightarrow 0 \end{aligned}$$

In the second case,

$$\begin{aligned} \frac{d(x'_n, y'_n)}{d'_n} &\leq \frac{d(x'_n, w'_n)}{d'_n} + \frac{d(w'_n, y'_n)}{d'_n} \\ &\leq \frac{d(o', x'_n)}{d'_n} \frac{d(x'_n, w'_n)}{d(x'_n, o')} + \frac{d(o', y'_n)}{d'_n} \frac{d(y'_n, w'_n)}{d(y'_n, o')} \\ &\leq \frac{d(o', x'_n)}{d'_n} \eta\left(\frac{d(x_n, w_n)}{d(x_n, o)}\right) + \frac{d(o', y'_n)}{d'_n} \eta\left(\frac{d(y_n, w_n)}{d(y_n, o)}\right) \\ &\leq \frac{d(o', x'_n)}{d'_n} \eta\left(\frac{d(x_n, w_n)}{d_n} \frac{d_n}{d(x_n, o)}\right) + \frac{d(o', y'_n)}{d'_n} \eta\left(\frac{d(y_n, w_n)}{d_n} \frac{d_n}{d(y_n, o)}\right) \\ &\rightarrow 0 \end{aligned}$$

Hence, in both cases  $x'_\omega = y'_\omega$ .

Let  $\{x_n\}$  be a sequence representing the origin  $o_\omega$ . If  $\{x_n\}$  is a bounded sequence, then by assumption on  $f$  also  $\{x'_n\}$  is bounded and therefore  $x'_\omega = o'_\omega$ .

If  $d(o, x_n)$  diverges, then  $\{x_n\}$  and  $\{o\}$  are  $u$ -separated. We may pick  $y_\omega \neq o_\omega$ , then by lemma 7,  $\{x_n\}, \{y_n\}, \{o\}$  are  $u$ -separated. Thus,

$$\begin{aligned} \frac{d(o', x'_n)}{d'_n} &= \frac{d(o', y'_n)}{d'_n} \frac{d(o', x'_n)}{d(o', y'_n)} \\ &\leq \frac{d(o', y'_n)}{d'_n} \eta\left(\frac{d(o, x_n)}{d(o, y_n)}\right) \\ &= \frac{d(o', y'_n)}{d'_n} \eta\left(\frac{d(o, x_n)}{d_n} \frac{d_n}{d(o, y_n)}\right) \\ &\rightarrow 0. \end{aligned} \tag{10}$$

Therefore  $o'_\omega = x'_\omega$ .

(3) The map  $g : \{x_n\} \rightarrow \{f(x_n)\}$  is quasi-symmetric and in particular continuous and injective.



Let  $\{w_n\}, \{x_n\}, \{y_n\}$  be sequences representing three distinct points  $w_\omega, x_\omega$  and  $y_\omega$  in  $X_\omega(d_n)$ . By lemma 7,  $\{w_n\}, \{x_n\}, \{y_n\}$  are pairwise  $u$  separated. Thus

$$\frac{d(x'_n, y'_n)}{d(x'_n, w'_n)} < \eta \left( \frac{d(x_n, y_n)}{d(x_n, w_n)} \right),$$

equivalently

$$\frac{d(x'_n, y'_n)}{d'_n} \frac{d'_n}{d(x'_n, w'_n)} < \eta \left( \frac{d(x_n, y_n)}{d_n} \frac{d_n}{d(x_n, w_n)} \right).$$

Taking the  $\omega$ -limit gives

$$\frac{d_\omega(x'_\omega, y'_\omega)}{d_\omega(x'_\omega, w'_\omega)} \leq \eta \left( \frac{d_\omega(x_\omega, y_\omega)}{d_\omega(x_\omega, w_\omega)} \right).$$

Hence, if  $g : \{x_n\} \mapsto \{f(x_n)\}$  is non-constant, then it is continuous and injective. Notice that  $\{o'\}$  and  $\{f(z_n)\}$  do not represent the same points, so  $g$  is indeed non-constant. ■

## 3 Examples

### 3.1 Sublinearly Lipschitz Equivalences

In [5] Y. Cornulier introduces *sublinearly Lipschitz maps*. A map  $f : X \rightarrow Y$  between metric spaces is a sublinearly Lipschitz map, if there exists an admissible function  $u : \mathbb{R}_+ \rightarrow \mathbb{R}$  such that

$$d(f(x), f(y)) \leq Cd(x, y) + C'u(|x| + |y|), \quad \forall x, y \in X,$$

for some constants  $C, C' > 0$ .

Two sublinearly Lipschitz maps  $f, f'$  are equivalent, if there exists an admissible function  $v$  and a constant  $C'' > 0$ , such that

$$d(f(x), f'(x)) \leq C''v(|x|)$$

for all  $x \in X$ .

Sublinearly Lipschitz maps between metric spaces form a category. Taking asymptotic cones yields a functor from the sublinearly Lipschitz category to the Lipschitz category. The sublinearly Lipschitz category is in some sense the maximal category with such a property.

The isomorphisms in the sublinearly Lipschitz category are called sublinearly Lipschitz equivalences, or SBE maps.

**Proposition 9.** *Every SBE-map  $f : X \rightarrow Y$  is an AM-map with linear  $\eta$ .*

**Proof:** If  $f$  is SBE, then  $f$  is bi-Lipschitz except at scales below some admissible function  $u$ . Indeed, let  $x, y \in X$  s.t.

$$d(x, y) \geq u(|x| + |y|).$$

Then

$$\begin{aligned} d(f(x), f(y)) &\leq cd(x, y) + Cu(|x| + |y|) \\ &\leq (c + C)d(x, y), \end{aligned} \tag{11}$$

and

$$\begin{aligned} d(f(x), f(y)) &\geq c'd(x, y) - C'u(|x| + |y|) \\ &\geq (c' - C')d(x, y) \end{aligned} \tag{12}$$

Hence, there exists  $D > 0$ , s.t. for all  $x, y > u$ ,

$$\frac{1}{D}d(x, y) \leq d(f(x), f(y)) \leq Dd(x, y).$$

In particular,  $f$  is an *AM*-map with linear  $\eta$ . ■

### 3.2 Assouad-type maps

Assouad's embedding theorem [1] [10] says that every snowflake  $X^\alpha = (X, d^\alpha)$ ,  $0 < \alpha < 1$ , of a doubling metric space admits a bi-Lipschitz embedding into some Euclidean space.

Clearly, the Assouad embedding of a doubling metric space into Euclidean space is an asymptotic-Möbius map.

In the following we construct an example of a Assouad type map from an infinite dimensional Heisenberg group into a Hilbert space. This construction follows straightforwardly from a construction by Lee and Naor in the finite dimensional case [9].

Let  $H$  be an infinite dimensional complex Hilbert space.  $H$  carries the symplectic form  $\Omega(a, b) = \text{Im}(\langle a, b \rangle)$ .

The *infinite dimensional Heisenberg group*  $\mathcal{H}_\Omega$ , is the set of tuples  $(a, t)$  with  $a \in H$ ,  $t \in \mathbb{R}$  and group law

$$(a, t)(a', t') = (a + a', t + t' + 2\Omega(a, a')).$$

Let  $G$  be a group with identity element  $e$ . A group seminorm on  $G$  is a function  $G \rightarrow [0, \infty)$  that satisfies  $N(g^{-1}) = N(g)$  for all  $g \in G$ ,  $N(gh) \leq N(g) + N(h)$  for all  $g, h \in G$  and  $N(e) = 0$ . If in addition  $N(g) = 0$  if and only if  $g = e$ , then  $N$  is a group norm on  $G$ .

The function

$$N(a, t) = \sqrt{\sqrt{\|a\|^4 + t^2} + \|a\|^2},$$

is a group norm on  $\mathcal{H}_\Omega$ . Indeed, if  $N_1$  and  $N_2$  are group seminorms, then  $\sqrt{N_1^2 + N_2^2}$  is a group seminorm. In this case  $N_1 = (\|a\|^4 + t^2)^{\frac{1}{4}}$  is the Koranyi-norm. Given  $N$ ,

$$d_N((a, t), (a', t')) = N((a, t)(a', t')^{-1})$$

is a right-invariant metric on  $\mathcal{H}_\Omega$ .

Let  $G$  be a group, a Hermitian kernel on  $G$  is a complex valued function  $K : G \times G \rightarrow \mathbb{C}$  that satisfies  $K(g, h) = \overline{K(h, g)}$  for all  $g, h \in G$ .

A Hermitian kernel on  $G$  is positive definite if

$$\sum_{i,j}^n K(g_i, g_j) c_i \overline{c_j} \geq 0$$

for all  $g_1, \dots, g_n \in G$  and for all complex numbers  $c_1, \dots, c_n \in \mathbb{C}$ , with equality if and only if the  $c_i$  vanish. A Hermitian kernel on  $G$  is conditionally negative definite if

$$\sum_{i,j}^n K(g_i, g_j) c_i \overline{c_j} \leq 0$$

for all  $g_1, \dots, g_n \in G$  and for all complex numbers  $c_1, \dots, c_n \in \mathbb{C}$  satisfying  $\sum_i^n c_i = 0$  and with equality if and only if the  $c_i$  vanish.

**Theorem 2** (Schönberg). *Let  $G$  be a group,  $K : G \times G \rightarrow \mathbb{R}$  a real valued kernel on  $G$  satisfying  $K(g, g) = 0 \forall g \in G$ . Then  $K$  is negative definite if and only if there exists a Hilbertspace  $H$  and a function  $T : G \rightarrow H$  such that  $\forall g, h \in G : K(g, h) = \|T(g) - T(h)\|^2$ .*

Thus, if  $d_N$  is negative definite, then there exists a Hilbert space  $H$  and an isometry  $T : (\mathcal{H}_\Omega, \sqrt{d_N}) \rightarrow H$ . In particular, the map  $T : (\mathcal{H}_\Omega, d_N) \rightarrow H$  is an  $AM$ -map.

A complex valued function  $F$  on a group  $G$  that satisfies  $F(g^{-1}) = \overline{F(g)}$  for all  $g \in G$  induces a Hermitian kernel on  $G$  by defining  $K_F(g, h) := F(gh^{-1})$ . The function  $F$  is said to be positive definite, if  $K_F$  is a positive definite Hermitian kernel on  $G$ .

**Lemma 10.** *For every  $\lambda \in \mathbb{R}$  the function*

$$\Phi_\lambda(a, t) = e^{-|\lambda| \|a\|^2 + i\lambda t}$$

*is positive definite on  $\mathcal{H}_\Omega$ .*

**Proof:** Indeed the function  $\Phi_\lambda$  satisfies  $\Phi_\lambda((a, t)^{-1}) = \overline{\Phi_\lambda((a, t))}$  and induces the kernel

$$K((a, s), (b, t)) := \exp(-|\lambda| \|a - b\|^2 + i\lambda(s - t - 2\Omega(a, b))).$$

The kernel can be rewritten as product of three exponentials

$$\exp(-|\lambda| (\|a\|^2 + \|b\|^2)) \exp(i\lambda(s - t)) \exp(2|\lambda| (\operatorname{Re}(\langle a, b \rangle) - i \operatorname{sign}(\lambda) \Omega(a, b))).$$

The product of positive definite kernels is positive definite. The first two factors are positive definite since for all  $c_1, \dots, c_n \in \mathbb{C}$ , the matrix  $(c_i \overline{c_j})_{i,j}$  is positive semi-definite. In particular also the matrix  $(\exp(i\lambda(s_i - t_j)) c_i \overline{c_j})_{i,j}$ . The exponential of any positive definite kernel is again a positive definite kernel ([3] Proposition 8.2.). Therefore the kernel  $K$  is positive definite if

$$\operatorname{Re}(\langle a, b \rangle) - i \operatorname{sign}(\lambda) \Omega(a, b)$$

is positive definite. This is  $\langle a, b \rangle$  if  $\lambda < 0$  and  $\overline{\langle a, b \rangle}$  if  $\lambda \geq 0$  and therefore clearly positive definite.  $\blacksquare$

**Proposition 11.**  $d_N : \mathcal{H}_\Omega \times \mathcal{H}_\Omega \rightarrow H$  is conditionally negative definite.

**Proof:** The existence of  $\frac{1}{2}$ -stable distributions implies that for all  $\epsilon > 0$ , there exists a non-negative integrable function  $\varphi_\epsilon : \mathbb{R} \rightarrow [0, \infty)$  s.t. its Fourier-transform  $\hat{\varphi}_\epsilon(t) = e^{-\epsilon\sqrt{|t|}}$ . Notice that  $\frac{1}{2\pi}\hat{\varphi}_\epsilon(x) = \varphi_\epsilon(x)$ .

By the lemma above,

$$F_\epsilon(a, t) = \int_{\mathbb{R}} e^{-|\lambda|\|a\|^2 + i\lambda t} \varphi_\epsilon(\lambda) d\lambda$$

is positive definite on  $\mathcal{H}_\Omega$ . Let

$$h_k(x) = \frac{k}{\pi} \frac{1}{k^2 + x^2}$$

denote the Cauchy-distribution with scale parameter  $k > 0$ . We rewrite

$$\begin{aligned} F_\epsilon(a, t) &= \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\lambda t} \hat{h}_{\|a\|^2}(\lambda) \hat{\varphi}_\epsilon(\lambda) d\lambda \\ &= \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\lambda t} \widehat{h_{\|a\|^2} * \varphi_\epsilon}(\lambda) d\lambda \\ &= (h_{\|a\|^2} * \hat{\varphi}_\epsilon)(t). \end{aligned} \quad (13)$$

Since  $F_\epsilon$  is positive definite,  $\frac{1-F_\epsilon}{\epsilon}$  is conditionally negative definite and

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \frac{1 - F_\epsilon(a, t)}{\epsilon} &= \lim_{\epsilon \rightarrow 0} \left[ h_{\|a\|^2} * \frac{1 - \hat{\varphi}_\epsilon}{\epsilon} \right] \\ &= \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}} \frac{1 - e^{-\epsilon\sqrt{|x|}}}{\epsilon} h_{\|a\|^2}(t-x) dx \\ &= \frac{\|a\|^2}{\pi} \int_{\mathbb{R}} \frac{\sqrt{|x|}}{\|a\|^4 + (t-x)^2} dx. \end{aligned} \quad (14)$$

We now show that for all  $r, t \in \mathbb{R}$

$$r^2 \int_{\mathbb{R}} \frac{\sqrt{|x|}}{r^4 + (t-x)^2} dx = \pi \sqrt{\sqrt{r^4 + t^2} + r^2}.$$

By change of variables  $x = r^2 y$  and  $s = t/r^2$ , the left-hand-side can be written as

$$\int_0^\infty \left( \frac{1}{1+(s-y)^2} + \frac{1}{1+(s+y)^2} \right) \sqrt{y} dy$$

This integral equals

$$\lim_{r \rightarrow 0} \lim_{R \rightarrow \infty} \frac{1}{2} \int_{C_{r,R}} \left( \frac{1}{1+(s-z)^2} + \frac{1}{1+(s+z)^2} \right) \sqrt{z} dz,$$

where  $C_{r,R}$  is the keyhole contour with a branch cut along the positive real axis. The integrand has simple poles at  $i \pm s$  and  $-i \pm s$ . With  $\text{Res}\left(\frac{\sqrt{z}}{1+(s-z)^2}, i+s\right) = \frac{\sqrt{i+s}}{2i}$ ,  $\text{Res}\left(\frac{\sqrt{z}}{1+(s-z)^2}, -i+s\right) = \frac{\sqrt{-i+s}}{-2i}$ ,  $\text{Res}\left(\frac{\sqrt{z}}{1+(s+z)^2}, i-s\right) = \frac{\sqrt{i-s}}{2i}$  and  $\text{Res}\left(\frac{\sqrt{z}}{1+(s+z)^2}, -i-s\right) = \frac{\sqrt{-i-s}}{-2i}$ . Now adding all the residues and multiplying by  $2\pi i$ , lets us evaluate the above equation as being

$$\frac{\pi}{2} (\sqrt{i+s} - \sqrt{-i+s} + \sqrt{i-s} - \sqrt{-i-s}).$$

This simplifies further to

$$\begin{aligned}
\pi \operatorname{Re}(\sqrt{i+s} + \sqrt{i-s}) &= \pi (\operatorname{Re}(\sqrt{i+s}) + \operatorname{Im}(\sqrt{i+s})) \\
&= \pi \left( \sqrt{\frac{\sqrt{1+s^2} + s}{2}} + \sqrt{\frac{\sqrt{1+s^2} - s}{2}} \right) \quad (15) \\
&= \pi \sqrt{\sqrt{1+s^2} + 1}
\end{aligned}$$

Thus we have seen, that the  $N$ -norm is a limit of conditionally negative definite functions on  $\mathcal{H}_\Omega$ , hence it is conditionally negative definite. ■

## 4 Applications to dimension theory

### 4.1 Dimension theory of finitely-generated groups

In 1993 M. Gromov introduced a notion of asymptotic dimension as a large scale analogue of Lebesgue's covering dimension [6]. The asymptotic dimension of a finitely generated group is a quasi-isometric invariant. Its most prominent application is due to Yu G., who showed that every finitely generated group with finite homotopy type and finite asymptotic dimension, satisfies the Novikov conjecture [15].

**Definition 12.** *Let  $X$  be a metric space. We say that the asymptotic dimension of  $X$  does not exceed  $n$  provided for every uniformly bounded open cover  $\mathcal{V}$  of  $X$  there is a uniformly bounded open cover  $\mathcal{U}$  of  $X$  of multiplicity  $\leq n+1$  so that  $\mathcal{V}$  refines  $\mathcal{U}$ . We write  $\operatorname{asdim} X = n$  if it is true that  $\operatorname{asdim} X \leq n$  and  $\operatorname{asdim} X \not\leq n-1$*

The asymptotic dimension of a finitely generated group  $\Gamma$  has several interesting implications. For example  $\operatorname{asdim} \Gamma = 0$  if and only if  $\Gamma$  is finite (Proposition 65. in [2]) and  $\operatorname{asdim} \Gamma = 1$  if and only if  $\Gamma$  is virtually free (Theorem 66. in [2]).

If  $G$  is a finitely generated nilpotent group, then its asymptotic cone  $G_\omega$  is a Carnot group [11] and the asymptotic dimension of  $G$  agrees with the topological dimension of its asymptotic cone [2].

**Theorem 3.** *Let  $G$  and  $G'$  be a finitely generated nilpotent groups and  $f : G \rightarrow G'$  an AM-map, then  $\operatorname{asdim}(G) \leq \operatorname{asdim}(G')$ . If  $G$  and  $G'$  are nilpotent Lie groups, then  $\dim(G) \leq \dim(G')$ . Furthermore, if  $\operatorname{asdim}(G) = \operatorname{asdim}(G')$ , then the asymptotic cones of  $G$  and  $G'$  are isomorphic graded Lie groups. Conversely, given nilpotent groups with isomorphic asymptotic cones, there exists an asymptotic-Möbius map between them.*

**Proof:** The map  $f$  induces a quasi-symmetric map  $g$  between asymptotic cones  $G_\omega$  and  $G'_\omega$  of  $G$  and  $G'$  respectively.

In particular  $g$  is an injective map from  $G_\omega$  to  $G'_\omega$  and therefore the topological dimensions increase. Since  $G$  and  $G'$  are nilpotent,  $\operatorname{asdim}(G) = \operatorname{top-dim}(G_\omega) \leq \operatorname{top-dim}(G'_\omega) = \operatorname{asdim}(G')$ .

The case for nilpotent Lie groups follows from the fact that for nilpotent Lie groups,  $\operatorname{top-dim}(G) = \operatorname{top-dim}(G_\omega)$ .

If  $\text{asdim}(G) = \text{asdim}(G')$ , the invariance of domain theorem implies that  $g$  is a homeomorphism. Since both  $g$  and  $g^{-1}$  are quasi-symmetric, they are a.e. differentiable by Pansu's theorem and the differential of  $g$  is at almost every point a graded isomorphism of groups [12]. ■

## 4.2 Dimension theory of CAT(0)-spaces

Heinrich Jung observed in early 1900, that the diameter of a set in Euclidean space  $\mathbb{R}^n$  is related to the radius of the minimum enclosing ball by an inequality, whose constant depends only on the dimension  $n$  [7]. Precisely, for any set  $A \subset \mathbb{R}^n$

$$r \leq d(A) \sqrt{\frac{n}{2(n+1)}}, \quad (16)$$

where  $r$  is the radius of the minimum enclosing ball of  $A$  and  $d(A)$  is the diameter of  $A$ .

Caprace and Lytchak observed [4], that a CAT(0)-space has *geometric dimension* less than  $n$  (as defined by B. Kleiner [8]), if and only if Jung's inequality (16) is true.

**Remark 13.** *It turns out that the geometric dimension of a CAT(0)-space  $X$  is equal to the maximal topological dimension of all compact subsets in  $X$  [8].*

The geometric dimension can be turned into a large-scale notion in the following manner. A CAT(0)-space  $X$  has telescopic dimension less than  $n$  if and only if for every  $\delta > 0$ , there exists a  $D > 0$  such that for every set  $A$  in  $X$  with diameter larger than  $D$ ,

$$r \leq \left( \delta + \sqrt{\frac{n}{2(n+1)}} \right) d(A). \quad (17)$$

Alternatively,

**Definition 14.** *A CAT(0)-space  $X$  has telescopic dimension  $\leq n$ , if every asymptotic cone has geometric dimension  $\leq n$ .*

We adopt the notation  $\text{tele-dim}(X)$  for the telescopic dimension of  $X$ .

**Theorem 4.** *Let  $X$  and  $Y$  be CAT(0) spaces. If there exists an AM-map  $f : X \rightarrow Y$ , then the telescopic dimension increases e.g.  $\text{tele-dim}(X) \leq \text{tele-dim}(Y)$ .*

The theorem follows straight forward from Theorem 1, Remark 13 and the fact that the topological dimension increases under injective continuous maps.

As a consequence, there cannot exist an AM-map from Euclidean buildings of finite rank  $r$  and thus telescopic dimension  $r$ , to a Euclidean building of lower rank.

If there exists an AM-map from a CAT(0) space to a hyperbolic metric space, then  $X$  is a hyperbolic metric space as well.

## References

- [1] ASSOUAD, P. Plongements lipschitziens dans  $\mathbb{R}^n$ . *Bulletin de la Société Mathématique de France* 111 (1983), 429 – 448.
- [2] BELL, G., AND DRANISHNIKOV, A. Asymptotic dimension. *Topology and its Applications* 155, 12 (2008), 1265–1296.
- [3] BENYAMINI, Y., AND LINDENSTRAUSS, J. *Geometric Nonlinear Functional Analysis*, vol. 1 of *American Mathematical Society colloquium publications*. American Mathematical Soc., 1998.
- [4] CAPRACE, P., AND LYTCHAK, A. At infinity of finite-dimensional  $\text{cat}(0)$  spaces. *Math. Ann.* 346 (2010), 1 – 21.
- [5] CORNULIER, Y. On sublinear bilipschitz equivalence of groups. *Annales scientifiques de l'École normale supérieure* 52 (02 2017).
- [6] GROMOV, M. *Asymptotic invariants of infinite groups*, vol. 2. Cambridge Univ. Press, 1993.
- [7] JUNG, H. Ueber die kleinste kugel, die eine räumliche figur einschliesst. *Journal für die reine und angewandte Mathematik (Crelles Journal)* 1901, 123 (1901), 241–257.
- [8] KLEINER, B. The local structure of length spaces with curvature bounded above. *Mathematische Zeitschrift* 231 (1999), 409 – 456.
- [9] LEE, J. R., AND NAOR, A. Lp metrics on the heisenberg group and the goemans-linial conjecture. In *2006 47th Annual IEEE Symposium on Foundations of Computer Science (FOCS'06)* (2006), pp. 99–108.
- [10] NAOR, A., AND NEIMAN, O. Assouad's theorem with dimension independent of the snowflaking. *Rev. Mat. Iberoam.* 28, 4 (2012), 1123 – 1142.
- [11] PANSU, P. Croissance des boules et des géodésiques fermées dans les nilvariétés. *Ergodic Theory and Dynamical Systems* 3, 3 (1983), 415–445.
- [12] PANSU, P. Métriques de carnot-caratheodory et quasiisométries des espaces symétriques de rang un. *Annals of Mathematics* 129, 1 (1989), 1–60.
- [13] PANSU, P. Large scale conformal maps. *Annales Scientifiques de l'École Normale Supérieure : publiées sous les auspices du Ministre de l'Instruction publique* 54, 4 (2021), 831 – 887.
- [14] VÄISÄLÄ, J. Quasi-möbius maps. *J. Analyse Math.* 44 (1984/85), 218 – 234.
- [15] YU, G. The novikov conjecture for groups with finite asymptotic dimension. *Annals of Mathematics* 147, 2 (1998), 325–355.

---

Orsay, September 30, 2021.

2020 *Mathematical Subject Classification*..

*Keywords and phrases*. Möbius geometry, large scale geometry.

GEORG GRÜTZNER  
UNIVERSITÉ PARIS-SACLAY, CNRS  
LABORATOIRE DE MATHÉMATIQUES D'ORSAY  
91405 ORSAY CÉDEX, FRANCE  
DATA SHAPE, CENTRE INRIA SACLAY  
91120 PALAISEAU, FRANCE  
`georg.gruetzner@universite-paris-saclay.fr`