



HAL
open science

Padé Approximation and Hypergeometric Functions: A Missing Link with the Spectrum of Delay-Differential Equations

Islam Boussaada, Guilherme Mazanti, Silviu-Iulian Niculescu

► **To cite this version:**

Islam Boussaada, Guilherme Mazanti, Silviu-Iulian Niculescu. Padé Approximation and Hypergeometric Functions: A Missing Link with the Spectrum of Delay-Differential Equations. MTNS 2022 - 25th International Symposium on Mathematical Theory of Networks and Systems, Sep 2022, Bayreuth, Germany. pp.206-211, 10.1016/j.ifacol.2022.11.053 . hal-03750884

HAL Id: hal-03750884

<https://hal.science/hal-03750884>

Submitted on 12 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Padé Approximation and Hypergeometric Functions: A Missing Link with the Spectrum of Delay-Differential Equations

Islam Boussaada ^{*,**} Guilherme Mazanti ^{*}
Silviu-Iulian Niculescu ^{*}

^{*} *Université Paris-Saclay, CNRS, CentraleSupélec, Inria, Laboratoire des Signaux et Systèmes, 91190, Gif-sur-Yvette, France (e-mails: {Islam.Boussaada, Guilherme.Mazanti, Silviu.Niculescu}@l2s.centralesupelec.fr).*

^{**} *Institut Polytechnique des Sciences Avancées (IPSA), 63 boulevard de Brandebourg, 94200 Ivry-sur-Seine, France*

Abstract: It is well known that rational approximation theory involves degenerate hypergeometric functions and, in particular, the Padé approximation of the exponential function is closely related to Kummer hypergeometric functions. Recently, in the context of the study of the exponential stability of the trivial solution of delay-differential equations, a new link between the degenerate hypergeometric function and the zeros distribution of the characteristic function associated with linear delay-differential equations was emphasized. Such a link allowed the characterization of a property of time-delay systems known as *multiplicity-induced-dominancy (MID)*, which opened a new direction in designing low-complexity controllers for time-delay systems by using a partial pole placement idea. Thanks to their relations to hypergeometric functions, we explore in this paper links between the spectrum of delay-differential equations and Padé approximations of the exponential function. This note exploits and further comments recent results from [I. Boussaada, G. Mazanti and S-I. Niculescu. 2022, Comptes Rendus. Mathématique] and [I. Boussaada, G. Mazanti and S-I. Niculescu. 2022, Bulletin des Sciences Mathématiques].

Keywords: Padé approximation, delay-differential equations, exponential stability, Kummer hypergeometric functions.

1. INTRODUCTION

On beyond of the arithmetic theory and the number theory, the theory of continued fractions is strongly related to the theory of analytic functions; it is involved in topics such as the study of definite integrals, power series and the summation of divergent series, see for instance Perron (1977); Wall (2018). Continuous fractions theory goes back to the pioneering work of T. S. Stieltjes mainly elaborated at the end of the 19th century. It appears that several links exist between continuous fractions theory with concepts and methods such as the well known Padé rational approximation and the ϵ -algorithm. As a matter of fact, the ϵ -algorithm is a transformation of the partial sums of a series into the corresponding Padé quotients or the corresponding continued fractions. In fact, the ϵ -algorithm is an old but still relevant way to improve the convergence rate of slowly convergent sequences, see, for instance, Graves-Morris et al. (2000) and references therein. Moreover, a strong connection between such an algorithm and the Padé approximation has been underlined by P. Wynn in the early 60s, Wynn (1966). Through his results on continuous fractions theory, Wynn's work covered several problems in analytic functions theory, such as the distribution of zeros of a class of special functions called the Kummer (de-

generate) hypergeometric functions or, equivalently, the *Wittaker hypergeometric functions*, see, e.g., Wynn (1973).

More contemporary contributions have pointed out to a missing link existing between the zeros of such a class of special functions with the spectrum distribution of an appropriate Delay-Differential Equation (DDE) including one pointwise delay, see for instance Mazanti et al. (2021a); Boussaada et al. (2022a). Interestingly, this link allowed deriving an exhaustive characterization of a property for DDE¹ called *Generic Multiplicity Induced Dominancy (GMID)*, see Boussaada et al. (2022b). Indeed, the GMID is a special case of a more general property called *Multiplicity Induced Dominancy (MID)*. Such a property consists of determining the conditions under which a given multiple complex zero of a quasipolynomial is dominant. It should be noted that the GMID property asserts that a DDE spectral value admitting the maximal multiplicity is necessarily the corresponding spectral abscissa. However, multiple roots with intermediate admissible multiplicities may or may not be dominant. Thanks to this property, a consistent control strategy is proposed in Boussaada et al. (2020); Balogh et al. (2022), which consists in assigning a root with an admissible multiplicity once appropriate

¹ in both retarded as well neutral cases

conditions guaranteeing its dominance have been established. Furthermore, the MID property may be used to tune standard controllers. For instance, in Ma et al. (2022) it is applied to the systematic tuning of the stabilizing PID controller of a first-order plant. Here, we aim at assigning dominant multiple real roots with admissible codimensions.

Beyond the reminder of the concepts and links discussed, the contribution of the present note is threefold. First, to highlight the existing links between roots of generic maximal multiplicity for a quasipolynomial, Kummer confluent geometric functions, and Padé approximations of the exponential function. Second, the use of Padé approximation allows simplifying some newly obtained analytic functions results, thus opening some new perspectives in a *partial pole placement* methodology for linear time-invariant DDE. Finally, it allows to slightly correct some of the results derived in Saff and Varga (1978) where the links between Kummer functions and Padé approximations are underlined.

The remaining of the paper is organized as follows: Section 2 recalls the definitions and properties of some concepts studied in the paper. Next, Section 3 is dedicated to the main result. Section 4 presents further comments on the distribution of the Kummer and the Whittaker zeros distribution. Some concluding remarks end the paper.

Notations: Throughout the paper, the following notations are used: \mathbb{N}^* denotes the set of positive integers and $\mathbb{N} = \mathbb{N}^* \cup \{0\}$. The set of all integers is denoted by \mathbb{Z} and, for $a, b \in \mathbb{R}$, we denote $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{Z}$, with the convention that $\llbracket a, b \rrbracket = \emptyset$ if $a > b$. For a complex number s , $\Re(s)$ and $\Im(s)$ denote its real and imaginary parts, respectively. The open left and right complex half-planes are the sets \mathbb{C}_- and \mathbb{C}_+ , respectively, defined by $\mathbb{C}_- = \{s \in \mathbb{C} \mid \Re(s) < 0\}$ and $\mathbb{C}_+ = \{s \in \mathbb{C} \mid \Re(s) > 0\}$. Given $k, n \in \mathbb{N}$ with $k \leq n$, the binomial coefficient $\binom{n}{k}$ is defined as $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ and this notation is extended to $k, n \in \mathbb{Z}$ by setting $\binom{n}{k} = 0$ when $n < 0$, $k < 0$, or $k > n$. For $\alpha \in \mathbb{C}$ and $k \in \mathbb{N}$, $(\alpha)_k$ is the *Pochhammer symbol* for the *ascending factorial*, defined inductively as $(\alpha)_0 = 1$ and $(\alpha)_{k+1} = (\alpha + k)(\alpha)_k$.

2. PREREQUISITES

This section provides a brief presentation of the definitions and results that shall be of use in the sequel concerning Padé rational approximation and degenerate hypergeometric functions.

2.1 On the Padé rational approximation

The Padé rational approximation can be seen as a generalization of the Taylor expansion of a function in which one wishes to approximate some function f around a given point (which is typically normalized to 0) by a rational function, i.e., by a function R expressed as $R = \frac{Q}{P}$, where Q and P are polynomials. We start this subsection by a precise definition of what is meant by Padé approximation in this paper.

Definition 2.1. Let $f : U \rightarrow \mathbb{C}$ be an analytic function defined on a neighborhood $U \subset \mathbb{C}$ of 0. Given $(m, n) \in \mathbb{N}^2$,

we say that a pair of polynomials (Q, P) with $\deg Q \leq m$ and $\deg P \leq n$ is a *Padé approximation of f of order (m, n)* if

$$P(z)f(z) - Q(z) = \mathcal{O}(z^{n+m+1}) \quad \text{as } z \rightarrow 0.$$

Classical results on Padé approximations can be found, for instance, in Baker and Graves-Morris (1996) and (Brezinski, 2002, Chapter 4), and we also refer to Brezinski (1991) for a historical presentation of this topic. For every $(m, n) \in \mathbb{N}^2$, the Padé approximation (Q, P) of an analytic function f defined on a neighborhood of 0 exists and is unique up to a multiplicative constant. If $n = 0$, the corresponding Padé approximation $(Q, 1)$ is such that Q coincides with the Taylor–Maclaurin expansion of f of order m .

Padé approximations are widely used in rational approximation theory (see, for instance, Szegő (1924); Dieudonné (1935); Saff and Varga (1978); Baratchart et al. (1995) and the references therein). In particular, several applications involve Padé approximations of the exponential function $z \mapsto e^z$ and some of the above works highlighted links between such approximations and *degenerate hypergeometric functions*. As reported in (Perron, 1977, p246), the Padé approximation of order (m, n) of the exponential function is the pair $(Q_{n,m}, P_{n,m})$ defined for $z \in \mathbb{C}$ by

$$\begin{cases} P_{n,m}(z) = \sum_{k=0}^n \frac{(n+m-k)!n!z^k}{k!m!(n-k)!}, \\ Q_{n,m}(z) = \sum_{k=0}^m \frac{(n+m-k)!(-z)^k}{k!(m-k)!}, \end{cases} \quad (1)$$

and the rational approximation $R_{n,m}(z) = \frac{Q_{n,m}(z)}{P_{n,m}(z)}$ satisfies

$$R_{n,m}(z) \xrightarrow[n, m \rightarrow \infty]{} e^z,$$

uniformly on compact subsets of \mathbb{C} . Further, as can be found in Perron (1977), the Padé remainder $e^z - R_{n,m}(z)$ satisfies

$$P_{n,m}(z)(e^z - R_{n,m}(z)) = \frac{z^{n+m+1}}{m!} \int_0^1 e^{tz}(1-t)^m t^n dt, \quad (2)$$

which, as detailed in Section 2.2 below, is closely related to Kummer hypergeometric functions (see for instance Saff and Varga (1978) and the references therein). We also highlight the fact that, from a control theory point of view, Padé approximations of the exponential function have been exploited in approximating delay systems via finite-dimensional systems (see, for instance, Glader et al. (1991) for a deeper discussion on the approximation of the parametric family of first-order stable delay systems).

Remark 2.2. The expressions of $P_{n,m}$ and $Q_{n,m}$ provided above use a different normalization with respect to the expressions in Perron (1977): we have chosen here the normalization consisting in requiring the denominator $P_{n,m}$ to be monic, i.e., the coefficient of the term z^n is 1.

2.2 Degenerate hypergeometric functions

We present in this section the definition and main properties of Kummer and Whittaker hypergeometric functions. The presentation in this section follows that of Boussaada

et al. (2022a), and we start by providing the definition and main properties of Kummer hypergeometric functions (for further details, see, for instance, Buchholz (1969); Erdélyi et al. (1981); Olver et al. (2010) and references therein). For an historical perspective, the reader is referred to Kampé de Fériet (1937).

Definition 2.3. Let $a, b \in \mathbb{C}$ and assume that b is not a nonpositive integer. The *Kummer confluent hypergeometric function* $\Phi(a, b, \cdot) : \mathbb{C} \rightarrow \mathbb{C}$ is the entire function defined for $z \in \mathbb{C}$ by the series

$$\Phi(a, b, z) = \sum_{k=0}^{\infty} \frac{(a)_k}{(b)_k} \frac{z^k}{k!}. \quad (3)$$

The series in (3) converges for every $z \in \mathbb{C}$, and the function $\Phi(a, b, \cdot)$ satisfies the *Kummer differential equation*

$$z \frac{\partial^2 \Phi}{\partial z^2}(a, b, z) + (b - z) \frac{\partial \Phi}{\partial z}(a, b, z) - a \Phi(a, b, z) = 0. \quad (4)$$

The equation (4) admits two linearly independent solutions, which sometimes are both called Kummer confluent hypergeometric functions. In the present paper, we are concerned only with the solution given by (3).

Kummer functions admit the integral representation

$$\Phi(a, b, z) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 e^{zt} t^{a-1} (1-t)^{b-a-1} dt, \quad (5)$$

for every $a, b, z \in \mathbb{C}$ such that $\Re(b) > \Re(a) > 0$, where Γ denotes the Gamma function. For all complex numbers a, b, z such that b is not a nonpositive integer, Kummer functions also satisfy the relation

$$\Phi(a, b, z) = e^z \Phi(b - a, b, -z), \quad (6)$$

and we have in addition

$$\Phi(a, b, z) = 1 + \mathcal{O}(z) \quad \text{as } z \rightarrow 0. \quad (7)$$

Kummer confluent hypergeometric functions have close links with Whittaker functions, defined as follows (see, e.g., Olver et al. (2010)).

Definition 2.4. Let $k, l \in \mathbb{C}$ and assume that $2l$ is not a negative integer. The *Whittaker function* $\mathcal{M}_{k,l}$ is the function defined for $z \in \mathbb{C}$ by

$$\mathcal{M}_{k,l}(z) = e^{-\frac{z}{2}} z^{\frac{1}{2}+l} \Phi\left(\frac{1}{2} + l - k, 1 + 2l, z\right). \quad (8)$$

If $\frac{1}{2} + l$ is not an integer, the function $\mathcal{M}_{k,l}$ is a multi-valued complex function with branch point at $z = 0$. Whenever $2l$ is not a negative integer, the nontrivial roots of $\mathcal{M}_{k,l}$ coincide with those of $\Phi(\frac{1}{2} + l - k, 1 + 2l, \cdot)$ and $\mathcal{M}_{k,l}$ satisfies the *Whittaker differential equation*

$$\varphi''(z) = \left(\frac{1}{4} - \frac{k}{z} + \frac{l^2 - \frac{1}{4}}{z^2} \right) \varphi(z). \quad (9)$$

Similarly to the Kummer differential equation (4), other solutions of the Whittaker differential equation (9) are also known as Whittaker functions in other works, but they will not be used in this paper. Notice also that, since $\mathcal{M}_{k,l}$ is a nontrivial solution of the second-order linear differential equation (9), any nontrivial root of $\mathcal{M}_{k,l}$ is necessarily simple. The roots of Whittaker functions satisfy the following immediate symmetry property.

Proposition 2.5. Let $k, l \in \mathbb{C}$ and assume that $2l$ is not a negative integer. If $z \in \mathbb{C} \setminus \{0\}$ is a nontrivial root of $\mathcal{M}_{k,l}$, then $-z$ is a root of $\mathcal{M}_{-k,l}$.

The following result from Boussaada et al. (2022a), whose proof relies on the Green–Hille transform of (9) (see, e.g., Hille (1922)), provides information on the location of nontrivial zeros of Whittaker functions.

Proposition 2.6. Let $k, l \in \mathbb{R}$ be such that $2l - 1 \geq 0$.

- (a) If $k = 0$, then all nontrivial roots z of $\mathcal{M}_{k,l}$ are purely imaginary.
- (b) If $k > 0$, then all nontrivial roots z of $\mathcal{M}_{k,l}$ satisfy $\Re(z) > 0$.
- (c) If $k < 0$, then all nontrivial roots z of $\mathcal{M}_{k,l}$ satisfy $\Re(z) < 0$.
- (d) If $k \neq 0$, then all nontrivial roots z of $\mathcal{M}_{k,l}$ satisfy

$$4k^2 \Im(z)^2 - (4(l^2 - k^2) - 1) \Re(z)^2 > 0. \quad (10)$$

Moreover, in all cases, all non-real roots z of $\mathcal{M}_{k,l}$ satisfy $|z| > \sqrt{4l^2 - 1}$.

As an immediate consequence of (8) and Proposition 2.6, we have the following result on the location of zeros of Kummer functions with real parameters, which was stated in Boussaada et al. (2022a).

Corollary 2.7. Let $a, b \in \mathbb{R}$ be such that $b \geq 2$.

- (a) If $b = 2a$, then all nontrivial roots z of $\Phi(a, b, \cdot)$ are purely imaginary.
- (b) If $b > 2a$, then all nontrivial roots z of $\Phi(a, b, \cdot)$ satisfy $\Re(z) > 0$.
- (c) If $b < 2a$, then all nontrivial roots z of $\Phi(a, b, \cdot)$ satisfy $\Re(z) < 0$.
- (d) If $b \neq 2a$, then all nontrivial roots z of $\Phi(a, b, \cdot)$ satisfy $(b - 2a)^2 \Im(z)^2 - (4a(b - a) - 2b) \Re(z)^2 > 0$.

Moreover, in all cases, all non-real roots z of $\Phi(a, b, \cdot)$ satisfy $|z| > \sqrt{b(b - 2)}$.

2.3 Delay-differential equations and their characteristic equations

Consider the linear time-invariant delay-differential equation

$$y^{(n)}(t) + \sum_{k=0}^{n-1} a_k y^{(k)}(t) + \sum_{k=0}^m \alpha_k y^{(k)}(t - \tau) = 0, \quad (11)$$

where $y(\cdot)$ is the real-valued unknown function, $\tau > 0$ is the delay, and $a_0, \dots, a_{n-1}, \alpha_0, \dots, \alpha_m$ are real coefficients. Such an equation is said to be of *retarded type* if $m < n$ (i.e., if the highest order of derivation appears only in the non-delayed term $y^{(n)}(t)$), or of *neutral type* if $m = n$. We refer to Hale and Verduyn Lunel (1993); Michiels and Niculescu (2014) for more information on delay-differential equations and for proofs of the major facts stated in the sequel.

Characterizing the stability of the zero solution of (11) in terms of the coefficients $a_0, \dots, a_{n-1}, \alpha_0, \dots, \alpha_m$ is in general a challenging question. Exponential stability can be characterized through the roots of the *characteristic function* of (11), which is the entire function $\Delta : \mathbb{C} \rightarrow \mathbb{C}$ defined for $s \in \mathbb{C}$ by

$$\Delta(s) = s^n + \sum_{k=0}^{n-1} a_k s^k + e^{-s\tau} \sum_{k=0}^m \alpha_k s^k. \quad (12)$$

More precisely, the zero solution of (11) is exponentially stable if and only if there exists $\gamma < 0$ such that, for every root s of Δ , we have $\Re(s) \leq \gamma$. Rightmost roots of Δ thus determine the asymptotic behavior of the system and its rate of exponential stability.

It should be mentioned that the function Δ from (12) is a particular case of a *quasipolynomial*. Quasipolynomials have been extensively studied in the literature (see, e.g., Bellman and Cooke (1963), Hale and Verduyn Lunel (1993)) and, in particular, a result presented in (Pólya and Szegő, 1998, Part Three, Problem 206.2) implies that any root s of Δ has multiplicity at most $\mathcal{D}_{PS} = n + m + 1$. The number \mathcal{D}_{PS} is known as the *degree* of Δ , and such a bound on the multiplicity of a root of Δ is known as *Pólya–Szegő bound*.

3. MAIN RESULTS

The main result of this paper is the following, which highlights the links between roots of maximal multiplicities of Δ , Kummer confluent hypergeometric functions, and Padé approximations of the exponential function. More precisely, our main result characterizes the case where Δ admits a root of maximal multiplicity \mathcal{D}_{PS} through a suitable Kummer hypergeometric function by using the results on the Padé approximations of the exponential function recalled in Section 2.1.

Theorem 3.1. Consider the quasipolynomial Δ given by (12) and let $s_0 \in \mathbb{R}$. Let P and Q be the polynomials defined by

$$P(s) = s^n + \sum_{k=0}^{n-1} a_k s^k, \quad Q(s) = \sum_{k=0}^m \alpha_k s^k$$

and define the quasipolynomial $\tilde{\Delta}$ and the polynomials \tilde{P} and \tilde{Q} by setting, for $z \in \mathbb{C}$,

$$\tilde{\Delta}(z) = \tau^n \Delta\left(s_0 + \frac{z}{\tau}\right), \quad (13)$$

$$\tilde{P}(z) = \tau^n P\left(s_0 + \frac{z}{\tau}\right), \quad (14)$$

$$\tilde{Q}(z) = e^{-s_0 \tau} \tau^n Q\left(s_0 + \frac{z}{\tau}\right) \quad (15)$$

Then the following assertions are equivalent.

- (a) The real number s_0 is a root of multiplicity $\mathcal{D}_{PS} = n + m + 1$ of Δ .
- (b) The pair of polynomials $(-\tilde{Q}, \tilde{P})$ forms a Padé approximation of order (m, n) of the exponential function $z \mapsto e^z$.
- (c) The quasipolynomial $\tilde{\Delta}$ is given by

$$\tilde{\Delta}(z) = \frac{n! z^{n+m+1}}{(n+m+1)!} \Phi(m+1, n+m+2, -z). \quad (16)$$

- (d) The coefficients $a_0, \dots, a_{n-1}, \alpha_0, \dots, \alpha_m$ of Δ are given by

$$\begin{cases} a_k = (-1)^{n-k} n! \sum_{j=k}^n \frac{\binom{j}{k} \binom{m+n-j}{m} s_0^{j-k}}{j! \tau^{n-j}} & \text{for } k \in \llbracket 0, n-1 \rrbracket, \\ \alpha_k = (-1)^{n-1} e^{s_0 \tau} \sum_{j=k}^m \frac{(-1)^{j-k} (m+n-j)! s_0^{j-k}}{k! (j-k)! (m-j)! \tau^{n-j}} & \text{for } k \in \llbracket 0, m \rrbracket. \end{cases} \quad (17)$$

In addition, if the above equivalent assertions are satisfied and $m \leq n$, then s_0 is the rightmost root of Δ , i.e., we have $\Re(s) \leq \Re(s_0)$ for every $s \in \mathbb{C}$ such that $\Delta(s) = 0$.

Proof. Clearly, thanks to the definition of $\tilde{\Delta}$, assertion (a) is equivalent to requiring 0 to be a root of multiplicity \mathcal{D}_{PS} of $\tilde{\Delta}$. Since $\tilde{\Delta}$ is an analytic function and \mathcal{D}_{PS} is the maximal possible multiplicity of any of its roots, the latter fact is equivalent to requiring that $\tilde{\Delta}(z) = \mathcal{O}(z^{n+m+1})$ as $z \rightarrow 0$. Using that $\tilde{\Delta}(z) = \tilde{P}(z) + e^{-z} \tilde{Q}(z)$, the latter is equivalent to

$$\tilde{P}(z)e^z + \tilde{Q}(z) = \mathcal{O}(z^{n+m+1}) \quad \text{as } z \rightarrow 0,$$

which, thanks to the definition of Padé approximation, is equivalent to (b).

Using the uniqueness of the Padé approximation of order (m, n) with a monic denominator, we deduce that (b) is equivalent to requiring that $-\tilde{Q} = Q_{n,m}$ and $\tilde{P} = P_{n,m}$, where $P_{n,m}$ and $Q_{n,m}$ are defined in (1), i.e., it is equivalent to requiring that, for every $z \in \mathbb{C}$,

$$\tilde{\Delta}(z) = P_{n,m}(z) - e^{-z} Q_{n,m}(z).$$

In particular, equivalence with (d) follows after straightforward but long computations by using (13) (see, e.g., (Mazanti et al., 2021a, Lemma 4.2) for detailed computations in the case $m = n-1$, which can be easily generalized to any m).

Finally, if (b) is satisfied, then, combining (2), (5), and (6), we get that

$$\begin{aligned} e^z \tilde{\Delta}(z) &= P_{n,m}(z)(e^z - R_{n,m}(z)) \\ &= \frac{n! z^{n+m+1}}{(n+m+1)!} \Phi(n+1, n+m+2, z) \\ &= e^z \frac{n! z^{n+m+1}}{(n+m+1)!} \Phi(m+1, n+m+2, -z), \end{aligned}$$

yielding (c). Conversely, if (c) holds, then, using (7) and (16), we deduce that

$$\tilde{P}(z)e^z + \tilde{Q}(z) = e^z \tilde{\Delta}(z) = \mathcal{O}(z^{n+m+1}),$$

yielding (b) thanks to the definition of Padé approximation.

Remark 3.2. The equivalence between items (a), (c), and (d) of Theorem 3.1 was already stated and proved in Boussaada et al. (2022b) by exploiting the integral representation (5). By exploiting the Padé approximation of the exponential function and in particular (2), Theorem 3.1 provides the additional equivalence with (b) and allows for a much simpler proof of the equivalences between (a)–(d).

Note that, by considering the first equation in (17) with $k = n-1$, one obtains the simple and interesting relation between s_0 , τ , and a_{n-1} given by

$$s_0 = -\frac{a_{n-1}}{n} - \frac{m+1}{\tau}. \quad (18)$$

As remarked in Boussaada et al. (2022b), if any of the equivalent assertions of Theorem 3.1 is satisfied, then s_0 is the unique real root of Δ , and, more precisely, it is the unique root of Δ on the horizontal strip $\{s \in \mathbb{C} \mid |\Im(s)| < \frac{2\pi}{\tau}\}$ of the complex plane. In addition, the assumption in Theorem 3.1 of requiring s_0 to be real is justified by the fact that nonreal roots of Δ cannot have a multiplicity

equal to the Pólya–Szegő bound \mathcal{D}_{PS} , since any root s_0 of Δ attaining the maximal multiplicity \mathcal{D}_{PS} necessarily satisfies (18), and thus it will be real since a_{n-1} is real. As an immediate consequence of Theorem 3.1, if (17) is satisfied for some $s_0 \in \mathbb{R}$, then the trivial solution of (11) is exponentially stable if and only if $a_{n-1} > -\frac{n(m+1)}{\tau}$.

Note that Theorem 3.1 provides necessary and sufficient conditions for a real number s_0 to be a root of maximal multiplicity of the quasipolynomial Δ from (12). The main result of this section states that, under those conditions, s_0 is necessarily a dominant root of Δ .

4. ADDITIONAL COMMENTS ON THE LOCATION OF ROOTS OF WHITTAKER FUNCTIONS

Proposition 2.6 is actually a corrected version of a result by G. E. Tsvetkov, (Tsvetkov, 1941, Theorem 7). In Tsvetkov (1941), the author provides statements of results concerning the location of zeros of Whittaker functions without the corresponding proofs, pointing only to the technique by E. Hille from Hille (1922). It turns out, as highlighted in (Boussaada et al., 2022a, Counterexample 3.1), that there are counterexamples to the statement of (Tsvetkov, 1941, Theorem 7). Proposition 2.6, proved in Boussaada et al. (2022a), uses the technique suggested by G. E. Tsvetkov in his paper, based on the Green–Hille transform, to provide a corrected version of (Tsvetkov, 1941, Theorem 7).

Some works in the literature have relied on (Tsvetkov, 1941, Theorem 7) to provide additional properties on the location of roots of Whittaker functions. This is the case, for instance, of (Saff and Varga, 1978, Proposition 3.2), which studies Whittaker functions motivated by the previously described links with Padé approximations of the exponential function. More precisely, the statement of (Saff and Varga, 1978, Proposition 3.2) is the following.

Proposition 4.1. (Saff and Varga (1978)). Let $k \in \mathbb{R}$, $l > 0$, and z be a nontrivial zero of $\mathcal{M}_{k,l}$.

- (a) If $k > 0$, then $\Re(z) > 2k$ and $\Im(z)^2 > \Re(z)^2 \frac{4l^2 - 4k^2 - 1}{4k^2}$.
- (b) If $k < 0$, then $\Re(z) < 2k$ and $\Im(z)^2 > \Re(z)^2 \frac{4l^2 - 4k^2 - 1}{4k^2}$.
- (c) If $k = 0$, then $\Re(z) = 0$ and $\Im(z)^2 > 4l^2 - 1$.

It turns out that the counterexample to (Tsvetkov, 1941, Theorem 7) from (Boussaada et al., 2022a, Counterexample 3.1) also provides a counterexample to items (a) and (b) of Proposition 4.1. For the sake of completeness, let us detail the counterexample.

Let $l > 0$ and take $k = l + \frac{3}{2}$. If z is a nontrivial root of $\mathcal{M}_{k,l}$, then, by (8), z is a nontrivial root of $\Phi(-1, 1+2l, \cdot)$. From (3), we have $\Phi(-1, 1+2l, z) = 1 - \frac{z}{1+2l}$, and its unique root is $z = 1+2l$. In particular, $\Re(z) = 1+2l < 2k = 3+2l$, and thus Proposition 4.1(a) is not verified. In addition, as a consequence of Proposition 2.5, $z = -1 - 2l$ is the unique nontrivial root of $\mathcal{M}_{-k,l}$ and, since $\Re(z) = -1 - 2l > -2k = -3 - 2l$, Proposition 4.1(b) is not verified.

In view of Proposition 4.1 and Proposition 2.6, a natural question is whether one may provide, in the case $k > 0$, a lower bound on the real part of nontrivial roots of $\mathcal{M}_{k,l}$ as a function of k . A numerical exploration of such a question was provided in Boussaada et al. (2022a) by considering the root $z = 1 + 2l$ of $\mathcal{M}_{k,l}$ for $k = l + \frac{3}{2}$. Note that

such a root exists whenever $l > -\frac{1}{2}$ and it is real and simple. Hence, for every $l > -\frac{1}{2}$, there exists an interval I_l containing $1 + 2l$ and a curve $k \mapsto z_l(k) \in \mathbb{R}$ defined on I_l such that, for every $k \in I_l$, $z_l(k)$ is a real root of $\mathcal{M}_{k,l}$ with $z_l(l + \frac{3}{2}) = 1 + 2l$. The article Boussaada et al. (2022a) provides the numerical computation of these curves for $l \in \{-\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$, which we reproduce in Figure 1. The black dots (k, z) correspond to $k = l + \frac{3}{2}$ and the root $z = 1 + 2l$.




Fig. 1. Real root $z(k)$ of $\mathcal{M}_{k,l}$ satisfying $z(l + \frac{3}{2}) = 1 + 2l$ for six different values of l .

An inspection of Figure 1 leads to the conjecture that the maximal interval I_l on which z_l is defined is $I_l = (l + \frac{1}{2}, +\infty)$ and that $z_l(k) \rightarrow 0$ as $k \rightarrow +\infty$ and $z_l(k) \rightarrow +\infty$ as $k \rightarrow l + \frac{1}{2}$. In particular, if this conjecture is true, then one cannot expect to correct Proposition 4.1(a) by replacing the term $2k$ by any function of k which remains lower bounded as $k \rightarrow +\infty$.

Finally, we notice the remarkable case depicted in Boussaada et al. (2022b), where one is able to characterize the zeros of the Whittaker function $\mathcal{M}_{0, n+\frac{1}{2}}$ for $n \in \mathbb{N}$ as $\{i\zeta \mid \zeta \in \Xi_n\}$, where Ξ_n is the set of $\zeta \in \mathbb{R}$ satisfying

$$\tan\left(\frac{\zeta}{2}\right) = \frac{\zeta \sum_{\ell=0}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^\ell \frac{(2n-2\ell-1)!}{(2\ell+1)!(n-2\ell-1)!} \zeta^{2\ell}}{\sum_{\ell=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^\ell \frac{(2n-2\ell)!}{(2\ell)!(n-2\ell)!} \zeta^{2\ell}}.$$

The particular case $n = 1$, $\Xi_1 = \left\{ \zeta \in \mathbb{R} \mid \tan\left(\frac{\zeta}{2}\right) = \frac{\zeta}{2} \right\}$, had already been identified in Mazanti et al. (2021b).

5. CONCLUDING REMARKS

This paper discusses the existing links between roots of generic maximal multiplicity for a quasipolynomial, Kum-

mer confluent geometric functions and Padé approximations of the exponential function. More precisely, we have shown that in the case of a real characteristic root having the maximum multiplicity of a simple quasipolynomial, the corresponding polynomials define a pair of Padé approximations. Moreover, the multiplicity of the root is equal to the Pólya–Szegő bound and it is dominant in the sense that it explicitly defines the spectral abscissa of the spectrum of the corresponding delay-differential equations. As a byproduct of the analysis, we have slightly corrected some of the results from the literature on the Padé approximation.

REFERENCES

- Baker, Jr., G.A. and Graves-Morris, P. (1996). *Padé approximants*, volume 59 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition.
- Balogh, T., Boussaada, I., Insperger, T., and Niculescu, S.I. (2022). Conditions for stabilizability of time-delay systems with real-rooted plant. *Internat. J. Robust Nonlinear Control*, 32(6), 3206–3224.
- Baratchart, L., Saff, E.B., and Wielonsky, F. (1995). Rational interpolation of the exponential function. *Canadian Journal of Mathematics*, 47(6), 1121–1147.
- Bellman, R. and Cooke, K.L. (1963). *Differential-difference equations*. Academic Press, New York.
- Boussaada, I., Mazanti, G., and Niculescu, S.I. (2022a). Some remarks on the location of non-asymptotic zeros of Whittaker and Kummer hypergeometric functions. *Bulletin des Sciences Mathématiques*, 174, 103093.
- Boussaada, I., Mazanti, G., and Niculescu, S.I. (2022b). The generic multiplicity-induced-dominancy property from retarded to neutral delay-differential equations: when delay-systems characteristics meet the zeros of Kummer functions. *C. R. Math. Acad. Sci. Paris*, 360, 349–369.
- Boussaada, I., Niculescu, S.I., El-Ati, A., Pérez-Ramos, R., and Trabelsi, K. (2020). Multiplicity-induced-dominancy in parametric second-order delay differential equations: analysis and application in control design. *ESAIM Control Optim. Calc. Var.*, 26, Paper No. 57, 34.
- Brezinski, C. (1991). *History of continued fractions and Padé approximants*, volume 12 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin.
- Brezinski, C. (2002). *Computational aspects of linear control*, volume 1 of *Numerical Methods and Algorithms*. Kluwer Academic Publishers, Dordrecht.
- Buchholz, H. (1969). *The confluent hypergeometric function with special emphasis on its applications*. Translated from the German by H. Lichtblau and K. Wetzel. Springer Tracts in Natural Philosophy, Vol. 15. Springer-Verlag New York Inc., New York.
- Dieudonné, J. (1935). Sur les zéros des polynômes-sections de e^x . *Bull. Sci. Math*, 70, 333–351.
- Erdélyi, A., Magnus, W., Oberhettinger, F., and Tricomi, F.G. (1981). *Higher transcendental functions. Vol. I*. Robert E. Krieger Publishing Co., Inc., Melbourne, Fla.
- Glader, C., Högnäs, G., Mäkilä, P.M., and Toivonen, H.T. (1991). Approximation of delay systems—a case study. *Internat. J. Control*, 53(2), 369–390.
- Graves-Morris, P., Roberts, D., and Salam, A. (2000). The epsilon algorithm and related topics. *Journal of Computational and Applied Mathematics*, 122(1-2), 51–80.
- Hale, J.K. and Verduyn Lunel, S.M. (1993). *Introduction to functional differential equations*, volume 99 of *Applied Mathematical Sciences*. Springer-Verlag, New York.
- Hille, E. (1922). Oscillation theorems in the complex domain. *Transactions of the American Mathematical Society*, 23(4), 350–385.
- Kampé de Fériet, J. (1937). La fonction hypergéométrique (in French). *Mémorial Sciences Math.*, Fasc. 85, 1–85.
- Ma, D., Boussaada, I., Chen, J., Bonnet, C., Niculescu, S.I., and Chen, J. (2022). PID control design for first-order delay systems via MID pole placement: Performance vs. robustness. *Automatica*, 137, 110102.
- Mazanti, G., Boussaada, I., and Niculescu, S.I. (2021a). Multiplicity-induced-dominancy for delay-differential equations of retarded type. *Journal of Differential Equations*, 286, 84–118.
- Mazanti, G., Boussaada, I., Niculescu, S.I., and Chitour, Y. (2021b). Effects of roots of maximal multiplicity on the stability of some classes of delay differential-algebraic systems: The lossless propagation case. In *Proceeding of 24th International Symposium on Mathematical Theory of Networks and Systems (MTNS 2021)*, IFAC-PapersOnLine. Cambridge, United Kingdom.
- Michiels, W. and Niculescu, S.I. (2014). *Stability, control, and computation for time-delay systems: An eigenvalue-based approach*, volume 27 of *Advances in Design and Control*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, second edition.
- Olver, F.W.J., Lozier, D.W., Boisvert, R.F., and Clark, C.W. (eds.) (2010). *NIST handbook of mathematical functions*. U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge.
- Perron, O. (1977). *Die Lehre von den Kettenbrüchen*. Springer.
- Pólya, G. and Szegő, G. (1998). *Problems and theorems in analysis. I*. Classics in Mathematics. Springer-Verlag, Berlin. Series, integral calculus, theory of functions, Translated from the German by Dorothee Aeppli, Reprint of the 1978 English translation.
- Saff, E.B. and Varga, R.S. (1978). On the zeros and poles of Padé approximants to e^z . III. *Numer. Math.*, 30(3), 241–266.
- Szegő, G. (1924). Über eine Eigenschaft der Exponentialreihe. *Sitzungsber. Berl. Math. Ges*, 23, 50–64.
- Tsvetkov, G.E. (1941). On the roots of Whittaker functions. *Dokl. Akad. Nauk SSSR*, 32(1), 10–12.
- Wall, H.S. (2018). *Analytic theory of continued fractions*. Courier Dover Publications.
- Wynn, P. (1966). On the convergence and stability of the epsilon algorithm. *SIAM Journal on Numerical Analysis*, 3(1), 91–122.
- Wynn, P. (1973). On the zeros of certain confluent hypergeometric functions. *Proc. Amer. Math. Soc.*, 40, 173–182.