



UNIVERSITÉ
DE MONTPELLIER



Transit Time index (TTi) as an adaptation of humification index to illustrate transit time differences in karst hydrosystems. Application to the karst springs of Fontaine de Vaucluse system (Southeastern France)


Leïla Serène¹, Naomi Mazzilli², Christelle Batiot-Guilhe¹, Christophe Emblanch², Milanka Babic², Julien Dupont², Roland Simler², Matthieu Blanc, Gérard Massonnat³

¹ HSM, Univ. Montpellier, CNRS, IMT, IRD, Montpellier, France

²UMR 1114 EMMAH (AU-INRAE), Université d'Avignon, 84000 Avignon, France

³TotalEnergies, CSTJF, Avenue Larribau, CEDEX 64018 Pau, France


Transit time issues in karsts



Fast flows in karst

- Complex and fast recharge through hierarchized network
 - Quantity and quality issues
- Essential to trace short transit time (< 6 months)

Transit time issues in karsts




Fast flows in karst

- Complex and fast recharge through hierarchized network
 - Quantity and quality issues
- Essential to trace short transit time (< 6 months)

Natural tracers


- Few available on the 0-6 months range
- High potential of natural organic matter (life duration < 6 months, *Batiot, 2002*)
- Use of Humification Index (HIX) by *Blondel et al. (2012)*

Transit time issues in karsts



Objective : Analyze the potential of fluorescent organic compounds as natural tracer of short transit time (<6 months)


Study site & sampling



Fontaine de Vaucluse karstic system

- Catchment of 1160 km² *Ollivier, 2020*
- Mean flow rate 23,3 m³/s *Cognard-Plancaq et al., 2006*
- Aquifer thickness up to 1500m
- Unsaturated zone average thickness 800m

Study site & sampling



Fontaine de Vaucluse karstic system


- Catchment of 1160 km² *Ollivier, 2020*
- Mean flow rate 23,3 m³/s *Cognard-Plancaq et al., 2006*
- Aquifer thickness up to 1500m
- Unsaturated zone average thickness 800m

Sampling test

- One year bi-monthly sampling (06-2020 to 10-2021)
- Major and trace elements, TOC, water stable isotopes, fluorescence of organic matter
- 4 springs, 1 losses, flows from unsaturated zone (LSBB)

Humification index (HIX) in brief

$HIX = H/L$ $HIX = H/(L+H)$	Light organic matter L	Heavy organic matter H
Organic matter type	Protein-like	Humic-like
Digestibility	high	low




Serène et al., 2022


Humification index (HIX) in brief

$HIX = H/L$ $HIX = H/(L+H)$	Light organic matter L	Heavy organic matter H
Organic matter type	Protein-like	Humic-like
Digestibility	high	low


Fluorescence measurement



Hypothetic soil




Biological activity



Time

Hypothetic soil biodegraded





Humification index (HIX) in brief

$HIX = H/L$ $HIX = H/(L+H)$	Light organic matter L	Heavy organic matter H
Organic matter type	Protein-like	Humic-like
Digestibility	high	low

Fluorescence measurement

H and L quantification

From HIX to transit time




Serène et al., 2022



Questioning the use of HIX in karst waters

Previous works

H and L windows calibrated
on soil water *Zsolnay (1999) and
Ohno (2002)*



Application to
groundwater
samples




Serène et al., 2022



Questioning the use of HIX in karst waters

Previous works

H and L windows calibrated on soil water Zsolnay (1999) and Ohno (2002)



Application to groundwater samples




Application results

- Mismatch of H and L windows with fluorescent peaks of groundwater samples
- Zsolnay (1999) and Ohno (2002) HIX formulas not applicable to groundwaters
- Blondel (2008) relation between HIX and transit time need to be rebuild

Serène et al., 2022

From the Humification Index to the Transit Time index




Identification of heavy & light compounds

- 3D matrix
- PARAFAC modeling

Serène et al., 2022

From the Humification Index to the Transit Time index



Humification Index (HIX)

- 2D spectra
- H and L windows

Transit Time index (TTi)


- 3D matrix
- PARAFAC modelling

$$\text{TTi} = \frac{\text{Humic-like}}{\text{Humic and protein like}} = \frac{H}{H+L}$$

High values → « long » transit time
 Low values → short transit time


Serène et al., 2022

Comparison of TTI between different springs



Serène et al., 2022


Comparison of TTI between different springs



- Increase of TTI with decrease of karstification degree
- Consistency of mean TTI with expected transit time

Serène et al., 2022

Comparison of TTI between different springs



- Increase of TTI with decrease of karstification degree
- Consistency of mean TTI with expected transit time
- FV TTI is similar to S.Nesque while it is the outlet of a wider system
- Limits of TTI usage in the case of mix with old waters (>> 6months)

Serène et al., 2022

Discussion, conclusions & perspectives

Results

Fluorescence of organic matter as a natural tracer of transit time ?

- Methodological biases in previous work with HIX (*Blondel, 2008*)
- Adaptation of HIX to groundwater usage thanks to PARAFAC modelling → TTi
- Consistency of mean TTi with expected transit time of studied springs

Serène et al., 2022

Discussion, conclusions & perspectives

Results

Fluorescence of organic matter as a natural tracer of transit time ?

- Methodological biases in previous work with HIX (*Blondel, 2008*)
- Adaptation of HIX to groundwater usage thanks to PARAFAC modelling → TTI
- Consistency of mean TTI with expected transit time of studied springs

Improving qualitative approach

Seasonal variations

One site scale

Little impact
(*Musadji et al., 2020*)

Several sites scale

Depending on hydro-pedoclimatic context

Anthropic activities

Fluorescent compounds selection thanks to PARAFAC modelling

Serène et al., 2022

Discussion, conclusions & perspectives

Results

Fluorescence of organic matter as a natural tracer of transit time ?

- Methodological biases in previous work with HIX (*Blondel, 2008*)
- Adaptation of HIX to groundwater usage thanks to PARAFAC modelling → TTi
- Consistency of mean TTi with expected transit time of studied springs

Improving qualitative approach

Seasonal variations

One site scale

Little impact
(*Musadji et al., 2020*)

Several sites scale

Depending on hydro-
pedoclimatic context

Anthropic activities

Fluorescent compounds selection thanks to
PARAFAC modelling

Developing quantitative approach

Biodigestibility in natural groundwaters? Artificial tracer test? Radiogenic isotopes?

Serène et al., 2022

Transit Time index (TTi) as an adaptation of humification index to illustrate transit time differences in karst hydrosystems. Application to the karst springs of Fontaine de Vaucluse system (Southeastern France)

Leïla Serène¹, Naomi Mazzilli², Christelle Batiot-Guilhe¹, Christophe Emblanch², Milanka Babic², Julien Dupont², Roland Simler², Matthieu Blanc, Gérard Massonnat³

Paper recently accepted for the journal HESS: <https://doi.org/10.5194/hess-2022-100>

Transit Time index (TTi)

- 3D matrix
- PARAFAC modelling

$$TTi = \frac{\text{Humic - like}}{\text{Humic and protein like}}$$

