



HAL
open science

Non-Alcoholic Steatohepatitis (NASH) and Liver Fibrosis: Molecular and Multicellular Control of Evolving Diseased States

Jérôme Eeckhoute

► **To cite this version:**

Jérôme Eeckhoute. Non-Alcoholic Steatohepatitis (NASH) and Liver Fibrosis: Molecular and Multicellular Control of Evolving Diseased States. *Cells*, 2022, 11 (16), pp.2551. 10.3390/cells11162551 . hal-03752605

HAL Id: hal-03752605

<https://hal.science/hal-03752605>

Submitted on 17 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Editorial

Non-Alcoholic Steatohepatitis (NASH) and Liver Fibrosis: Molecular and Multicellular Control of Evolving Diseased States

Jérôme Eeckhoutte 

U1011-EGID, Institut Pasteur de Lille, CHU Lille, Inserm, Univ. Lille, F-59000 Lille, France; jerome.eeckhoutte@inserm.fr

Non-alcoholic fatty liver disease (NAFLD), the most common chronic liver disease, has emerged as a major threat to public health [1]. Indeed, obesity and type 2 diabetes are independent risk factors of NAFLD for which increases contribute to the devastating consequences of the metabolic disease pandemic [2]. Importantly, NAFLD presents with a range of conditions initiated by relatively benign hepatic steatosis, which can evolve to non-alcoholic steatohepatitis (NASH); fibrosis; and ultimately, cirrhosis [3]. Advanced stages promote hepatocellular carcinoma and/or liver failure [1]. In this context, NASH has been the fastest growing indication for liver transplantation in Western countries and is expected to become the leading cause of liver transplant in the United States [4,5].

NASH is characterized by an escalation in the disease complexity, where suffering hepatocytes trigger inflammation and, secondarily, liver fibrosis [1–3]. These processes are increasingly recognized as being driven by dynamic and coordinated changes in the complexity and activities of the various liver cell populations [6]. Indeed, in addition to hepatocytes and immune cells, resident or recruited, the role exerted by cholangiocytes, (myo)fibroblasts, and endothelial cells is also now being analyzed in detail (e.g., [7–9]). In this context, our understanding of the molecular and multicellular mechanisms underlying the various NAFLD stages is still limited and defining how/why the disease evolves to more advanced stages has remained challenging. For instance, the kinetic, spatial organization, and functional hierarchy of changes in cellular states and activities largely remain to be described together with the identity and integration of intercellular signals involved. At the individual cell-type level, molecular (mal)adaptations accompanying the disease progression, including transcriptomic dysregulation, also need to be further characterized. Indeed, diagnosis and therapeutic treatment of NASH and fibrosis remain controversial and of limited efficacy [10–12] and would greatly benefit from a greater understanding of the molecular mechanisms at play in the multicellular processes defining advanced NAFLD stages. The purpose of this Special Issue is to highlight recent findings in those areas, enlightening how NASH and its associated liver fibrosis develop through coordinated modulation of the activities of different liver cell-types.

Funding: Work at Inserm U1011 is supported by the Agence Nationale de la Recherche (ANR) grants “HSCreg” (ANR-21-CE14-0032), “PreciNASH” (ANR-16-RHUS-0006-PreciNASH), and “European Genomic Institute for Diabetes” E.G.I.D (ANR-10-LABX-0046), a French State fund managed by ANR under the frame program Investissements d’Avenir I-SITE ULNE/ANR-16-IDEX-0004 ULNE and by Fondation Recherche Médicale (FRM grants EQU202203014645 and EQU202203014650).

Conflicts of Interest: The author declares no conflict of interest.



Citation: Eeckhoutte, J. Non-Alcoholic Steatohepatitis (NASH) and Liver Fibrosis: Molecular and Multicellular Control of Evolving Diseased States. *Cells* **2022**, *11*, 2551. <https://doi.org/10.3390/cells11162551>

Received: 10 August 2022

Accepted: 15 August 2022

Published: 17 August 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

References

1. Loomba, R.; Friedman, S.L.; Shulman, G.I. Mechanisms and disease consequences of nonalcoholic fatty liver disease. *Cell* **2021**, *184*, 2537–2564. [[CrossRef](#)] [[PubMed](#)]
2. Haas, J.T.; Francque, S.; Staels, B. Pathophysiology and Mechanisms of Nonalcoholic Fatty Liver Disease. *Annu. Rev. Physiol.* **2016**, *78*, 181–205. [[CrossRef](#)] [[PubMed](#)]
3. Schwabe, R.F.; Tabas, I.; Pajvani, U.B. Mechanisms of Fibrosis Development in NASH. *Gastroenterology* **2020**, *158*, 1913–1928. [[CrossRef](#)] [[PubMed](#)]
4. Wong, R.J.; Aguilar, M.; Cheung, R.; Perumpail, R.B.; Harrison, S.A.; Younossi, Z.M.; Ahmed, A. Nonalcoholic steatohepatitis is the second leading etiology of liver disease among adults awaiting liver transplantation in the United States. *Gastroenterology* **2015**, *148*, 547–555. [[CrossRef](#)] [[PubMed](#)]
5. Younossi, Z.M.; Stepanova, M.; Ong, J.; Trimble, G.; AlQahtani, S.; Younossi, I.; Ahmed, A.; Racila, A.; Henry, L. Nonalcoholic Steatohepatitis Is the Most Rapidly Increasing Indication for Liver Transplantation in the United States. *Clin. Gastroenterol. Hepatol.* **2021**, *19*, 580–589. [[CrossRef](#)] [[PubMed](#)]
6. Wallace, S.J.; Tacke, F.; Schwabe, R.F.; Henderson, N.C. Understanding the cellular interactome of non-alcoholic fatty liver disease. *JHEP Rep.* **2022**, *4*, 100524. [[CrossRef](#)] [[PubMed](#)]
7. Williams, M.; Bonnardel, J.; Haest, B.; Vanderborght, B.; Wagner, C.; Remmerie, A.; Bujko, A.; Martens, L.; Thone, T.; Browaeys, R.; et al. Spatial proteogenomics reveals distinct and evolutionarily conserved hepatic macrophage niches. *Cell* **2022**, *185*, 379–396.e338. [[CrossRef](#)] [[PubMed](#)]
8. Terkelsen, M.K.; Bendixen, S.M.; Hansen, D.; Scott, E.A.H.; Moeller, A.F.; Nielsen, R.; Mandrup, S.; Schlosser, A.; Andersen, T.L.; Sorensen, G.L.; et al. Transcriptional dynamics of hepatic sinusoid-associated cells after liver injury. *Hepatology* **2020**, *72*, 2119–2133. [[CrossRef](#)] [[PubMed](#)]
9. Haas, J.T.; Vonghia, L.; Mogilenko, D.A.; Verrijken, A.; Molendi-Coste, O.; Fleury, S.; Deprince, A.; Nikitin, A.; Woittrain, E.; Ducrocq-Geoffroy, L.; et al. Transcriptional Network Analysis Implicates Altered Hepatic Immune Function in NASH development and resolution. *Nat. Metab.* **2019**, *1*, 604–614. [[CrossRef](#)] [[PubMed](#)]
10. Rinella, M.E.; Loomba, R.; Caldwell, S.H.; Kowdley, K.; Charlton, M.; Tetri, B.; Harrison, S.A. Controversies in the Diagnosis and Management of NAFLD and NASH. *Gastroenterol. Hepatol.* **2014**, *10*, 219–227.
11. Leslie, M. The liver’s weighty problem. *Science* **2015**, *349*, 18–20. [[CrossRef](#)]
12. Friedman, S.L.; Pinzani, M. Hepatic Fibrosis 2022: Unmet Needs and a Blueprint for the Future. *Hepatology* **2022**, *75*, 473–488. [[CrossRef](#)] [[PubMed](#)]