



HAL
open science

Solutions of the LPD equation and multi-parametric rogue waves

Pierre Gaillard

► **To cite this version:**

Pierre Gaillard. Solutions of the LPD equation and multi-parametric rogue waves. 2022. hal-03755079

HAL Id: hal-03755079

<https://hal.science/hal-03755079>

Preprint submitted on 21 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Solutions of the LPD equation and multi-parametric rogue waves

Pierre Gaillard,

Université de Bourgogne Franche Comté,
Institut de mathématiques de Bourgogne,
9 avenue Alain Savary BP 47870
21078 Dijon Cedex, France :
E-mail : Pierre.Gaillard@u-bourgogne.fr

Abstract

Quasi-rational solutions to the Lakshmanan Porsezian Daniel equation are presented. We construct explicit expressions of these solutions for the first orders depending on real parameters.

We study the patterns of these configurations in the (x, t) plane in function of the different parameters.

We observe in the case of order 2, three rogue waves which move according to the two parameters.

In the case of order 3, six rogue waves are observed with specific configurations moving according to the four parameters.

Key Words : Lakshmanan Porsezian Daniel equation, quasi-rational solutions.

PACS numbers :

33Q55, 37K10, 47.10A-, 47.35.Fg, 47.54.Bd

1 Introduction

We consider the Lakshmanan Porsezian Daniel (LPD) equation in the following normalization

$$iu_t + u_{xxxx} + 8|u|^2 u_{xx} + 2u^2 \bar{u}_{xx} + 6u_x^2 \bar{u} + 4u|u_x|^2 + 6|u|^4 u = 0, \quad (1)$$

where the subscripts mean the partial derivatives.

There are a different models that describe the dynamics of soliton propagation. One of the most important models is the nonlinear Lakshmanan Porsezian Daniel (LPD) equation. This equation with spatio-temporal dispersion as well as group velocity dispersion, is used to get and describe solitons. This model was introduced originally in the context of Heisenberg spin chain [1]. Solitons

are some of the main focal points in the field of mathematical physics, optical fibers and nonlinear optics in particular. These solutions are self-reinforcing single waves, which move at a constant velocity while maintaining its shape; they are relevant because of their localized and stable nature. The LPD equation has been studied in the polarization-preserving fibers context or in the context of the birefringent fibers [2]; it has been used in the study of the propagation of periodic ultrashort pulses in the optical fibers [3]. Many mathematical methods have used to study this equation; we can mention the following ones. For instance, the method of undetermined coefficients is applied in [4] to look for bright, dark and singular soliton solutions. The extended trial equation method has been used to construct bright solitons, dark solitons, periodic solitary wave, rational and elliptic solutions in [5]. The semi inverse variational principle has been employed in [6] to construct bright solitons with Kerr and power laws of nonlinearity. One can also quote the extended Jacobi elliptic function approach [7] to obtain dark and singular optical solitons; the $\exp(-\phi(\epsilon))$ -expansion method in [8]; the Lie symmetry analysis in [9]; the Riccati equation method has been applied to get dark and bright soliton solutions of (1) in [10]; the sine-Gordon expansion method has been used to get analytical solutions of the LPD equation in [11]; the modified extended direct algebraic method [12] and expansion method [13].

2 Quasi-rational solutions of order 1 to the Lakshmanan Porsezian Daniel equation

Theorem 2.1 *The function $v(x, t)$ defined by*

$$v(x, t) = \left(1 - 4 \frac{1 + 24it}{1 + 4x^2 + 576t^2}\right) e^{6it} \quad (2)$$

is a solution to the Lakshmanan Porsezian Daniel equation (1)

(3)




Figure 1. Solution of order 1 to the equation (1).

3 Quasi-rational solutions of order 2 to the Lakshmanan Porsezian Daniel equation

Theorem 3.1 *The function $v(x, t)$ defined by*

$$v(x, t) = \left(1 - 12 \frac{n(x, t)}{d(x, t)}\right) \exp(6it + 2a_1) \quad (4)$$

with

$$n(x, t) = (2x - 12b_1)^4 + 6((4a_1 + 24t)^2 + 1)(2x - 12b_1)^2 - 192b_1(2x - 12b_1) + 5(4a_1 + 24t)^4 + 18(4a_1 + 24t)^2 + 384t(4a_1 + 24t) - 3 + i((4a_1 + 24t)(2x - 12b_1)^4 + 2((4a_1 + 24t)^3 - 12a_1 - 168t)(2x - 12b_1)^2 - 192(4a_1 + 24t)b_1(2x - 12b_1) + (4a_1 + 24t)^5 + 2(4a_1 + 24t)^3 + 192t(4a_1 + 24t)^2 - 60a_1 - 552t)$$

and

$$d(x, t) = ((2x - 12b_1)^2 + (4a_1 + 24t)^2 + 1)^3 + 192b_1(2x - 12b_1)^3 - 24((4a_1 + 24t)^2 + 48t(4a_1 + 24t) - 1)(2x - 12b_1)^2 - 576((4a_1 + 24t)^2 + 1)b_1(2x - 12b_1) + 24(4a_1 + 24t)^4 + 384t(4a_1 + 24t)^3 + 96(4a_1 + 24t)^2 + 3456t(4a_1 + 24t) + 36864t^2 + 9216b_1^2 + 8$$

is a solution to the Lakshmanan Porsezian Daniel equation (1).




Figure 2. Solution of order 1 to the equation (1); to the left $a_1 = 0$, $b_1 = 0$; in the center $a_1 = 1$, $b_1 = 0$; to the right $a_1 = 10$, $b_1 = 0$.

A very fast evolution of the structure of the solution is observed when the parameter a_1 grows; it evolves toward three peaks.




Figure 3. Solution of order 1 to the equation (1); to the left $a_1 = 0$, $b_1 = 0.1$; in the center $a_1 = 0$, $b_1 = 1$; to the right $a_1 = 0$, $b_1 = 10$.

As in the case of the parameter a_1 , an evolution of the structure of the solution toward three peaks is observed when the parameter b_1 grows.

4 Quasi-rational solutions of order 3 to the Lakshmanan Porsezian Daniel equation

Theorem 4.1 *The function $v(x, t)$ defined by*

$$v(x, t) = \left(1 - 24 \frac{n(x, t)}{d(x, t)}\right) e^{6i\alpha t} \quad (5)$$

with

$$\begin{aligned} n(x, t) = & 1024x^{10} + 3840(1 + 576t^2)x^8 + 64(210 - 218880t^2 + 16588800t^4)x^6 + \\ & 675 + 16(-1584000t^2 - 315187200t^4 + 13377208320t^6 - 450)x^4 + 4(5424537600t^4 + \\ & 80263249920t^6 + 4953389137920t^8 - 1555200t^2(5 - 3072t^2) - 675 - 19353600t^2)x^2 + \\ & 1977915801600t^6 + 107323431321600t^8 + 697437190619136t^{10} + 149299200t^4(-17 + \\ & 3072t^2) + 2388787200t^4 + 259200t(16t - 16384t^3) + 388800t^2(-3 + 4096t^2) + \\ & 23500800t^2 + i(24576tx^{10} + 256(-840t + 69120t^3)x^8 + 64(240t - 4147200t^3 + \\ & 79626240t^5)x^6 + 16(-22809600t^3 - 3264675840t^5 + 45864714240t^7 + 10800t(-3 - \\ & 1024t^2) + 14400t)x^4 + 4(51836682240t^5 - 275188285440t^7 + 13209037701120t^9 - \\ & 12441600t^3(7 - 3072t^2) + 16200t(7 + 4096t^2) + 345600t + 88473600t^3)x^2 + \\ & 716636160t^5(-107 + 3072t^2) + 3110400t^2(-176t - 16384t^3) - 3110400t^3(11 + \\ & 20480t^2) - 16200t(-7 + 22528t^2) - 265420800t^3 - 138549657600t^5 - 2155641569280t^7 + \\ & 224553640919040t^9 + 1521681143169024t^{11} + 151200t) \end{aligned}$$

and

$$\begin{aligned} d(x, t) = & 2024 + 256(-207360t^2 + 120)x^8 + 64(3778560t^2 - 291962880t^4 + \\ & 2320)x^6 + 16(-5255331840t^4 - 91729428480t^6 + 138240t^2(56 - 11520t^2) - \\ & 26265600t^2 + 3360)x^4 + 4(5870683422720t^6 + 26418075402240t^8 - 79626240t^4(-326 - \\ & 34560t^2) + 276480t^2(-76 + 311040t^2) + 57330892800t^4 - 1555200t(96t - 16384t^3) + \end{aligned}$$

$12144 + 8294400 t^2)x^2 + 15288238080 t^6(191 + 6912 t^2) + 298598400 t^3(368 t - 16384 t^3) + 79626240 t^4(599 - 57600 t^2) - 388800 t(-496 t - 65536 t^3) + 13824 t^2(3881 + 7257600 t^2) + (1 + 4 x^2 + 576 t^2)^6 + 223948800 t^2 + 33973862400 t^4 + 4127824281600 t^6 + 937841676779520 t^8 + 12680676193075200 t^{10}$
is a solution to the (LPD) equation (1).

We can give also the solutions to the Lakshmanan Porsezian Daniel equation depending on 4 real parameters. But, because of the length of the expression, we only give it in the appendix.

We give patterns of the modulus of the solutions in function of the parameters a_1, a_2, b_1, b_2 .




Figure 4. Solution of order 3 to the equation (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 0$; in the center $a_1 = 1, b_1 = 0, a_2 = 0, b_2 = 0$; to the right $a_1 = 10, b_1 = 0, a_2 = 10, b_2 = 0$; with $\alpha = \beta = 1$.




Figure 5. Solution of order 3 to the equation (1); to the left $a_1 = 0, b_1 = 0.1, a_2 = 0, b_2 = 0$; in the center $a_1 = 0, b_1 = 1, a_2 = 0, b_2 = 0$; to the right $a_1 = 0, b_1 = 5, a_2 = 0, b_2 = 0$; with $\alpha = \beta = 1$.




Figure 6. Solution of order 3 to the equation (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0.5, b_2 = 0$; in the center $a_1 = 0, b_1 = 0, a_2 = 1, b_2 = 0$; to the right $a_1 = 2, b_1 = 0, a_2 = 5, b_2 = 0$; with $\alpha = \beta = 1$.




Figure 7. Solution of order 3 to the equation (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 0.1$; in the center $a_1 = 0, b_1 = 10, a_2 = 0, b_2 = 1$; to the right $a_1 = 10, b_1 = 0, a_2 = 0, b_2 = 5$; with $\alpha = \beta = 1$.

In the evolution of the structure of the solutions according to the different parameters, we note the appearance of triangles with six peaks with more at least speed according to the parameters.

5 Conclusion

Quasi-rational solutions to the Lakshmanan Porsezian Daniel equation have been given for the first orders depending on several real parameters.

In the case of order 2, the evolution of the structure of the solutions toward three peaks is observed when parameters a_1 or b_1 grow.

In the same way, in the case of order 3, the evolution of the structure of the solutions toward six peaks is observed when parameters a_1, a_2, a_1, a_2 grow.

These solutions have to be compared with these constructed by the author in

the case of the NLS or mKdV equations [19, 21, 25, 59, 65].

We can cite some recent works about this equation. In [11], new travelling wave solutions to the (LPD) equation with Kerr nonlinearity are built using Bäcklund transformation method based on Riccati equation, Kudryashov method and a new auxiliary ordinary differential equation (ODE). In [69], bright, dark, singular, kink and periodic optical solitons solutions of the LPD equation are constructed by improved $\tan \psi(\eta)/2$ -method. In [70], three images of nonlinearity to the fractional LPD equation in birefringent fibers are investigated. In particular, the new bright, periodic wave and singular optical soliton solutions are constructed via the $m + G'/G$ expansion method. In [71], new analytical solutions to the LPD equation is presented by Jacobi elliptic functions. It would be relevant to continue the study of this article to determine other configurations in particular, those concerning no longer triangles but rings as in the case of the NLS equation.

References

- [1] M. Lakshmanan, K. Porsezian, M. Daniel, Effect of discreteness on the continuum limit of the Heisenberg spin chain, *Phys. Lett. A*, V. 133, N. 9, 483488, 1998
- [2] Y. Yyldyrym, E. Topkara, A. Biswas, H. Triki, M. Ekici, P. Guggilla, S. Khan, M.R. Belic, Cubicquartic optical soliton perturbation with LakshmananPorsezianDaniel model by sine-Gordon equation approach, *J. Opt.*, V. 50, N. 2, 322329, 2021
- [3] Y.L. Ye, D.D. Hou, C. Cheng, S.H. Chen, Rogue wave solutions of the vector LakshmananPorsezianDaniel equation, *Phys. Lett. A*, V. 384, N. 11, 126226, 2020
- [4] J. Vega-Guzman, R.T. Alqahtani, Q. Zhou, M.F. Mahmood, S.P. Moshokoa, M.Z. Ullah, A. Biswas, M. Belic, Optical solitons for Lakshmanan Porsezian Daniel model with spatiotemporal dispersion using the method of undetermined coefficients, *Optik*, V. 144, 115-123, 2017
- [5] J. Manafian, M. Foroutan, A. Guzali, Applications of the ETEM for obtaining optical soliton solutions for the Lakshmanan-Porsezian-Daniel model, *The European Physical Journal Plus*, V. 132, N. 11, 1-22, 2017
- [6] R.T. Alqahtani, M.M. Babatin, A. Biswas, Bright optical solitons for Lakshmanan- Porsezian-Daniel model by semi-inverse variational principle, *Optik*, V. 154, 109-114, 2018
- [7] A. Biswas, M. Ekici, A. Sonmezoglu, H. Triki, F.B. Majida, Q. Zhouf, S.P. Moshokoac, M. Mirzazadeh, M. Belic, Optical solitons with LakshmananPorsezianDaniel model using a couple of integration schemes, *Optik*, V. 158, 705711, 2018

- [8] A. Biswas, Y. Yyldyrym, E. Yasar, Q. Zhou, S.P. Moshokoac, M. Belic, Optical solitons for LakshmananPorsezianDaniel model by modified simple equation method, *Optik*, V. 160, 2432, 2018
- [9] A. Bansala, A. Biswasb, H. Triki, Q. Zhou, S.P. Moshokoad, M. Belic, Optical solitons and group invariant solutions to Lakshmanan Porsezian Daniel model in optical fibers and PCF, *Optik*, V. 160, 8691, 2018
- [10] A.A. AlQarni et al., Optical solitons for Lakshmanan Porsezian Daniel model by Riccati equation approach, *Optik*, V. 182, 922-929, 2019
- [11] H. Rezaazadeh et al., Optical solitons of Lakshmanan Porsezian Daniel model with a couple of nonlinearities, *Optik*, V. 164, 414-423, 2018
- [12] M.B. Hubert, et al., Optical solitons with Lakshmanan Porsezian Daniel model by modified extended direct algebraic method, *Optik*, V. 162, 228-236, 2018
- [13] S. Arshed et al., Optical solitons in birefringent fibers for Lakshmanan Porsezian Daniel model using $\exp(-i\phi)$ -expansion method, *Optik*, V. 172, 651-656, 2018
- [14] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. **161**, 2002
- [15] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. **68**, 77-90, 2004
- [16] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, 2009
- [17] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 404409-1-16, 2009
- [18] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. **185**, 247-258, 2010
- [19] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. **44**, 435204-1-15, 2011
- [20] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. **13**, N. 2, 71-153, 2012
- [21] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. **54**, 013504-1-32, 2013

- [22] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, 2013
- [23] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, 2013
- [24] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, 2013
- [25] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, 2013
- [26] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013
- [27] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013
- [28] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [29] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [30] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014
- [31] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [32] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [33] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [34] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [35] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, 2015

- [36] P. Gaillard, Other $2N-2$ parameters solutions to the NLS equation and $2N+1$ highest amplitude of the modulus of the N -th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [37] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, N. **5**, 346-364, 2015
- [38] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, 2016
- [39] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. **10**,1-7, 2016
- [40] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys*, V. **65**, N. **2**, 136-144, 2016
- [41] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **367**, 1-5, 2016
- [42] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. **10**, 83-89, 2016
- [43] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. **189**, N. **1**, 1440-1449, 2016
- [44] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [45] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. **3**, 130-139, 2017
- [46] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N2$ parameters, *Int. Jour. of Appl. Sci. And Math.*, V. **4**, N. **3**, 60-70, 2017
- [47] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, *New Hor. in Math. Phys.*, V. **1**, N. **1**, 26-31, 2017
- [48] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. **2**, 92-98, 2017

- [49] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 9, 19-37, 2017
- [50] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 11, 24-30, 2017
- [51] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, *Int. Jour. of Inn. In Sci. and Math.*, V. **6**, N. 1, 14-19, 2018
- [52] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. **196**, N. 2, 1174-1199, 2018
- [53] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. **2018**, 1-18, 2018
- [54] P. Gaillard, Families of Solutions of Order 5 to the Johnson Equation Depending on 8 Parameters, *NHIMP*, V. **2**, N. 4, 53-61, 2018
- [55] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. **1141**, 012102-1-10, 2018
- [56] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. , 109-112, 2019
- [57] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. **2**, N. 4, 1-4, 2019
- [58] P. Gaillard, Multi-parametric families of solutions of order N to the Boussinesq and KP equations and the degenerate rational case, *UJMA*, V. **3**, N. 2, 44-52, 2020
- [59] P. Gaillard, The mKdV equation and multi-parameters rational solutions, *Wave Motion*, V. **100**, 102667-1-9, 2021
- [60] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, *Jour. Of Geom. And Phys.*, V. **161**, 104059-1-12, 2021
- [61] P. Gaillard, $2N$ parameter solutions to the Burgers equation, *Jour. Of Appl. Nonlin. Dyn.*, V. **1**, N. **1**, 1-6, 2021
- [62] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. **3**, N. 3, 14-21, 2021

- [63] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. **11**, N. 4, 2102-2113, 2021
- [64] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. **17**, N. 6, 27-34, 2021
- [65] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. **107**, 102824-1-11, 2021
- [66] P. Gaillard, Multiparametric Rational Solutions of Order N to the KPI Equation and the Explicit Case of Order 3, *Arch. Of cur. Res. Int.*, V. **21**, N . 6, 58-71, 2021
- [67] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. **108**, 102828-1-9, 2021
- [68] H. Rezazadeh, D. Kumar, A. Neirameh, M. Eslami, M. Mirzazadeh, Applications of three methods for obtaining optical soliton solutions for the LPD model with Kerr law nonlinearity, *Pramana J?Phys.*, V. **89**, 1-11, 2020
- [69] G. Akram, M. Sadaf, M. Dawood, D. Baleanu, Optical solitons for LakshmananPorsezianDaniel equation with Kerr law non-linearity using improved tan expansion technique, *Res. In Phys.*, V. **29**, 104758-1-13, 2021
- [70] H.F. Ismael, H.M. Baskonius, H. Bulut, Abundant novel solutions of the conformable Lakshmanan-Porsezian-Daniel model, *Disc. And Cont. Dun. Syst. Ser. S*, V. **14**, N. 7, 2311-2333, 2021
- [71] H. Yopez-Martinez, H. Rezazadeh, M. Inc, M.Ali-Akinlar, J.F. Gomez-Aguilar, Analytical solutions to the fractional LakshmananPorsezianDaniel model, *Opt. And Quant. Elec.*, V. **54**, 1-41, 2021

Appendix

Solution of order 3 to the (LPD) equation depending on 4 real parameters :
The function $v(x, t)$ defined by

$$v(x, t) = \left(1 - 24 \frac{n(x, t)}{d(x, t)}\right) e^{i(2a_1 + 6t + 20b_2)} \quad (6)$$

with

$$\begin{aligned} n(x, t) = & 675 + (2x + 12b_1 + 60a_2)^{10} + 2190(4a_1 + 24t + 120b_2)^6 + 27000(-8b_1 - 80a_2)^2 + 91800(-16t - 160b_2)^2 + 495(4a_1 + 24t + 120b_2)^8 + 11(4a_1 + 24t + 120b_2)^{10} + 88473600a_2^2 + 353894400b_2^2 + 15(1 + (4a_1 + 24t + 120b_2)^2)(2x + 12b_1 + 60a_2)^8 + (210 - 60(4a_1 + 24t + 120b_2)^2 + 50(4a_1 + 24t + 120b_2)^4 + 480(4a_1 + 24t + 120b_2)(-16t - 160b_2))(2x + 12b_1 + 60a_2)^6 + (-720(4a_1 + 24t + 120b_2)^2(-8b_1 - 80a_2) + 5760b_1 - 11520a_2)(2x + 12b_1 + 60a_2)^5 + (450(4a_1 + 24t + 120b_2)^2 - 150(4a_1 + 24t + 120b_2)^4 + 70(4a_1 + 24t + 120b_2)^6 + 1200(4a_1 + 24t + 120b_2)^8 - 150(4a_1 + 24t + 120b_2)^4 + 70(4a_1 + 24t + 120b_2)^6 + 1200(4a_1 + 24t + 120b_2)^8) \end{aligned}$$

$$\begin{aligned}
& 120 b_2)^3(-16 t-160 b_2)-450+5400(-8 b_1-80 a_2)^2-1800(-16 t-160 b_2)^2+ \\
& 3600(4 a_1+24 t+120 b_2)(-16 t-224 b_2)(2 x+12 b_1+60 a_2)^4+(-2400(4 a_1+ \\
& 24 t+120 b_2)^4(-8 b_1-80 a_2)+28800(4 a_1+24 t+120 b_2)(-8 b_1-80 a_2)(-16 t- \\
& 160 b_2)-57600 b_1-806400 a_2-7200(4 a_1+24 t+120 b_2)^2(-16 b_1-128 a_2))(2 x+ \\
& 12 b_1+60 a_2)^3+(6750(4 a_1+24 t+120 b_2)^4+420(4 a_1+24 t+120 b_2)^6+45(4 a_1+ \\
& 24 t+120 b_2)^8-2700(4 a_1+24 t+120 b_2)^2(5+4(-8 b_1-80 a_2)^2-12(-16 t- \\
& 160 b_2)^2)-675-10800(-8 b_1-80 a_2)^2-10800(-16 t-160 b_2)^2+21600(4 a_1+ \\
& 24 t+120 b_2)(-32 t-384 b_2)-7200(4 a_1+24 t+120 b_2)^3(-32 t-128 b_2))(2 x+ \\
& 12 b_1+60 a_2)^2+(-1680(4 a_1+24 t+120 b_2)^6(-8 b_1-80 a_2)-28800(4 a_1+ \\
& 24 t+120 b_2)^3(-8 b_1-80 a_2)(-16 t-160 b_2)-10800(4 a_1+24 t+120 b_2)^2(-8 b_1- \\
& 272 a_2)-86400 b_1-1209600 a_2+43200(-8 b_1-80 a_2)^3+43200(-8 b_1-80 a_2)(-16 t- \\
& 160 b_2)^2+3600(4 a_1+24 t+120 b_2)^4(-8 b_1+80 a_2)-86400(4 a_1+24 t+120 b_2)((-8 b_1- \\
& 80 a_2)(-16 t-160 b_2)+32(-16 t-160 b_2)a_2-64(-8 b_1-80 a_2)b_2))(2 x+12 b_1+ \\
& 60 a_2)-720(4 a_1+24 t+120 b_2)^7(-16 t-160 b_2)+450(4 a_1+24 t+120 b_2)^4(-17+ \\
& 28(-8 b_1-80 a_2)^2+12(-16 t-160 b_2)^2)-3600(4 a_1+24 t+120 b_2)^3(-48 t- \\
& 1376 b_2)-720(4 a_1+24 t+120 b_2)^5(-272 t-3168 b_2)+10800(4 a_1+24 t+ \\
& 120 b_2)(16 t+224 b_2+4(-8 b_1-80 a_2)^2(-16 t-160 b_2)+4(-16 t-160 b_2)^3)+ \\
& 675(4 a_1+24 t+120 b_2)^2(-3+16(-8 b_1-80 a_2)^2+16(-16 t-160 b_2)^2- \\
& 4096(-8 b_1-80 a_2)a_2-8192(-16 t-160 b_2)b_2)-2764800(-8 b_1-80 a_2)a_2- \\
& 11059200(-16 t-160 b_2)b_2+i(151200 t-5529600(-8 b_1-80 a_2)(-16 t-160 b_2)a_2+ \\
& 64800(-16 t-160 b_2)^3-870(4 a_1+24 t+120 b_2)^7+25(4 a_1+24 t+120 b_2)^9+(4 a_1+ \\
& 24 t+120 b_2)^{11}-21600(-8 b_1-80 a_2)^2(-16 t-160 b_2)+(4 a_1+24 t+120 b_2)(2 x+ \\
& 12 b_1+60 a_2)^{10}+(-60 a_1-840 t-6600 b_2+5(4 a_1+24 t+120 b_2)^3)(2 x+12 b_1+ \\
& 60 a_2)^8+(-600 a_1+240 t+58800 b_2-140(4 a_1+24 t+120 b_2)^3+10(4 a_1+ \\
& 24 t+120 b_2)^5+240(4 a_1+24 t+120 b_2)^2(-16 t-160 b_2))(2 x+12 b_1+60 a_2)^6+ \\
& (-240(4 a_1+24 t+120 b_2)^3(-8 b_1-80 a_2)-1440(-8 b_1-80 a_2)(-16 t-160 b_2)+ \\
& 720(4 a_1+24 t+120 b_2)(-8 b_1-176 a_2))(2 x+12 b_1+60 a_2)^5+(-450(4 a_1+ \\
& 24 t+120 b_2)^3-210(4 a_1+24 t+120 b_2)^5+10(4 a_1+24 t+120 b_2)^7+300(4 a_1+ \\
& 24 t+120 b_2)^4(-16 t-160 b_2)+450(4 a_1+24 t+120 b_2)(-3+12(-8 b_1-80 a_2)^2- \\
& 4(-16 t-160 b_2)^2)+14400 t+259200 b_2+1800(4 a_1+24 t+120 b_2)^2(-16 t- \\
& 224 b_2))(2 x+12 b_1+60 a_2)^4+(-480(4 a_1+24 t+120 b_2)^5(-8 b_1-80 a_2)+ \\
& 14400(4 a_1+24 t+120 b_2)^2(-8 b_1-80 a_2)(-16 t-160 b_2)+7200(4 a_1+24 t+ \\
& 120 b_2)(-8 b_1-48 a_2)-2400(4 a_1+24 t+120 b_2)^3(-16 b_1-128 a_2)-14400(-8 b_1- \\
& 80 a_2)(-16 t-160 b_2)-460800(-16 t-160 b_2)a_2+921600(-8 b_1-80 a_2)b_2)(2 x+ \\
& 12 b_1+60 a_2)^3+(1710(4 a_1+24 t+120 b_2)^5-60(4 a_1+24 t+120 b_2)^7+5(4 a_1+ \\
& 24 t+120 b_2)^9-900(4 a_1+24 t+120 b_2)^3(7+4(-8 b_1-80 a_2)^2-12(-16 t- \\
& 160 b_2)^2)+675(4 a_1+24 t+120 b_2)(7+16(-8 b_1-80 a_2)^2+16(-16 t-160 b_2)^2)+ \\
& 345600 t+4492800 b_2-21600(-8 b_1-80 a_2)^2(-16 t-160 b_2)-21600(-16 t- \\
& 160 b_2)^3+691200(4 a_1+24 t+120 b_2)^2 b_2-1800(4 a_1+24 t+120 b_2)^4(-64 t- \\
& 448 b_2))(2 x+12 b_1+60 a_2)^2+(-240(4 a_1+24 t+120 b_2)^7(-8 b_1-80 a_2)- \\
& 7200(4 a_1+24 t+120 b_2)^4(-8 b_1-80 a_2)(-16 t-160 b_2)+10800(4 a_1+24 t+ \\
& 120 b_2)(-24 b_1-400 a_2+4(-8 b_1-80 a_2)^3+4(-8 b_1-80 a_2)(-16 t-160 b_2)^2)+ \\
& 3600(4 a_1+24 t+120 b_2)^3(-24 b_1-176 a_2)+720(4 a_1+24 t+120 b_2)^5(-56 b_1- \\
& 400 a_2)+21600(-8 b_1-80 a_2)(-16 t-160 b_2)+1382400(-16 t-160 b_2)a_2- \\
& 2764800(-8 b_1-80 a_2)b_2-43200(4 a_1+24 t+120 b_2)^2((-8 b_1-80 a_2)(-16 t- \\
& 160 b_2)+32(-16 t-160 b_2)a_2-64(-8 b_1-80 a_2)b_2))(2 x+12 b_1+60 a_2)-
\end{aligned}$$

$$\begin{aligned}
& 90(4a_1 + 24t + 120b_2)^8(-16t - 160b_2) + 90(4a_1 + 24t + 120b_2)^5(-107 + \\
& 28(-8b_1 - 80a_2)^2 + 12(-16t - 160b_2)^2) + 5400(4a_1 + 24t + 120b_2)^2(-176t - \\
& 2464b_2 + 4(-8b_1 - 80a_2)^2(-16t - 160b_2) + 4(-16t - 160b_2)^3) - 120(4a_1 + 24t + \\
& 120b_2)^6(-80t - 1248b_2) + 900(4a_1 + 24t + 120b_2)^4(-464t - 4000b_2) - 225(4a_1 + \\
& 24t + 120b_2)^3(11 + 80(-8b_1 - 80a_2)^2 + 80(-16t - 160b_2)^2 + 4096(-8b_1 - \\
& 80a_2)a_2 + 8192(-16t - 160b_2)b_2) - 675(4a_1 + 24t + 120b_2)(-7 + 56(-8b_1 - \\
& 80a_2)^2 + 88(-16t - 160b_2)^2 - 4096(-8b_1 - 80a_2)a_2 - 131072a_2^2 - 524288b_2^2) + \\
& 5529600(-8b_1 - 80a_2)^2b_2 - 5529600(-16t - 160b_2)^2b_2 + 1857600b_2)
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & 2024 + 518400(-16t - 160b_2)^4 + 356400(-8b_1 - 80a_2)^2 + 874800(-16t - \\
& 160b_2)^2 + 3720(4a_1 + 24t + 120b_2)^8 + 120(4a_1 + 24t + 120b_2)^{10} + 530841600a_2^2 + \\
& 2123366400b_2^2 + (1 + (2x + 12b_1 + 60a_2)^2 + (4a_1 + 24t + 120b_2)^2)^6 + 518400(-8b_1 - \\
& 80a_2)^4 + (-1440(4a_1 + 24t + 120b_2)^4 + 720(4a_1 + 24t + 120b_2)^5(-16t - 160b_2) + \\
& 240(4a_1 + 24t + 120b_2)^2(56 + 135(-8b_1 - 80a_2)^2 - 45(-16t - 160b_2)^2) + \\
& 32400(4a_1 + 24t + 120b_2)(-16t - 288b_2) + 7200(4a_1 + 24t + 120b_2)^3(-48t - \\
& 544b_2) + 3360 + 32400(-8b_1 - 80a_2)^2 - 54000(-16t - 160b_2)^2 + 2764800(-8b_1 - \\
& 80a_2)a_2 + 5529600(-16t - 160b_2)b_2)(2x + 12b_1 + 60a_2)^4 + (-960(4a_1 + 24t + \\
& 120b_2)^6(-8b_1 - 80a_2) + 57600(4a_1 + 24t + 120b_2)^3(-8b_1 - 80a_2)(-16t - \\
& 160b_2) - 43200(4a_1 + 24t + 120b_2)^2(-24b_1 - 272a_2) - 7200(4a_1 + 24t + \\
& 120b_2)^4(-48b_1 - 448a_2) - 345600b_1 - 5529600a_2 - 86400(-8b_1 - 80a_2)^3 - \\
& 86400(-8b_1 - 80a_2)(-16t - 160b_2)^2 + 172800(4a_1 + 24t + 120b_2)((-8b_1 - \\
& 80a_2)(-16t - 160b_2) - 32(-16t - 160b_2)a_2 + 64(-8b_1 - 80a_2)b_2))(2x + 12b_1 + \\
& 60a_2)^3 + (13440(4a_1 + 24t + 120b_2)^6 + 240(4a_1 + 24t + 120b_2)^8 - 240(4a_1 + \\
& 24t + 120b_2)^4(-326 + 45(-8b_1 - 80a_2)^2 - 135(-16t - 160b_2)^2) + 480(4a_1 + \\
& 24t + 120b_2)^2(-76 + 135(-8b_1 - 80a_2)^2 + 1215(-16t - 160b_2)^2) - 129600(4a_1 + \\
& 24t + 120b_2)^3(-32t - 256b_2) - 12960(4a_1 + 24t + 120b_2)^5(-32t - 256b_2) - \\
& 64800(4a_1 + 24t + 120b_2)(96t + 1280b_2 + 4(-8b_1 - 80a_2)^2(-16t - 160b_2) + \\
& 4(-16t - 160b_2)^3) + 12144 - 97200(-8b_1 - 80a_2)^2 + 32400(-16t - 160b_2)^2 + \\
& 530841600a_2^2 - 33177600(-16t - 160b_2)b_2 + 2123366400b_2^2)(2x + 12b_1 + 60a_2)^2 + \\
& (-360(4a_1 + 24t + 120b_2)^8(-8b_1 - 80a_2) - 17280(4a_1 + 24t + 120b_2)^5(-8b_1 - \\
& 80a_2)(-16t - 160b_2) - 1440(4a_1 + 24t + 120b_2)^6(-8b_1 - 240a_2) + 32400(4a_1 + \\
& 24t + 120b_2)^4(-8b_1 + 112a_2) + 64800(4a_1 + 24t + 120b_2)^2(40b_1 + 752a_2 + \\
& 4(-8b_1 - 80a_2)^3 + 4(-8b_1 - 80a_2)(-16t - 160b_2)^2) - 777600(4a_1 + 24t + \\
& 120b_2)((-8b_1 - 80a_2)(-16t - 160b_2) + 64(-16t - 160b_2)a_2 - 128(-8b_1 - \\
& 80a_2)b_2) - 172800(4a_1 + 24t + 120b_2)^3(3(-8b_1 - 80a_2)(-16t - 160b_2) + 32(-16t - \\
& 160b_2)a_2 - 64(-8b_1 - 80a_2)b_2) + 648000b_1 + 8553600a_2 + 259200(-8b_1 - \\
& 80a_2)^3 + 1296000(-8b_1 - 80a_2)(-16t - 160b_2)^2 - 33177600(-8b_1 - 80a_2)^2a_2 + \\
& 33177600(-16t - 160b_2)^2a_2 - 132710400(-8b_1 - 80a_2)(-16t - 160b_2)b_2)(2x + \\
& 12b_1 + 60a_2) + 120(-8b_1 - 80a_2)(2x + 12b_1 + 60a_2)^9 - 120(4a_1 + 24t + \\
& 120b_2)^9(-16t - 160b_2) + 80(4a_1 + 24t + 120b_2)^6(191 + 63(-8b_1 - 80a_2)^2 + \\
& 27(-16t - 160b_2)^2) - 2160(4a_1 + 24t + 120b_2)^5(-240t - 4576b_2) + 21600(4a_1 + \\
& 24t + 120b_2)^3(368t + 3488b_2 + 4(-8b_1 - 80a_2)^2(-16t - 160b_2) + 4(-16t - \\
& 160b_2)^3) - 1440(4a_1 + 24t + 120b_2)^7(-80t - 864b_2) + 240(4a_1 + 24t + 120b_2)^4(599 + \\
& 135(-8b_1 - 80a_2)^2 - 225(-16t - 160b_2)^2 - 11520(-8b_1 - 80a_2)a_2 - 23040(-16t - \\
& 160b_2)b_2) - 16200(4a_1 + 24t + 120b_2)(-496t - 6240b_2 + 80(-8b_1 - 80a_2)^2(-16t - \\
& 160b_2) + 16(-16t - 160b_2)^3 + 4096(-8b_1 - 80a_2)(-16t - 160b_2)a_2 - 4096(-8b_1 -
\end{aligned}$$

$$\begin{aligned}
& 80 a_2)^2 b_2 + 4096 (-16 t - 160 b_2)^2 b_2 + 24 (4 a_1 + 24 t + 120 b_2)^2 (3881 + 12150 (-8 b_1 - \\
& 80 a_2)^2 + 28350 (-16 t - 160 b_2)^2 + 691200 (-8 b_1 - 80 a_2) a_2 + 22118400 a_2^2 + \\
& 88473600 b_2^2) + 1036800 (-8 b_1 - 80 a_2)^2 (-16 t - 160 b_2)^2 + 46080 a_2 (2 x + 12 b_1 + \\
& 60 a_2)^7 - 24883200 (-8 b_1 - 80 a_2) a_2 - 82944000 (-16 t - 160 b_2) b_2 + (-120 (4 a_1 + \\
& 24 t + 120 b_2)^2 + 360 (4 a_1 + 24 t + 120 b_2) (-16 t - 160 b_2) + 120) (2 x + 12 b_1 + \\
& 60 a_2)^8 + (480 (4 a_1 + 24 t + 120 b_2)^2 - 240 (4 a_1 + 24 t + 120 b_2)^4 + 960 (4 a_1 + 24 t + \\
& 120 b_2)^3 (-16 t - 160 b_2) + 2320 + 2160 (-8 b_1 - 80 a_2)^2 + 5040 (-16 t - 160 b_2)^2 - \\
& 1440 (4 a_1 + 24 t + 120 b_2) (-64 t - 960 b_2)) (2 x + 12 b_1 + 60 a_2)^6 + (-720 (4 a_1 + \\
& 24 t + 120 b_2)^4 (-8 b_1 - 80 a_2) - 17280 (4 a_1 + 24 t + 120 b_2) (-8 b_1 - 80 a_2) (-16 t - \\
& 160 b_2) + 4320 (4 a_1 + 24 t + 120 b_2)^2 (-8 b_1 - 176 a_2) + 51840 b_1 + 103680 a_2) (2 x + \\
& 12 b_1 + 60 a_2)^5
\end{aligned}$$

is a solution to the Lakshmanan Porsezian Daniel equation (1).