



**HAL**  
open science

## **Impact of Hypoxia and Hadrontherapy on Radioresistant Chondrosarcoma Cancer Stem Cells**

Antoine Gilbert, Julie Laurence, Marilyne Guillamin, Siamak Haghdoost, Samuel Valable, François Chevalier

► **To cite this version:**

Antoine Gilbert, Julie Laurence, Marilyne Guillamin, Siamak Haghdoost, Samuel Valable, et al.. Impact of Hypoxia and Hadrontherapy on Radioresistant Chondrosarcoma Cancer Stem Cells. Proceeding of the ERRS 2021, Nov 2021, CAEN, France. pp.93. <hal-03757408>

**HAL Id: hal-03757408**

**<https://hal.science/hal-03757408v1>**

Submitted on 22 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

---

# Impact of Hypoxia and Hadrontherapy on Radioresistant Chondrosarcoma Cancer Stem Cells

Antoine Gilbert<sup>\*1</sup>, Julie Laurence<sup>1</sup>, Marilyne Guillamin<sup>2</sup>, Siamak Haghdoost<sup>1</sup>, Samuel Valable<sup>3</sup>, and Francois Chevalier<sup>1</sup>

<sup>1</sup>Centre de recherche sur les Ions, les MATériaux et la Photonique – Centre National de la Recherche Scientifique : UMR6252, Ecole Nationale Supérieure d'Ingénieurs de Caen, Normandie Université, Commissariat à l'énergie atomique et aux énergies alternatives : DRF/IRAMIS, Université de Caen Normandie – CIMAP - UMR 6252, 6, Boulevard du Maréchal Juin 14050 Caen cedex 4, France

<sup>2</sup>Université de Caen Normandie – Icore – Esplanade de la Paix - CS 14032 - 14032 CAEN Cedex 5, France

<sup>3</sup>Imagerie et Stratégies Thérapeutiques des pathologies Cérébrales et Tumorales – Université de Caen Normandie, Normandie Université, Commissariat à l'énergie atomique et aux énergies alternatives, Centre National de la Recherche Scientifique : UMR6030 – GIP CYCERON Campus Jules Horowitz Bd Henri Becquerel BP 5229 14074 CAEN, France

## Abstract

Chondrosarcoma is a malignant tumor characterized by an abundant extracellular matrix, which is responsible of a hypoxic microenvironment. Hypoxia can lead to resistance of conventional X-rays treatment by the absence of oxygen, essential for the formation of Reactive Oxygen Species (ROS) associated with cellular toxicity. Hypoxia is also related to a possible stabilization of subpopulation of Cancer Stem Cells (CSCs) possibly implicated in radioresistance [1]. Hadrontherapy is expected to induce more damage under a hypoxia environment, according to the characteristics of ions, but the effect of hadrons on the CSCs needs to be further analyzed.

For this study, the impact of the different irradiations qualities (X-rays, protons and carbon ions) under low (1%) and high (20%) oxygen conditions were analyzed on 2 chondrosarcoma cell lines: L835 and Oums27 [2], respectively muted on IDH1 and p53. Cell survival was evaluated by clonogenic assay and the impact of treatments on CSC subpopulation was studied using an ALDH activity test by flow cytometry and a tumorsphere assay.

As expected, hypoxia decreased X-rays efficacy on both chondrosarcoma cell lines. This effect was less evident with proton and carbon ions irradiations, excepted in case of the L835 cells irradiated with carbon ions. Without irradiation, hypoxia increased ALDH-high cells only with Oums27 cells. After a high dose (3Gy) of carbon ions, an increased of the ALDH-high population was observed with Oums27 cells, in both oxygen conditions. Using L835 cells, an increase of the ALDH-high population was also observed with hypoxic culture condition, independently of the dose (1 to 3Gy).

This preliminary study shows a differential impact of hypoxia on chondrosarcoma cells survival after X-rays, protons and carbon ions irradiations. The corresponding radioresistance

---

\*Speaker

could be linked with a CSC subpopulation.

1. Vares G, Ahire V, Sunada S, Ho Kim E, Sai S, Chevalier F, et al. *Radiother Oncol.* 2020;150:253–61.
2. Chevalier F, Hamdi DH, Lepleux C, Temelie M, Nicol A, Austray JB, et al. *Technol Cancer Res Treat.* 2019;18:153303381987130.