

## Neurons membrane potential beats to the rhythm of breath

Maxime Juventin, Baptiste Girin, Mickaël Zbili, Nicolas Fourcaud-Trocmé, S. Garcia, Nathalie Buonviso, Corine Amat

#### ▶ To cite this version:

Maxime Juventin, Baptiste Girin, Mickaël Zbili, Nicolas Fourcaud-Trocmé, S. Garcia, et al.. Neurons membrane potential beats to the rhythm of breath. FENS (Federation of European Neuroscience Societies ) 2020, 2020, Virtual forum, France. hal-03758974

## HAL Id: hal-03758974 https://hal.science/hal-03758974

Submitted on 30 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Juventin M., Girin B., Zbili M., Fourcaud-Trocmé N., Garcia S., Buonviso N., and Amat C.

Lyon Neuroscience Research Center, team : Olfaction : from Coding to Memory, CNRS UMR5292 – INSERM U1028 – University Lyon 1


### Introduction

Respiratory rhythm has recently emerged as a key element in the cortical dynamics, since an increasing number of studies have shown a global entrainment of LFP activities by the respiratory rhythm. This observation has been done in rodents as well in humans, and in a large set of brain areas (cortical and limbic areas). Since this question has been tackled in non-olfactory structures almost exclusively through LFP analysis, the mechanisms underlying such respiratory modulation cannot be fully understood. To carry on understanding, we performed intracellular and LFP co-recordings in four non-olfactory brain areas : Hippocampus (H), and the somatosensory (S1), primary visual (V1) and prelimbic cortices (pL), in anesthetized rats.

## Methods


Experiments were performed on freely breathing anesthetized rats. Intracellular activity (or **Vm**) and LFP were co-recorded. For Vm, sharp electrodes filled with a solution of 2M potassium acetate (Resistance 45-120 Mohms) were used. Respiration was monitored using a pressure sensor placed before the right nostril.

# Result 1 : Respiratory-rhythm (**RR**) can entrain neuron activity in widespread brain areas




## Result 2 : Brain-wide quantification of Vm RR oscillations

To precisely quantify RR entrainement of **Vm oscillations**, we developped a detection method using respiration triggered waveform Vm (4 respi. cycles sliding windows), and compared to surrogates. This algorhythm finally led to a RR encoding of Vm for each respiratory cycles.







**Brain-wide** : proportion of RR entrained cells

100

75


50


25

pL S1

V1 H

Proportion of cells





To what extent brain areas are sensitive to RR?

## Conclusions

Our results suggest that a respiratory drive of cellular membrane potential exists in a large brain network, influencing temporal discharge pattern. This drive is not unique but varies according to the brain area, LFP dynamics and probably other parameters that remain to be explored. Result 3 : Vm and LFP entrainement are partly coordinated

We encoded with the same detection method both Vm and LFP. Then we computed the probability for Vm to be RR-entrained as a function of LFP dynamics.

