



Evidence of an additional center of apple domestication in Iran, with contributions from the Caucasian crab apple *Malus orientalis* Uglitzk. to the cultivated apple gene pool. Running head: Apple domestication in the Caucasus and Iran

Hamid Bina, Hamed Yousefzadeh, Anthony Venon, Carine Remoué, Agnès Rousselet, Matthieu Falque, Shadab Faramarzi, Chen Xilong, Jarkyn Samanchina, David Gill, et al.

► **To cite this version:**

Hamid Bina, Hamed Yousefzadeh, Anthony Venon, Carine Remoué, Agnès Rousselet, et al.. Evidence of an additional center of apple domestication in Iran, with contributions from the Caucasian crab apple *Malus orientalis* Uglitzk. to the cultivated apple gene pool. Running head: Apple domestication in the Caucasus and Iran. *Molecular Ecology*, In press, 31 (21), pp.5581-5601. <10.1111/mec.16667>. <hal-03759285>

HAL Id: hal-03759285

<https://hal.science/hal-03759285v1>

Submitted on 24 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 Title: Evidence of an additional center of apple domestication in Iran, with contributions from the
2 Caucasian crab apple *Malus orientalis* Uglitzk. to the cultivated apple gene pool.

3
4 **Running head:** Apple domestication in the Caucasus and Iran

5
6 Bina Hamid¹, Yousefzadeh Hamed², Venon Anthony³, Remoué Carine³, Rousselet Agnès³, Falque
7 Matthieu³, Faramarzi Shadab⁴, Xilong Chen³, Jarkyn Samanchina⁵, David Gill⁵, Akylai Kabaeva⁵,
8 Giraud Tatiana⁶, Hossainpour Batool⁷, Abdollahi Hamid⁸, Gabrielyan Ivan⁹, Nersesyan Anush¹⁰,
9 Cornille A.³

10
11 ¹ Department of Forestry, Tarbiat Modares University, Noor, Iran

12 ² Department of Environmental science, Biodiversity Branch, Natural resources faculty, Tarbiat
13 Modares University, Noor, Iran

14 ³ Université Paris-Saclay, INRAE, CNRS, AgroParisTech, GQE - Le Moulon, 91190 Gif-sur-
15 Yvette, France

16 ⁴ Department of Plant Production and Genetics, Faculty of Agriculture, Razi University,
17 Kermanshah, Iran

18 ⁵ Fauna & Flora International, Cambridge, UK

19 ⁶ Ecologie Systematique Evolution, Universite Paris-Saclay, CNRS, AgroParisTech, 91190 Gif-
20 sur-Yvette, France

21 ⁷ Iranian Research Organization for Science and Technology (IROST), Institute of Agriculture,
22 Department of Agriculture, Tehran, Iran

23 ⁸ Temperate Fruits Research Center, Horticultural Sciences Research Institute, Agricultural
24 Research, Education and Extension Organization (AREEO), Karaj, Iran

25 ⁹ Department of Palaeobotany, A. Takhtajyan Institute of Botany, Armenian National Academy
26 of Sciences, Acharyan Str.1, 0040 Yerevan, Armenia

27 ¹⁰ Department of Conservation of Genetic Resources of Armenian Flora, A. Takhtajyan Institute
28 of Botany, Armenian National Academy of Sciences, Acharyan Str.1, 0040 Yerevan, Armenia

29
30
31 Corresponding authors: amandine.cornille@cnrs.fr, h.yousefzadeh@modares.ir

Abstract

Divergence processes in crop-wild fruit tree complexes in pivotal regions for plant domestication such as the Caucasus and Iran remain little studied. We investigated anthropogenic and natural divergence processes in apples in these regions using 26 microsatellite markers amplified in 550 wild and cultivated samples. We found two genetically distinct cultivated populations in Iran that are differentiated from *Malus domestica*, the standard cultivated apple worldwide. Coalescent-based inferences showed that these two cultivated populations originated from specific domestication events of *Malus orientalis* in Iran. We found evidence of substantial wild-crop and crop-crop gene flow in the Caucasus and Iran, as has been described in apple in Europe. In addition, we identified seven genetically differentiated populations of wild apple (*M. orientalis*), not introgressed by the cultivated apple. Niche modeling combined with genetic diversity estimates indicated that these wild populations likely resulted from range changes during past glaciations. This study identifies Iran as a key region in the domestication of apple and *M. orientalis* as an additional contributor to the cultivated apple gene pool. Domestication of the apple tree therefore involved multiple origins of domestication in different geographic locations and substantial crop-wild hybridization, as found in other fruit trees. This study also highlights the impact of climate change on the natural divergence of a wild fruit tree and provides a starting point for apple conservation and breeding programs in the Caucasus and Iran.

Key words: apple, Caucasus, Iran, crop-wild gene flow, domestication, fruit tree, climate, introgression.

Introduction

Crop-wild complexes, *i.e.*, groups of crops and their associated wild species that exchange genes (Anderson, 2005), provide good models for understanding how anthropogenic and natural factors shape population divergence in the presence of gene flow. Indeed, crops are the result of a recent anthropogenic divergence process, *i.e.*, domestication, which began around 10,000 years ago, and which has often been followed by subsequent crop-wild gene flow (Besnard, Terral, & Cornille, 2018; Brandenburg et al., 2017; Chen et al., 2019; Cornille, Giraud, Smulders, Roldán-Ruiz, & Gladieux, 2014; Cornille et al., 2012; Diez et al., 2015; Flowers et al., 2019; Gaut, Díez, & Morrell, 2015). Conversely, wild species allow the study of natural divergence over a longer timescale. Indeed, wild species have often undergone shifts in their distribution following past climate changes associated with glacial periods, and range contraction has often led to population differentiation and divergence (Excoffier, Foll, & Petit, 2009; Hewitt, 1990; Hewitt, 1996; Jezkova, Olah-Hemmings, & Riddle, 2011; Petit, Bialozyt, Garnier-Géré, & Hampe, 2004; Schmitt, 2007). Understanding the evolutionary processes shaping the natural and anthropogenic divergence of crop-wild complexes is not just an academic exercise: it will also help assess the future status of wild resources. Because of the socio-economic importance of crop plants, protecting the wild relatives of crops, beyond the need for preserving biodiversity (Bacles & Jump, 2011), will allow us to manage the genetic resources for future breeding strategies in the face of global changes (*e.g.*, climate change, emerging diseases) (Bailey-Serres, Parker, Ainsworth, Oldroyd, & Schroeder, 2019; Castañeda-Álvarez et al., 2016; Zhang, Mittal, Leamy, Barazani, & Song, 2017).

Fruit trees present several historical and biological features that make them fascinating models for investigating anthropogenic and natural divergence with gene flow. Several fruit tree crops are found across the world and sometimes occur in sympatry with their wild relatives (Besnard, Terral, & Cornille, 2018; Cornille et al., 2019; Cornille, Giraud, Smulders, Roldán-Ruiz, & Gladieux, 2014; Delplancke et al., 2011; Liu et al., 2019; Miller & Gross, 2011). Fruit trees are also characterized by high levels of gene flow during divergence, which is to be expected considering the typical life history traits of trees (Cornille et al., 2013; Cornille, Gladieux, & Giraud, 2013; Oddou-Muratorio & Klein, 2008; Petit & Hampe, 2006). Population genetics studies of natural divergence processes associated with the last glacial maximum in Europe, North America and Asia in wind-dispersed trees (*e.g.*, *Abies*, *Pinus*, *Fraxinus*, *Quercus*, *Betula* (Lascoux, Palmé, Cheddadi, & Latta, 2004; Petit et al., 2004)) and animal-dispersed trees (Cornille et al., 2013;

George et al., 2015) showed that there were high levels of gene flow between populations and that trees had high dispersal capabilities. These studies also located single (Bai & Spitkovsky, 2010; Tian, Li, Ji, Zhang, & Luo, 2009; Zeng et al., 2011) or multiple (Qiu, Wang, Liu, Shen, & Tang, 2011; Tian et al., 2009) glacial refugia where most temperate tree species persisted during the last glacial maximum, and from which populations recolonized higher or lower latitudes during the Holocene post-glacial expansion (Giesecke, Brewer, Finsinger, Leydet, & Bradshaw, 2017). Population genetics and genomics studies also revealed the prominent role of gene flow during the anthropogenic divergence of fruit trees. Domestication of several emblematic fruit tree crops such as grape and apple occurred with substantial crop-crop and wild-crop gene flow and without a bottleneck (Arroyo-García et al., 2006; Cornille et al., 2012; Decroocq et al., 2016; Diez et al., 2015; Duan et al., 2017; Liu et al., 2019; Meyer, Duval, & Jensen, 2012; Myles et al., 2011). These studies thus revealed that the domestication of fruit trees displays different patterns from that of annuals which can be explained by the long lifespan, long juvenile phase and self-incompatibility system of trees (Fuller, 2018; Gaut et al., 2015). However, studies of natural and anthropogenic divergence processes in crop-wild fruit tree complexes are still scarce in the geographic regions that were pivotal in the divergence history of these complexes.

The Caucasus ecoregion harbors a remarkable concentration of both economically important plants and their wild relatives, in particular wheat, rye, barley and also fruit trees including walnut, apricot and apple (Asanidze, Akhalkatsi, Henk, Richards, & Volk, 2014; Gabrielian & Zohary, 2004; Vavilov, 1926, 1992; Yousefzadeh, Hosseinzadeh Colagar, Tabari, Sattarian, & Assadi, 2012). This region covers Georgia, Armenia, Azerbaijan, the North Caucasian part of the Russian Federation, the northeastern part of Turkey and the Hyrcanian Mixed Forests region in northwestern Iran (Nakhutsrishvili, Zazanashvili, Batsatsashvili, & Montalvo, 2015; Zazanashvili et al., 2020). Two Pleistocene refugia for temperate plants are recognized in this region (Bina, Yousefzadeh, Ali, & Esmailpour, 2016; Yousefzadeh et al., 2012): the Colchis refugium in the catchment basin of the Black Sea (in the Western Caucasus) and the Hyrcanian refugium at the southern edge of the Caspian Sea. Glacial refugia are known to harbor higher levels of species and genetic diversity (Hewitt, 2004), and this is the case for the Colchis and Hyrcanian refugia. Furthermore, it has been suggested that Iran, with its close proximity to Central Asia - the center of origin of emblematic fruit trees - and its historic position on the Silk Trade Routes, is a possible secondary center of domestication of apple, grape and apricot (Decroocq et al., 2016;

Liang et al., 2019; Liu et al., 2019). However, inferences of the natural and anthropogenic divergence history of wild-crop fruit tree complexes in the Caucasus have been limited by the small number of samples (Decroocq et al., 2016; Liu et al., 2019) and/or genetic markers investigated so far (Amirchakhmaghi et al., 2018; Asanidze et al., 2014; Cornille et al., 2013; Gharghani et al., 2010; Myles et al., 2011; Volk & Cornille, 2019; Vouillamoz et al., 2006).

The history of domestication of the apple tree in the Caucasus and Iran remains to be resolved. Some authors suggest that the local apple cultivars from several regions of the Caucasus originated from *M. orientalis* (Forsline, Aldwinckle, Dickson, Luby, & Hokanson, 2003; Langenfeld, 1991; Schmitt, 2007) and not from the Central Asian wild apple, *M. sieversii*, the progenitor of *M. domestica* (Cornille et al., 2019; Cornille, Giraud, Smulders, Roldán-Ruiz, & Gladieux, 2014; Duan et al., 2017, 2017; Harris, Robinson, & Juniper, 2002; Migicovsky et al., 2021), or may even be *M. orientalis* grown in gardens and orchards. The Caucasian crab apple, *Malus orientalis* Uglitzk., is an endemic wild apple species in the Caucasus. More specifically, it is found in the Western Caucasus (*i.e.*, the southern part of Russia, northern Anatolia and northwestern Turkey), Georgia, Armenia, the mountainous belt in northern Iran, the Hyrcanian Forests (Büttner, 2001; Rechinger, 1964) and the Zagros Forests in eastern and central Iran (Browicz, 1969; Rechinger, 1964) (Figure 1). This species displays high phenotypic diversity across its distribution range where it occurs as scattered individuals in natural forests or at high altitude in rocky mountains (Fischer & Schmidt, 1938; Rechinger, 1964). The Caucasian crab apple has a high resistance to pests and diseases (Büttner, 2001) and its fruit, of high quality, are variable in size (2-4 cm) and color (green to yellowish) (Cornille et al., 2014). *Malus orientalis* has been present and cultivated in Iran (formerly Persia) long before *M. domestica*, the standard cultivated apple worldwide, was introduced (Abdollahi, 2021; Curtis et al., 2005; Fallahi, Colt, Fallahi, & Chun, 2002; Fallahi, Fallahi, & Mahdavi, 2020; Gharghani et al., 2010; Gharghani et al., 2009; Spengler, 2019). Translations of cuneiform scripts from Persepolis and other historical documents that are related to the Sassanid and Parthian dynasties revealed that edible apples were grown in Iran in the form of dwarf apples three centuries before Christ (BC) (Abdollahi, 2021; Curtis et al., 2005; Fallahi et al., 2002). Cultivation of *M. orientalis* in Persian gardens became more widespread after the invasion and subjugation of large portions of the Persian Empire (334 BC) by Alexander the Great (Bowe, 2004). Dwarf apples became popular in gardens and paradise gardens (derived from the Persian term ‘*pardis*’) around 60 BC in Iran (Fallahi et al., 2002). Apple orchards were

cultivated more widely in Iran during the Safavid period (1501-1736), and apple cultivation has continued in Iran until now (Fallahi et al., 2002, 2020). Nowadays, the fruit of several local cultivars that are unique to Iran are harvested in the Caucasus for stewing and processed as juice and other beverages (cider, wine), jelly, syrup, jam and vinegar (Amirchakhmaghi et al., 2018; Büttner, 2001). *Malus domestica* is also currently grown in various regions of the Caucasus, but to a much lesser extent than the local cultivars (Forsline et al., 2003; Gharghani et al., 2010; Gharghani et al., 2009; Langenfeld, 1991; Schmitt, 2007). Therefore, there may have been additional events of domestication of the apple tree in the Caucasus, from *M. orientalis*, or *M. domestica*, or apple domestication in the Caucasus may be in its early stage, with the cultivation of *M. orientalis* individuals in orchards. *Malus orientalis* may also be at the origin of wild-to-crop introgressions in the same way as *M. sylvestris*, the European crab apple, contributed to the *M. domestica* gene pool (Cornille et al., 2012; Gharghani et al., 2009); hybrid cultivated trees would then occur in gardens and orchards in Iran. The origin of the cultivated apple tree in the Caucasus and Iran therefore remains unresolved despite the large consumption of the apple fruit in these regions.

Reciprocally, gene flow from cultivated trees may have led to the presence of hybrid trees in the wild, or even to feral trees. In France, the occurrence of orchards surrounding wild apple populations have directly impacted the levels of crop-to-wild introgression of the European wild apple populations (Cornille et al., 2015). However, the extent of crop-wild gene flow in apples in the Caucasus and Iran has just begun to be investigated. A study suggested that *M. orientalis* only made a minor contribution to Mediterranean *M. domestica* cultivars (Cornille et al., 2012), but lacked in-depth investigation. A population genetics study revealed low levels of crop-to-wild gene flow from *M. domestica* to *M. orientalis* in natural forests of Armenia, Turkey and Russia (Cornille et al., 2013). Population genetic diversity and structure analyses of *M. orientalis* populations from the Western Caucasus and South Caucasus identified three differentiated populations: one in Turkey, one in Armenia and one in Russia (Cornille et al., 2013). On a smaller geographic scale, an east-west genetic subdivision was found across the Hyrcanian Forests in Iran, with five main populations showing admixture (Amirchakhmaghi et al., 2018). However, we still lack a comprehensive view (beyond Armenia) of the genetic diversity and structure of *M. orientalis* to understand its natural divergence history and its contribution to local cultivars.

Here, we therefore investigate anthropogenic and natural divergence processes in apples from the Caucasus and the extent of gene flow during divergence. A total of 550 apple trees, comprising local cultivated and wild apple trees from the Caucasus, *M. domestica* apple cultivars and *M. sieversii*, were genotyped using 26 microsatellite markers. In addition, the Siberian wild apple *Malus baccata* was used as an outgroup. From the analysis of this comprehensive genetic dataset, combined with ecological niche modeling approaches, we addressed the following questions: 1) What is the population genetic structure among Caucasian wild and cultivated apples, *M. domestica* and *M. sieversii*, and what are their genetic relationships and levels of genetic diversity? 2) What is the extent of crop-crop, crop-wild and wild-wild gene flow in apple trees in the Caucasus? Are there feral or hybrid trees in the wild? Do humans cultivate wild or hybrid trees? 3) Is *M. orientalis* an additional contributor to the domestication history of cultivated apple? 4) Did *M. orientalis* experience past range contraction and expansion associated with the last glacial maximum?

Materials and methods

Sampling, DNA extraction and microsatellite genotyping

Microsatellite genotyping data for *M. orientalis*, *M. sieversii* from Kazakhstan, *M. domestica* and *M. baccata* were from previously published studies, including: 207 *M. orientalis* individuals from Turkey, Armenia and Russia (23 sites, Tables S1 and S2, (Cornille et al., 2013; Cornille et al., 2012)), four apple cultivars from Armenia, 40 “pure” European cultivated *M. domestica* individuals (*i.e.*, not introgressed by *M. sylvestris*) (Tables S1 and S2, (Cornille et al., 2013, 2012)), 20 *M. sieversii* individuals from Kazakhstan (Cornille et al., 2013) and 22 *M. baccata* individuals from Russia (Cornille et al., 2012). We collected 257 new samples in 2017 in Iran and in 2018 in Kyrgyzstan (*M. sieversii*) for this study: 167 *M. orientalis* individuals from the Hyrcanian Forests and the Zagros region in Iran (Table S1), 48 local Iranian apple cultivars from the *Seed and Plant Improvement Institute* (Karaj, Iran) (Table S1) and 42 *M. sieversii* individuals from Kyrgyzstan. Note that for 18 of the 48 local Iranian samples, fruit size was measured (Table S1). Collections meet the requirements of the recently enacted Nagoya protocol on access to genetic resources and the fair and equitable sharing of benefits. Thus, a total of 550 individuals were analyzed, comprising 374 *M. orientalis*, 48 Iranian and four Armenian apple cultivars, 40 European apple

cultivars belonging to *M. domestica*, 62 *M. sieversii* (from Kyrgyzstan and Kazakhstan) and 22 *M. baccata* individuals (details are provided in Table S1).

DNA from the new samples ($N = 257$) was extracted from dried leaves with the NucleoSpin plant II DNA extraction kit (Macherey & Nagel, Düren, Germany®) following the manufacturer's instructions. Multiplex microsatellite PCR amplifications were performed with a multiplex PCR kit (Qiagen Inc.®) for 26 microsatellite markers as previously described (Cornille et al., 2012; Patocchi, Frei, Frey, & Kellerhals, 2009). Note that on each DNA plate, we included three controls, *i.e.*, one sample of *M. orientalis*, one of *M. sieversii* and one of *M. domestica* for which data were already available (Cornille et al., 2013). Genotypes of the controls were compared with the 2013 dataset. We retained only multilocus genotypes for which $< 20\%$ of the data was missing. The suitability of these markers for population genetics analyses has been demonstrated in previous studies (Cornille et al., 2013; Cornille, Gladieux, & Giraud, 2013; Cornille et al., 2012).

Bayesian inferences of population structure, genetic differentiation among wild and cultivated apples, and genetic diversity estimates

We investigated the population structure of wild and cultivated apples with the individual-based Bayesian clustering methods implemented in STRUCTURE 2.3.3 (Pritchard, Stephens, & Donnelly, 2000). STRUCTURE uses Markov chain Monte Carlo (MCMC) simulations to infer the proportion of ancestry of genotypes from K distinct clusters. The underlying algorithms attempt to minimize deviations from Hardy–Weinberg within clusters and linkage disequilibrium among loci. We ran STRUCTURE from $K = 1$ to $K = 15$. Based on 10 repeated runs of MCMC sampling from 500,000 iterations after a burn-in of 50,000 steps, we determined the amount of additional information explained by increasing K using the ΔK statistic (Evanno, Regnaut, & Goudet, 2005) as implemented in the online post-processing software Structure Harvester (Earl, 2012). However, ΔK provides statistical support for the strongest but not the finest population structure (Puechmaille, 2016). Natural populations can display hierarchical genetic structure, with fine-scale population structure. It has been shown that interpreting K values below the one corresponding to the finest substructure can lead to misinterpretations, especially with uneven sampling among populations (Puechmaille, 2016). Indeed, simulations have shown that STRUCTURE is not reliable for identifying groups that are not at Hardy-Weinberg equilibrium, *i.e.*, when the K is too low (Cunningham et al., 2020; Kalinowski, 2011; Puechmaille, 2016). For instance, when the

number of clusters is suboptimal, STRUCTURE does not always group the genetically closest clusters into the same cluster. The problem seems to be caused when the program is forced to cluster individuals into an inappropriately small number of clusters (Cullingham et al., 2020; Kalinowski, 2011; Puechmaille, 2016). A more powerful method for identifying the best K value is to look at the bar plots and choose the highest K value for which no clusters is only represented by admixed individuals, indicating that we have reached the highest K value for which new genuine clusters could be delimited (Chen et al., 2022; Cornille et al., 2013, 2012; Cornille et al., 2015; Fontaine, Gladieux, Hood, & Giraud, 2013; Vercken et al., 2010). The K value we therefore considered corresponded to the finest one, which can be higher than the K value of the strongest population structure level identified by ΔK .

We ran STRUCTURE for the whole dataset ($N = 550$) to investigate the population genetic structure among the Caucasian crab apple *M. orientalis*, the Caucasian and Iranian cultivated apples, *M. sieversii*, *M. domestica* and *M. baccata*. We further explored the genetic variation and differentiation among the genetic groups detected with STRUCTURE using three different methods. First, we ran a principal component analysis (PCA) for all individuals with the *dudi.pca* function from the *ade4* R package (Jombart & Ahmed, 2011). For the PCA, individuals that were assigned to a given cluster with a membership coefficient ≥ 0.85 were color-coded according to the respective color of each cluster, and admixed individuals (*i.e.*, individuals with a membership coefficient to any given cluster < 0.85) were colored in gray. We chose this threshold based on the distribution of the maximum membership coefficients inferred with STRUCTURE (see results). Second, we generated a neighbor-net with Splitstree v4 (Huson, 1998; Huson & Scornavacca, 2012), using the PCA color code. Third, we explored the relationships among populations identified with STRUCTURE (*i.e.*, clusters of individuals with a membership coefficient ≥ 0.85 to a given cluster) with a neighbor joining (NJ) tree (Huson, 1998; Huson & Scornavacca, 2012). The NJ tree and the neighbor-net tree were built using Nei's standard genetic distance (Nei, 1987) computed among individuals or populations with the Populations software v1.2.31 (<https://bioinformatics.org/populations/>).

We computed descriptive population genetic estimates for each population (*i.e.*, each cluster inferred with STRUCTURE, excluding admixed individuals with a membership coefficient < 0.85). We calculated allelic richness (A_R) and private allelic richness (A_P) with ADZE (Szpiech, Jakobsson, & Rosenberg, 2008) using standardized sample sizes of $N_{ADZE} = 7$ (one individual x

two chromosomes), corresponding to the minimal number of observations across populations. Heterozygosity (expected and observed), Weir and Cockerham F -statistics and deviations from Hardy–Weinberg equilibrium were calculated with Genepop v4.2 (Raymond & Rousset, 1995; Rousset, 2008).

Identification of crop-wild hybrids in the Caucasus and historical wild-wild gene flow in the Caucasian crab apple, *M. orientalis*

To assess the extent of crop-wild gene flow in the Caucasus, we removed *M. sieversii* and *M. baccata* from the dataset (resulting in a dataset with $N = 466$, Table S2) and ran STRUCTURE with the same parameters as above. We defined hybrids resulting from crop-to-wild introgression as *M. orientalis* trees assigned to the *M. domestica* or the Iranian or Armenian cultivated gene pools with a membership coefficient > 0.10 . We defined hybrids resulting from wild-to-crop introgression as cultivars assigned to any of the wild gene pools with a membership coefficient > 0.10 . We chose this threshold based on the distribution of the maximum membership coefficients inferred with STRUCTURE (see results).

After removing crop-wild hybrids, we estimated the extent of historical wild-wild gene flow in *M. orientalis* in the Caucasus using two methods. First, we tested whether there was a significant isolation-by-distance (IBD) pattern. We computed the correlation between $F_{ST}/(1-F_{ST})$ and the natural algorithm of geographic distance between geographical sites with SPAGeDI 1.5 (Hardy & Vekemans, 2002). Second, for each population, we computed the Nason’s kinship coefficient F_{ij} between pairs of individuals i and j (Loiselle, Sork, Nason, & Graham, 1995) with SPAGeDI 1.5 (Hardy & Vekemans, 2002), and regressed F_{ij} against the natural logarithm of geographic distance, $\ln(d_{ij})$, to obtain the regression slope b . We permuted the spatial position of individuals 9,999 times to test whether there was a significant spatial genetic structure between sites. We then calculated the Sp statistic, defined as $Sp = -bLd/(1-F_N)$, where F_N is the mean F_{ij} between neighboring individuals (Vekemans & Hardy, 2004), and $-bLd$ is the regression slope of F_{ij} against $\ln(d_{ij})$. A low Sp implies low spatial population structure, which suggests high historical gene flow and/or high effective population size.

We investigated the directionality of crop-wild, wild-wild and crop-crop gene flow (*i.e.*, immigration rate m_{ji} , the proportion of immigrants in population i that arrive from population j) over the past few generations with a Bayesian assignment method implemented in BAYESASS

v3.0 (Wilson & Rannala, 2003). As recommended by Wilson and Rannala (Wilson & Rannala, 2003), we run 3×10^6 iterations, sampled every 2,000 iterations with a discarded burn-in of 10^6 iterations, delta values were adjusted to 0.12 to ensure that chain swapping occurred in approximately 50% of the total iterations. We repeated five times the analysis with different random number seeds to evaluate consistency of results.

Approximate Bayesian computation to reconstruct the domestication history of apple

We used approximate Bayesian computation to test whether Iranian cultivated apples (see results below, we removed Armenian cultivated apples as they were represented by only four samples) were derived from the same domestication event as *M. domestica*. We used the ABC method based on a machine learning tool named “random forest” (ABC-RF) to perform model selection and parameter estimates (Estoup et al., 2018; Pudlo et al., 2016; Raynal et al., 2019). In brief, this method creates a “forest” of bootstrapped decision trees that ranks scenarios based on the summary statistics of the datasets. Some simulations are not used to build the trees and can thus be used to cross-validate the analysis by computing a “prior error rate”. This approach allows the comparison of complex demographic models (Pudlo et al., 2016) by comparing groups of scenarios with a specific type of evolutionary event with other groups with different types of evolutionary events instead of considering all scenarios separately (Estoup et al., 2018).

Using the ABC-RF framework, we compared different scenarios of domestication of Iranian cultivars, *i.e.*, with an origin from *M. domestica*, *M. sieversii* from Kazakhstan or *M. orientalis*. Populations were defined as the clusters detected with STRUCTURE with the whole dataset ($N = 550$), removing putative hybrid individuals (*i.e.*, individuals with a membership coefficient < 0.85 to any given cluster), and excluding *M. baccata* and Armenian cultivars (see Results part). We removed putative hybrids in order to retrace the divergence history and historical gene flow among populations; more recent admixture events being detectable directly from the STRUCTURE bar plots. We assumed bidirectional gene flow among wild and cultivated apple populations. The model parameters used were: the divergence time between X and Y populations (T_{X-Y}), the effective population size of population X (N_{E-X}) and the migration rate per generation between X and Y populations (m_{X-Y}). Prior values for divergence time were drawn from the log-uniform distribution bounded between the distributions used in the approximate Bayesian computations and are given in Table S3.

For all models, microsatellite datasets were simulated for 14 out of the 26 markers that had perfect repeats (Ch01h01, Ch01h10, Ch02c06, Ch02d08, Ch05f06, Ch01f02, Hi02c07, Ch02c09, Ch03d07, Ch04c07, Ch02b03b, MS06g03, Ch04e03, Ch02g01 (Cornille et al., 2013; Cornille et al., 2012; Patocchi, Fernández-Fernández, et al., 2009). We checked that the population structure inferred with 14 microsatellite markers did not differ significantly from the inferences obtained with 26 SSR markers (data not shown). We assumed a generalized stepwise model of microsatellite evolution (Slatkin, 1995). Mutation rates were allowed to vary across loci, with locus-specific mutation rates where μ is the mutation rate per generation, with a log-uniform prior distribution for μ (10^{-4} , 10^{-3}) (Table S3).

We used ABCtoolbox (Wegmann, Leuenberger, Neuenschwander, & Excoffier, 2010) with fastsimcoal 2.5 (Excoffier & Foll, 2011) to simulate datasets, using model parameters drawn from prior distributions (Table S3). We performed 8,000 simulations per scenario. For each simulation, we calculated three summary statistics per population with arlsumstats v 3.5 (Excoffier and Lischer 2010): H , the mean heterozygosity across loci, GW , the mean Garza-Williamson statistic over populations (Garza & Williamson, 2001) and the pairwise F_{ST} between populations (Weir & Cockerham, 1984).

We used the abcrf v.1.7.0 R statistical package (Pudlo et al., 2016) to carry out the ABC-RF analysis. This analysis provides a classification vote that represents the number of times a scenario is selected as the best one among n trees in the constructed random forest. For each ABC step, we selected the scenario, or the group of scenarios, with the highest number of classification votes as the best scenario, or best group of scenarios, among a total of 500 classification trees (Breiman, 2001). We computed the posterior probabilities and prior error rates (*i.e.*, the probability of choosing a wrong group of scenarios when drawing model index and parameter values from the priors of the best scenario) over 10 replicate analyses (Estoup et al., 2018) for each ABC step. We also checked visually that the simulated models were compatible with the observed dataset by projecting the simulated and the observed datasets onto the first two linear discriminant analysis (LDA) axes (Pudlo et al., 2016), and by checking that the observed dataset fell within the clouds of simulated datasets. We then calculated parameter inferences using the final selected model. The median and 90% confidence interval (CI, 5%-95%) are given for each model parameter estimate. Note that the ABC-RF approach includes the model checking step that was performed *a posteriori* in previous ABC methods.

Spatial pattern of genetic diversity in the Caucasian crab apple

We investigated spatial patterns of diversity in “pure” *M. orientalis*. To this aim, we excluded the crop-to-wild hybrids detected in the second STRUCTURE analysis (*i.e.*, excluding *M. baccata* and *M. sieversii*), as well as *M. domestica* and the Iranian and Armenian cultivars. Spatial patterns of genetic diversity in “pure” *M. orientalis* were visualized by mapping the variation across space (A_R) at 36 sites (*i.e.*, geographic locations for which at least five individuals were successfully genotyped for each marker, Table S2) with the geometry-based inverse distance weighted interpolation in QGIS (Quantum GIS, GRASS, SAGA GIS). We calculated allelic richness (A_R) and private allelic richness (A_P) per site with ADZE (Szpiech et al., 2008) using standardized sample sizes of $N_{ADZE} = 6$ (one individual x two chromosomes), corresponding to the minimal number of observations across sites.

Species distribution modeling

The BIOMOD2 R package (Thuiller, Georges, Engler & Breiner, 2016) was used to project past and present distributions of *M. orientalis* following the species distribution modeling methods of Leroy *et al.* (2014). A set of 19 bioclimatic variables from WorldClim.org was used in addition to monthly temperature and precipitation values. Climate data were obtained for past conditions from the last glacial maximum and for the current period between 1960 and 1990. The climate projection at the 2.5-minute spatial resolution from the CCSM4 global climate model was used (<https://www.worldclim.org/data/worldclim21.html#>), as we previously showed that it was the most accurate for apple trees (Cornille et al., 2013). Past and present distributions were projected using three modeling algorithms: a generalized linear model (GLM), a generalized additive model (GAM) and artificial neural networks (ANN).

The location of 339 “pure” *M. orientalis* trees (*i.e.*, individuals assigned to a wild apple gene pool with a membership coefficient > 0.9 , see results from the second STRUCTURE analysis) provided the longitude and latitude coordinates. Duplicate data points were removed, resulting in 57 presence points for *M. orientalis* (Table S4). We did not have absence data so we randomly selected pseudo-absences to serve as “absence” points for the model, and weighted presence and absence points equally as per Barbet-Massin *et al.* (2012). Models were calibrated using the set of bioclimatic variables and model evaluation was calculated with Jaccard’s indices. Ensemble model

forecasting was completed by pulling the average trend of the three modeling algorithms and retaining only the uncorrelated bioclimatic variables with a Pearson correlation threshold greater than 0.75 (Table S5). The model was run again using only variables with high predictive power.

Results

Clear population genetic structure among Caucasian wild and cultivated apples, *M. domestica* and *M. sieversii*

The ΔK statistic indicated that the strongest level of population subdivision was at $K = 3$ (Figures S1 and S2). At $K = 3$, a genetic cluster encompassed *M. orientalis* from Turkey, Russia, Armenia and *M. baccata* samples, a second cluster encompassed *M. sieversii* from Central Asia (Kazakhstan and Kyrgyzstan), and a third cluster comprised *M. domestica* and Iranian cultivars. However, several lines of evidence indicated that $K = 3$ did not correspond to a biologically relevant population structure. Indeed, *M. baccata* is known to be highly genetically differentiated from *M. orientalis*, *M. sieversii* and *M. domestica* ((Cornille et al., 2014, 2012) and see the PCA below), yet it clustered with *M. orientalis* at $K=3$. In addition, we observed multiple ΔK peaks, up to $K=12$ (Figure S1a) and further genetic subdivisions were revealed for $K > 3$ (Figure S2), with well-delimited and biologically meaningful clusters. For example, a population of *M. sieversii* admixed at $K=3$ formed a well-separated cluster without admixture at $K=12$ (the light blue cluster at $K=12$ in Figure S1). We therefore visually examined the bar plots (Figure S2) for K values that had ΔK peaks (Figure S1), and chose from these K values the highest one for which all clusters had fully assigned individuals, indicating the finest level of genetic subdivision. At $K = 12$, STRUCTURE identified twelve well-delimited clusters corresponding to species and/or geographic regions (Figures 1 and S2). We therefore considered these twelve clusters as the most relevant genetic structure to investigate the domestication history of the apple tree.

Among these twelve clusters were two distinct genetic clusters of *M. sieversii*, one from Kazakhstan and one from Kyrgyzstan (in two shades of light blue, respectively, Figure 1), and a specific genetic cluster of *M. baccata* (orange red). We identified seven distinct genetic groups of *M. orientalis*: a genetic group from the Western Caucasus (Russia, Turkey and northwestern Armenia; orange), a central Armenian group (blue), a southern Armenian group (brown), four genetic groups in Iran corresponding to two gene pools spread across the Zagros Forests (including

samples from the Lorestan province in light green and from the Kurdistan province in red), and two gene pools (pink and purple) spread across the Hyrcanian Forests (Figure 1).

The *M. domestica* apple cultivars formed a specific genetic group (yellow) that was well separated from the wild *M. orientalis* and the Iranian and Armenian cultivars. The Iranian apple cultivars formed two genetic clusters: one that included only cultivars (dark green) and another (purple) that included cultivated trees and wild individuals from the Hyrcanian Forests. We also detected Iranian cultivated trees that were highly admixed between the Iranian cultivated dark green cluster, the Iranian purple cluster, the *M. domestica* yellow cluster and with two other clusters (red and orange); the latter two included several wild *M. orientalis* individuals from the Zagros Forests in the Kurdistan province in Iran and the Western Caucasus, respectively (Figures 1 and S2). The four Armenian cultivated apple trees fell in the blue and orange clusters, which also included *M. orientalis* trees.

Genetic variation and differentiation among wild and cultivated apples

We assigned individuals with a membership coefficient > 0.85 (Figure S3) to a given cluster to the corresponding 12 populations to assess the genetic variation and differentiation among wild and cultivated apples.

The PCA showed that *M. baccata* was highly differentiated from the other genetic groups (Figure S4). The PCA without *M. baccata* showed that the European (yellow) and the two Iranian cultivated genetic clusters (green and purple) formed well differentiated gene pools (Figure 2a and S5). *Malus sieversii* from Kazakhstan was closer to the Armenian and Iranian wild apple than *M. sieversii* from Kyrgyzstan. The neighbor-net (Figure 2b) confirmed that the apple cultivars (*M. domestica* and the purple and dark green Iranian cultivar clusters) were distinct from the wild populations, with the exception of trees from the wild Hyrcanian purple population and *M. sieversii* from Kazakhstan; trees of *M. domestica* and *M. sieversii* from Kazakhstan, and of the wild and cultivated Hyrcanian purple population, were indeed intermingled in the neighbor net (Figure 2b). *Malus orientalis* from the Western Caucasus (orange) and Central and Southern Armenian (brown and blue) were closer to each other than to the Iranian wild apples (light green, red and pink), which clustered together. Most of the NJ tree nodes were not statistically supported, except three nodes that showed a sister relationship of 1) the two cultivated Iranian populations (green and purple), 2) *M. sieversii* from Kyrgyzstan and the Lorestan population, and, 3) among the three Armenian

populations (Figure S6). The close relationship between *M. domestica* and *M. sieversii* was not statistically supported but was demonstrated in previous studies (Cornille et al., 2019; Cornille, Giraud, Smulders, Roldán-Ruiz, & Gladieux, 2014; Duan et al., 2017, 2017; Harris, Robinson, & Juniper, 2002; Migicovsky et al., 2021).

The two cultivated apple populations from Iran were genetically highly differentiated (mean F_{ST} = 0.24, Table S6), and had lower levels of genetic diversity and fewer private alleles than *M. domestica* ($P < 0.01$, Tables 1 and S7); the purple gene pool displayed the lowest level of genetic diversity and the lowest number of private alleles. The close genetic relationship between apple trees sampled in the Hyrcanian Forests and cultivars from the purple gene pool (Figures 1 and 2), together with the lower levels of genetic diversity in these populations (Table 1), suggest that the trees sampled in the Hyrcanian Forests assigned to the purple genetic group may be feral. Alternatively, the cultivated trees from the purple gene pool may represent the first step of apple domestication in Iran, *i.e.*, wild trees cultivated by humans.

STRUCTURE analysis, PCA and neighbor net thus revealed insights into the history of apples in the Caucasus. First, cultivated apples in Iran (dark green and purple clusters) may have resulted from additional domestication events, distinct from that of *M. domestica*, as their respective clusters are not the most closely related in the PCA and the neighbor net. These additional domestication events could have involved *M. orientalis* as a progenitor in Iran; alternatively, cultivated apples in Iran could be derived from *M. sieversii* or *M. domestica* with subsequent local gene flow from *M. orientalis*. Second, the full membership of a substantial number of cultivated trees to genetic clusters of Iranian and Armenian *M. orientalis* suggests that wild trees are grown in orchards for consumption without any strong domestication process and/or that feral individuals occur (*e.g.*, the purple genetic cluster may represent a cultivated group that is also found in the wild as feral or an early domestication step). Third, the high level of admixture in several Iranian cultivated apple trees with wild *M. orientalis* gene pools indicates substantial wild-crop gene flow. Fourth, the spatial population structure of *M. orientalis* in the Caucasus may result from past range contraction and expansion associated with the last glacial maximum. We tested these hypotheses below. First, we estimated the extent of crop-wild and wild-wild gene flow in apple trees in the Caucasus (*i.e.*, excluding *M. baccata* and *M. sieversii*). Second, we inferred the anthropogenic divergence history of the Iranian cultivated apple using a coalescent-based inference method for assessing the probability of different domestication scenarios. Third, we investigated

both spatial genetic distribution of genetic diversity and modeled the past and present climatic niches of *M. orientalis* to test whether this species underwent past contraction and expansion in the Caucasus during past glaciations.

Substantial crop-wild, crop-crop and wild-wild gene flow in apples in the Caucasus

The second STRUCTURE analysis, focused on the Caucasus, revealed the same genetic clustering for wild apples and *M. domestica* at $K = 9$ (Figures S1 and S7) as in the previous analysis ($K = 12$) (Figures 1 and 2). At $K = 9$, we detected 150 apple genotypes that could be considered hybrids (*i.e.*, individuals assigned to a gene pool with a membership coefficient < 0.9 , this cut-off being chosen on the basis of the distribution of the cumulated membership coefficients for each individual at $K = 9$, Figure S8); these 150 hybrids represented 32% of the total dataset (Table 2). The Iranian cultivars had the highest proportion of hybrids (67%), mostly admixed with the wild and cultivated gene pools from Iran, but also with the *M. domestica* gene pool. Hybrids of the wild Armenian apple were mostly an admixture of the wild Armenian gene pools (*i.e.*, Western, Central and Southern), suggesting local gene flow between crop and wild populations.

We removed the 150 hybrids and all apple cultivars (Tables 2 and S2) and focused on gene flow within the “pure” Caucasian crab apple *M. orientalis*. We detected a significant but weak isolation by distance pattern across the Caucasus ($P < 0.001$, $R^2 = 0.07$, Figure S9), suggesting a high level of gene flow among the sampled geographic sites. We estimated Sp values for populations with at least five sampling sites and 20 individuals, *i.e.*, the Hyrcanian (pink) and the Central Armenian (blue) *M. orientalis* populations. Sp values were low but significant ($Sp_{\text{Hyrcanian_pink}} = 0.0076$, $Sp_{\text{Central_blue}} = 0.0027$, $P < 0.001$) suggesting a high level of historical gene flow within populations. However, the Sp value was higher for the Iranian population than for the Armenian population, suggesting a lower level of historical gene flow within the Hyrcanian (pink) population than the Central Armenian (blue) population. Contemporary estimates of gene flow further confirmed the occurrence of wild-to-wild, crop-wild and crop-crop gene flow in the Caucasus (Table S8 and Figure S10). Our results therefore are in agreement with the observations obtained above for the full dataset, suggesting substantial crop-crop, crop-wild and wild-wild gene flow in apples in Iran and the Caucasus.

Additional domestication events from *M. orientalis* in Iran inferred with ABC-RF

We defined the populations used in the ABC framework from the clusters detected with STRUCTURE at $K=12$ for 550 wild and cultivated apple accessions, excluding admixed individuals (*i.e.*, with a membership coefficient < 0.85 to any given cluster, Figure S3, as recent gene flow can easily be seen from visual inspection of the bar plots). We excluded *M. baccata* as it was the most genetically differentiated species (Figure S4 and Table S6), and the Armenian cultivars as they were represented by only four individuals. We included two *M. sieversii* populations (from Kazakhstan and Kyrgyzstan), the cultivated (green and purple) populations from Iran, the standard *M. domestica* population and the seven populations of *M. orientalis* (four populations from Iran and three populations from Armenia, Table 1). We also simulated an ancestral population. In total, we therefore simulated thirteen populations (Figures S11 and S12).

To keep the number of scenarios tractable, we fixed the divergence histories of the wild apple populations as previously known and inferred from the trees (Figures 2b and S6), and assumed bidirectional gene flow among populations which is congruent with the high levels of crop-wild and crop-crop gene flow shown above. We built eight groups of scenarios to test whether the two Iranian cultivated apple populations diverged from the same wild population or whether they diverged from different populations (Figures S11 and S12). In order to assess the origin of the wild purple population, these eight groups of scenarios were simulated under two alternative hypotheses (Figures S11 and S12, respectively): the wild purple population was either assumed to have diverged from the ancestral population (Figure S11), or the Iranian wild and crop purple populations were assumed to be sister groups (Figure S12). A total of 46 scenarios was therefore simulated, representing 16 groups of hypotheses (Figures S11 and S12).

For each step of the ABC-RF approach, the projection of the reference table and the observed datasets onto the two LDA axes that explained most of the variance of the summary statistics showed that the observed data fell within the distribution of the simulated summary statistics (Figure S13), forming distinct clouds for each scenario or groups of scenarios. Visual inspection of the LDA plots indicated that we had enough power to discriminate and select scenarios; results were subsequently validated by the ABC-RF inferences presented below.

We used a five-step nested ABC-RF approach. In the first step, we tested the divergence history of the purple wild and cultivated populations. All ten replicate ABC-RF analyses supported a sister relationship between the cultivated and wild purple populations (groups of scenarios 9 to 16, Figure S11, an average of 282 votes out of the 500 RF-trees; posterior probabilities = 65%,

prior error rate = 0.98%, Table S9). In a second step, we compared the remaining groups (groups 9 to 16) to reconstruct the history of the Iranian cultivated apple populations. All ten replicate ABC-RF analyses supported that the two Iranian cultivated apple populations (green and purple) diverged from the same wild apple population (groups 9 to 16, an average of 296 votes out of the 500 RF-trees; posterior probabilities = 82%, prior error rate = 1.1%, Table S10). In the third step, we compared the remaining groups (groups 9 to 12) to test whether these Iranian cultivated apple populations diverged from *M. sieversii*, *M. domestica* or *M. orientalis*. All ten replicate ABC-RF analyses supported that the two Iranian cultivated apple populations originated from *M. orientalis* (groups 11 and 12, an average of 290 votes out of the 500 RF-trees; posterior probabilities = 79%, prior error rate = 0.2%, Table S11). In the fourth step, we compared the remaining groups (groups 11 and 12) to test whether the two Iranian cultivated apple populations were derived from *M. orientalis* from Iran or from Armenia. All ten replicate ABC-RF analyses supported that the two Iranian cultivated apple populations originated from *M. orientalis* from Iran (group 11, an average of 316 votes out of the 500 RF-trees; posterior probabilities = 92%, prior error rate = 0.10%, Table S12). For step five, we compared the last three remaining individual scenarios to select the final model. All ten replicate ABC-RF analyses supported that the two Iranian cultivated apple populations originated from the Iranian red *M. orientalis* population (sc45, an average of 316 votes out of the 500 RF-trees; posterior probabilities = 90%, prior error rate = 0.11%, Table S13). This nested ABC approach avoids comparing too many complex models with too many parameters, and is more powerful than testing all scenarios individually to disentangle the main evolutionary events characterizing demography and divergence (Estoup et al., 2018).

ABC-RF inferences therefore provided support for additional apple domestication events in Iran, from the red *M. orientalis* population (Figures 3 and S12). The time of domestication was inferred to be *ca.* 826 years ago (confidence interval (CI): 130-5,030) for the green Iranian crop population and *ca.* 803 ya (CI: 160-3,969) for the purple crop population, respectively; by contrast *M. domestica* was inferred to have diverged from *M. sieversii* *ca.* 3,898 ya (CI: 270-7,670) (Table S14, Figure 3). The divergence time between the wild and the crop purple populations was estimated to be 418 ya (CI: 100-2,500). Note that the confidence intervals were relatively high for some parameters (*e.g.*, migration rates) as illustrated by the high normalized mean absolute error values (NMAE). High posterior probabilities and low *prior* error rates for model choice

nevertheless indicate high support for the final model, even if some parameters such as migration rates cannot be precisely estimated.

Range expansion and contraction of the wild apple *M. orientalis* in the Caucasus associated with the past glaciations

We investigated the spatial variation of genetic diversity and used ecological niche modeling to test the existence of past contraction and expansion of the range of the wild apple in the Caucasus. After removing the 150 crop-wild hybrids identified from the second STRUCTURE analysis, we found a significant positive correlation between longitude and allelic richness (Figure S14, average adjusted $R^2 = 0.66$, $P < 0.0001$) and a significant negative correlation between latitude and allelic richness (Figure S14, average adjusted $R^2 = -0.43$, $P < 0.001$). We also found that the Western (orange) population had the highest level of allelic richness (Tables 1 and S15 and Figure 4). The level of allelic richness was also significantly lower in the wild apple populations from the Zagros Forests, *i.e.*, Lorestan (light green) and Kurdistan (red), than in the other wild populations (Tables 1 and S7). The Western Caucasus may therefore have been a glacial refugium in the past. In addition to high levels of genetic diversity in the west, across northeastern Turkey and the Lesser Caucasus mountains in Armenia, we observed local hotspots of genetic diversity in the Hyrcanian Forests and the High Caucasus mountains (Figure 4), suggesting that these mountainous regions may have been potential glacial refugia.

Ecological niche modeling further indicated past contraction and expansion of the *M. orientalis* range. Model performance as assessed with AUC and TSS was high (Table S15), indicating that the ANN, GLM and GAM algorithms fitted the data well (Allouche, Tsoar, & Kadmon, 2006; Fieldings & Bell, 1997; Monserud & Leemans, 1992). The following six bioclimatic variables were found to have high predictive power: mean diurnal range temperature (bio2), temperature seasonality (bio4), minimum temperature of the wettest quarter (bio8), minimum temperature of the driest quarter (bio9), annual precipitation (bio12) and precipitation of the coldest quarter (bio19). These bioclimatic variables were used to calibrate the models to predict the past and present distribution of *M. orientalis*. The MIROC model (Figure 4) predicted that the areas suitable for *M. orientalis* during the LGM contracted to the western Lesser Caucasus and northeastern Turkey along the Black Sea and into the Colchis region, and also in the eastern part of the Hyrcanian Forests, near Azerbaijan, in agreement with the genetic data (Figure 4). The

climatic model therefore suggested that populations of the Caucasian wild apple *M. orientalis* may have been maintained in at least two glacial refugia.

Discussion

This study provides insights into the natural and anthropogenic divergence history of apples in hotspots of crop diversity, the Caucasus and Iran. Our results reveal that the evolution of the domesticated apple involved an additional wild species, *M. orientalis*. We identified two distinct cultivated gene pools in Iran that were well differentiated from the standard *M. domestica* apple cultivars and were derived from the same wild Iranian *M. orientalis* population. In addition, several cultivated apple genotypes from the Caucasus were found to belong to the Caucasian *M. orientalis* gene pool, suggesting that local farmers use the Caucasian crabapple for cultivation, which may represent the early stages of domestication. *Malus orientalis* has also contributed to the Caucasian cultivated apple germplasm through wild-to-crop introgression; a similar process has been previously described in apples in Europe (Cornille et al., 2014, 2012). Conversely, crop-to-wild gene flow, which has been reported in Europe (Cornille et al., 2015), was also detected in the Caucasus. We also showed that « pure » *M. orientalis* in this region displays a clear spatial genetic structure with at least seven populations. The combination of niche modeling and population genetics approaches suggested that these populations resulted from range contraction and expansion associated with the past glaciations. Iran is therefore an additional center of apple domestication, where distinct cultivated gene pools evolved from their progenitor *M. orientalis*. *Malus orientalis* has also contributed to cultivated gene pools in the Caucasus and Iran through wild-to-crop gene flow. Our results thereby reinforce the view that apple tree domestication has involved multiple wild apple species from multiple geographic regions, a pattern that has also been found in apricot and pear trees (Groppi, Liu, Cornille, Decroocq, & Decroocq, 2021; Liu et al., 2019; Wu et al., 2018). We also found evidence of substantial hybridization between domestic and wild forms, and this has also been described in other fruit trees (Groppi et al., 2021; Liu et al., 2019; Wu et al., 2018). In addition, this study highlights the impact of climate change on the natural divergence of a wild fruit tree and provides a starting point for apple conservation and breeding programs in the Caucasus and Iran.

Additional events of apple domestication from *M. orientalis* in Iran

The occurrence of cultivated populations in Iran which are genetically differentiated from the cultivated apple *M. domestica* and are derived from *M. orientalis*, suggests that Iran is an additional center of apple domestication. The existence of two Iranian cultivated groups (green and purple), genetically closely related, suggests that they diverged from the same wild population, perhaps corresponding to two successive domestication steps, or represent separate domestication events from the same progenitor. We tested these hypotheses and inferred more precisely the Iranian cultivated apple domestication history using coalescent-based methods combined with approximate Bayesian computation. This confirmed that the two Iranian cultivated populations diverged from a *M. orientalis* population in Iran, most likely from the Kurdistan province (red population). Thus, despite the spread of the cultivated apple *M. domestica* along the Silk Trade Routes that crossed Iran and the South Caucasus to reach Turkey (Canepa, 2010; Spengler, 2019), specific domestication events in this region have resulted in local cultivated apple gene pools. It is still not clear, however, whether these two cultivated apple populations represent *de novo* domestication events. Distinct genetic ancestries may not reflect independent *de novo* domestication, but may instead represent a single domestication event with multiple origins (Choi et al., 2017; Gros-Balthazard & Flowers, 2021). The occurrence of independent domestication events in many crop species is a source of ongoing debate (Besnard et al., 2018; Choi et al., 2017; Gros-Balthazard & Flowers, 2021). In apricot and pear trees, there is evidence of independent domestication events in Europe and Asia, with distinct cultivated gene pools of different ancestries which display different genomic regions showing footprints of selection (Groppi et al., 2021; Liu et al., 2019; Wu et al., 2018). Here, our data shows that there are distinct cultivated apple populations, with different geographic origins and which are derived from distinct ancestral gene pools. Microsatellite markers however do not allow to investigate history of selection in populations and have lower power than nucleotide sequences to infer demographic history as mutation models for microsatellite markers are less accurate than for nucleotide sequences (Putman & Carbone, 2014). The low bootstrap support for some of the nodes of the NJ tree and conflicting results with the ABC inferences may be due to the limited power of microsatellite markers to resolve relationships among genetically close populations that exchanged gene flow. The resolution of NJ tree nodes can indeed depends on the number of loci (Koskinen, Hirvonen, Landry, & Primmer, 2004; Takezaki & Nei, 2008), and when admixture among populations occurred (Cavalli-Sforza L. Luca, 1998), as seen in the apple tree. While inferences with ABC coalescent simulate

admixture and gene flow among populations, NJ tree does not. ABC inferences are thus more powerful to infer the history of apple domestication, as actually shown by high posterior probabilities and low prior error rates. Sequencing genomes of Iranian and European varieties will likely give a more accurate view of the domestication history of apple in the Caucasus and Iran. First, it would help to get more genetic markers and resolve the relationships among apple populations (X. Chen et al., 2022). Second, the existence of distinct genomic regions under selection in the European and Iranian cultivated gene pools, for instance related to apple taste or size, would be a hallmark of multiple *de novo* domestication events, as shown in apricot and pear trees (Groppi et al., 2021; Wu et al., 2018). In addition, a broader view of the domestication history of the apple tree will require additional sampling of *M. sylvestris*, which is another contributor to the *M. domestica* gene pool (Cornille et al., 2014, 2012), as well as of local varieties across Eurasia, which are missing in this study.

Wild-to-crop gene flow from *M. orientalis* to local cultivars and cultivation of the local wild apple in the Caucasus

The Caucasian crab apple has considerably contributed to the Caucasian apple germplasm through wild-to-crop introgression. We found evidence of substantial wild-crop and crop-crop gene flow in the Caucasus. Indeed, we found that 41.6% of Iranian cultivars were introgressed by local wild apple gene pools or were an admixture of two cultivated gene pools. This extensive wild-to-crop and crop-crop gene flow is strikingly similar to the pattern documented in apples in Europe. *Malus sylvestris* has been shown to be a significant contributor to the *M. domestica* gene pool through recurrent and recent hybridization and introgression events ever since the cultivated apple was introduced in Europe by the Greeks around 1,500 years ago (Cornille et al., 2012). Conversely, substantial crop-to-wild gene flow has been reported from *M. domestica* to *M. sylvestris* (Cornille et al., 2015). Similarly, we found many crop-to-wild hybrids (*M. orientalis* introgressed with *M. domestica*) in the forests of Armenia and Iran. Extensive gene flow has been found during the domestication of other fruit trees (Arroyo-García et al., 2006; Cornille et al., 2012; Decroocq et al., 2016; Diez et al., 2015; Duan et al., 2017; Liu et al., 2019; Meyer, Duval, et al., 2012; Myles et al., 2011). The evolutionary consequences of crop-to-wild gene flow remains unclear in fruit trees (Feurtey, Cornille, Shykoff, Snirc, & Giraud, 2017); the extent to which crop-to-wild gene flow may threaten the Caucasian crab apple remains to be tested.

Our findings also suggest that farmers in the Caucasus and Iran grow *M. orientalis* rather than *M. domestica*. Indeed, the four Armenian cultivars share their gene pools with the Western (orange) and Central (blue) Caucasian crabapple populations, and some Iranian cultivars are fully assigned to wild populations (red and purple) in Iran. Coalescent-based methods suggest that the wild purple population found in the forest is feral or results from a recent domestication event. It is difficult to discriminate between these two hypotheses using only genotypes, as both predict a sister relationship between the wild and cultivated purple populations. Individual isolated trees belonging to the purple population are found in natural mountainous areas close to fruit trees grafted on *Crataegus*, although they are not themselves cultivated (personal observation. H. Yousefzadeh). In addition, the wild purple population found in the forest had a lower level of genetic diversity than the other wild Iranian populations. Altogether, this may suggest that this population has recently escaped from cultivation. Like domestication, feralization can be seen as a process that is accompanied by admixture and introgression, and can be accompanied by a range of genetic, phenotypic and demographic changes (Mabry, Rowan, Pires, & Decker, 2021). Feral populations have also been documented in other fruit trees, including olive (Besnard et al., 2018), almond (Balaguer-Romano et al., 2021) and apricot (Spengler, Chang, & Tourtellotte, 2013). However, the trees found in the Hyrcanian Forests could also be the result of a first domestication step, as wild apples are widely used for cultivation in Iran (personal communication H. Yousefzadeh and from our results). Additional phenotypic data to identify shifts in phenotypic traits associated with domestication or feralization are required to further disentangle these hypotheses. The cultivation of wild genotypes (here, from the red or purple populations) would not be surprising as *M. orientalis* can grow in mountainous areas, is highly resistant to pests, diseases and drought (Amirchakhmaghi et al., 2018; Büttner, 2001; Höfer et al., 2013; Volk et al., 2008), and has high-quality fruits that have several features that are intermediate between those of *M. sylvestris* and *M. sieversii* (Cornille et al., 2014). The use of local wild apples has also been documented in Europe for specific purposes at different times in history (Tardío, Arnal, & Lázaro, 2020).

The natural divergence history of the Caucasian wild apple was shaped by the past glaciations

The climatic variations since the last glacial maximum, along with the landscape features of the Caucasus, have likely shaped the population structure and diversity of the Caucasian wild apple.

We identified seven populations of *M. orientalis* in the Caucasus and Iran: one highly genetically differentiated population in the Western Caucasus (Turkey, Russia and northwestern Armenia), two in Armenia (a southern and a Central population) and four in Iran, including two in the Zagros Forest (one in the Kurdistan province and one in the Lorestan province) and two in the Hyrcanian Forests bordering the southern Caspian Sea. These wild apple populations likely arose from isolation in several refugia during the last glacial maximum. This hypothesis is supported by the observation of a large hotspot of genetic diversity located in the Western Caucasus, and several local hotspots of genetic diversity in Armenia and the Hyrcanian Forests (Zazanashvili et al., 2020). Ecological niche modeling further supported the existence of strong contractions in the range of *M. orientalis* in the Western Caucasus bordering the Black Sea (including the Colchis region), as well as in the Lesser Caucasus and in some parts of the Hyrcanian Forests. Additional samples from the Western Caucasus are required to confirm this hypothesis. These glacial refugia have been described in relation to other species (Parvizi et al., 2019). Indeed, two refugia are recognized in the Caucasus (Tarkhnishvili et al. 2012; Yousefzadeh et al. 2012; Bina et al. 2016; Aradhya et al. 2017): a major forest refugium between the western Lesser Caucasus and northeastern Turkey (including the Colchis region in the catchment basin of the Black Sea) and the Hyrcanian refugium at the southern edge of the Caucasus. Further sampling of *M. orientalis* in the far Western and Eastern Caucasus and genotyping with the same microsatellite markers are needed to uncover the role of these two refugia in the divergence history of *M. orientalis*.

We also found that the natural divergence history of the Caucasian wild apple involved gene flow across the Caucasus. The weak but significant isolation-by-distance pattern further supports the existence of substantial gene flow among wild apple populations in the Caucasus. Widespread gene flow during divergence associated with the last glacial maximum has been documented for another wild apple relative *M. sylvestris* (Cornille et al., 2013). Calculation of the *Sp* parameter within the largest populations revealed high levels of historical gene flow within populations. *Sp* can also be used to compare the dispersal capacities of *M. orientalis* with that of other plants (Cornille et al., 2013; Cornille, Gladieux, & Giraud, 2013; Vekemans & Hardy, 2004). The Caucasian wild apple showed dispersal capacities that were similar to previous estimates in other wild apple species and lower than that of wind-dispersed trees. Wild apples can disperse over kilometers (Cornille et al., 2015; Feurtey, Cornille, Shykoff, Snirc, & Giraud, 2017). The spatial population structure was somewhat stronger in Iran than in Armenia, suggesting lower levels of

gene flow in the Hyrcanian population. In addition to having a stronger genetic structure, the Iranian populations had lower genetic diversity than the Armenian populations, especially the Zagros and Kurdistan populations. In Iran, traditional animal husbandry is a widespread practice (Soofi et al., 2018). Such intensive farming environments may lead to forest fragmentation and may impact wild apple populations, which form low density populations. The future of Iranian wild apple populations, especially in the south where genetic diversity is low, may depend on our ability to protect them through conservation programs. However, *M. orientalis* in the Caucasian ecoregion is highly diverse and further investigations and additional sampling are necessary, as well as a better assessment of its species status and genetic relationship with *M. sieversii*. Here, the microsatellite markers yielded a neighbor-net with intermingled *M. sieversii* and *M. orientalis* from Iran, but the microsatellite markers may lack power to discriminate closely related genetic clusters, as discussed above. A better understanding of the functional genetic diversity and of the ecological relationships of wild apples in their ecosystem are needed for developing and implementing effective conservation genetic strategies in this region (Teixeira & Huber, 2021), and is crucial to better understand crop domestication. This question needs to be resolved urgently, as *M. sieversii* is endangered across its distribution (Omasheva et al., 2017; Zhang, Li, & Li, 2018).

Conclusion

We identified Iran as a key center in the evolution and domestication of apple, and *M. orientalis* as a contributor to the evolutionary history of cultivated apples in Iran and in the Caucasus. We also provided insights into the processes underlying the natural divergence of this emblematic wild species and identified several populations that could be the target of conservation programs. Our study revealed the role of gene flow and human practices in natural and anthropogenic divergence processes of an emblematic fruit tree in the Caucasus and Iran. Our results are consistent with those reported for other woody perennials, including apricot (Groppi et al., 2021; Liu et al., 2019), olive (Besnard et al., 2018; Diez et al., 2015), pear (Volk & Cornille, 2019; Wu et al., 2018), and date palm (Flowers et al., 2019). This study also supports the view that the domestication of fruit trees has often been a geographically diffuse and protracted process, involving multiple, geographically disparate origins of domestication, as also reported for example in apricots and pears (Groppi et al., 2021; Wu et al., 2018).

Acknowledgements

We thank the Franco-Iranian Campus France program « Gundhishapur » 2016-2018, the Institut Diversité Écologie et Évolution du Vivant (IDEEV), ATIP-Avenir for funding, and the ANR JCJC “PLEASURE”. We thank Bolotbek Tagaev (Sustainable Livelihoods Coordinator of FFI-Kyrgyzstan) for sampling and prospection, Fauna & Flora International and more specifically the Global Trees Campaign (GTC) Program. We also thank Adrien Falce, Olivier Langella and Benoit Johannet for help and support on the INRAE-Génétique Quantitative et Evolution- Le Moulon lab cluster and the genotyping platform GENTYANE INRA UMR 1095. We thank the INRAE MIGALE bioinformatics platform (<http://migale.jouy.inra.fr>) for providing help and support, in particular Véronique Martin, Eric Montaubon and Valentin Loux. We also thanks Céline Bellard for her advices for ecological niche modeling analyses.

Data Availability

SSR data are available on the ZENODO repository 10.5281/zenodo.6981530. Scripts are available https://forgemia.inra.fr/amandine.cornille/caucasian_apple_domestication_ssr

Author Contributions

AC, HY conceived and designed the experiments; AC, HY obtained funding; HB, HY, SF, HBa, IG, AN, AC, JS, DG, AK sampled the material; AV, CR, AR, MF performed the molecular work; AC, HB analyzed the data; AC, HB: wrote the original draft and preparation of the figures; AC, HB, HY, TG, XC, IG, AN and all co-authors: gave critical inputs in final draft and revisions.

References

- Abdollahi, H. (2021). An illustrated review on manifestation of pome fruit germplasm in the historic miniatures of ancient Persia. *Genetic Resources and Crop Evolution*, 68(7), 2775–2791. doi: 10.1007/s10722-021-01244-y
- Allouche, O., Tsoar, A., & Kadmon, R. (2006). Assessing the accuracy of species distribution models: Prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, 43(6), 1223–1232.
- Amirchakhmaghi, N., Yousefzadeh, H., Hosseinpour, B., Espahbodi, K., Aldaghi, M., & Cornille, A. (2018). First insight into genetic diversity and population structure of the Caucasian wild apple (*Malus orientalis* Uglitzk.) in the Hyrcanian forest (Iran) and its resistance to apple scab and powdery mildew. *Genetic Resources and Crop Evolution*, 65(4), 1255–1268. doi: 10.1007/s10722-018-0611-z
- Anderson, E., . (2005). *Plants, man, and life: Illustrated*. Mineola, N.Y.: Dover Publications. /z-wcorg/.
- Arroyo-García, R., Ruiz-García, L., Bolling, L., Ocete, R., López, M. A., Arnold, C., ... Martinez-Zapater, J. M. (2006). Multiple origins of cultivated grapevine (*Vitis vinifera* L. ssp. *Sativa*) based on chloroplast DNA polymorphisms. *Molecular Ecology*, 15(12), 3707–3714. doi: 10.1111/j.1365-294X.2006.03049.x
- Asanidze, Z., Akhalkatsi, M., Henk, A. D., Richards, C. M., & Volk, G. M. (2014). Genetic relationships between wild progenitor pear (*Pyrus* L.) species and local cultivars native to Georgia, South Caucasus. *Flora: Morphology, Distribution, Functional Ecology of Plants*, 209(9), 504–512. doi: 10.1016/j.flora.2014.06.013
- Bacles, C. F. E., & Jump, A. S. (2011). Taking a tree's perspective on forest fragmentation genetics. *Trends in Plant Science*, 16(1), 13–18. doi: 10.1016/j.tplants.2010.10.002

842 Bai, X. N., & Spitkovsky, A. (2010). Uncertainties of modeling gamma-ray pulsar light curves
 843 using vacuum dipole magnetic field. *Astrophysical Journal*, 715(2), 1270–1281. doi:
 844 10.1088/0004-637X/715/2/1270

845 Bailey-Serres, J., Parker, J. E., Ainsworth, E. A., Oldroyd, G. E. D., & Schroeder, J. I. (2019).
 846 Genetic strategies for improving crop yields. *Nature*, 575(7781), 109–118. doi:
 847 10.1038/s41586-019-1679-0

848 Balaguer-Romano, R., Barea-Marquez, A., Ocaña-Calahorra, F. J., Gomez, J. M., Schupp, E. W.,
 849 Zhang, J., & de Casas, R. R. (2021). The potential role of synzoochory in the
 850 naturalization of almond tree. *Basic and Applied Ecology*, 50, 97–106. doi:
 851 10.1016/j.baae.2020.11.004

852 Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absence for
 853 species distribution models: How, where and how many? *Methods in Ecology and*
 854 *Evolution*, in press(3), 327–338. doi: <https://doi.org/10.1111/j.2041-210X.2011.00172.x>

855 Besnard, G., Terral, J. F., & Cornille, A. (2018). On the origins and domestication of the olive: A
 856 review and perspectives. *Annals of Botany*, 121(3), 385–403. doi: 10.1093/aob/mcx145

857 Bina, H., Yousefzadeh, H., Ali, S. S., & Esmailpour, M. (2016). Phylogenetic relationships,
 858 molecular taxonomy, biogeography of *Betula*, with emphasis on phylogenetic position of
 859 Iranian populations. *Tree Genetics and Genomes*, 12(5). doi: 10.1007/s11295-016-1037-4

860 Bowe, Patrick. (2004). *Gardens of the Roman world*. Los Angeles: J. Paul Getty Museum. /z-
 861 wcorg/.

862 Brandenburg, J. T., Mary-Huard, T., Rigauill, G., Hearne, S. J., Corti, H., Joets, J., ... Tenailon,
 863 M. I. (2017). Independent introductions and admixtures have contributed to adaptation of
 864 European maize and its American counterparts. *PLoS Genetics*, 13(3), 1–30. doi:
 865 10.1371/journal.pgen.1006666

866 Breiman, L. (2001). Random Forests. *Machine Learning*, 45(1), 5–32. doi:
867 10.1023/A:1010933404324

868 Browicz, K. (1969). Amygdalus. *Flora Iranica*, 66, 166–168.

869 Büttner, R. (2001). Malus. *Hanelt P, Institute of Plant Genetics and Crop Plant Research (Eds)*
870 *Mansfelds Encyclopedia of Agricultural and Horticultural Crops*, 471–482.

871 Canepa, M. (2010). Distant displays of power: Understanding Cross-Cultural Interaction Among
872 the Elites of Rome, Sasanian Iran, and Sui-Tang China. *Ars Orientalis*, 38, 121–154.
873 JSTOR. Retrieved from JSTOR.

874 Castañeda-Álvarez, N. P., Khoury, C. K., Achicanoy, H. A., Bernau, V., Dempewolf, H.,
875 Eastwood, R. J., ... Toll, J. (2016). Global conservation priorities for crop wild relatives.
876 *Nature Plants*, 2(4), 16022. doi: 10.1038/nplants.2016.22

877 Cavalli-Sforza L. Luca. (1998). The Chinese Human Genome Diversity Project. *Proceedings of*
878 *the National Academy of Sciences*, 95(20), 11501–11503. doi: 10.1073/pnas.95.20.11501

879 Chen, J., Li, L., Milesi, P., Jansson, G., Berlin, M., Karlsson, B., ... Lascoux, M. (2019).
880 Genomic data provide new insights on the demographic history and the extent of recent
881 material transfers in Norway spruce. *Evolutionary Applications*, 12(8), 1539–1551. doi:
882 10.1111/eva.12801

883 Chen, X., Cornille, A., An, H., Xing, L., Ma, J., Zhao, C., ... Dong, Z. (2022). The East Asian
884 wild apples, *Malus baccata* (L.) Borkh and *Malus hupehensis* (Pamp.) Rehder., are
885 additional contributors to the genomes of cultivated European and Chinese varieties.
886 *Molecular Ecology*, 0, 1–15. doi: 10.1111/mec.16485

887 Choi, J. Y., Platts, A. E., Fuller, D. Q., Hsing (邢禹依), Y.-I., Wing, R. A., & Purugganan, M. D.
888 (2017). The Rice Paradox: Multiple Origins but Single Domestication in Asian Rice.
889 *Molecular Biology and Evolution*, 34(4), 969–979. doi: 10.1093/molbev/msx049

890 Cornille, A., Antolín, F., Garcia, E., Vernesi, C., Fietta, A., Brinkkemper, O., ... Roldán-Ruiz, I.
891 (2019). A Multifaceted Overview of Apple Tree Domestication. *Trends in Plant Science*,
892 24(8), 770–782. doi: <https://doi.org/10.1016/j.tplants.2019.05.007>

893 Cornille, A., Giraud, T., Bellard, C., Tellier, A., Le Cam, B., Smulders, M. J. M., ... Gladieux, P.
894 (2013). Post-glacial recolonization history of the European crabapple (*Malus sylvestris*
895 Mill.), a wild contributor to the domesticated apple. *Molecular Ecology*, 22(8), 2249–
896 2263.

897 Cornille, A., Gladieux, P., Smulders, M. J. M., Roldán-Ruiz, I., Laurens, F., Le Cam, B., ...
898 Giraud, T. (2012). New insight into the history of domesticated apple: Secondary
899 contribution of the European wild apple to the genome of cultivated varieties. *PLoS*
900 *Genet*, 8(5), e1002703.

901 Cornille, Amandine, Feurtey, A., Gélín, U., Ropars, J., Misvanderbrugge, K., Gladieux, P., &
902 Giraud, T. (2015). Anthropogenic and natural drivers of gene flow in a temperate wild
903 fruit tree: A basis for conservation and breeding programs in apples. *Evolutionary*
904 *Applications*, 8(4), 373–384. doi: 10.1111/eva.12250

905 Cornille, Amandine, Giraud, T., Smulders, M. J. M., Roldán-Ruiz, I., & Gladieux, P. (2014). The
906 domestication and evolutionary ecology of apples. *Trends in Genetics*, 30(2), 57–65. doi:
907 10.1016/j.tig.2013.10.002

908 Cornille, Amandine, Gladieux, P., & Giraud, T. (2013). Crop-to-wild gene flow and spatial
 909 genetic structure in the closest wild relatives of the cultivated apple. *Evolutionary*
 910 *Applications*, 6(5), 737–748. doi: 10.1111/eva.12059
 911 Cornille, Amandine, Gladieux, P., Smulders, M. J. M., Roldán-Ruiz, I., Laurens, F., Le Cam, B.,
 912 ... Giraud, T. (2012). New insight into the history of domesticated apple: Secondary
 913 contribution of the European wild apple to the genome of cultivated varieties. *PLoS*
 914 *Genetics*, 8(5). doi: 10.1371/journal.pgen.1002703
 915 Cullingham, C. I., Miller, J. M., Peery, R. M., Dupuis, J. R., Malenfant, R. M., Gorrell, J. C., &
 916 Janes, J. K. (2020). Confidently identifying the correct K value using the ΔK method:
 917 When does $K = 2$? *Molecular Ecology*, 29(5), 862–869. doi: 10.1111/mec.15374
 918 Curtis, J., Tallis, N., Andre-Salvini, B., André-Salvini, B., Museum, B., & Armbruster, B. (2005).
 919 *Forgotten Empire: The World of Ancient Persia*. University of California Press. Retrieved
 920 from <https://books.google.fr/books?id=kJnaKu9DdNEC>
 921 Decroocq, S., Cornille, A., Tricon, D., Babayeva, S., Chague, A., Eyquard, J. P., ... Decroocq, V.
 922 (2016). New insights into the history of domesticated and wild apricots and its
 923 contribution to Plum pox virus resistance. *Molecular Ecology*, 25(19), 4712–4729. doi:
 924 10.1111/mec.13772
 925 Delplancke, M., Alvarez, N., Espíndola, A., Joly, H., Benoit, L., Brouck, E., & Arrigo, N. (2011).
 926 Gene flow among wild and domesticated almond species: Insights from chloroplast and
 927 nuclear markers. *Evolutionary Applications*, 5(4), 317-329.
 928 Diez, C. M., Trujillo, I., Martinez-Urdiroz, N., Barranco, D., Rallo, L., Marfil, P., & Gaut, B. S.
 929 (2015). Olive domestication and diversification in the Mediterranean Basin. *New*
 930 *Phytologist*, 206(1), 436–447. doi: 10.1111/nph.13181

931 Duan, N., Bai, Y., Sun, H., Wang, N., Ma, Y., Li, M., ... Chen, X. (2017). Genome re-
 932 sequencing reveals the history of apple and supports a two-stage model for fruit
 933 enlargement. *Nature Communications*, 8(1). doi: 10.1038/s41467-017-00336-7

934 Earl, D. A. (2012). STRUCTURE HARVESTER: a website and program for visualizing
 935 STRUCTURE output and implementing the Evanno method. *Conservation Genetics*
 936 *Resources*, 4(2), 359–361.

937 Estoup, A. A., Raynal, L., Verdu, P., & Marin, J.-M. (2018). Model choice using Approximate
 938 Bayesian Computation and Random Forests: Analyses based on model grouping to make
 939 inferences about the genetic history of Pygmy human populations. *Journal de La Societe*
 940 *Française de Statistique*, 159(3), 167–190.

941 Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals
 942 using the software STRUCTURE: a simulation study. *Molecular Ecology*, 14(8), 2611–
 943 2620.

944 Excoffier, L., & Foll, M. (2011). fastsimcoal: A continuous-time coalescent simulator of genomic
 945 diversity under arbitrarily complex evolutionary scenarios. *Bioinformatics*, 27(9), 1332–
 946 1334. doi: 10.1093/bioinformatics/btr124

947 Excoffier, L., Foll, M., & Petit, R. J. (2009). Genetic Consequences of Range Expansions. *Annual*
 948 *Review of Ecology, Evolution, and Systematics*, 40(1), 481–501. doi:
 949 10.1146/annurev.ecolsys.39.110707.173414

950 Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: A new series of programs to
 951 perform population genetics analyses under Linux and Windows. *Molecular Ecology*
 952 *Resources*, 10(3), 564–567. doi: 10.1111/j.1755-0998.2010.02847.x

953 Fallahi, E., Colt, W. M., Fallahi, B., & Chun, I.-J. (2002). The Importance of Apple Rootstocks
 954 on Tree Growth, Yield, Fruit Quality, Leaf Nutrition, and Photosynthesis with an

955 Emphasis on 'Fuji'. *HortTechnology Horttech*, 12(1), 38–44. doi:

956 10.21273/HORTTECH.12.1.38

957 Fallahi, E., Fallahi, P., & Mahdavi, S. (2020). Ancient Urban Gardens of Persia: Concept,

958 History, and Influence on Other World Gardens. *HortTechnology Hortte*, 30(1), 6–12.

959 doi: 10.21273/HORTTECH04415-19

960 Feurtey, A., Cornille, A., Shykoff, J. A., Snirc, A., & Giraud, T. (2017a). Crop-to-wild gene flow

961 and its fitness consequences for a wild fruit tree: Towards a comprehensive conservation

962 strategy of the wild apple in Europe. *Evolutionary Applications*, 10(2), 180–188. PMC

963 (PMC5253423). doi: 10.1111/eva.12441

964 Feurtey, A., Cornille, A., Shykoff, J. A., Snirc, A., & Giraud, T. (2017b). Crop-to-wild gene flow

965 and its fitness consequences for a wild fruit tree: Towards a comprehensive conservation

966 strategy of the wild apple in Europe. *Evolutionary Applications*, 10(2), 180–188. PMC

967 (PMC5253423). doi: 10/f9wm3v

968 Fieldings, A. H., & Bell, J. F. (1997). A review of methods for the assessment of prediction errors

969 in conservation presence/absence models. *Environmental Conservation*, 24, 38–49.

970 Fischer, A., & Schmidt, M. (1938). Wilde Kern-und Steinobstarten, ihre Heimat und ihre

971 Bedeutung für die Entstehung der Kultursorten und die Züchtung. *Der Züchter*, 10(6),

972 157–167.

973 Flowers, J. M., Hazzouri, K. M., Gros-Balthazard, M., Mo, Z., Koutroumpa, K., Perrakis, A., ...

974 Purugganan, M. D. (2019). Cross-species hybridization and the origin of North African

975 date palms. *Proceedings of the National Academy of Sciences of the United States of*

976 *America*, 116(5), 1651–1658. doi: 10.1073/pnas.1817453116

977 Fontaine, M. C., Gladieux, P., Hood, M. E., & Giraud, T. (2013). History of the invasion of the
 978 anther smut pathogen on *Silene latifolia* in North America. *New Phytologist*, 198(3), 946–
 979 956. doi: 10.1111/nph.12177
 980 Forsline, P. L., Aldwinckle, H. S., Dickson, E. E., Luby, J. J., & Hokanson, S. C. (2003). *Of Wild*
 981 *Apples of Central Asia* (Vol. 29).
 982 Fuller, D. Q. (2018). Long and attenuated: Comparative trends in the domestication of tree fruits.
 983 *Vegetation History and Archaeobotany*, 27(1), 165–176. doi: 10.1007/s00334-017-0659-2
 984 Gabrielian, E. T., & Zohary, D. (2004). Wild relatives of food crops native to Armenia and
 985 Nakhichevan. *Flora Mediterranea*, 14, 5–80.
 986 Garza, J. C., & Williamson, E. G. (2001). Detection of reduction in population size using data
 987 from microsatellite loci. *Molecular Ecology*, 10(2), 305–318. doi: 10.1046/j.1365-
 988 294X.2001.01190.x
 989 Gaut, B. S., Díez, C. M., & Morrell, P. L. (2015). Genomics and the Contrasting Dynamics of
 990 Annual and Perennial Domestication. *Trends in Genetics*, 31(12), 709–719. doi:
 991 10.1016/j.tig.2015.10.002
 992 George, J.-P., Konrad, H., Collin, E., Thevenet, J., Ballian, D., Idzajt, M., ... Geburek, T.
 993 (2015). High molecular diversity in the true service tree (*Sorbus domestica*) despite
 994 rareness: Data from Europe with special reference to the Austrian occurrence. *Annals of*
 995 *Botany*, 115(7), 1105–1115. doi: 10.1093/aob/mcv047
 996 Gharghani, A., Zamani, Z., Talaie, A., Fattahi, R., Hajnajari, H., Oraguzie, N. C., ... Gardiner, S.
 997 E. (2010). The Role of Iran (Persia) in Apple (*Malus × domestica* Borkh.) Domestication,
 998 Evolution and Migration via the Silk Trade Route. *Acta Horticulturae*, (859), 229–236.
 999 Gharghani, Ali, Zamani, Z., Talaie, A., Oraguzie, N. C., Fatahi, R., Hajnajari, H., ... Gardiner, S.
 1000 E. (2009). Genetic identity and relationships of Iranian apple (*Malus × domestica* Borkh.)

1001 cultivars and landraces, wild *Malus* species and representative old apple cultivars based
 1002 on simple sequence repeat (SSR) marker analysis. *Genetic Resources and Crop Evolution*,
 1003 56(6), 829–842.

1004 Giesecke, T., Brewer, S., Finsinger, W., Leydet, M., & Bradshaw, R. H. W. (2017). Patterns and
 1005 dynamics of European vegetation change over the last 15,000 years. *Journal of*
 1006 *Biogeography*, 44(7), 1441–1456. doi: 10.1111/jbi.12974

1007 Groppi, A., Liu, S., Cornille, A., Decroocq, S., & Decroocq, D. (2021). Population genomics of
 1008 apricots unravels domestication history and adaptive events. *Nature Communications*, 12,
 1009 3956.

1010 Gros-Balthazard, M., & Flowers, J. M. (2021). A Brief History of the Origin of Domesticated
 1011 Date Palms. In J. M. Al-Khayri, S. M. Jain, & D. V. Johnson (Eds.), *The Date Palm*
 1012 *Genome, Vol. 1: Phylogeny, Biodiversity and Mapping* (pp. 55–74). Cham: Springer
 1013 International Publishing. doi: 10.1007/978-3-030-73746-7_3

1014 Hardy, O. J., & Vekemans, X. (2002). SPAGeDi: A versatile computer program to analyse spatial
 1015 genetic structure at the individual or population levels. *Molecular Ecology Notes*, 2(4),
 1016 618–620.

1017 Harris, S. A., Robinson, J. P., & Juniper, B. E. (2002). Genetic clues to the origin of the apple.
 1018 *Trends Genet*, 18(8), 426–430.

1019 Hewitt, G. M. (1990). Divergence and speciation as viewed from an insect hybrid zone.
 1020 *Canadian Journal of Zoology*, 68(8), 1701–1715. doi: 10.1139/z90-251

1021 Hewitt, G. M. (2004). Genetic consequences of climatic oscillations in the Quaternary.
 1022 *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*,
 1023 359(1442), 183–195. doi: 10.1098/rstb.2003.1388

1024 Hewitt, Godfrey M. (1996). Some genetic consequences of ice ages, and their role in divergence
 1025 and speciation. *Biological Journal of the Linnean Society*, 58(3), 247–276. doi:
 1026 10.1006/bijl.1996.0035

1027 Höfer, M., Flachowsky, H., Hanke, M.-V., Semënov, V., Šlâvas, A., Bandurko, I., ... Alexanian,
 1028 S. (2013). Assessment of phenotypic variation of *Malus orientalis* in the North Caucasus
 1029 region. *Genetic Resources and Crop Evolution*, 60(4), 1463–1477. doi: 10.1007/s10722-
 1030 012-9935-2

1031 Huson, D. H. (1998). SplitsTree: Analyzing and visualizing evolutionary data. *Bioinformatics*
 1032 (*Oxford, England*), 14(1), 68–73.

1033 Huson, D. H., & Scornavacca, C. (2012). Dendroscope 3: An interactive tool for rooted
 1034 phylogenetic trees and networks. *Systematic Biology*, 61(6), 1061–1067.

1035 Jezkova, T., Olah-Hemmings, V., & Riddle, B. R. (2011). Niche shifting in response to warming
 1036 climate after the last glacial maximum: Inference from genetic data and niche assessments
 1037 in the chisel-toothed kangaroo rat (*Dipodomys microps*). *Global Change Biology*, 17(11),
 1038 3486–3502. doi: 10.1111/j.1365-2486.2011.02508.x

1039 Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide
 1040 SNP data. *Bioinformatics*, 27(21), 3070–3071. doi: 10.1093/bioinformatics/btr521

1041 Kalinowski, S. T. (2011). The computer program STRUCTURE does not reliably identify the
 1042 main genetic clusters within species: Simulations and implications for human population
 1043 structure. *Heredity*, 106(4), 625–632.

1044 Koskinen, M. T., Hirvonen, H., Landry, P. A., & Primmer, C. R. (2004). The benefits of
 1045 increasing the number of microsatellites utilized in genetic population studies: An
 1046 empirical perspective. *Hereditas*, 141(1), 61–67. doi: 10.1111/j.1601-5223.2004.01804.x

1047 Langenfeld, W. T. (1991). Apple trees. *Morphological Evolution, Phylogeny, Geography and*
 1048 *Systematics. Riga (Zinatne), 232.*

1049 Lascoux, M., Palmé, A. E., Cheddadi, R., & Latta, R. G. (2004). Impact of Ice Ages on the
 1050 genetic structure of trees and shrubs. *Philosophical Transactions of the Royal Society B:*
 1051 *Biological Sciences, 359(1442), 197–207.* doi: 10.1098/rstb.2003.1390

1052 Leroy, B., Bellard, C., Dubos, N., Colliot, A., Vasseur, M., Courtial, C., ... Ysnel, F. (2014).
 1053 Forecasted climate and land use changes, and protected areas: The contrasting case of
 1054 spiders. *Diversity and Distributions, 20(6), 686–697.* doi: 10.1111/ddi.12191

1055 Liang, Z., Duan, S., Sheng, J., Zhu, S., Ni, X., Shao, J., ... Dong, Y. (2019). Whole-genome
 1056 resequencing of 472 *Vitis* accessions for grapevine diversity and demographic history
 1057 analyses. *Nature Communications, 10(1), 1190.* doi: 10.1038/s41467-019-09135-8

1058 Liu, S., Cornille, A., Decroocq, S., Tricon, D., Chague, A., Eyquard, J., ... Decroocq, V. (2019).
 1059 The complex evolutionary history of apricots: Species divergence, gene flow and multiple
 1060 domestication events. In *Molecular Ecology*. doi: 10.1111/mec.15296

1061 Loiselle, B. A., Sork, V. L., Nason, J., & Graham, C. (1995). Spatial genetic structure of a
 1062 tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of*
 1063 *Botany, 82(11), 1420–1425.*

1064 Mabry, M. E., Rowan, T. N., Pires, J. C., & Decker, J. E. (2021). Feralization: Confronting the
 1065 Complexity of Domestication and Evolution. *Trends in Genetics, 37(4), 302–305.* doi:
 1066 10.1016/j.tig.2021.01.005

1067 Meyer, R. S., Duval, A. E., & Jensen, H. R. (2012). Patterns and processes in crop domestication:
 1068 An historical review and quantitative analysis of 203 global food crops. *New Phytologist,*
 1069 *196(1), 29–48.* doi: 10.1111/j.1469-8137.2012.04253.x

1070 Migicovsky, Z., Gardner, K. M., Richards, C., Chao, C. T., Schwaninger, H. R., Fazio, G., ...
 1071 Myles, S. (2021). Genomic consequences of apple improvement. *Horticulture Research*,
 1072 8(1), 1–13. doi: 10.1038/s41438-020-00441-7
 1073 Miller, A. J., & Gross, B. L. (2011). From forest to field: Perennial fruit crop domestication.
 1074 *American Journal of Botany*, 98(9), 1389–1414. doi: 10.3732/ajb.1000522
 1075 Monserud, R. A., & Leemans, R. (1992). Comparing global vegetation maps with the Kappa
 1076 statistic. *Ecological Modelling*, 62(4), 275–293.
 1077 Myles, S., Boyko, A. R., Owens, C. L., Brown, P. J., Grassi, F., Aradhya, M. K., ... Buckler, E.
 1078 S. (2011). Genetic structure and domestication history of the grape. *Proceedings of the*
 1079 *National Academy of Sciences of the United States of America*, 108(9), 3530–3535. doi:
 1080 10.1073/pnas.1009363108
 1081 Nakhutsrishvili, G., Zazanashvili, N., Batsatsashvili, K., & Montalvo, CS. (2015). Colchic and
 1082 Hyrcanian forests of the Caucasus: Similarities, differences and conservation status. *Flora*
 1083 *Mediterranea*, 25, 185–192.
 1084 Nei, M. (1987). *Molecular Evolutionary Genetics*. Columbia University Press. Retrieved from
 1085 <https://books.google.es/books?id=UhRSsLkjxDgC>
 1086 Oddou-Muratorio, S., & Klein, E. K. (2008). Comparing direct vs. Indirect estimates of gene
 1087 flow within a population of a scattered tree species. *Molecular Ecology*, 17(11), 2743–
 1088 2754.
 1089 Omasheva, M. Y., Flachowsky, H., Ryabushkina, N. A., Pozharskiy, A. S., Galiakparov, N. N., &
 1090 Hanke, M.-V. (2017). To what extent do wild apples in Kazakhstan retain their genetic
 1091 integrity? *Tree Genetics & Genomes*, 13(3), 52. doi: 10.1007/s11295-017-1134-z
 1092 Parvizi, E., Keikhosravi, A., Naderloo, R., Solhjoui-Fard, S., Sheibak, F., & Schubart, C. D.
 1093 (2019). Phylogeography of *Potamon ibericum* (Brachyura: Potamidae) identifies

1094 Quaternary glacial refugia within the Caucasus biodiversity hot spot. *Ecology and*
 1095 *Evolution*, 9(8), 4749–4759. doi: 10.1002/ece3.5078
 1096 Patocchi, A., Fernández-Fernández, F., Evans, K., Gobbin, D., Rezzonico, F., Boudichevskaia,
 1097 A., ... van de Weg, W. (2009). Development and test of 21 multiplex PCRs composed of
 1098 SSRs spanning most of the apple genome. *Tree Genetics and Genomes*, 5(1), 211–223.
 1099 Patocchi, A., Frei, A., Frey, J. E., & Kellerhals, M. (2009). Towards improvement of marker
 1100 assisted selection of apple scab resistant cultivars: *Venturia inaequalis* virulence surveys
 1101 and standardization of molecular marker alleles associated with resistance genes.
 1102 *Molecular Breeding*, 24(4), 337–347. doi: 10.1007/s11032-009-9295-6
 1103 Petit, R. J., Bialozyt, R., Garnier-Géré, P., & Hampe, A. (2004). Ecology and genetics of tree
 1104 invasions: From recent introductions to Quaternary migrations. *Forest Ecology and*
 1105 *Management*, 197(1–3), 117–137. doi: 10.1016/j.foreco.2004.05.009
 1106 Petit, R. J., & Hampe, A. (2006). Some Evolutionary Consequences of Being a Tree. *Annual*
 1107 *Review of Ecology, Evolution, and Systematics*, 37(1), 187–214. doi:
 1108 10.1146/annurev.ecolsys.37.091305.110215
 1109 Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using
 1110 multilocus genotype data. *Genetics*, 155(2), 945–959.
 1111 Pudlo, P., Marin, J.-M., Estoup, A., Cornuet, J.-M., Gautier, M., & Robert, C. P. (2016). Reliable
 1112 ABC model choice via random forests. *Bioinformatics*, 32(6), 859–866. doi:
 1113 10.1093/bioinformatics/btv684
 1114 Puechmaille, S. J. (2016). The program structure does not reliably recover the correct population
 1115 structure when sampling is uneven: Subsampling and new estimators alleviate the
 1116 problem. *Molecular Ecology Resources*, 16(3), 608–627. doi: 10.1111/1755-0998.12512

1117 Putman, A. I., & Carbone, I. (2014). Challenges in analysis and interpretation of microsatellite
 1118 data for population genetic studies. *Ecology and Evolution*, 4(22), 4399–4428. doi:
 1119 10.1002/ece3.1305

1120 Qiu, J., Wang, L., Liu, M., Shen, Q., & Tang, J. (2011). An efficient and simple protocol for a
 1121 PdCl₂-ligandless and additive-free Suzuki coupling reaction of aryl bromides.
 1122 *Tetrahedron Letters*, 52(48), 6489–6491. doi: 10.1016/j.tetlet.2011.09.115

1123 Raymond, M., & Rousset, F. (1995). An exact test for population differentiation. *Evolution*,
 1124 49(6), 1280–1283.

1125 Raynal, L., Marin, J.-M., Pudlo, P., Ribatet, M., Robert, C. P., & Estoup, A. (2019). ABC random
 1126 forests for Bayesian parameter inference. *Bioinformatics*, 35(10), 1720–1728. doi:
 1127 10.1093/bioinformatics/bty867

1128 Rechinger, K. H. (1964). Flora Iranica, Akademische Druck-und Verlagsanstalt Graz. *University*
 1129 *of Tehran, Iran*, 549.

1130 Rousset, F. (2008). genepop'007: A complete re-implementation of the genepop software for
 1131 Windows and Linux. *Molecular Ecology Resources*, 8(1), 103–106.

1132 Schmitt, T. (2007). Molecular biogeography of Europe: Pleistocene cycles and postglacial trends.
 1133 *Frontiers in Zoology*, 4, 1–13. doi: 10.1186/1742-9994-4-11

1134 Slatkin, M. (1995). A measure of population subdivision based on microsatellite allele
 1135 frequencies. *Genetics*, 139(1), 457–462. PubMed (7705646). doi:
 1136 10.1093/genetics/139.1.457

1137 Soofi, M., Ghoddousi, A., Zeppenfeld, T., Shokri, S., Soufi, M., Jafari, A., ... Waltert, M. (2018).
 1138 Livestock grazing in protected areas and its effects on large mammals in the Hyrcanian
 1139 forest, Iran. *Biological Conservation*, 217, 377–382. doi: 10.1016/j.biocon.2017.11.020

1140 Spengler, R.N. (2019). *Fruit from the Sands: The Silk Road Origins of the Foods We Eat*.
 1141 University of California Press. Retrieved from
 1142 <https://books.google.fr/books?id=fr6SDwAAQBAJ>
 1143 Spengler, Robert N., Chang, C., & Tourtellotte, P. A. (2013). Agricultural production in the
 1144 Central Asian mountains: Tuzusai, Kazakhstan (410–150 b.c.). *Journal of Field*
 1145 *Archaeology*, 38(1), 68–85. doi: 10.1179/0093469012Z.0000000000037
 1146 Spengler, Robert Nicholas. (2019). Origins of the Apple: The Role of Megafaunal Mutualism in
 1147 the Domestication of Malus and Rosaceous Trees. *Frontiers in Plant Science*, 10. doi:
 1148 10.3389/fpls.2019.00617
 1149 Szpiech, Z. A., Jakobsson, M., & Rosenberg, N. A. (2008). ADZE: a rarefaction approach for
 1150 counting alleles private to combinations of populations. *Bioinformatics*, 24(21), 2498–
 1151 2504.
 1152 Takezaki, N., & Nei, M. (2008). Empirical Tests of the Reliability of Phylogenetic Trees
 1153 Constructed With Microsatellite DNA. *Genetics*, 178(1), 385–392. doi:
 1154 10.1534/genetics.107.081505
 1155 Tardío, J., Arnal, A., & Lázaro, A. (2020). Ethnobotany of the crab apple tree (*Malus sylvestris*
 1156 (L.) Mill., Rosaceae) in Spain. *Genetic Resources and Crop Evolution*. doi:
 1157 10.1007/s10722-020-01026-y
 1158 Teixeira, J. C., & Huber, C. D. (2021). The inflated significance of neutral genetic diversity in
 1159 conservation genetics. *Proceedings of the National Academy of Sciences*, 118(10),
 1160 e2015096118. doi: 10.1073/pnas.2015096118
 1161 Thuiller, W., Georges, D., Engler, R., & Breiner, F. (2016). *Biomod2: Ensemble platform for*
 1162 *species distribution modeling. R package version 3.3-7*.

1163 Tian, F., Li, B., Ji, B., Zhang, G., & Luo, Y. (2009). Identification and structure-activity
 1164 relationship of gallotannins separated from *Galla chinensis*. *LWT - Food Science and*
 1165 *Technology*, 42(7), 1289–1295. doi: 10.1016/j.lwt.2009.03.004

1166 Vavilov, N. I. (1926). Studies on the origin of cultivated plants. *Trudy Byuro. Prikl. Bot.*, 16,
 1167 139–245.

1168 Vavilov, N. I. (1992). *Origin and geography of cultivated plants*. Cambridge: Cambridge
 1169 University Press. CABDirect.

1170 Vekemans, X., & Hardy, O. J. (2004a). New insights from fine-scale spatial genetic structure
 1171 analyses in plant populations. *Molecular Ecology*, 13(4), 921–935.

1172 Vekemans, X., & Hardy, O. J. (2004b). New insights from fine-scale spatial genetic structure
 1173 analyses in plant populations. *Molecular Ecology*, 13(4), 921–935. doi: 10.1046/j.1365-
 1174 294X.2004.02076.x

1175 Vercken, E., Fontaine, M. C., Gladieux, P., Hood, M. E., Jonot, O., & Giraud, T. (2010). Glacial
 1176 refugia in pathogens: European genetic structure of anther smut pathogens on *Silene*
 1177 *latifolia* and *Silene dioica*. *PLoS Pathogens*, 6(12), e1001229.

1178 Volk, G. M., & Cornille, A. (2019). Genetic Diversity and Domestication History in *Pyrus*. In
 1179 *The Pear Genome* (pp. 51–62). Springer.

1180 Volk, G. M., Richards, C. M., Reilley, A. A., Henk, A. D., Reeves, P. A., Forsline, P. L., &
 1181 Aldwinckle, H. S. (2008). Genetic diversity and disease resistance of wild *Malus*
 1182 *orientalis* from Turkey and Southern Russia. *Journal of the American Society for*
 1183 *Horticultural Science*, 133(3), 383–389.

1184 Vouillamoz, J. F., McGovern, P. E., Ergul, A., Söylemezoğlu, G., Tevzadze, G., Meredith, C. P.,
 1185 & Grando, M. S. (2006). Genetic characterization and relationships of traditional grape

1186 cultivars from Transcaucasia and Anatolia. *Plant Genetic Resources*, 4(2), 144–158. doi:
 1187 10.1079/pgr2006114
 1188 Wegmann, D., Leuenberger, C., Neuenschwander, S., & Excoffier, L. (2010). ABCtoolbox: A
 1189 versatile toolkit for approximate Bayesian computations. *BMC Bioinformatics*, 11(1), 116.
 1190 doi: 10.1186/1471-2105-11-116
 1191 Weir, B. S., & Cockerham, C. C. (1984). Estimating F-Statistics for the analysis of population
 1192 structure. *Evolution*, 38(6), 1358–1370.
 1193 Wilson, G. A., & Rannala, B. (2003). Bayesian Inference of Recent Migration Rates Using
 1194 Multilocus Genotypes. *Genetics*, 163(3), 1177–1191. doi: 10.1093/genetics/163.3.1177
 1195 Wu, J., Wang, Y., Xu, J., Korban, S. S., Fei, Z., Tao, S., ... Zhang, S. (2018). Diversification and
 1196 independent domestication of Asian and European pears. *Genome Biology*, 19(1), 77. doi:
 1197 10.1186/s13059-018-1452-y
 1198 Yousefzadeh, H., Hosseinzadeh Colagar, A., Tabari, M., Sattarian, A., & Assadi, M. (2012).
 1199 Utility of ITS region sequence and structure for molecular identification of *Tilia* species
 1200 from Hyrcanian forests, Iran. *Plant Systematics and Evolution*, 298(5), 947–961. doi:
 1201 10.1007/s00606-012-0604-x
 1202 Zazanashvili, N., Sanadiradze, G., Garforth, M., Bitsadze, M., Manvelyan, K., Askerov, E., ...
 1203 Devranoğlu Tavşel, S. (2020). *Ecoregional Conservation Plan (ECP) For The Caucasus*
 1204 *2020 Edition*. WWF, KfW, Tbilisi.
 1205 Zeng, G., Zhang, J., Chen, Y., Yu, Z., Yu, M., Li, H., ... Hu, C. (2011). Relative contributions of
 1206 archaea and bacteria to microbial ammonia oxidation differ under different conditions
 1207 during agricultural waste composting. *Bioresour Technol*, 102(19), 9026–9032. doi:
 1208 10.1016/j.biortech.2011.07.076

Zhang, H., Mittal, N., Leamy, L. J., Barazani, O., & Song, B. H. (2017). Back into the wild—
 Apply untapped genetic diversity of wild relatives for crop improvement. *Evolutionary
 Applications*, 10(1), 5–24. doi: 10.1111/eva.12434

Zhang, H.-X., Li, H.-Y., & Li, Y.-X. (2018). Identifying evolutionarily significant units for
 conservation of the endangered *Malus sieversii* using genome-wide RADseq data. *Nordic
 Journal of Botany*, 36(7), e01733. doi: 10.1111/njb.01733

Figures and Tables

**Figure 1. Population genetic structure in cultivated and wild apples from the Caucasus and
 Iran, *Malus domestica*, *Malus sieversii* and *Malus baccata* based on 26 microsatellite markers.**

a. Spatial population genetic structure inferred with STRUCTURE at $K = 12$ ($N = 550$); the map
 represents membership proportions averaged over each geographic site for the Caucasian crabapple
M. orientalis ($N = 374$, 43 sites across Turkey, Russia, Armenia and Iran). In the bottom right
 corner, the mean membership proportions for the apple cultivars from Armenia ($N = 4$), Iran ($N =$
 48) and Europe (*M. domestica*, $N = 40$). The size of the pie charts is proportional to the number of
 samples per site. b. STRUCTURE bar plot ($N = 550$) at $K = 12$ showing 12 distinct genetic clusters.
 Each vertical line represents an individual. Colors represent the inferred ancestry from K ancestral
 genetic clusters. Sites are grouped by country for the wild apple samples (*i.e.*, Turkey, Russia,
 Armenia, Iran) and *M. sieversii* (*i.e.*, Kazakhstan and Kyrgyzstan), apple cultivars are grouped
 according to their origin: Armenia ($N = 4$), Iran ($N = 48$) and *M. domestica* ($N = 40$). Countries
 (Kazakhstan, Kyrgyzstan, Armenia) and/or main regions in the Caucasus (the Western Caucasus,
i.e., Turkey and Russia, Zagros and Hyrcanian Forests, Central and Southern Armenia) are shown
 on the map. Reference samples from previously published studies of each species are: *M. orientalis*
 from the Western Caucasus and Central and Southern Armenia, *M. domestica* (European cultivars),
M. sieversii from Kazakhstan (Cornile et al., 2013) and *M. baccata* (Cornille et al., 2012).

1237

1238 **Figure 2. Genetic variation and differentiation among cultivated and wild apples from the Caucasus, *Malus domestica*, *Malus***
1239 ***sieversii* and *Malus baccata* based on 26 microsatellite markers.** a. Principal component analysis after removing the outgroup *M.*
1240 *baccata* ($N = 530$), with the respective total variation explained by each of the two first components. b. Neighbor-net representing the
1241 genetic relationships among wild and cultivated individuals inferred with STRUCTURE at $K = 12$. Colors correspond to the genetic
1242 groups inferred with STRUCTURE at $K=12$, and admixed samples are in grey.

1243

1244 **Figure 3. Most likely scenario of domestication of cultivated apples in Iran using random-forest approximate Bayesian**
1245 **computation (ABC-RF) combined with coalescent-based simulations (SC45, Figure S12).** Population names correspond to the ones
1246 detected with STRUCTURE for $K = 12$, excluding admixed individuals (*i.e.*, individuals with a membership coefficient < 0.85 to any
1247 given cluster), *M. baccata* was excluded from the inferences. Bidirectional gene flow between populations was assumed. For clarity,
1248 only estimates of divergence time of the cultivated populations are provided, other parameters estimates are provided in Table S13.

1249

1250

1

2 **Figure 4. Spatial diversity and past contraction and expansion of *Malus orientalis* across the Caucasus. a.** Spatial genetic
3 diversity (allelic richness) at 36 sites ($N = 339$). **b. and c.** Ensemble forecasting of the three different algorithms (ANN, GLM and
4 GAM) predicting the current and last glacial maximum (LGM) distribution range of suitable areas for *M. orientalis*, respectively. The
5 probabilities of being a suitable habitat are given in the legend. The Colchis and Hyrcanian regions are shown on the maps.

1 **Table 1. Genetic diversity estimates for wild and cultivated apple populations detected with STRUCTURE at $K = 12$** ($N = 424$,
2 *i.e.*, individuals with a membership coefficient < 0.85 to any given cluster were excluded from the analysis). Note that the purple cluster
3 was split between cultivated and wild samples. Thus, samples were partitioned into 13 populations, including 10 wild and three cultivated
4 apple populations.
5

| Wild or cultivated | Species | Country of origin | Population | N | H_O | H_E | F_{IS} | $A_R (G=7)$ | $A_P (G=7)$ |
|--------------------|-------------------------|-------------------|------------------------|-----|-------|-------|---------------------|-------------|-------------|
| wild | <i>Malus orientalis</i> | Armenia | Western (orange) | 20 | 0.82 | 0.86 | 0.05** | 5.04±0.13 | 0.97±0.10 |
| | | | Central (blue) | 109 | 0.78 | 0.79 | 0.01 ^{NS} | 4.03±0.14 | 0.60±0.08 |
| | | | Southern (brown) | 29 | 0.76 | 0.79 | 0.03 ^{NS} | 4.38±0.17 | 0.63±0.12 |
| | | Iran | Lorestan (light green) | 10 | 0.84 | 0.46 | -0.82 *** | 2.03±0.12 | 0.23±0.06 |
| | | | Kurdestan (red) | 12 | 0.75 | 0.71 | -0.05 * | 3.71±0.15 | 0.50±0.09 |
| | | | Hyrceanian (pink) | 88 | 0.71 | 0.76 | 0.07 *** | 4.12±0.18 | 0.65±0.10 |
| | | | Hyrceanian (purple) | 19 | 0.76 | 0.63 | -0.21 *** | 3.24±0.13 | 0.21±0.03 |
| | <i>Malus sieversii</i> | Kazakhstan | (light blue) | 40 | 0.74 | 0.76 | 0.03 * | 4.15±0.16 | 0.99±0.13 |
| | | Kyrkyzstan | (cyan) | 17 | 0.70 | 0.73 | 0.04 * | 3.47±0.14 | 0.30±0.06 |
| | <i>Malus baccata</i> | Russia | (light red) | 17 | 0.52 | 0.55 | 0.06 ^{NS} | 3.23±0.19 | 1.08±0.14 |
| cultivated | <i>Malus domestica</i> | Mostly Europe | European cultivars | 38 | 0.78 | 0.78 | -0.00 ^{NS} | 4.27±0.09 | 0.85±0.10 |

| | | | | | | | | | |
|-------|---|------|-----------------------------------|-----|------|------|------------------|-----------|-----------|
| | ? | Iran | Iranian cultivars (purple) | 18 | 0.77 | 0.61 | -0.26 *** | 3.08±0.13 | 0.10±0.02 |
| | | | Iranian cultivars (dark green) | 7 | 0.71 | 0.65 | -0.08 * | 3.39±0.13 | 0.52±0.09 |
| TOTAL | | | | 424 | | | | | |

1

2 *N*: number of individuals assigned to a focal cluster with a membership coefficient > 0.85; *H_O* and *H_E*: observed and expected
3 heterozygosity; *F_{IS}*: inbreeding coefficient; *A_R*: mean allelic richness across loci, corrected by the rarefaction method, estimated for a
4 sample size of 7; *A_P*: number of private alleles, corrected by the rarefaction method, estimated for a sample size of 7; *: 0.05 < *P* < 0.01;
5 ***: *P* < 0.001; NS: non-significant.

6

1 **Table 2. Distribution of hybrids (*i.e.*, individuals with a membership coefficient < 0.90 to any given genetic cluster, as inferred**
2 **with STRUCTURE for $K = 9$) in cultivated and wild apple in the Caucasus ($N = 466$, 26 microsatellite markers).**

| Gene pool | | Crop | | | | Wild | | | |
|--|-----------------------------------|------------------------|----------------------|-------------------------------------|--------------------------------------|--------------------------|-----------------------|-----------------------|-----------------------|
| | | Armenia $N_{tot}=3$ | Iran $N_{tot}=48$ | <i>M. domestica</i> $N_{tot}=40$ | | Armenia $N_{tot}=196$ | Iran $N_{tot}=167$ | Russia $N_{tot}=5$ | Turkey $N_{tot}=6$ |
| <i>Malus domestica</i> | mean introgression rate | 0 | 0.13 | 0.66 | | 0.04 | 0.02 | 0 | 0 |
| | N (%) over the total | 0 | 11 (23%) | 2 (5%) | | 4 (2%) | 5 (2.9%) | 0 | 0 |
| Wild and cultivated Hyrcanian (purple) | mean introgression rate | 0 | 0.36 | 0.33 | | 0 | 0.16 | 0 | 0 |
| | N (%) | 0 | 18 (37.5%) | 2 (5%) | | 0 | 13 (7.8%) | 0 | 0 |
| Cult. Iran (dark_green) | mean introgression rate | 0 | 0.17 | 0 | | 0.05 | 0.06 | 0 | 0 |
| | N (%) | 0 | 14 (30%) | 0 | | 11 (5.6%) | 8 (4.7%) | 0 | 0 |
| | Ncrop-crop hybrids (and %) | 0 | 43 (89.5%) | 4 (1%) | Ncrop-to-wild hybrids (and %) | 15 (7.6%) | 26 (18.1%) | 0 | 0 |
| Hyrcanian (pink) | mean introgression rate | 0 | 0.05 | 0 | | 0 | 0.41 | 0 | 0 |
| | N (%) | 0 | 6 (12.5%) | 0 | | 0 | 34 (20.3%) | 0 | 0 |
| Lorestan (light green) | mean introgression rate | 0 | 0.01 | 0 | | 0 | 0.04 | 0 | 0 |
| | N (%) | 0 | 2 (4%) | 0 | | 0 | 6 (3.6%) | 0 | 0 |
| Wild Kurdistan (red) | mean introgression rate | 0 | 0.17 | 0 | | 0 | 0.27 | 0 | 0 |
| | N (%) | 0 | 12 (35%) | 0 | | 0 | 30 (17.9%) | 0 | 0 |
| Western (orange) | mean introgression rate | 0.65 | 0.08 | 0 | | 0.28 | 0.02 | 0.49 | 0.84 |
| | N (%) | 1 (33%) | 3 (6.2%) | 0 | | 33 (16.8%) | 4 (2.3%) | 1 (20%) | 2 (33%) |
| Southern (brown) | mean introgression rate | 0 | 0.02 | 0 | | 0.17 | 0.02 | 0.35 | 0 |
| | N (%) | 0 | 2 (4.1%) | 0 | | 13 (6.6%) | 2 (1.1%) | 1 (20%) | 0 |

| | | | | | | | | | | |
|----------------|--|----------|-------------------|---------------|---|-------------------|-----------------|----------------|----------------|----------------------|
| Central (blue) | mean introgression rate | 0.32 | 0.02 | 0 | | 0.46 | 0.01 | 0 | 0.11 | |
| | N (%) | 1 (33%) | 1 (2%) | 0 | | 46 (23.4%) | 1 (0.5%) | 0 | 1 (16.6%) | |
| | Nwild-to-crop hybrids (and %) | 2 (66%) | 26 (54.1%) | 0 | Nwild-to- wild hybrids (and %) | 92 (46.9%) | 77 (46.1%) | 2 (40%) | 3 (50%) | |
| | Total number of hybrids (and %) | 1 | 32 (66%) | 2 (5%) | | 58 (28%) | 54 (32%) | 1 (20%) | 2 (33%) | 150 (32%) |

1 *N_{tot}*: total number of samples in each group (*i.e.*, cultivated or wild, from different regions); *N* (and %): number of hybrids assigned to
2 each gene pool, and the respective percentage over the total number of samples from each group; mean introgression rate: mean
3 membership coefficient to this gene pool. Note that some admixed trees were assigned to several gene pools with a membership
4 coefficient < 0.90, therefore, the total number of hybrids associated with each cluster (TOTAL) is given on the last line of the table. We
5 also showed the distribution of crop-crop, crop-to-wild, wild-to-crop and wild-to-wild hybrids.

