



HAL
open science

About the unification type of $K + \text{Box Box false}$

Philippe Balbiani, Cigdem Gencer, Maryam Rostamigiv, Tinko Tinchev

► **To cite this version:**

Philippe Balbiani, Cigdem Gencer, Maryam Rostamigiv, Tinko Tinchev. About the unification type of $K + \text{Box Box false}$. *Annals of Mathematics and Artificial Intelligence*, 2021, 90, pp.481-497. hal-03762105

HAL Id: hal-03762105

<https://hal.science/hal-03762105>

Submitted on 26 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

About the unification type of $\mathbf{K} + \Box\Box\perp^*$

Philippe Balbiani^a Çiğdem Gencer^{a,b} Maryam Rostamigiv^a Tinko Tinchev^c

^aToulouse Institute of Computer Science Research
CNRS — Toulouse University, Toulouse, France

^bFaculty of Arts and Sciences

Istanbul Aydın University, Istanbul, Turkey

^cFaculty of Mathematics and Informatics

Sofia University St. Kliment Ohridski, Sofia, Bulgaria

Abstract

The unification problem in a propositional logic is to determine, given a formula φ , whether there exists a substitution σ such that $\sigma(\varphi)$ is in that logic. In that case, σ is a unifier of φ . When a unifiable formula has minimal complete sets of unifiers, it is either infinitary, finitary, or unitary, depending on the cardinality of its minimal complete sets of unifiers. Otherwise, it is nullary. In this paper, we prove that in modal logic $\mathbf{K} + \Box\Box\perp$, unifiable formulas are either finitary, or unitary.

Keywords

Propositional modal logics. Locally tabular modal logics. Unification problem. Unification types.

1 Introduction

The unification problem in a propositional logic is to determine, given a formula φ , whether there exists a substitution σ such that $\sigma(\varphi)$ is in that logic. In that case, σ is a unifier of φ . We say that a set of unifiers of a unifiable formula φ is complete if for all unifiers σ of φ , there exists a unifier τ of φ in that set such that τ is more general than σ ¹. Now, an important question is to determine whether a given unifiable formula has minimal complete sets of unifiers [1]. When such sets exist, they all have the same cardinality. In that case, a unifiable formula is either infinitary, or finitary, or unitary, depending whether its complete sets of unifiers are either infinite, or finite, or with cardinality 1. Otherwise, the formula is nullary. The importance of the unification problem lies in its connection with the admissibility problem. In a consistent propositional logic \mathbf{L} , unification is reducible to non-admissibility, seeing that the unifiability in \mathbf{L} of a formula φ is equivalent to the non-admissibility in \mathbf{L} of the inference rule $\frac{\varphi}{\perp}$. As observed by Ghilardi [15], when \mathbf{L} has a decidable membership problem and \mathbf{L} is either unitary, or finitary, algorithms for computing minimal complete sets of unifiers in \mathbf{L} can be used as a key component of algorithms for solving the admissibility problem in \mathbf{L} , seeing that the admissibility in \mathbf{L} of an inference rule $\frac{\varphi_1, \dots, \varphi_p}{\psi}$ is equivalent to the inclusion in \mathbf{L} of the set $\{\sigma(\psi) : \sigma \in \Sigma\}$, where Σ is an arbitrary minimal complete set of unifiers of $\varphi_1 \wedge \dots \wedge \varphi_p$ in \mathbf{L} .

Within the context of the unification problem in a propositional logic, we distinguish between elementary unification and unification with constants. In unification with constants, some variables (called constants) are never replaced by formulas when one applies a substitution whereas in elementary unification, all variables are likely to be replaced. About the unification type of modal logics², it is known

*Corresponding author: Philippe Balbiani. E-mail address: philippe.balbiani@irit.fr.

¹A substitution σ is more general than a substitution τ in a propositional logic if there exists a substitution ν such that for all variables x , $\nu(\sigma(x)) \leftrightarrow \tau(x)$ is in that logic.

²In this paper, all modal logics are normal. We follow the same conventions as in [9, 10, 23] for talking about them: $\mathbf{S5}$ is the least modal logic containing the formulas usually denoted \mathbf{T} , 4 and 5, \mathbf{KD} is the least modal logic containing the formula usually denoted \mathbf{D} , etc.

that \mathbf{KT} , \mathbf{KD} and \mathbf{KB} are nullary [2, 3, 5], $\mathbf{KD45}$ and $\mathbf{K45}$ are unitary [4, 8, 17, 20], $\mathbf{Alt}_1 + \Box^d\perp$ (the least modal logic containing \mathbf{Alt}_1 and $\Box^d\perp$) is unitary for each $d \geq 2$ [6], $\mathbf{S5}$ and $\mathbf{S4.3}$ are unitary [11, 12, 13], transitive modal logics like $\mathbf{K4}$ and $\mathbf{S4}$ are finitary [15, 19], \mathbf{K} is nullary [21] and $\mathbf{K4D1}$ (the least modal logic containing $\mathbf{K4}$ and $\Box(\Box x \rightarrow y) \vee \Box(\Box y \rightarrow x)$) is unitary [22], the type of \mathbf{KT} , \mathbf{KD} and \mathbf{KB} having only been obtained within the context of unification with constants and the type of $\mathbf{Alt}_1 + \Box^d\perp$ having only been obtained within the context of elementary unification. About the unification type of \mathbf{Alt}_1 and its extensions, the line of reasoning determining in [3, 5] the unification type (nullary) of \mathbf{KD} within the context of unification with constants can be adapted to $\mathbf{Alt}_1 + \Diamond\top$ whereas the line of reasoning determining in [21] the unification type (nullary) of \mathbf{K} has been adapted to \mathbf{Alt}_1 [7]. In this paper, within the context of elementary unification, we prove that in $\mathbf{K} + \Box\Box\perp$ (the least modal logic containing $\Box\Box\perp$), unifiable formulas are either finitary, or unitary³.

2 A preliminary result

Let S be a finite set. We write $\|S\|$ for the cardinality of S . If S is nonempty then for all equivalence relations \sim on S , for all $\alpha \in S$, $[\alpha]$ denotes the equivalence class of α modulo \sim and for all $T \subseteq S$, T/\sim denotes the quotient set of T modulo \sim . Our first result, Proposition 1, is used later in Section 6. Its proof is presented in an Annex along with the proofs of all other results asserted in this paper.

Proposition 1. *Let T be a finite set. If S is nonempty then for all equivalence relations \sim on S , $\|S/\sim\| \leq \|T\| \leq \|S\|$ iff there exists a surjective function f from S to T such that for all $\alpha, \beta \in S$, if $f(\alpha) = f(\beta)$ then $\alpha \sim \beta$.*

3 Syntax

Let \mathbf{VAR} be a countably infinite set of *variables* (with typical members denoted x, y , etc). Let (x_1, x_2, \dots) be an enumeration of \mathbf{VAR} without repetitions. Let $n \geq 1$. The set \mathbf{FOR}_n of all n -formulas (with typical members denoted φ, ψ , etc) is inductively defined by:

- $\varphi, \psi ::= x_i \mid \perp \mid \neg\varphi \mid (\varphi \vee \psi) \mid \Box\varphi$.

We adopt the standard rules for omission of the parentheses. The connectives $\top, \wedge, \rightarrow$ and \leftrightarrow are defined by the usual abbreviations. We have also a connective \Diamond which is defined by

- $\Diamond\varphi ::= \neg\Box\neg\varphi$.

For all $\varphi \in \mathbf{FOR}_n$, we respectively write “ φ^0 ” and “ φ^1 ” to mean “ $\neg\varphi$ ” and “ φ ”. An n -substitution is a couple (k, σ) where $k \geq 1$ and σ is a homomorphism from \mathbf{FOR}_n to \mathbf{FOR}_k . Let \mathbf{SUB}_n be the set of all n -substitutions. From now on,

we write “ \mathbf{L}_2 ” to mean “ $\mathbf{K} + \Box\Box\perp$ ”.

The standard axiomatization of \mathbf{L}_2 consists of the following axioms and rules of proof:

- all propositional tautologies,
- $\Box(x_i \rightarrow x_j) \rightarrow (\Box x_i \rightarrow \Box x_j)$,
- $\Box\Box\perp$,

³From now on, “... iff ...” means “... if and only if ...” and “... not-iff ...” means “... if and only if not ...”.

- modus ponens,
- uniform substitution,
- generalization: given φ , prove $\square\varphi$.

As is well-known, \mathbf{L}_2 is the modal logic of directed graphs where there is no path of length 3. The generated subgraphs of such directed graphs are therefore tree-like Kripke frames of depth at most 1. They constitute the semantic basis of \mathbf{L}_2 — tree-like Kripke models of depth at most 1 — developed in Section 4. In the meantime, let us consider the equivalence relation \equiv_n on \mathbf{FOR}_n defined by:

- $\varphi \equiv_n \psi$ iff $\varphi \leftrightarrow \psi \in \mathbf{L}_2$.

Proposition 2. \equiv_n possesses finitely many equivalence classes.

In other words, \mathbf{L}_2 is *locally tabular*. Locally tabular modal logics possess interesting properties. In particular, in terms of decidability [24, 25, 26]. See also [10, Chapter 12] and [14]. The equivalence relation \simeq_n on \mathbf{SUB}_n is defined by:

- $(k, \sigma) \simeq_n (l, \tau)$ iff for all $i \in \{1, \dots, n\}$, $\sigma(x_i) \leftrightarrow \tau(x_i) \in \mathbf{L}_2$.

The preorder \preceq_n on \mathbf{SUB}_n is defined by:

- $(k, \sigma) \preceq_n (l, \tau)$ iff there exists a k -substitution (m, v) such that for all $i \in \{1, \dots, n\}$, $v(\sigma(x_i)) \leftrightarrow \tau(x_i) \in \mathbf{L}_2$.

4 Semantics

Let $n \geq 1$. An n -tuple of bits (denoted α, β , etc) is a function from $\{1, \dots, n\}$ to $\{0, 1\}$. Such function should be understood as a propositional valuation of the variables x_1, \dots, x_n : for all $i \in \{1, \dots, n\}$, if $\alpha_i = 0$ then it is interpreted to mean “ x_i is false” else it is interpreted to mean “ x_i is true”. Let \mathbf{BIT}_n be the set of all n -tuples of bits. An n -model is a structure of the form (α, S) where $\alpha \in \mathbf{BIT}_n$ and $S \subseteq \mathbf{BIT}_n$. Such structure should be understood as a tree-like Kripke model of depth at most 1: α is the valuation of its root node and S is the set of the valuations of its non-root nodes. Let \mathbf{MOD}_n be the set of all n -models. We say that an n -model (α, S) is *degenerated* if $S = \emptyset$. Let $\mathbf{MOD}_n^{\text{deg}}$ be the set of all degenerated n -models. Notice that $\|\mathbf{MOD}_n^{\text{deg}}\| = 2^n$. Notice also that for all sets S of n -tuples of bits, $S \times \{\emptyset\}$ is a set of degenerated n -models. The binary relation \models_n of n -satisfiability between \mathbf{MOD}_n and \mathbf{FOR}_n is defined as expected:

- $(\alpha, S) \models_n x_i$ iff $\alpha_i = 1$,
- $(\alpha, S) \not\models_n \perp$,
- $(\alpha, S) \models_n \neg\varphi$ iff $(\alpha, S) \not\models_n \varphi$,
- $(\alpha, S) \models_n \varphi \vee \psi$ iff either $(\alpha, S) \models_n \varphi$, or $(\alpha, S) \models_n \psi$,
- $(\alpha, S) \models_n \square\varphi$ iff for all $\beta \in S$, $(\beta, \emptyset) \models_n \varphi$.

As a result,

- $(\alpha, S) \models_n \diamond\varphi$ iff there exists $\beta \in S$ such that $(\beta, \emptyset) \models_n \varphi$.

Obviously, for all $\alpha \in \mathbf{BIT}_n$ and for all $\varphi \in \mathbf{FOR}_n$, $(\alpha, \emptyset) \models_n \Box\varphi$ and $(\alpha, \emptyset) \not\models_n \Diamond\varphi$. We say that $\varphi \in \mathbf{FOR}_n$ is *n-valid* if for all $(\alpha, S) \in \mathbf{MOD}_n$, $(\alpha, S) \models_n \varphi$. The soundness and the completeness of \mathbf{L}_2 with respect to this concept of validity is well-known. This is what Proposition 3 is about.

Proposition 3. *For all $\varphi \in \mathbf{FOR}_n$, $\varphi \in \mathbf{L}_2$ iff φ is n-valid.*

For all $\alpha \in \mathbf{BIT}_n$, the *n*-formula

$$\bullet \bar{x}^\alpha = \bigwedge \{x_i^{\alpha_i} : i \in \{1, \dots, n\}\}$$

exactly characterizes the propositional valuation of the variables x_1, \dots, x_n represented by α . This is what Proposition 4 is about.

Proposition 4. *For all $(\alpha, S) \in \mathbf{MOD}_n$ and for all $\beta \in \mathbf{BIT}_n$, $\alpha = \beta$ iff $(\alpha, S) \models_n \bar{x}^\beta$.*

For all $(\alpha, S) \in \mathbf{MOD}_n$, the *n*-formula

$$\bullet \mathbf{for}_n(\alpha, S) = \bar{x}^\alpha \wedge \Box \bigvee \{\bar{x}^\gamma : \gamma \in S\} \wedge \bigwedge \{\Diamond \bar{x}^\gamma : \gamma \in S\}$$

exactly characterizes the tree-like Kripke model of depth at most 1 represented by (α, S) . This is what Proposition 5 is about.

Proposition 5. *For all $(\alpha, S), (\beta, T) \in \mathbf{MOD}_n$, $(\alpha, S) = (\beta, T)$ iff $(\alpha, S) \models_n \mathbf{for}_n(\beta, T)$.*

As we know, an *n*-substitution (k, σ) is a homomorphism from \mathbf{FOR}_n to \mathbf{FOR}_k . Taking into account the duality between *n*-formulas and *n*-models, Propositions 6 and 7 tell us how to associate a function $g_{(k, \sigma)}$ from \mathbf{MOD}_k to \mathbf{MOD}_n to any *n*-substitution (k, σ) .

Proposition 6. *Let $(k, \sigma) \in \mathbf{SUB}_n$. For all $(\alpha, S) \in \mathbf{MOD}_k$, there exists $(\beta, T) \in \mathbf{MOD}_n$ such that $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T))$.*

Proposition 7. *Let $(k, \sigma) \in \mathbf{SUB}_n$. Let $(\alpha, S) \in \mathbf{MOD}_k$. For all $(\beta, T), (\gamma, U) \in \mathbf{MOD}_n$, if $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T))$ and $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\gamma, U))$ then $(\beta, T) = (\gamma, U)$.*

For all $(k, \sigma) \in \mathbf{SUB}_n$, let $g_{(k, \sigma)}$ be the function from \mathbf{MOD}_k to \mathbf{MOD}_n such that for all $(\alpha, S) \in \mathbf{MOD}_k$,

$$\bullet g_{(k, \sigma)}(\alpha, S) \text{ is the } (\beta, T) \in \mathbf{MOD}_n \text{ such that } (\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T)).$$

For all $(k, \sigma) \in \mathbf{SUB}_n$, notice that by Propositions 6 and 7, $g_{(k, \sigma)}$ is well-defined. Moreover, for all $(k, \sigma) \in \mathbf{SUB}_n$, for all $(\alpha, S) \in \mathbf{MOD}_k$ and for all $(\beta, T) \in \mathbf{MOD}_n$, if $g_{(k, \sigma)}(\alpha, S) = (\beta, T)$ then

- for all $\gamma \in S$, there exists $\delta \in T$ such that $g_{(k, \sigma)}(\gamma, \emptyset) = (\delta, \emptyset)$,
- for all $\delta \in T$, there exists $\gamma \in S$ such that $g_{(k, \sigma)}(\gamma, \emptyset) = (\delta, \emptyset)$.

See Proposition 8 below. Obviously, the above conditions are very similar to the forward condition and backward condition of bounded morphisms usually considered in modal logic [9, Definition 2.10]. This motivates the following definition. For all $k \geq 1$, a *(k, n)-morphism* is a function f from \mathbf{MOD}_k to \mathbf{MOD}_n such that for all $(\alpha, S) \in \mathbf{MOD}_k$ and for all $(\beta, T) \in \mathbf{MOD}_n$, if $f(\alpha, S) = (\beta, T)$ then

- for all $\gamma \in S$, there exists $\delta \in T$ such that $f(\gamma, \emptyset) = (\delta, \emptyset)$,
- for all $\delta \in T$, there exists $\gamma \in S$ such that $f(\gamma, \emptyset) = (\delta, \emptyset)$.

Proposition 8. *For all $(k, \sigma) \in \mathbf{SUB}_n$, $g_{(k, \sigma)}$ is a (k, n) -morphism.*

However, the morphisms described here should not be mistaken for the bounded morphisms. In particular, in the above definition, there is no condition related to the propositional valuations of the variables. For all $(k, \sigma) \in \mathbf{SUB}_n$, the best we can say about the propositional valuations of the variables concerns $g_{(k, \sigma)}$ and is contained in the following result.

Proposition 9. *For all $(k, \sigma) \in \mathbf{SUB}_n$ and for all $(\alpha, S), (\beta, T) \in \mathbf{MOD}_k$, if $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$ then for all $i \in \{1, \dots, n\}$, $(\alpha, S) \models_k \sigma(x_i)$ iff $(\beta, T) \models_k \sigma(x_i)$.*

Nevertheless, it is not particularly surprising that we have the following results.

Proposition 10. *Let $k \geq 1$. Let f be a (k, n) -morphism. Let $(\beta, T) \in \mathbf{MOD}_k$ and $(\gamma, U) \in \mathbf{MOD}_n$. If $f(\beta, T) = (\gamma, U)$ then the image by f of $T \times \{\emptyset\}$ is equal to $U \times \{\emptyset\}$. Moreover, $T = \emptyset$ iff $U = \emptyset$.*

Proposition 11. *Let $k \geq 1$. Let f be a (k, n) -morphism. Let $(\beta, T) \in \mathbf{MOD}_k$ and $(\gamma, U) \in \mathbf{MOD}_n$. If the following conditions hold then $f(\beta, T) = (\gamma, U)$:*

- $f(\beta, T) \models_n \bar{x}^\gamma$,
- for all $\delta \in T$, there exists $\epsilon \in U$ such that $f(\delta, \emptyset) = (\epsilon, \emptyset)$,
- for all $\epsilon \in U$, there exists $\delta \in T$ such that $f(\delta, \emptyset) = (\epsilon, \emptyset)$.

5 Unification

Let $n \geq 1$. An n -unifier of $\varphi \in \mathbf{FOR}_n$ is an n -substitution (k, σ) such that $\sigma(\varphi) \in \mathbf{L}_2$. We say that $\varphi \in \mathbf{FOR}_n$ is n -unifiable if there exists an n -unifier of φ . We say that a set Σ of n -unifiers of an n -unifiable $\varphi \in \mathbf{FOR}_n$ is n -complete if for all n -unifiers (k, σ) of φ , there exists $(l, \tau) \in \Sigma$ such that $(l, \tau) \preceq_n (k, \sigma)$. We say that an n -complete set Σ of n -unifiers of an n -unifiable $\varphi \in \mathbf{FOR}_n$ is *minimal* if for all n -complete sets Δ of n -unifiers of φ , if $\Delta \subseteq \Sigma$ then $\Delta = \Sigma$. As is well-known, for all $\varphi \in \mathbf{FOR}_n$, if φ is n -unifiable then for all minimal n -complete sets Σ, Δ of n -unifiers of φ , Σ and Δ have the same cardinality. Then, an important question is the following: when $\varphi \in \mathbf{FOR}_n$ is n -unifiable, is there a minimal n -complete set of n -unifiers of φ ? When the answer is “yes”, how large is this set? For all n -unifiable $\varphi \in \mathbf{FOR}_n$, we say that:

- φ is n -nullary if there exists no minimal complete set of unifiers of φ ,
- φ is n -infinite if there exists a minimal complete set of unifiers of φ with infinite cardinality,
- φ is n -finite if there exists a minimal complete set of unifiers of φ with finite cardinality ≥ 2 ,
- φ is n -unitary if there exists a minimal complete set of unifiers of φ with cardinality 1.

Obviously, considered as an n -formula, $\diamond x_1 \rightarrow \square x_1$ is n -unifiable. Indeed, let (n, v_\perp) and (n, v_\top) be the n -substitutions defined by:

- $v_\perp(x_1) = \perp$ and $v_\top(x_1) = \top$,
- for all $i \in \{2, \dots, n\}$, $v_\perp(x_i) = x_i$ and $v_\top(x_i) = x_i$.

Obviously, $v_\perp(\diamond x_1 \rightarrow \square x_1) \in \mathbf{L}_2$ and $v_\top(\diamond x_1 \rightarrow \square x_1) \in \mathbf{L}_2$. Hence, (n, v_\perp) and (n, v_\top) are n -unifiers of $\diamond x_1 \rightarrow \square x_1$. Moreover,

Proposition 12. *The n -unifiable n -formula $\diamond x_1 \rightarrow \square x_1$ is n -finite.*

For all n -unifiable $\varphi \in \mathbf{FOR}_n$ and for all $\pi \geq 1$, we say that φ is n - π -reasonable if for all n -unifiers (k, σ) of φ , if $k \geq \pi$ then there exists an n -unifier (l, τ) of φ such that $(l, \tau) \preceq_n (k, \sigma)$ and $l \leq \pi$. The idea behind the concept of reasonableness is simple: a unifiable formula is reasonable when a bounded set of variables suffices to express the set of all its unifiers. Since \mathbf{L}_2 is locally tabular, it is not particularly surprising that we have the following result.

Proposition 13. *Let $\varphi \in \mathbf{FOR}_n$ be n -unifiable and $\pi \geq 1$. If φ is n - π -reasonable then φ is either n -finitary, or n -unitary.*

As a result, in order to prove that n -unifiable n -formulas are either n -finitary, or n -unitary, it suffices to prove that n -unifiable n -formulas are n - n -reasonable. This is what Proposition 14 asserts below.

6 Main result

Let $n \geq 1$. Our aim is now to prove that n -unifiable n -formulas do not require more variables than the variables x_1, \dots, x_n in order to express their unifiers. Let $\varphi \in \mathbf{FOR}_n$. Suppose φ is n -unifiable. Let (k, σ) be an n -unifier of φ such that $k \geq n$. To achieve our aim, it suffices to construct an n -unifier (n, τ) of φ such that $(n, \tau) \preceq_n (k, \sigma)$. The construction of (n, τ) is based on the definition of a surjective (k, n) -morphism f such that for all $(\alpha, S), (\beta, T) \in \mathbf{MOD}_k$, if $f(\alpha, S) = f(\beta, T)$ then $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$. See Lemmas 4, 5 and 6 below. The argument leading to the definition of f is based on a sequence of combinatorial facts which proofs require twice the use of Proposition 1. See Lemmas 1, 2 and 3 below. Now, let us start. Let \sim_k be the equivalence relation on \mathbf{MOD}_k defined by:

- $(\alpha, S) \sim_k (\beta, T)$ iff $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$.

Lemma 1. 1. $\|\mathbf{MOD}_k^{\text{deg}} / \sim_k\| \leq \|\mathbf{MOD}_n^{\text{deg}}\|$,

- 2. $\|\mathbf{MOD}_n^{\text{deg}}\| \leq \|\mathbf{MOD}_k^{\text{deg}}\|$.

Hence, by Proposition 1 and Lemma 1, there exists a surjective function f^{deg} from $\mathbf{MOD}_k^{\text{deg}}$ to $\mathbf{MOD}_n^{\text{deg}}$ such that for all $(\alpha, \emptyset), (\beta, \emptyset) \in \mathbf{MOD}_k^{\text{deg}}$, if $f^{\text{deg}}(\alpha, \emptyset) = f^{\text{deg}}(\beta, \emptyset)$ then $(\alpha, \emptyset) \sim_k (\beta, \emptyset)$.

Lemma 2. *For all nonempty sets S, T of k -tuples of bits, if the images by f^{deg} of $S \times \{\emptyset\}$ and $T \times \{\emptyset\}$ are equal then the images by $g_{(k, \sigma)}$ of $S \times \{\emptyset\}$ and $T \times \{\emptyset\}$ are equal.*

For all nonempty sets E of n -tuples of bits, let

- $f^\circ(E)$ be the set of all $(\alpha, S) \in \mathbf{MOD}_k \setminus \mathbf{MOD}_k^{\text{deg}}$ such that the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$,
- $f^\bullet(E)$ be the set of all $(\alpha, S) \in \mathbf{MOD}_n \setminus \mathbf{MOD}_n^{\text{deg}}$ such that $S = E$.

Notice that for all nonempty sets E of n -tuples of bits, since f^{deg} is surjective, $\|f^\circ(E)\| \geq 2^k$. Notice also that for all nonempty sets E of n -tuples of bits, $\|f^\bullet(E)\| = 2^n$.

Lemma 3. *For all nonempty sets E of n -tuples of bits,*

- 1. $\|f^\circ(E) / \sim_k\| \leq \|f^\bullet(E)\|$,
- 2. $\|f^\bullet(E)\| \leq \|f^\circ(E)\|$.

Thus, for all nonempty sets E of n -tuples of bits, by Proposition 1 and Lemma 3, there exists a surjective function f^E from $f^\circ(E)$ to $f^\bullet(E)$ such that for all $(\alpha, S), (\beta, T) \in f^\circ(E)$, if $f^E(\alpha, S) = f^E(\beta, T)$ then $(\alpha, S) \sim_k (\beta, T)$. Let f be the function from \mathbf{MOD}_k to \mathbf{MOD}_n such that for all $(\alpha, \emptyset) \in \mathbf{MOD}_k^{\text{deg}}$,

- $f(\alpha, \emptyset) = f^{\text{deg}}(\alpha, \emptyset)$

and for all $(\alpha, S) \in \text{MOD}_k \setminus \text{MOD}_k^{\text{deg}}$, E being the nonempty set of n -tuples of bits such that the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$,

- $f(\alpha, S) = f^E(\alpha, S)$.

Lemma 4. f is surjective.

Lemma 5. f is a (k, n) -morphism.

Lemma 6. For all $(\alpha, S), (\beta, T) \in \text{MOD}_k$, if $f(\alpha, S) = f(\beta, T)$ then $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$.

The reader is invited to notice how Proposition 1 has been used — twice — in the argument leading to the definition of f . Let $(n, \tau), (k, \nu)$ be the n -substitutions defined by:

- $\tau(x_i) = \bigvee \{ \text{for}_n(f(\alpha, S)) : (\alpha, S) \in \text{MOD}_k \text{ is such that } (\alpha, S) \models_k \sigma(x_i) \}$ where $i \in \{1, \dots, n\}$,
- $\nu(x_i) = \bigvee \{ \text{for}_k(\alpha, S) : (\alpha, S) \in \text{MOD}_k \text{ is such that } f(\alpha, S) \models_n x_i \}$ where $i \in \{1, \dots, n\}$.

In Lemmas 7–10 below, we prove interesting properties of these n -substitutions: Lemma 7 is used for showing that (n, τ) is an n -unifier of φ , Lemma 8 is used in the proof of Lemma 9, Lemma 9 is used in the proof of Lemma 10 and Lemma 10 is used for showing that $(n, \tau) \preceq_n (k, \sigma)$.

Lemma 7. Let $\psi \in \text{FOR}_n$. For all $(\beta, T) \in \text{MOD}_n$, the following conditions are equivalent: (i) there exists $(\alpha, S) \in \text{MOD}_k$ such that $f(\alpha, S) = (\beta, T)$ and $(\alpha, S) \models_k \sigma(\psi)$; (ii) for all $(\alpha, S) \in \text{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\psi)$; (iii) $(\beta, T) \models_n \tau(\psi)$.

Lemma 8. For all $(\beta, T) \in \text{MOD}_k$ and for all $i \in \{1, \dots, n\}$, $(\beta, T) \models_k \nu(x_i)$ iff $f(\beta, T) \models_n x_i$.

Lemma 9. For all $(\beta, T) \in \text{MOD}_k$ and for all $(\gamma, U) \in \text{MOD}_n$, $f(\beta, T) = (\gamma, U)$ iff $(\beta, T) \models_k \nu(\text{for}_n(\gamma, U))$.

Lemma 10. For all $(\beta, T) \in \text{MOD}_k$ and for all $i \in \{1, \dots, n\}$, $(\beta, T) \models_k \nu(\tau(x_i))$ iff $(\beta, T) \models_k \sigma(x_i)$.

Since (k, σ) is an n -unifier of φ , $\sigma(\varphi) \in \mathbf{L}_2$. Thus, by Proposition 3, $\sigma(\varphi)$ is k -valid, i.e. for all $(\alpha, S) \in \text{MOD}_k$, $(\alpha, S) \models_k \sigma(\varphi)$. Consequently, by Lemma 7, for all $(\beta, T) \in \text{MOD}_n$, $(\beta, T) \models_n \tau(\varphi)$, i.e. $\tau(\varphi)$ is n -valid. Hence, by Proposition 3, $\tau(\varphi) \in \mathbf{L}_2$. Thus, (n, τ) is an n -unifier of φ . Since by Lemma 10, $(n, \tau) \preceq_n (k, \sigma)$, φ is n - n -reasonable. Since $\varphi \in \text{FOR}_n$ is arbitrary, this proves the following result.

Proposition 14. For all $\varphi \in \text{FOR}_n$, if φ is n -unifiable then φ is n - n -reasonable.

Propositions 13 and 14, lead us to the following result.

Theorem 1. For all $\varphi \in \text{FOR}_n$, if φ is n -unifiable then φ is either n -finitary, or n -unitary.

In the light of Proposition 12 and Theorem 1, we therefore conclude that \mathbf{L}_2 is finitary.

7 Conclusion

In this paper, within the context of elementary unification, we have proved Theorem 1 asserting that in \mathbf{L}_2 , unifiable formulas are either finitary, or unitary. We believe that in the line of reasoning leading to Theorem 1, the main properties of \mathbf{L}_2 are the ones given in Propositions 2, 6 and 7. Proposition 2 says that \mathbf{L}_2 is locally tabular — it is used in the proof of Proposition 13. For all $n \geq 1$, Propositions 6

and 7 tell us how to associate a function $g_{(k,\sigma)}$ from \mathbf{MOD}_k to \mathbf{MOD}_n to any n -substitution (k, σ) — they are used in the proof of Proposition 14. Notice that Theorem 1 is an immediate consequence of Propositions 13 and 14. However, since \mathbf{L}_2 is locally tabular, the reader may think that the line of reasoning leading to Theorem 1 is unnecessarily complicated. In other respect, since the unification type is a categorical invariant, the reader may think that it is possible to directly work in the category of all finite models of \mathbf{L}_2 . Indeed, the categorical approach to the unification problem in propositional logic is powerful [16]. Nevertheless, the fact that the question of the unification type of \mathbf{L}_2 has not been solved before indicates that things may not be so simple.

Here are open questions: (i) determine the unification type of the locally tabular modal logic $\mathbf{K} + \Box^d\perp$ for each $d \geq 3$, (ii) determine the unification types of the locally tabular modal logics studied in [24, 25, 26], (iii) determine the unification types of the modal logics \mathbf{KB} , \mathbf{KD} and \mathbf{KT} . We conjecture that the modal logics mentioned in (i) and (ii) are either finitary, or unitary for elementary unification. As for the modal logics considered in (iii), it is only known that \mathbf{KD} and \mathbf{KT} are not unitary within the context of elementary unification and \mathbf{KB} , \mathbf{KD} and \mathbf{KT} are nullary within the context of unification with parameters [2, 3, 5]. We conjecture that the modal logics mentioned in (iii) are finitary within the context of elementary unification. And of course there is also the open question of the decidability of the unification problem in the modal logics considered in (i), (ii) and (iii). When the modal logics mentioned in (i) and (ii) are decidable, their local tabularity will probably imply the decidability of their unification problem. Concerning the modal logics considered in (iii), the decidability of their unification problem remains a mystery within the context of unification with parameters⁴.

Funding

The preparation of this paper has been supported by the Project *DN02/15/19.12.2016* of *Bulgarian Science Fund* and the Programme *Professeurs invités 2018* of *Université Paul Sabatier*.

Acknowledgement

We are indebted to Silvio Ghilardi for his suggestion to consider the question of the unification type of \mathbf{L}_2 . We also make a point of thanking for their feedback the referees of all preliminary versions of this paper: their useful suggestions have been essential for improving its correctness and its readability.

References

- [1] BAADER, F. and S. GHILARDI, ‘Unification in modal and description logics’, *Logic Journal of the IGPL* **19** (2011) 705–730.
- [2] BALBIANI, P., ‘Remarks about the unification type of several non-symmetric non-transitive modal logics’, *Logic Journal of the IGPL* **27** (2019) 639–658.
- [3] BALBIANI, P. and Ç. GENCER, ‘ \mathbf{KD} is nullary’, *Journal of Applied Non-Classical Logics* **27** (2017) 196–205.
- [4] BALBIANI, P., and Ç. GENCER, ‘Unification in epistemic logics’, *Journal of Applied Non-Classical Logics* **27** (2017) 91–105.
- [5] BALBIANI, P., and Ç. GENCER, ‘About the unification type of modal logics between \mathbf{KB} and \mathbf{KTB} ’, *Studia Logica* **108** (2020) 941–966.
- [6] BALBIANI, P., Ç. GENCER, M. ROSTAMIGIV and T. TINCHEV, ‘About the unification types of the modal logics determined by classes of deterministic frames’, arXiv:2004.07904v1 [cs.LO].

⁴As is well-known, the unification problem in \mathbf{KB} , \mathbf{KD} and \mathbf{KT} is in \mathbf{NP} within the context of elementary unification.

- [7] BALBIANI, P. and T. TINCHEV, ‘Unification in modal logic \mathbf{Alt}_1 ’, In: *Advances in Modal Logic*, College Publications (2016) 117–134.
- [8] BALBIANI, P., and T. TINCHEV, ‘Elementary unification in modal logic $\mathbf{KD45}$ ’, *Journal of Applied Logics* **5** (2018) 301–317.
- [9] BLACKBURN, P., M. DE RIJKE and Y. VENEMA, *Modal Logic*, Cambridge University Press (2001).
- [10] CHAGROV, A., and M. ZAKHARYASCHEV, *Modal Logic*, Oxford University Press (1997).
- [11] DZIK, W., ‘Unitary unification of $\mathbf{S5}$ modal logics and its extensions’, *Bulletin of the Section of Logic* **32** (2003) 19–26.
- [12] DZIK, W., *Unification Types in Logic*, Wydawnictwo Uniwersytetu Slaskiego (2007).
- [13] DZIK, W. and P. WOJTYLAK, ‘Projective unification in modal logic’, *Logic Journal of the IGPL* **20** (2012) 121–153.
- [14] GABBAY, D. and V. SHEHTMAN, ‘Products of modal logics, part 1’, *Logic Journal of the IGPL* **6** (1998) 73–146.
- [15] GHILARDI, S., ‘Best solving modal equations’, *Annals of Pure and Applied Logic* **102** (2000) 183–198.
- [16] GHILARDI, S., ‘Unification, finite duality and projectivity in varieties of Heyting algebras’, *Annals of Pure and Applied Logic* **127** (2004) 99–115.
- [17] GHILARDI, S. and L. SACCHETTI, ‘Filtering unification and most general unifiers in modal logic’, *Journal of Symbolic Logic* **69** (2004) 879–906.
- [18] GORANKO, V. and M. OTTO, ‘Model theory of modal logic’, In: *Handbook of Modal Logic*, Elsevier (2007) 249–329.
- [19] IEMHOFF, R., ‘A syntactic approach to unification in transitive reflexive modal logics’, *Notre Dame Journal of Formal Logic* **57** (2016) 233–247.
- [20] JEŘÁBEK, E., ‘Logics with directed unification’, In: *Algebra and Coalgebra meet Proof Theory*, Workshop at Utrecht University (2013).
- [21] JEŘÁBEK, E., ‘Blending margins: the modal logic \mathbf{K} has nullary unification type’, *Journal of Logic and Computation* **25** (2015) 1231–1240.
- [22] KOST, S., ‘Projective unification in transitive modal logics’, *Logic Journal of the IGPL* **26** (2018) 548–566.
- [23] KRACHT, M., *Tools and Techniques in Modal Logic*, Elsevier (1999).
- [24] MIYAZAKI, Y., ‘Normal modal logics containing \mathbf{KTB} with some finiteness conditions’, In: *Advances in Modal Logic*, College Publications (2004) 171–190.
- [25] NAGLE, M. and S. THOMASON, ‘The extensions of the modal logic $\mathbf{K5}$ ’, *Journal of Symbolic Logic* **50** (1985) 102–109.
- [26] SHAPIROVSKY, I. and V. SHEHTMAN, ‘Local tabularity without transitivity’, In: *Advances in Modal Logic*, College Publications (2016) 520–534.

Annex

Proof of Proposition 1: Suppose S is nonempty. Let \sim be an equivalence relation on S .

Suppose $\|S/\sim\| \leq \|T\| \leq \|S\|$. Let h be a function from S/\sim to S such that for all $\alpha \in S$, $h([\alpha]) \in [\alpha]$, i.e. h is a function selecting an element in each equivalence class modulo \sim ⁵. Obviously, h is injective. Let $S_0 = \{h([\alpha]) : \alpha \in S\}$. Since h is injective, $\|S/\sim\| = \|S_0\|$. Since $\|S/\sim\| \leq \|T\|$, $\|S_0\| \leq \|T\|$. Let T_0 be a subset of T such that $\|T_0\| = \|S_0\|$. Let f_0 be a one-to-one correspondence between S_0 and T_0 . Let $T_1 = T \setminus T_0$. Notice that T_0 and T_1 make a partition of T . Since $\|T\| \leq \|S\|$ and $\|T_0\| = \|S_0\|$, $\|T_1\| \leq \|S \setminus S_0\|$. Let S_1 be a subset of $S \setminus S_0$ such that $\|S_1\| = \|T_1\|$. Let f_1 be a one-to-one correspondence between S_1 and T_1 . Let $S_2 = (S \setminus S_0) \setminus S_1$. Let f_2 be the function from S_2 to T such that for

⁵Since S is finite, the proof of the existence of \sim does not require the use of the axiom of choice.

all $\alpha \in S_2$, $f_2(\alpha) = f_0(h([\alpha]))$. Let f be the function from S to T defined by $f|_{S_0} = f_0$, $f|_{S_1} = f_1$ and $f|_{S_2} = f_2$.

Claim. f is surjective.

Proof. Let $\beta \in T$. We consider the following cases.

Case $\beta \in T_0$. Since f_0 is one-to-one, let $\alpha \in S_0$ be such that $f_0(\alpha) = \beta$. Thus, $\alpha \in S$. Moreover, $f(\alpha) = f_0(\alpha)$. Since $f_0(\alpha) = \beta$, $f(\alpha) = \beta$.

Case $\beta \in T_1$. Since f_1 is one-to-one, let $\alpha \in S_1$ be such that $f_1(\alpha) = \beta$. Hence, $\alpha \in S$. Moreover, $f(\alpha) = f_1(\alpha)$. Since $f_1(\alpha) = \beta$, $f(\alpha) = \beta$.

Claim. For all $\alpha, \beta \in S$, if $f(\alpha) = f(\beta)$ then $\alpha \sim \beta$.

Proof. Let $\alpha, \beta \in S$ be such that $f(\alpha) = f(\beta)$. We consider the following cases.

Case $\alpha \in S_0$ and $\beta \in S_0$. Consequently, $f(\alpha) = f_0(\alpha)$ and $f(\beta) = f_0(\beta)$. Since $f(\alpha) = f(\beta)$, $f_0(\alpha) = f_0(\beta)$. Since f_0 is one-to-one, $\alpha = \beta$. Thus, $\alpha \sim \beta$.

Case $\alpha \in S_0$ and $\beta \in S_1$. Consequently, $f(\alpha) = f_0(\alpha)$ and $f(\beta) = f_1(\beta)$. Since $f(\alpha) = f(\beta)$, $f_0(\alpha) = f_1(\beta)$. Since $f_0(\alpha) \in T_0$ and $f_1(\beta) \in T_1$, T_0 and T_1 do not make a partition of T : a contradiction.

Case $\alpha \in S_0$ and $\beta \in S_2$. Hence, $f(\alpha) = f_0(\alpha)$ and $f(\beta) = f_2(\beta)$. Since $f(\alpha) = f(\beta)$, $f_0(\alpha) = f_2(\beta)$. Thus, $f_0(\alpha) = f_0(h([\beta]))$. Since f_0 is one-to-one, $\alpha = h([\beta])$. Since $h([\beta]) \in [\beta]$, $\alpha \in [\beta]$. Consequently, $\alpha \sim \beta$.

Case $\alpha \in S_1$ and $\beta \in S_1$. Hence, $f(\alpha) = f_1(\alpha)$ and $f(\beta) = f_1(\beta)$. Since $f(\alpha) = f(\beta)$, $f_1(\alpha) = f_1(\beta)$. Since f_1 is one-to-one, $\alpha = \beta$. Thus, $\alpha \sim \beta$.

Case $\alpha \in S_1$ and $\beta \in S_2$. Hence, $f(\alpha) = f_1(\alpha)$ and $f(\beta) = f_2(\beta)$. Since $f(\alpha) = f(\beta)$, $f_1(\alpha) = f_2(\beta)$. Thus, $f_1(\alpha) = f_0(h([\beta]))$. Since $f_1(\alpha) \in T_1$ and $f_0(h([\beta])) \in T_0$, T_0 and T_1 do not make a partition of T : a contradiction.

Case $\alpha \in S_2$ and $\beta \in S_2$. Hence, $f(\alpha) = f_2(\alpha)$ and $f(\beta) = f_2(\beta)$. Since $f(\alpha) = f(\beta)$, $f_2(\alpha) = f_2(\beta)$. Consequently, $f_0(h([\alpha])) = f_0(h([\beta]))$. Since f_0 is one-to-one, $h([\alpha]) = h([\beta])$. Since $h([\alpha]) \in [\alpha]$ and $h([\beta]) \in [\beta]$, $[\alpha] \cap [\beta] \neq \emptyset$. Thus, $\alpha \sim \beta$.

Suppose f is a surjective function from S to T such that for all $\alpha, \beta \in S$, if $f(\alpha) = f(\beta)$ then $\alpha \sim \beta$. For the sake of the contradiction, suppose either $\|S/\sim\| > \|T\|$, or $\|T\| > \|S\|$. Since f is surjective, $\|T\| \leq \|S\|$. Since either $\|S/\sim\| > \|T\|$, or $\|T\| > \|S\|$, $\|S/\sim\| > \|T\|$. Let $p \in \mathbb{N}$ and $\beta^1, \dots, \beta^p \in S$ be such that $p > \|T\|$ and for all $q, r \in \mathbb{N}$, if $1 \leq q, r \leq p$ and $q \neq r$ then $\beta^q \not\sim \beta^r$. Hence, for all $q, r \in \mathbb{N}$, if $1 \leq q, r \leq p$ and $q \neq r$ then $f(\beta^q) \neq f(\beta^r)$. Thus, $p \leq \|T\|$: a contradiction.

Proof of Proposition 2: From [9, Proposition 2.29] and from the fact that for all $\varphi \in \mathbf{FOR}_n$, there exists $\psi \in \mathbf{FOR}_n$ such that $\deg_n(\psi) < 2$ and $\varphi \leftrightarrow \psi \in \mathbf{L}_2$.

Proof of Proposition 3: From [9, Proposition 2.6], from [9, Lemma 4.21] and from the fact that for all maximal consistent sets w of n -formulas, there exists $(\alpha, S) \in \mathbf{MOD}_n$ such that the submodel of the

canonical model of \mathbf{L}_2 generated by w is isomorphic to (α, S) .

Proof of Proposition 4: Let $(\alpha, S) \in \mathbf{MOD}_n$ and $\beta \in \mathbf{BIT}_n$. For the sake of the contradiction, suppose $\alpha = \beta$ not-iff $(\alpha, S) \models_n \bar{x}^\beta$. Hence, either $\alpha = \beta$ and $(\alpha, S) \not\models_n \bar{x}^\beta$, or $\alpha \neq \beta$ and $(\alpha, S) \models_n \bar{x}^\beta$. In the former case, for all $i \in \{1, \dots, n\}$, $\alpha_i = \beta_i$. Thus, for all $i \in \{1, \dots, n\}$, $(\alpha, S) \models_n x_i^{\beta_i}$. Consequently, $(\alpha, S) \models_n \bar{x}^\beta$: a contradiction. In the latter case, let $i \in \{1, \dots, n\}$ be such that $\alpha_i \neq \beta_i$. Hence, $(\alpha, S) \not\models_n x_i^{\beta_i}$. Thus, $(\alpha, S) \not\models_n \bar{x}^\beta$: a contradiction.

Proof of Proposition 5: From [18, Theorem 32], from the fact that for all $(\alpha, S) \in \mathbf{MOD}_n$, $\mathbf{for}_n(\alpha, S)$ characterizes n -models modulo bisimulation and from the fact that for all $(\alpha, S), (\beta, T) \in \mathbf{MOD}_n$, if (α, S) and (β, T) are bisimilar then $(\alpha, S) = (\beta, T)$.

Proof of Proposition 6: Let $(\alpha, S) \in \mathbf{MOD}_k$. Let β be the n -tuple of bits such that for all $i \in \{1, \dots, n\}$, if $(\alpha, S) \not\models_k \sigma(x_i)$ then $\beta_i = 0$ else $\beta_i = 1$. Let T be the least set of n -tuples of bits such that for all $\gamma \in S$, there exists $\delta \in T$ such that for all $i \in \{1, \dots, n\}$, if $(\gamma, \emptyset) \not\models_k \sigma(x_i)$ then $\delta_i = 0$ else $\delta_i = 1$. By induction on $\varphi \in \mathbf{FOR}_n$, the reader may easily verify that $(\alpha, S) \models_k \sigma(\varphi)$ iff $(\beta, T) \models_n \varphi$. Since by Proposition 5, $(\beta, T) \models_n \mathbf{for}_n(\beta, T)$, $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T))$.

Proof of Proposition 7: Let $(\beta, T), (\gamma, U) \in \mathbf{MOD}_n$. Suppose $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T))$ and $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\gamma, U))$. Hence, $(\alpha, S) \models_k \sigma(\bar{x}^\beta)$ and $(\alpha, S) \models_k \sigma(\bar{x}^\gamma)$. Thus, $\beta = \gamma$. Let $\beta' \in T$ be arbitrary. Since $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T))$, let $\alpha' \in S$ be such that $(\alpha', \emptyset) \models_k \sigma(\bar{x}^{\beta'})$. Since $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\gamma, U))$, let $\gamma' \in U$ be such that $(\alpha', \emptyset) \models_k \sigma(\bar{x}^{\gamma'})$. Since $(\alpha', \emptyset) \models_k \sigma(\bar{x}^{\beta'})$, $\beta' = \gamma'$. Consequently, $\beta' \in U$. Since β' is arbitrary, $T \subseteq U$. Reciprocally, the reader may easily verify that $U \subseteq T$. Hence, $T = U$. Since $\beta = \gamma$, $(\beta, T) = (\gamma, U)$.

Proof of Proposition 8: Let $(k, \sigma) \in \mathbf{SUB}_n$. For the sake of the contradiction, suppose $g_{(k, \sigma)}$ is not a (k, n) -morphism. Hence, let $(\alpha, S) \in \mathbf{MOD}_k$ and $(\beta, T) \in \mathbf{MOD}_n$ be such that $g_{(k, \sigma)}(\alpha, S) = (\beta, T)$ — and therefore $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\beta, T))$ — and either there exists $\gamma \in S$ such that for all $\delta \in T$, $g_{(k, \sigma)}(\gamma, \emptyset) \neq (\delta, \emptyset)$, or there exists $\delta \in T$ such that for all $\gamma \in S$, $g_{(k, \sigma)}(\gamma, \emptyset) \neq (\delta, \emptyset)$. In the former case, let $\delta' \in T$ be such that $(\gamma, \emptyset) \models_k \sigma(\bar{x}^{\delta'})$. Thus, $g_{(k, \sigma)}(\gamma, \emptyset) \neq (\delta', \emptyset)$. Since $(\gamma, \emptyset) \models_k \sigma(\bar{x}^{\delta'})$, $g_{(k, \sigma)}(\gamma, \emptyset) = (\delta', \emptyset)$: a contradiction. In the latter case, let $\gamma' \in S$ be such that $(\gamma', \emptyset) \models_k \sigma(\bar{x}^\delta)$. Consequently, $g_{(k, \sigma)}(\gamma', \emptyset) \neq (\delta, \emptyset)$. Since $(\gamma', \emptyset) \models_k \sigma(\bar{x}^\delta)$, $g_{(k, \sigma)}(\gamma', \emptyset) = (\delta, \emptyset)$: a contradiction.

Proof of Proposition 9: Let $(k, \sigma) \in \mathbf{SUB}_n$ and $(\alpha, S), (\beta, T) \in \mathbf{MOD}_k$. Suppose $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$. Hence, let $(\gamma, U) \in \mathbf{MOD}_n$ be such that $g_{(k, \sigma)}(\alpha, S) = (\gamma, U)$ and $g_{(k, \sigma)}(\beta, T) = (\gamma, U)$. Thus, $(\alpha, S) \models_k \sigma(\mathbf{for}_n(\gamma, U))$ and $(\beta, T) \models_k \sigma(\mathbf{for}_n(\gamma, U))$. Consequently, $(\alpha, S) \models_k \sigma(\bar{x}^\gamma)$ and $(\beta, T) \models_k \sigma(\bar{x}^\gamma)$. Hence, for all $i \in \{1, \dots, n\}$, $(\alpha, S) \models_k \sigma(x_i)$ iff $(\beta, T) \models_k \sigma(x_i)$.

Proof of Proposition 10: For the sake of the contradiction, suppose $f(\beta, T) = (\gamma, U)$ and the image by f of $T \times \{\emptyset\}$ is not equal to $U \times \{\emptyset\}$. Hence, either the image by f of $T \times \{\emptyset\}$ is not included in $U \times \{\emptyset\}$, or the image by f of $T \times \{\emptyset\}$ does not include to $U \times \{\emptyset\}$. In the former case, let $\delta \in T$ be such that $f(\delta, \emptyset) \notin U \times \{\emptyset\}$. Since f is a (k, n) -morphism and $f(\beta, T) = (\gamma, U)$, let $\epsilon' \in U$ be such that $f(\delta, \emptyset) = (\epsilon', \emptyset)$. Thus, $f(\delta, \emptyset) \in U \times \{\emptyset\}$: a contradiction. In the latter case, let $\epsilon \in U$ be such that $(\epsilon, \emptyset) \notin f(T \times \{\emptyset\})$. Since f is a (k, n) -morphism and $f(\beta, T) = (\gamma, U)$, let $\delta' \in T$ be such that $f(\delta', \emptyset) = (\epsilon, \emptyset)$. Consequently, $(\epsilon, \emptyset) \in f(T \times \{\emptyset\})$: a contradiction.

Proof of Proposition 11: For the sake of the contradiction, suppose $f(\beta, T) \neq (\gamma, U)$, $f(\beta, T) \models_n \bar{x}^\gamma$, for all $\delta \in T$, there exists $\epsilon \in U$ such that $f(\delta, \emptyset) = (\epsilon, \emptyset)$ and for all $\epsilon \in U$, there exists $\delta \in T$ such that $f(\delta, \emptyset) = (\epsilon, \emptyset)$. Let $(\gamma', U') \in \mathbf{MOD}_n$ be such that $f(\beta, T) = (\gamma', U')$. Since $f(\beta, T) \models_n \bar{x}^\gamma$, by Pro-

position 4, $\gamma=\gamma'$. Since $f(\beta, T)\neq(\gamma, U)$ and $f(\beta, T)=(\gamma', U')$, $U\neq U'$. Hence, either $U\not\subseteq U'$, or $U\not\supseteq U'$. In the former case, let $\epsilon'\in U$ be such that $\epsilon'\notin U'$. Since for all $\epsilon\in U$, there exists $\delta\in T$ such that $f(\delta, \emptyset)=(\epsilon, \emptyset)$, let $\delta'\in T$ be such that $f(\delta', \emptyset)=(\epsilon', \emptyset)$. Since f is a (k, n) -morphism and $f(\beta, T)=(\gamma', U')$, $\epsilon'\in U'$: a contradiction. In the latter case, let $\epsilon''\in U'$ be such that $\epsilon''\notin U$. Since f is a (k, n) -morphism and $f(\beta, T)=(\gamma', U')$, let $\delta''\in T$ be such that $f(\delta'', \emptyset)=(\epsilon'', \emptyset)$. Since for all $\delta\in T$, there exists $\epsilon\in U$ such that $f(\delta, \emptyset)=(\epsilon, \emptyset)$, $\epsilon''\in U$: a contradiction.

Proof of Proposition 12: Let (n, σ) and (n, τ) be the n -substitutions defined by:

- $\sigma(x_1)=\square\perp \vee x_1$ and $\tau(x_1)=\diamond\top \wedge x_1$,
- for all $i\in\{2, \dots, n\}$, $\sigma(x_i)=x_i$ and $\tau(x_i)=x_i$.

Obviously, $\sigma(\diamond x_1 \rightarrow \square x_1)\in\mathbf{L}_2$ and $\tau(\diamond x_1 \rightarrow \square x_1)\in\mathbf{L}_2$. Hence, (n, σ) and (n, τ) are n -unifiers of $\diamond x_1 \rightarrow \square x_1$. In order to prove that $\diamond x_1 \rightarrow \square x_1$ is n -finitary, it suffices to prove that $\{(n, \sigma), (n, \tau)\}$ is a minimal n -complete set of n -unifiers of $\diamond x_1 \rightarrow \square x_1$.

n -completeness of $\{(n, \sigma), (n, \tau)\}$: Let (k, v) be an arbitrary n -unifier of $\diamond x_1 \rightarrow \square x_1$. Thus, $v(\diamond x_1 \rightarrow \square x_1)\in\mathbf{L}_2$. By using the semantics of \mathbf{L}_2 , it follows that either $\square\perp \rightarrow v(x_1)\in\mathbf{L}_2$, or $v(x_1) \rightarrow \diamond\top\in\mathbf{L}_2$. Indeed, for the sake of the contradiction, suppose neither $\square\perp \rightarrow v(x_1)\in\mathbf{L}_2$, nor $v(x_1) \rightarrow \diamond\top\in\mathbf{L}_2$. Consequently, by Proposition 3, neither $\square\perp \rightarrow v(x_1)$ is k -valid, nor $v(x_1) \rightarrow \diamond\top$ is k -valid. Hence, let $(\alpha, \emptyset), (\beta, \emptyset)\in\mathbf{MOD}_k^{\text{deg}}$ be such that $(\alpha, \emptyset)\not\models_k v(x_1)$ and $(\beta, \emptyset)\models_k v(x_1)$. Let $\gamma\in\mathbf{BIT}_k$. Since $(\alpha, \emptyset)\not\models_k v(x_1)$ and $(\beta, \emptyset)\models_k v(x_1)$, $(\gamma, \{\alpha, \beta\})\not\models_k v(\diamond x_1 \rightarrow \square x_1)$. Thus, $v(\diamond x_1 \rightarrow \square x_1)$ is not k -valid. Consequently, by Proposition 3, $v(\diamond x_1 \rightarrow \square x_1)\notin\mathbf{L}_2$: a contradiction. In the former case where $\square\perp \rightarrow v(x_1)\in\mathbf{L}_2$, it follows immediately that $v(\sigma(x_1))\equiv_k v(x_1)$. Hence, $(n, \sigma)\preceq_n(k, v)$. In the latter case where $v(x_1) \rightarrow \diamond\top\in\mathbf{L}_2$, it follows immediately that $v(\tau(x_1))\equiv_{\mathbf{L}_2} v(x_1)$. Thus, $(n, \tau)\preceq_n(k, v)$.

Minimality of $\{(n, \sigma), (n, \tau)\}$: For the sake of the contradiction, suppose $\{(n, \sigma), (n, \tau)\}$ is not minimal. Consequently, either $(n, \sigma)\preceq_n(n, \tau)$, or $(n, \tau)\preceq_n(n, \sigma)$. In the former case, there exists an n -substitution (n, v) such that $v(\sigma(x_1))\equiv_n \tau(x_1)$. Hence, $\square\perp \vee v(x_1)\equiv_n \diamond\top \wedge x_1$. In the latter case, there exists a substitution (n, v) such that $v(\tau(x_1))\equiv_n \sigma(x_1)$. Thus, $\diamond\top \wedge v(x_1)\equiv_n \square\perp \vee x_1$. In both cases, $\square\perp \rightarrow \diamond\top\in\mathbf{L}_2$. Consequently, $\diamond\top\in\mathbf{L}_2$: a contradiction.

Proof of Proposition 13: Suppose φ is n - π -reasonable. Let Σ be the set of all n -unifiers of φ . Notice that Σ is n -complete. Let Σ' be the set of n -substitutions obtained from Σ by keeping only the n -substitutions (k, σ) such that $k\leq\pi$. Since φ is n - π -reasonable and Σ is n -complete, Σ' is n -complete. Let Σ'' be the set of n -substitutions obtained from Σ' by keeping only one representative of each equivalence class modulo \simeq_n . Since Σ' is n -complete, Σ'' is n -complete. Moreover, by Proposition 2, Σ'' is finite. Hence, either φ is n -finitary, or φ is n -unitary.

Proof of Lemma 1: (1) For the sake of the contradiction, suppose $\|\mathbf{MOD}_k^{\text{deg}}/\sim_k\|>\|\mathbf{MOD}_n^{\text{deg}}\|$. Let $p\in\mathbb{N}$ and $(\alpha^1, \emptyset), \dots, (\alpha^p, \emptyset)\in\mathbf{MOD}_k^{\text{deg}}$ be such that $p>\|\mathbf{MOD}_n^{\text{deg}}\|$ and for all $q, r\in\mathbb{N}$, if $1\leq q, r\leq p$ and $q\neq r$ then $(\alpha^q, \emptyset)\not\sim_k(\alpha^r, \emptyset)$. Hence, for all $q, r\in\mathbb{N}$, if $1\leq q, r\leq p$ and $q\neq r$ then $g_{(k, \sigma)}(\alpha^q, \emptyset)\neq g_{(k, \sigma)}(\alpha^r, \emptyset)$. Since $g_{(k, \sigma)}$ is a (k, n) -morphism, let $\beta^1, \dots, \beta^p\in\mathbf{BIT}_n$ be such that $g_{(k, \sigma)}(\alpha^1, \emptyset)=(\beta^1, \emptyset)$, \dots , $g_{(k, \sigma)}(\alpha^p, \emptyset)=(\beta^p, \emptyset)$. Since for all $q, r\in\mathbb{N}$, if $1\leq q, r\leq p$ and $q\neq r$ then $g_{(k, \sigma)}(\alpha^q, \emptyset)\neq g_{(k, \sigma)}(\alpha^r, \emptyset)$, for all $q, r\in\mathbb{N}$, if $1\leq q, r\leq p$ and $q\neq r$ then $\beta^q\neq\beta^r$. Thus, $p\leq 2^n$. Since $\|\mathbf{MOD}_n^{\text{deg}}\|=2^n$, $p\leq\|\mathbf{MOD}_n^{\text{deg}}\|$: a contradiction.

(2) Since $\|\mathbf{MOD}_n^{\text{deg}}\|=2^n$, $\|\mathbf{MOD}_k^{\text{deg}}\|=2^k$ and $k\geq n$, $\|\mathbf{MOD}_n^{\text{deg}}\|\leq\|\mathbf{MOD}_k^{\text{deg}}\|$.

Proof of Lemma 2: Let S, T be nonempty sets of k -tuples of bits. Suppose the images by f^{deg} of $S \times \{\emptyset\}$ and $T \times \{\emptyset\}$ are equal. For the sake of the contradiction, suppose the images by $g_{(k,\sigma)}$ of $S \times \{\emptyset\}$ and $T \times \{\emptyset\}$ are not equal. Since $g_{(k,\sigma)}$ is a (k, n) -morphism, let $(\gamma, \emptyset) \in \mathbf{MOD}_n^{\text{deg}}$ be such that either (γ, \emptyset) is in the image by $g_{(k,\sigma)}$ of $S \times \{\emptyset\}$ without being in the image by $g_{(k,\sigma)}$ of $T \times \{\emptyset\}$, or (γ, \emptyset) is in the image by $g_{(k,\sigma)}$ of $T \times \{\emptyset\}$ without being in the image by $g_{(k,\sigma)}$ of $S \times \{\emptyset\}$. Without loss of generality, suppose (γ, \emptyset) is in the image by $g_{(k,\sigma)}$ of $S \times \{\emptyset\}$ without being in the image by $g_{(k,\sigma)}$ of $T \times \{\emptyset\}$. Hence, let $\alpha \in S$ be such that $g_{(k,\sigma)}(\alpha, \emptyset) = (\gamma, \emptyset)$. Since the images by f^{deg} of $S \times \{\emptyset\}$ and $T \times \{\emptyset\}$ are equal, let $\beta \in T$ be such that $f^{\text{deg}}(\alpha, \emptyset) = f^{\text{deg}}(\beta, \emptyset)$. Thus, $(\alpha, \emptyset) \sim_k (\beta, \emptyset)$. Consequently, $g_{(k,\sigma)}(\alpha, \emptyset) = g_{(k,\sigma)}(\beta, \emptyset)$. Since $g_{(k,\sigma)}(\alpha, \emptyset) = (\gamma, \emptyset)$, (γ, \emptyset) is in the image by $g_{(k,\sigma)}$ of $T \times \{\emptyset\}$: a contradiction.

Proof of Lemma 3: Let E be a nonempty set of n -tuples of bits.

(1) For the sake of the contradiction, suppose $\|f^\circ(E)/\sim_k\| > \|f^\bullet(E)\|$. Let $p \in \mathbb{N}$ and $(\alpha^1, S_1), \dots, (\alpha^p, S_p) \in \mathbf{MOD}_k \setminus \mathbf{MOD}_k^{\text{deg}}$ be such that $p > \|f^\bullet(E)\|$, the images by f^{deg} of $S_1 \times \{\emptyset\}, \dots, S_p \times \{\emptyset\}$ are equal to $E \times \{\emptyset\}$ and for all $q, r \in \mathbb{N}$, if $1 \leq q, r \leq p$ and $q \neq r$ then $(\alpha^q, S_q) \not\sim_k (\alpha^r, S_r)$. Hence, for all $q, r \in \mathbb{N}$, if $1 \leq q, r \leq p$ and $q \neq r$ then $g_{(k,\sigma)}(\alpha^q, S_q) \neq g_{(k,\sigma)}(\alpha^r, S_r)$. Since $g_{(k,\sigma)}$ is a (k, n) -morphism and the images by f^{deg} of $S_1 \times \{\emptyset\}, \dots, S_p \times \{\emptyset\}$ are equal to $E \times \{\emptyset\}$, let $\beta^1, \dots, \beta^p \in \mathbf{BIT}_n$ be such that $g_{(k,\sigma)}(\alpha^1, S_1) = (\beta^1, E), \dots, g_{(k,\sigma)}(\alpha^p, S_p) = (\beta^p, E)$. Since for all $q, r \in \mathbb{N}$, if $1 \leq q, r \leq p$ and $q \neq r$ then $g_{(k,\sigma)}(\alpha^q, S_q) \neq g_{(k,\sigma)}(\alpha^r, S_r)$, for all $q, r \in \mathbb{N}$, if $1 \leq q, r \leq p$ and $q \neq r$ then $\beta^q \neq \beta^r$. Thus, $p \leq 2^n$. Since $\|f^\bullet(E)\| = 2^n$, $p \leq \|f^\bullet(E)\|$: a contradiction.

(2) Since $k \geq n$, $\|f^\circ(E)\| \geq 2^k$ and $\|f^\bullet(E)\| = 2^n$, $\|f^\bullet(E)\| \leq \|f^\circ(E)\|$.

Proof of Lemma 4: Let $\beta \in \mathbf{BIT}_n$ and T be a set of n -tuples of bits. We consider the following cases.

Case $(\beta, T) \in \mathbf{MOD}_n^{\text{deg}}$. Since f^{deg} is surjective, let $\alpha \in \mathbf{BIT}_k$ be such that $f^{\text{deg}}(\alpha, \emptyset) = (\beta, \emptyset)$. Hence, $f(\alpha, \emptyset) = (\beta, T)$.

Case $(\beta, T) \in \mathbf{MOD}_n \setminus \mathbf{MOD}_n^{\text{deg}}$. Thus, $(\beta, T) \in f^\bullet(T)$. Since f^T is surjective, let $(\alpha, S) \in f^\circ(T)$ be such that $f^T(\alpha, S) = (\beta, T)$. Consequently, $(\alpha, S) \in \mathbf{MOD}_k \setminus \mathbf{MOD}_k^{\text{deg}}$ and the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $T \times \{\emptyset\}$. Hence, $f(\alpha, S) = f^T(\alpha, S)$. Since $f^T(\alpha, S) = (\beta, T)$, $f(\alpha, S) = (\beta, T)$.

Proof of Lemma 5: For the sake of the contradiction, suppose f is not a (k, n) -morphism. Hence, let $(\alpha, S) \in \mathbf{MOD}_k$ and $(\beta, T) \in \mathbf{MOD}_n$ be such that $f(\alpha, S) = (\beta, T)$ and either there exists $\gamma' \in S$ such that for all $\delta' \in T$, $f(\gamma', \emptyset) \neq (\delta', \emptyset)$, or there exists $\delta'' \in T$ such that for all $\gamma'' \in S$, $f(\gamma'', \emptyset) \neq (\delta'', \emptyset)$. In the former case, $S \neq \emptyset$. Thus, $(\alpha, S) \in \mathbf{MOD}_k \setminus \mathbf{MOD}_k^{\text{deg}}$ and, E being the nonempty set of n -tuples of bits such that the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$, $f(\alpha, S) = f^E(\alpha, S)$. Since $f(\alpha, S) = (\beta, T)$, $f^E(\alpha, S) = (\beta, T)$. Consequently, $T = E$. Since the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$, the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $T \times \{\emptyset\}$. Hence, let $\delta \in T$ be such that $f(\gamma', \emptyset) = (\delta, \emptyset)$. Since for all $\delta' \in T$, $f(\gamma', \emptyset) \neq (\delta', \emptyset)$, $f(\gamma', \emptyset) \neq (\delta, \emptyset)$: a contradiction. In the latter case, $T \neq \emptyset$. Since $f(\alpha, S) = (\beta, T)$, $S \neq \emptyset$. Thus, $(\alpha, S) \in \mathbf{MOD}_k \setminus \mathbf{MOD}_k^{\text{deg}}$ and, E being the nonempty set of n -tuples of bits such that the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$, $f(\alpha, S) = f^E(\alpha, S)$. Since $f(\alpha, S) = (\beta, T)$, $f^E(\alpha, S) = (\beta, T)$. Consequently, $T = E$. Since the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$, the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $T \times \{\emptyset\}$. Hence, let $\gamma \in S$ be such that $f(\gamma, \emptyset) = (\delta'', \emptyset)$. Since for all $\gamma'' \in S$, $f(\gamma'', \emptyset) \neq (\delta'', \emptyset)$, $f(\gamma, \emptyset) \neq (\delta'', \emptyset)$: a contradiction.

Proof of Lemma 6: Let $(\alpha, S), (\beta, T) \in \mathbf{MOD}_k$. Suppose $f(\alpha, S) = f(\beta, T)$. We consider the following cases.

Case $S = \emptyset$. Since $f(\alpha, S) = f(\beta, T)$, by Lemma 5, $T = \emptyset$. Since $S = \emptyset$, $(\alpha, S), (\beta, T) \in \mathbf{MOD}_k^{\text{deg}}$. Hence, $f(\alpha, S) = f^{\text{deg}}(\alpha, S)$ and $f(\beta, T) = f^{\text{deg}}(\beta, T)$. Since $f(\alpha, S) = f(\beta, T)$, $f^{\text{deg}}(\alpha, S) = f^{\text{deg}}(\beta, T)$. Thus, $(\alpha, S) \sim_k (\beta, T)$. Consequently, $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$.

Case $S \neq \emptyset$. Since $f(\alpha, S) = f(\beta, T)$, by Lemma 5, $T \neq \emptyset$. Since $S \neq \emptyset$, $(\alpha, S), (\beta, T) \in \mathbf{MOD}_k \setminus \mathbf{MOD}_k^{\text{deg}}$. Hence, $f(\alpha, S) = f^E(\alpha, S)$ and $f(\beta, T) = f^F(\beta, T)$, E being the nonempty set of n -tuples of bits such that the image by f^{deg} of $S \times \{\emptyset\}$ is equal to $E \times \{\emptyset\}$ and F being the nonempty set of n -tuples of bits such that the image by f^{deg} of $T \times \{\emptyset\}$ is equal to $F \times \{\emptyset\}$. Since $f(\alpha, S) = f(\beta, T)$, $f^E(\alpha, S) = f^F(\beta, T)$. Thus, $E = F$. Since $f^E(\alpha, S) = f^F(\beta, T)$, $(\alpha, S) \sim_k (\beta, T)$. Consequently, $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\beta, T)$.

Proof of Lemma 7: By induction on ψ . Let $(\beta, T) \in \mathbf{MOD}_n$. We consider the following cases.

Case $\psi = x_i$ for some $i \in \{1, \dots, n\}$. **(i) \Rightarrow (ii)** Suppose $f(\alpha, S) = (\beta, T)$ and $(\alpha, S) \models_k \sigma(x_i)$ for some $(\alpha, S) \in \mathbf{MOD}_k$. Let $(\alpha', S') \in \mathbf{MOD}_k$. Suppose $f(\alpha', S') = (\beta, T)$. Since $f(\alpha, S) = (\beta, T)$, $f(\alpha, S) = f(\alpha', S')$. Hence, $g_{(k, \sigma)}(\alpha, S) = g_{(k, \sigma)}(\alpha', S')$. Thus, by Proposition 9, $(\alpha, S) \models_k \sigma(x_i)$ iff $(\alpha', S') \models_k \sigma(x_i)$. Since $(\alpha, S) \models_k \sigma(x_i)$, $(\alpha', S') \models_k \sigma(x_i)$.

(ii) \Rightarrow (iii) Suppose for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(x_i)$. Since f is surjective, let $(\alpha', S') \in \mathbf{MOD}_k$ be such that $f(\alpha', S') = (\beta, T)$. Since for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(x_i)$, $(\alpha', S') \models_k \sigma(x_i)$. Consequently, $\text{for}_n(f(\alpha', S'))$ is one of the disjuncts of $\tau(x_i)$. Since $f(\alpha', S') = (\beta, T)$, by Proposition 5, $(\beta, T) \models_n \tau(x_i)$.

(iii) \Rightarrow (i) Suppose $(\beta, T) \models_n \tau(x_i)$. Hence, let $(\alpha, S) \in \mathbf{MOD}_k$ be such that $(\alpha, S) \models_k \sigma(x_i)$ and $(\beta, T) \models_n \text{for}_n(f(\alpha, S))$. Thus, by Proposition 5, $f(\alpha, S) = (\beta, T)$.

Case $\psi = \perp$. **(i) \Rightarrow (ii)** Obviously, the condition “ $f(\alpha, S) = (\beta, T)$ and $(\alpha, S) \models_k \sigma(\perp)$ for some $(\alpha, S) \in \mathbf{MOD}_k$ ” cannot hold.

(ii) \Rightarrow (iii) Since f is surjective, the condition “for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\perp)$ ” cannot hold.

(iii) \Rightarrow (i) Obviously, the condition “ $(\beta, T) \models_n \tau(\perp)$ ” cannot hold.

Case $\psi = \neg\chi$. **(i) \Rightarrow (ii)** Suppose $f(\alpha, S) = (\beta, T)$ and $(\alpha, S) \models_k \sigma(\neg\chi)$ for some $(\alpha, S) \in \mathbf{MOD}_k$. Let $(\alpha', S') \in \mathbf{MOD}_k$. Suppose $f(\alpha', S') = (\beta, T)$. For the sake of the contradiction, suppose $(\alpha', S') \not\models_k \sigma(\neg\chi)$. Consequently, $(\alpha', S') \models_k \sigma(\chi)$. Since $f(\alpha, S) = (\beta, T)$ and $f(\alpha', S') = (\beta, T)$, by induction hypothesis, $(\alpha, S) \models_k \sigma(\chi)$. Hence, $(\alpha, S) \not\models_k \sigma(\neg\chi)$: a contradiction.

(ii) \Rightarrow (iii) Suppose for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\neg\chi)$. For the sake of the contradiction, suppose $(\beta, T) \not\models_n \tau(\neg\chi)$. Thus, $(\beta, T) \models_n \tau(\chi)$. Since f is surjective, let $(\alpha', S') \in \mathbf{MOD}_k$ be such that $f(\alpha', S') = (\beta, T)$. Since for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\neg\chi)$, $(\alpha', S') \models_k \sigma(\neg\chi)$. Consequently, $(\alpha', S') \not\models_k \sigma(\chi)$. Since $f(\alpha', S') = (\beta, T)$, by induction hypothesis, $(\beta, T) \not\models_n \tau(\chi)$: a contradiction.

(iii) \Rightarrow (i) Suppose $(\beta, T) \models_n \tau(\neg\chi)$. Since f is surjective, let $(\alpha, S) \in \mathbf{MOD}_k$ be such that $f(\alpha, S) = (\beta, T)$. For the sake of the contradiction, suppose $(\alpha, S) \not\models_k \sigma(\neg\chi)$. Hence, $(\alpha, S) \models_k \sigma(\chi)$. Since $f(\alpha, S) = (\beta, T)$, by induction hypothesis, $(\beta, T) \models_n \tau(\chi)$. Thus, $(\beta, T) \not\models_n \tau(\neg\chi)$: a contradiction.

Case $\psi = \chi \vee \theta$. (i) \Rightarrow (ii) Suppose $f(\alpha, S) = (\beta, T)$ and $(\alpha, S) \models_k \sigma(\chi \vee \theta)$ for some $(\alpha, S) \in \mathbf{MOD}_k$. Let $(\alpha', S') \in \mathbf{MOD}_k$. Suppose $f(\alpha', S') = (\beta, T)$. For the sake of the contradiction, suppose $(\alpha', S') \not\models_k \sigma(\chi \vee \theta)$. Consequently, neither $(\alpha', S') \models_k \sigma(\chi)$, nor $(\alpha', S') \models_k \sigma(\theta)$. Since $f(\alpha, S) = (\beta, T)$ and $f(\alpha', S') = (\beta, T)$, by induction hypothesis, neither $(\alpha, S) \models_k \sigma(\chi)$, nor $(\alpha, S) \models_k \sigma(\theta)$. Hence, $(\alpha, S) \not\models_k \sigma(\chi \vee \theta)$: a contradiction.

(ii) \Rightarrow (iii) Suppose for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\chi \vee \theta)$. For the sake of the contradiction, suppose $(\beta, T) \not\models_n \tau(\chi \vee \theta)$. Thus, neither $(\beta, T) \models_n \tau(\chi)$, nor $(\beta, T) \models_n \tau(\theta)$. Since f is surjective, let $(\alpha', S') \in \mathbf{MOD}_k$ be such that $f(\alpha', S') = (\beta, T)$. Since for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\chi \vee \theta)$, $(\alpha', S') \models_k \sigma(\chi \vee \theta)$. Consequently, either $(\alpha', S') \models_k \sigma(\chi)$, or $(\alpha', S') \models_k \sigma(\theta)$. Since $f(\alpha', S') = (\beta, T)$, by induction hypothesis, either $(\beta, T) \models_n \tau(\chi)$, or $(\beta, T) \models_n \tau(\theta)$: a contradiction.

(iii) \Rightarrow (i) Suppose $(\beta, T) \models_n \tau(\chi \vee \theta)$. Since f is surjective, let $(\alpha, S) \in \mathbf{MOD}_k$ be such that $f(\alpha, S) = (\beta, T)$. For the sake of the contradiction, suppose $(\alpha, S) \not\models_k \sigma(\chi \vee \theta)$. Hence, neither $(\alpha, S) \models_k \sigma(\chi)$, nor $(\alpha, S) \models_k \sigma(\theta)$. Since $f(\alpha, S) = (\beta, T)$, by induction hypothesis, neither $(\beta, T) \models_n \tau(\chi)$, nor $(\beta, T) \models_n \tau(\theta)$. Thus, $(\beta, T) \not\models_n \tau(\chi \vee \theta)$: a contradiction.

Case $\psi = \square\chi$. (i) \Rightarrow (ii) Suppose there exists $(\alpha, S) \in \mathbf{MOD}_k$ such that $f(\alpha, S) = (\beta, T)$ and $(\alpha, S) \models_k \sigma(\square\chi)$. Let $(\alpha', S') \in \mathbf{MOD}_k$. Suppose $f(\alpha', S') = (\beta, T)$. For the sake of the contradiction, suppose $(\alpha', S') \not\models_k \sigma(\square\chi)$. Consequently, let $\gamma' \in S'$ be such that $(\gamma', \emptyset) \not\models_k \sigma(\chi)$. Since f is a (k, n) -morphism and $f(\alpha', S') = (\beta, T)$, let $\delta \in T$ be such that $f(\gamma', \emptyset) = (\delta, \emptyset)$. Since f is a (k, n) -morphism and $f(\alpha, S) = (\beta, T)$, let $\gamma \in S$ be such that $f(\gamma, \emptyset) = (\delta, \emptyset)$. Since $(\gamma', \emptyset) \not\models_k \sigma(\chi)$ and $f(\gamma', \emptyset) = (\delta, \emptyset)$, by induction hypothesis, $(\gamma, \emptyset) \not\models_k \sigma(\chi)$. Hence, $(\alpha, S) \not\models_k \sigma(\square\chi)$: a contradiction.

(ii) \Rightarrow (iii) Suppose for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\square\chi)$. For the sake of the contradiction, suppose $(\beta, T) \not\models_n \tau(\square\chi)$. Thus, let $\delta \in T$ be such that $(\delta, \emptyset) \not\models_n \tau(\chi)$. Since f is surjective, let $(\alpha', S') \in \mathbf{MOD}_k$ be such that $f(\alpha', S') = (\beta, T)$. Since for all $(\alpha, S) \in \mathbf{MOD}_k$, if $f(\alpha, S) = (\beta, T)$ then $(\alpha, S) \models_k \sigma(\square\chi)$, $(\alpha', S') \models_k \sigma(\square\chi)$. Since f is a (k, n) -morphism and $f(\alpha', S') = (\beta, T)$, let $\gamma' \in S'$ be such that $f(\gamma', \emptyset) = (\delta, \emptyset)$. Since $(\delta, \emptyset) \not\models_n \tau(\chi)$, by induction hypothesis, $(\gamma', \emptyset) \not\models_k \sigma(\chi)$. Consequently, $(\alpha', S') \not\models_k \sigma(\square\chi)$: a contradiction.

(iii) \Rightarrow (i) Suppose $(\beta, T) \models_n \tau(\square\chi)$. Since f is surjective, let $(\alpha, S) \in \mathbf{MOD}_k$ be such that $f(\alpha, S) = (\beta, T)$. For the sake of the contradiction, suppose $(\alpha, S) \not\models_k \sigma(\square\chi)$. Hence, let $\gamma \in S$ be such that $(\gamma, \emptyset) \not\models_k \sigma(\chi)$. Since f is a (k, n) -morphism and $f(\alpha, S) = (\beta, T)$, let $\delta \in T$ be such that $f(\gamma, \emptyset) = (\delta, \emptyset)$. Since $(\gamma, \emptyset) \not\models_k \sigma(\chi)$, by induction hypothesis, $(\delta, \emptyset) \not\models_n \tau(\chi)$. Thus, $(\beta, T) \not\models_n \tau(\square\chi)$: a contradiction.

Proof of Lemma 8: Let $(\beta, T) \in \mathbf{MOD}_k$ and all $i \in \{1, \dots, n\}$. For the sake of the contradiction, suppose either $(\beta, T) \models_k \nu(x_i)$ and $f(\beta, T) \not\models_n x_i$, or $(\beta, T) \not\models_k \nu(x_i)$ and $f(\beta, T) \models_n x_i$. In the former case, by definition of ν , let $(\alpha, S) \in \mathbf{MOD}_k$ be such that $f(\alpha, S) \models_n x_i$ and $(\beta, T) \not\models_k \text{for}_k(\alpha, S)$. Hence, by Proposition 5, $(\beta, T) = (\alpha, S)$. Since $f(\alpha, S) \models_n x_i$, $f(\beta, T) \not\models_n x_i$: a contradiction. In the latter case, by definition of ν , $\text{for}_k(\beta, T)$ is one of the disjuncts of $\nu(x_i)$. Since by Proposition 5, $(\beta, T) \models \text{for}_k(\beta, T)$, $(\beta, T) \not\models_k \nu(x_i)$: a contradiction.

Proof of Lemma 9: Let $(\beta, T) \in \mathbf{MOD}_k$ and $(\gamma, U) \in \mathbf{MOD}_n$. For the sake of the contradiction, suppose either $f(\beta, T) = (\gamma, U)$ and $(\beta, T) \not\models_k \nu(\mathbf{for}_n(\gamma, U))$, or $f(\beta, T) \neq (\gamma, U)$ and $(\beta, T) \models_k \nu(\mathbf{for}_n(\gamma, U))$. In the former case, since by Proposition 4, $(\gamma, U) \models_n \bar{x}^\gamma$, $f(\beta, T) \models_n \bar{x}^\gamma$. Hence, by Lemma 8, $(\beta, T) \models_k \nu(\bar{x}^\gamma)$. Since f is a (k, n) -morphism and by Proposition 4, for all $\gamma''', \gamma'''' \in U$, $(\gamma''', \emptyset) \models_n \bar{x}^{\gamma''''}$ iff $\gamma''' = \gamma''''$, for all $\beta' \in T$, there exists $\gamma' \in U$ such that $f(\beta', \emptyset) \models_n \bar{x}^{\gamma'}$ and for all $\gamma'' \in U$, there exists $\beta'' \in T$ such that $f(\beta'', \emptyset) \models_n \bar{x}^{\gamma''}$. Thus, by Lemma 8, for all $\beta' \in T$, there exists $\gamma' \in U$ such that $(\beta', \emptyset) \models_k \nu(\bar{x}^{\gamma'})$ and for all $\gamma'' \in U$, there exists $\beta'' \in T$ such that $(\beta'', \emptyset) \models_k \nu(\bar{x}^{\gamma''})$. Since $(\beta, T) \models_k \nu(\bar{x}^\gamma)$, $(\beta, T) \models_k \nu(\mathbf{for}_n(\gamma, U))$: a contradiction. In the latter case, $(\beta, T) \models_k \nu(\bar{x}^\gamma)$. Moreover, for all $\beta' \in T$, there exists $\gamma' \in U$ such that $(\beta', \emptyset) \models_k \nu(\bar{x}^{\gamma'})$ and for all $\gamma'' \in U$, there exists $\beta'' \in T$ such that $(\beta'', \emptyset) \models_k \nu(\bar{x}^{\gamma''})$. Consequently, by Lemma 8, for all $\beta' \in T$, there exists $\gamma' \in U$ such that $f(\beta', \emptyset) \models_n \bar{x}^{\gamma'}$ and for all $\gamma'' \in U$, there exists $\beta'' \in T$ such that $f(\beta'', \emptyset) \models_n \bar{x}^{\gamma''}$. Since f is a (k, n) -morphism and by Proposition 4, for all $\gamma''', \gamma'''' \in U$, $(\gamma''', \emptyset) \models_n \bar{x}^{\gamma''''}$ iff $\gamma''' = \gamma''''$, for all $\beta' \in T$, there exists $\gamma' \in U$ such that $f(\beta', \emptyset) = (\gamma', \emptyset)$ and for all $\gamma'' \in U$, there exists $\beta'' \in T$ such that $f(\beta'', \emptyset) = (\gamma'', \emptyset)$. Since $(\beta, T) \models_k \nu(\bar{x}^\gamma)$, by Proposition 11, $f(\beta, T) = (\gamma, U)$: a contradiction.

Proof of Lemma 10: Let $(\beta, T) \in \mathbf{MOD}_k$ and $i \in \{1, \dots, n\}$. For the sake of the contradiction, suppose either $(\beta, T) \models_k \nu(\tau(x_i))$ and $(\beta, T) \not\models_k \sigma(x_i)$, or $(\beta, T) \not\models_k \nu(\tau(x_i))$ and $(\beta, T) \models_k \sigma(x_i)$. In the former case, by definition of τ , let $(\alpha, S) \in \mathbf{MOD}_k$ be such that $(\alpha, S) \models_k \sigma(x_i)$ and $(\beta, T) \models_k \nu(\mathbf{for}_n(f(\alpha, S)))$. Hence, by Lemma 9, $f(\beta, T) = f(\alpha, S)$. Thus, $g_{(k, \sigma)}(\beta, T) = g_{(k, \sigma)}(\alpha, S)$. Consequently, by Proposition 9, $(\beta, T) \models_k \sigma(x_i)$ iff $(\alpha, S) \models_k \sigma(x_i)$. Since $(\alpha, S) \models_k \sigma(x_i)$, $(\beta, T) \models_k \sigma(x_i)$: a contradiction. In the latter case, by definition of τ , $\mathbf{for}_n(f(\beta, T))$ is one of the disjuncts of $\tau(x_i)$. Since by Lemma 9, $(\beta, T) \models_k \nu(\mathbf{for}_n f(\beta, T))$, $(\beta, T) \models_k \nu(\tau(x_i))$: a contradiction.