



HAL
open science

2D-Block Geminals: a non 1-orthogonal and non 0-seniority model with reduced computational complexity

Patrick Cassam-Chenaï, Thomas Perez, Davide Accomasso

► **To cite this version:**

Patrick Cassam-Chenaï, Thomas Perez, Davide Accomasso. 2D-Block Geminals: a non 1-orthogonal and non 0-seniority model with reduced computational complexity. 2022. hal-03763128v1

HAL Id: hal-03763128

<https://hal.science/hal-03763128v1>

Preprint submitted on 1 Sep 2022 (v1), last revised 3 Jan 2023 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2D-Block Geminals: a non 1-orthogonal and non 0-seniority model with reduced computational complexity

Patrick Cassam-Chenai, Thomas Perez¹ and Davide Accomasso²

¹*Université Côte d'Azur, LJAD, UMR 7351, 06100 Nice, France*

²*Dipartimento di Chimica e Chimica Industriale, Università di Pisa, via Moruzzi 13, 56124 Pisa, Italy*

We present a new geminal product wave function ansatz where the geminals are not constrained to be strongly orthogonal nor to be of seniority zero. Instead, we introduce weaker orthogonality constraints between geminals which significantly lower the computational effort, without sacrificing the indistinguishability of the electrons. That is to say, the electron pairs corresponding to the geminals are not fully distinguishable, and their product has still to be antisymmetrized according to the Pauli principle to form a *bona fide* electronic wave function. Our geometrical constraints translate into simple equations involving the traces of products of our geminal matrices. In the simplest non-trivial model, a set of solutions is given by block-diagonal matrices where each block is of size 2×2 and consists of either a Pauli matrix or a normalized diagonal matrix, multiplied by a complex parameter to be optimized. With this simplified ansatz for geminals, the number of terms in the calculation of the matrix elements of quantum observables is considerably reduced. A proof of principle is reported and confirms that the ansatz is more accurate than strongly orthogonal geminal products while remaining computationally affordable.

Keywords:

Strongly correlated electron pairs, antisymmetric geminal products, Pauli matrices

Suggested running head:

2D-Block Geminals

All correspondance to be send to P. Cassam-Chenai,

cassam@unice.fr,

I. INTRODUCTION

The accurate, *ab initio* calculation of the electronic energy levels of a molecular system is still a challenging problem with exponential complexity with respect to the number of particles. This is particularly true for the so-called “strongly correlated” systems, where many Slater determinants have to be included in the wave function to achieve a qualitatively correct description of the molecular state. In such systems, it has been established that better ansätze can be based on an electron pair model, that is to say, an antisymmetric product of two-electron wave functions, called “geminals”.

However, without further restrictions, such a model has still a factorial computational cost with respect to the number of electronic orbitals and its applicability is therefore limited to small systems. Reducing the scaling of the computational cost to a polynomial one is an active research field¹⁻³. We postpone to the next section a review of the most popular proposals. Here, we just note that they either enforce the “strong orthogonality” constraint which breaks the indistinguishability of electron pairs, and is thus at odds with the Pauli principle, or have been limited to seniority zero systems, that is to say, to systems with no unpaired electronic configuration.

In the past, we have introduced generalized orthogonality constraints between geminals to reduce the computational effort, without sacrificing the indistinguishability of the electrons nor assuming a given seniority^{4,5}. In other words, the electron pairs corresponding to the geminals were not fully distinguishable, so that geminal products had still to be antisymmetrized according to the Pauli principle to form *bona fide* electronic wave functions, and each geminal combined linearly electronic configurations of mixed seniorities. The aim of the present work is to pursue our exploration of orthogonality constraints to improve the computational affordability of antisymmetric product of geminals (APG).

Recall that an APG wave function can be described in an alternative and practical

way, by a set of matrices, one for each geminal. Our geometrical constraints translated in terms of these geminal matrices, into simple equations involving the traces of their products. Here, we will impose further geometrical constraints whose solutions will lead us to consider block-diagonal matrices, where each block is at most of size 2×2 . These blocks consist essentially in a Pauli or a diagonal matrix, multiplied by a complex parameter to be optimized. With this simple ansatz for geminals, the number of terms in the calculation of quantum observable matrix elements is considerably reduced.

The paper is organized as follows: In the next section we present an overview of existing APG-based methods. Then, we obtain an overlap formula for the general APG case, which exhibits its full complexity and serves us as a starting point to elaborate new constraints able to make it more amenable to practical computations. So, in the following section, we propose to enforce the so-called “permutationally invariant 2-orthogonality constraints” and show how we arrive at the 2D-block geminal model. Finally, we obtain the necessary quantities to implement our ansatz and provide a proof of principle that it is able to give strictly lower energies than strongly orthogonal geminal products. In the last section, we conclude on the prospect of our method.

II. ANTISYMMETRIZED PRODUCT OF GEMINALS (APG) ANSÄTZE: AN OVERVIEW

Many APG models have appeared in the literature. We provide here a quick overview to help the reader in situating the model proposed in the following sections. We recall in Appendix A, the generalized concept of seniority, and the definition of p -orthogonality, because these two notions make the constraints generally imposed on APG wave functions amenable to a synthetic presentation.

A. General APG ansatz

A general APG can be written as

$$\Phi = g^1 \wedge g^2 \wedge \cdots \wedge g^p \quad (1)$$

with

$$g^r = \sum_{i=1}^{2m} \sum_{j>i}^{2m} \lambda_{i,j}^r \varphi_i \wedge \varphi_j \quad . \quad (2)$$

The φ_i are a basis set but not necessarily spin-adapted, that is to say, they can be linear combination involving both spin α and spin β spin-orbitals. Each geminal is parametrized by $m(2m - 1)$ independent scalar (real or complex) numbers $\lambda_{i,j}^r$.

However, even in the spin-adapted case, optimizing such ansatz is extremely costly from a computational standpoint⁶ and has rarely been attempted^{5,7}. All popular geminal ansätze can be seen as APG with additional constraints. The latter can be divided into “intra-geminal” constraints, bearing on the spin-orbitals of the geminal expansion Eq.(2), and “inter-geminal” constraints, imposing certain relationships between the geminals of the product Eq.(1).

B. Intra-geminal constraints

1. *APsetG ansatz:*

A first approach consists in partitioning the spin-orbital basis into two subsets of cardinal m , $\mathcal{H} = \mathcal{H}_\alpha \oplus \mathcal{H}_\beta$, ($\dim \mathcal{H}_\alpha = \dim \mathcal{H}_\beta = m$), and to expand the geminals on two-electron basis functions of generalized seniority equal to 2 with respect to this partition:

$$g_{\text{set}}^r = \sum_{i=1}^m \sum_{j=1}^m \lambda_{i,j}^r \varphi_{2i-1} \wedge \varphi_{2j} \quad , \quad (3)$$

where the subscript “set” hints to the fact that we have two disjoint sets of identical cardinal. One can choose for example to gather all spin-orbitals of spin α in the first set and all those of spin β in the second set. Then, one has wave functions that are eigenfunctions of \hat{S}_z associated with the eigenvalue 0.

Geminals of this type are parametrized by m^2 independent scalar numbers.

The Grassmann product of those geminals is called an APsetG (“Antisymmetric Product of set-divided Geminals”)⁸ wave function:

$$\Phi_{\text{APsetG}} = \bigwedge_{r=1}^k g_{\text{set}}^r \quad . \quad (4)$$

2. *APIG ansatz:*

An alternative idea is to partition the spin-orbital basis set into m shells of the form $(\varphi_{2i-1}, \varphi_{2i})$ and to build geminals out of seniority 0 two-electron functions exclusively:

$$g_{\text{I}}^r = \sum_{i=1}^m \lambda_i^r \varphi_{2i-1} \wedge \varphi_{2i} \quad , \quad (5)$$

where the subscript “I” stands for “Interacting”, as the Grassmann product of those geminals is known as the APIG wave function (“Antisymmetric Product of Interacting Geminals”)⁸

$$\Phi_{\text{APIG}} = \bigwedge_{r=1}^k g_{\text{sen0}}^r \quad . \quad (6)$$

Each geminals in the product is parametrized by m independent scalar numbers.

Note that, if the shells of the partition are such that $\forall i$, φ_{2i-1} and φ_{2i} are spin-orbitals of the same spatial part and of opposite spin, one retrieve the usual concept of seniority,

and the APIG wave functions are those considered by Silver⁹ and revived recently by Limacher¹⁰. In this case a more natural notation is:

$$g_I^r = \sum_{i=1}^m \lambda_i^r \varphi_i \wedge \bar{\varphi}_i \quad . \quad (7)$$

Most of other APG ansätze in use are particular cases of the APIG ansatz, hence of seniority 0. Let us mention the antisymmetric product of rank-two geminals, APr2G, where the coefficients are constrained to assume the form^{11,12}: $\lambda_i^r = \frac{1}{a_r \epsilon_i - b_i \lambda_r}$, $a_r, \epsilon_i, b_i, \lambda_r \in \mathbb{C}$. When these numbers are derived as eigenstate coefficients of reduced Bardeen-Cooper-Schrieffer (BCS) model Hamiltonians¹³, one obtains the so-called Richardson–Gaudin (RG) geminals which are currently under active development.^{1,14}

C. Inter-geminal constraints

1. *AP1roG ansatz*:

Another important subcase of the APIG model consists in imposing the following additional constraint: each geminal is only allowed to have a nonzero coefficient on a single pair of spin-orbitals (numbered from 1 to k) pertaining to an Hartree-Fock type reference wave function (with k doubly occupied orbitals):

$$g_{1ro}^r = \varphi_r \wedge \bar{\varphi}_r + \sum_{i=k+1}^m \lambda_i^r \varphi_i \wedge \bar{\varphi}_i \quad , \quad (8)$$

where the subscript “1ro” stands for “1-reference orbital”. Note that the constraint has an intra-geminal character but has also an inter-geminal one, since two different geminals must have different doubly occupied reference orbitals. Each 1ro-geminal is parametrized by $m - k$ independent scalar numbers.

The Grassmann product of those geminals is called the “AP1roG” wave function (“Antisymmetric Product of 1-reference orbital Geminals”)^{12,15–19}:

$$\Phi_{\text{AP1roG}} = \bigwedge_{r=1}^k g_{1\text{ro}}^r \quad . \quad (9)$$

2. *APSG ansatz:*

The 1-orthogonality constraint (see Appendix VII B) between the geminals in the Grassmann product has often been added as an inter-geminal condition. One obtains in this way an APSG (“Antisymmetrized Product of Strongly-orthogonal Geminals”) wave functions^{9,20–28}.

The 1-orthogonality constraint reduces drastically the computational cost, however, it is usually too stringent to achieve chemical accuracy.

3. *AGP ansatz:*

At the other extreme, one can take all k geminals, not just non-orthogonal but even equal. Then, one obtains a product wave function called “AGP” for “Antisymmetrized Geminal Power”²⁹. Dropping the subscript and denoting g the geminal involved, we have:

$$\Phi_{\text{AGP}} = g^{\wedge k} = \underbrace{g \wedge \dots \wedge g}_{k \text{ times}} \quad . \quad (10)$$

4. *GMFCI p -orthogonal ansätze:*

The p -orthogonality constraint, $1 \leq p \leq \text{inf}(n_1, n_2)$ between an n_1 -electron wave function and an n_2 -electron one (see Appendix VII B), can be tuned from strong orthogonality

($p = 1$), to “weak orthogonality” ($p = \inf(n_1, n_2)$), to restrict variational freedom. Note that in the case ($n_1 = n_2 = n$) weak orthogonality ($p = n$) is just the usual orthogonality between n -electron wave functions.

Two such constraints in between the APSG and AGP extreme cases, have been considered in the past for Geminal Mean Field Configuration Interaction (GMFCI) wave functions⁵: (i) 2-orthogonality between every pairs of geminals (g^{r_1}, g^{r_2}) of the APG; (ii) 2-orthogonality between any geminal g^r and any Grassmann product of the remaining ones $\bigwedge_{s \neq r} g^s$.

5. *Various post-treatments:*

There have been many proposals to improve APSG or AP1roG wave functions by adding perturbative corrections^{30–35}. Other attempts have consisted in applying linearized coupled cluster corrections to APSG or AP1roG reference wave functions^{36,37}. However, these improvements were at the expense of the simplicity of the antisymmetrized geminal product model.

Many recent works aim at improving the AGP ansatz. A promising method called “CJAGP” for “Cluster-Jastrow Antisymmetrized Geminal Power”^{38,39}, or its “JAGP” particular case when atomic orbitals are not optimized^{40–42}, have been proposed by Neuscamman and can achieve polynomial cost, but with a large prefactor. Beside the latter approach qualified as “subtractive” according to Neuscamman’s interesting terminology, AGPs were also considered as reference wave functions for more classical “additive” approaches^{43,44}.

Note also that, very recently the AGP ansatz⁴⁵ and an unitary version of AP1roG⁴⁶ have been ported to quantum computers.

In order to propose new constraints to enforce on the APG wave functions, we find appropriate to return first to the general case in the next section.

III. GENERAL APG OVERLAP FORMULA

In general, in quantum chemistry, the observables of interest such as the Hamiltonian, contain at most two-particle operators. So, the computational cost of an electronic wave function method is essentially given by the cost of calculating overlap integrals.

In this section, we provide a general formula, proved in Appendix VII C for the overlap between two wave functions which are products of geminals, i.e. to compute $\langle \Phi_1 \wedge \dots \wedge \Phi_n | \Phi'_1 \wedge \dots \wedge \Phi'_n \rangle$, where the Φ_k 's and the Φ'_k 's are general ($S_z = 0$)-geminals.

We will use a spin-adapted orthonormal basis set for the one-electron Hilbert space \mathcal{H} (of finite dimension $2m$), that is made of two subsets $(\varphi_i)_{1 \leq i \leq m}$ and $(\bar{\varphi}_i)_{1 \leq i \leq m}$, the latter being, respectively, spin α and spin β spin-orbitals of the same set of orbitals.

We associate with each geminal, Φ_k , a matrix of size $m \times m$, denoted C_k , defined as follows:

$$\forall k \in \{1, \dots, n\}, C_k = \left(\langle \Phi_k | \varphi_i \wedge \bar{\varphi}_j \rangle \right)_{\substack{1 \leq i \leq m \\ 1 \leq j \leq m}} \quad , \quad (11)$$

so that:

$$\forall k \in \{1, \dots, n\}, \Phi_k = \sum_{1 \leq i, j \leq m} (C_k)_{i,j} \varphi_i \wedge \bar{\varphi}_j \quad . \quad (12)$$

Then the scalar product of two geminals can be expressed as the trace of the product of two matrices:

$$\langle \Phi_i | \Phi_j \rangle = \text{tr}(C_i^\dagger C_j) \quad . \quad (13)$$

Note that, the matrix associated with a singlet geminal, will be symmetric (even if we work with complex numbers), and that of a triplet geminal, will be skew-symmetric.

Let Φ'_k 's be geminals whose associated matrices are denoted by C'_k 's. As proved in Appendix VII C the overlap between the wave functions $\Psi_e = \Phi_1 \wedge \dots \wedge \Phi_n$ and $\Psi'_e = \Phi'_1 \wedge \dots \wedge \Phi'_n$ is given by the formula:

$$\langle \Psi_e | \Psi'_e \rangle = \langle \Phi_1 \wedge \dots \wedge \Phi_n | \Phi'_1 \wedge \dots \wedge \Phi'_n \rangle = \sum_{\substack{0 \leq N_{n,0}, \dots, N_{n,n} \leq n \\ \sum_{k=0}^n N_{n,k} = \sum_{k=0}^n k N_{n,k} = n}} (-1)^{N_{n,0}} \sum_{\sigma, \sigma' \in \mathfrak{S}_n} \prod_{k=1}^n \frac{T_{N_{n,0}, \dots, N_{n,k-1}}^{N_{n,k}}(j, \sigma, \sigma')}{k^{N_{n,k}} N_{n,k}!} \quad (14)$$

with:

$$T_{N_{n,0}, \dots, N_{n,k-1}}^{N_{n,k}}(j, \sigma, \sigma') = \text{tr} \left[C^\dagger_{\sigma \left(\sum_{p=0}^{k-1} p N_{n,p} + (j-1)k + 1 \right)} C'_{\sigma' \left(\sum_{p=0}^{k-1} p N_{n,p} + (j-1)k + 1 \right)} \dots C^\dagger_{\sigma \left(\sum_{p=0}^{k-1} p N_{n,p} + jk \right)} C'_{\sigma' \left(\sum_{p=0}^{k-1} p N_{n,p} + jk \right)} \right] \quad (15)$$

The external sum is on the "partitions of the integer n ", that is, on decompositions of n into a sum of (possibly repeated) natural integers. Each such partition is entirely determined by a set of integers $(N_{n,k})_{k=1, \dots, n}$, which correspond to the numbers of times integer k is found in the decomposition. That is to say, we can write:

$$n = \sum_{k=1}^n k N_{n,k} = \sum_{k=0}^n k N_{n,k} \quad . \quad (16)$$

The number $N_{n,0}$, which is introduced in the second equality, extends the definition to $k = 0$ and depends linearly on the others $N_{n,k}$'s:

$$N_{n,0} = n - \sum_{k=1}^n N_{n,k} \quad , \quad (17)$$

It plays a particularly important rôle in the formula, as it determines the sign factor of

each term.

The internal sum on permutations gives *a priori* $(n!)^2$ terms. However, there are redundancies due to the invariance of a trace by circular permutation of its matrices, and by permutations of the traces of the same form, themselves. So, for a given partition of n the number of *a priori* distinct products of traces, to be calculated, reduces to

$$\frac{(n!)^2}{\prod_{k=1}^n k^{N_{n,k}} N_{n,k}!}.$$

These products of traces are decomposed into double products, as follows:

The most external product runs over an index k which determines the number of matrices (k matrices C_i and k matrices C'_j) to be multiplied within each of the $N_{n,k}$ trace factors (provided $N_{n,k} \neq 0$) of the most internal product on the j -index.

Example ($n = 3$) : We compute $\langle \Phi_1 \wedge \Phi_2 \wedge \Phi_3 | \Phi'_1 \wedge \Phi'_2 \wedge \Phi'_3 \rangle = A_1 + A_2 + A_3$, as follow:

$$\begin{aligned} \blacktriangleright 3=0+0+3: N_{3,0} = 2, N_{3,1} = 0, N_{3,2} = 0, N_{3,3} = 1 &\rightsquigarrow A_1 = \sum_{\sigma, \sigma' \in \mathfrak{S}_3} \frac{\text{tr} \left[C_{\sigma(1)}^\dagger C'_{\sigma'(1)} C_{\sigma(2)}^\dagger C'_{\sigma'(2)} C_{\sigma(3)}^\dagger C'_{\sigma'(3)} \right]}{3}. \\ \blacktriangleright 3=0+1+2: N_{3,0} = 1, N_{3,1} = 1, N_{3,2} = 1, N_{3,3} = 0 &\rightsquigarrow A_2 = - \sum_{\sigma, \sigma' \in \mathfrak{S}_3} \frac{\text{tr} \left[C_{\sigma(1)}^\dagger C'_{\sigma'(1)} \right] \text{tr} \left[C_{\sigma(2)}^\dagger C'_{\sigma'(2)} C_{\sigma(3)}^\dagger C'_{\sigma'(3)} \right]}{2}. \\ \blacktriangleright 3=1+1+1: N_{3,0} = 0, N_{3,1} = 3, N_{3,2} = 0, N_{3,3} = 0 &\rightsquigarrow A_3 = \sum_{\sigma, \sigma' \in \mathfrak{S}_3} \frac{\text{tr} \left[C_{\sigma(1)}^\dagger C'_{\sigma'(1)} \right] \text{tr} \left[C_{\sigma(2)}^\dagger C'_{\sigma'(2)} \right] \text{tr} \left[C_{\sigma(3)}^\dagger C'_{\sigma'(3)} \right]}{6}. \end{aligned}$$

Formula Eqs.(14),(15) can easily be checked in the particular case of an AGP against Eq.(8) of Ref.⁴⁷. In this case, all geminal matrices are equal to one and the same matrix, say C_0 : $\forall k, C_k = C'_k = C_0$. So, the sum over permutations give just a factor $(n!)^2$. Moreover, by choosing the ϕ_i to be the natural orbital of the geminal function, we can make this matrix diagonal, without loss of generality: $\forall i, j, (C_0)_{i,j} = \delta_{i,j} \eta_i$. So, all traces reduce to traces of powers of C_0 , and $\text{tr} \left[C_0^p \right] = \sum_{i=1}^m \eta_i^p$. In the above example, we

obtain

$$\langle \Phi_1 \wedge \Phi_2 \wedge \Phi_3 | \Phi_1 \wedge \Phi_2 \wedge \Phi_3 \rangle = 36 \times \left(\frac{\sum_{i=1}^m \eta_i^6}{3} - \frac{\sum_{i=1}^m \eta_i^2 \times \sum_{j=1}^m \eta_j^4}{2} + \frac{\sum_{i=1}^m \eta_i^2 \times \sum_{j=1}^m \eta_j^2 \times \sum_{k=1}^m \eta_k^2}{6} \right). \quad (18)$$

The coefficient of terms of the form η_i^6 is found to be $\frac{1}{3} - \frac{1}{2} + \frac{1}{6} = 0$, and that of terms $\forall i \neq j, \eta_i^2 \eta_j^4$ is $-\frac{1}{2} + 3 \times \frac{1}{6} = 0$, so we are left with terms of the form $\eta_i^2 \eta_j^2 \eta_k^2$ with i, j, k all distinct, arising from the last term of Eq.(18) only, with a combinatorial factor $3!$. Hence the final result:

$$\langle \Phi_1 \wedge \Phi_2 \wedge \Phi_3 | \Phi_1 \wedge \Phi_2 \wedge \Phi_3 \rangle = 36 \times \left(\sum_{1 \leq i < j < k \leq m} \eta_i^2 \eta_j^2 \eta_k^2 \right). \quad (19)$$

which is the expected result, since in Ref.⁴⁷, both bra and ket AGP's are normalized by a factor $n!$, thus eliminating the factor $36 = (3!)^2$. A similar path can be followed to retrieve the formula for an arbitrary n .

At the other extreme, we can check Eqs.(14),(15) for the APSG case. The 1-orthogonality constraint imposed to the geminals amounts to setting

$$\forall i \neq j \quad C_i^\dagger C_j' = 0. \quad (20)$$

So, only one partition of n in the external sum gives a nonzero contribution, namely $n = \underbrace{1 + \dots + 1}_{n \text{ times}}$. Moreover, this constraint also imposes $\sigma = \sigma'$ i.e. transforms the double sum on permutations into a single sum. The latter just gives a factor $N_{n,1}! = n!$ that cancels out with the redundancy factor in the denominator. Therefore, we obtain the expected formula:

$$\langle \Phi_1 \wedge \dots \wedge \Phi_n | \Phi_1' \wedge \dots \wedge \Phi_n' \rangle = \text{tr} \left[C_1^\dagger C_1' \right] \text{tr} \left[C_2^\dagger C_2' \right] \dots \text{tr} \left[C_n^\dagger C_n' \right]. \quad (21)$$

IV. COMPUTATIONAL COST REDUCTION BY ORTHOGONALITY CONSTRAINTS

The number of partitions of n giving the number of terms in the external sum of Eq.(18), grows asymptotically as $\frac{1}{4n\sqrt{3}}\exp\left(\pi\sqrt{\frac{2n}{3}}\right)$ when n goes to infinity. So, even if one disregards the internal sum over non-redundant permutations, the cost of calculating overlaps in the general case is rapidly prohibitive. Hence, the necessity of introducing constraints such as those reviewed in Section II. Here, we investigate permutationally invariant 2-orthogonality constraints already mentioned in conclusion of Ref.⁵.

A. permutationally invariant 2-orthogonality constraints

We have seen in the previous section how choosing all geminals parallel (actually equal) to one another or all 1-orthogonal simplifies the overlap formula. Our aim is to propose less drastic inter-geminal orthogonality constraint, intermediate between the AGP (parallel) and the APSG (1-orthogonality) constraints.

Let us consider wave functions which are products of singlet or triplet geminals (i.e. with symmetric or antisymmetric associated matrices). We want to impose to the geminals Φ_k 's what we call "permutationally invariant 2-orthogonality constraints":

$$\forall i, j, k \in \{1, \dots, n\} \text{ distinct, } \begin{cases} \langle \Phi_i | \Phi_j \rangle = 0 \\ \Phi_k \leftarrow (\Phi_i \wedge \Phi_j) = 0 \end{cases} \quad (22)$$

(the interior product by Φ_k , noted " $\Phi_k \leftarrow$ ", plays in first quantization the same rôle as annihilation in second quantization) or in terms of matrices:

$$\forall i, j, k \in \{1, \dots, n\} \text{ distinct, } \begin{cases} \text{tr}(C_i^\dagger C_j) = 0 \\ C_i C_k^\dagger C_j + C_j C_k^\dagger C_i = 0 \end{cases} \quad (23)$$

Clearly Eq.(20) implies Eq.(23) so 1-orthogonality is at least as strong as permutationally invariant 2-orthogonality. However, the question arises whether there actu-

ally exists geminals which are permutationally invariant 2-orthogonal without being 1-orthogonal i.e. whether 1-orthogonality is strictly stronger than permutationally invariant 2-orthogonality.

For $m = 1$, there can only be one nonzero C_i satisfying the 2-orthogonal condition, so 1-orthogonality is also satisfied. To answer the question positively, let us consider the case $m = 2$, so that the C_i 's are 2×2 -matrices. In this limit case, $\forall i, j, k$ distinct, $\Phi_i \wedge \Phi_j \neq 0 \implies \Phi_i \wedge \Phi_j \propto \varphi_1 \wedge \varphi_2 \wedge \bar{\varphi}_1 \wedge \bar{\varphi}_2 \implies \Phi_k = 0$. So, to have at least three distinct matrices, we necessarily have $\forall i \neq j, \Phi_i \wedge \Phi_j = 0$. A general study⁴⁸ shows that one can have at most three singlet solutions of the form:

$$\begin{aligned} C_1 &= \begin{pmatrix} e^{-i\tau} & 0 \\ 0 & e^{i\tau} \end{pmatrix} z_1 \quad ; \quad C_2 = \begin{pmatrix} \rho e^{-i\tau} & 1 \\ 1 & -\rho e^{i\tau} \end{pmatrix} z_2 \quad ; \\ C_3 &= \begin{pmatrix} -e^{-i\tau} & \rho \\ \rho & e^{i\tau} \end{pmatrix} z_3 \quad , \quad \text{with} \quad \begin{cases} \rho \in \mathbb{R}_+ \\ \tau \in [0, \pi[\end{cases} \quad , \end{aligned} \quad (24)$$

and $z_1, z_2, z_3 \in \mathbb{C}^*$ unimportant factors. To this set one can add at most one triplet geminal matrix. A convenient choice corresponding to $\rho = \tau = 0, z_1 = z_2 = 1, z_3 = -1$ gives the set:

$$I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad ; \quad \sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad ; \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad i\sigma_y = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad , \quad (25)$$

where σ_x, σ_y and σ_z are the well-known Pauli matrices. We will call the set $\{I, \sigma_x, i\sigma_y, \sigma_z\}$ the "4-type set". One can trivially verify that the associated geminals are not 1-orthogonal.

Another non-maximal but useful set of solutions to the permutationally invariant 2-orthogonality conditions in dimension 2 is, $\{G_\theta, \sigma_x, i\sigma_y\}$, where:

$$G_\theta = \begin{pmatrix} \sqrt{2} \sin \theta & 0 \\ 0 & \sqrt{2} \cos \theta \end{pmatrix}, \quad \text{with } \theta \in [0, \pi[\quad (26)$$

whose associated geminal is:

$$\Phi_{G_\theta} = \sqrt{2} (\sin \theta \varphi_1 \wedge \bar{\varphi}_1 + \cos \theta \varphi_2 \wedge \bar{\varphi}_2) \quad \text{with } \|\Phi_{G_\theta}\| = \sqrt{2} \quad . \quad (27)$$

Note that by playing with θ , one can retrieve special matrices of interest:

$$\left\{ \begin{array}{l} G_0 = \sqrt{2} E_{22} \\ G_{\frac{\pi}{4}} = I_2 \\ G_{\frac{\pi}{2}} = \sqrt{2} E_{11} \\ G_{\frac{3\pi}{4}} = \sigma_z \end{array} \right. , \quad (28)$$

where $E_{ij} = (\delta_{ki} \delta_{lj})_{kl}$. It is important to also consider such "3-type sets" in order to break the degeneracy of the one-electron reduced density matrix (1RDM) eigenvalues. Note that all matrix types are either diagonal or anti-diagonal, so we partition them into two sets $\mathcal{D} = \{I_2, \sigma_z, G_\theta\}$ and $\bar{\mathcal{D}} = \{\sigma_x, i\sigma_y\}$.

One could study in the same fashion general solutions for increasing m -values. However, we prefer in the present article to limit ourselves to m -dimensional matrices built as block-diagonal matrices, each block being either a 1-dimensional (1D) block (1-orthogonal part corresponding to a geminal $\varphi_i \wedge \bar{\varphi}_i$) or a 2-dimensional (2D) block constituted of a 2×2 -matrix (whose elements correspond to geminal basis functions $\varphi_i \wedge \bar{\varphi}_i, \varphi_i \wedge \bar{\varphi}_{i+1}, \varphi_{i+1} \wedge \bar{\varphi}_i, \varphi_{i+1} \wedge \bar{\varphi}_{i+1}$) belonging to the 4-type set or to a 3-type set for some value of θ to be optimized.

Note that the permutationally invariant 2-orthogonality conditions further impose that

$$\forall i, j, k \in \{1, \dots, m\} \text{ distinct, } \begin{cases} \sum_{l \text{ such that } B_i^l = B_j^l} \overline{\lambda}_i^l \lambda_j^l + \sum_{l \text{ such that } B_{\theta_l, i}^l = B_{\theta_l, j}^l} \overline{\lambda}_{\theta_l, i}^l \lambda_{\theta_l, j}^l = 0 \\ \lambda_i^l \overline{\lambda}_k^l \lambda_j^l = 0 \text{ as soon as } \exists \{i', j'\} \subset \{i, j, k\}, B_{i'}^l = B_{j'}^l \end{cases} . \quad (30)$$

For practical reason, we choose to rather enforce the following sufficient set of conditions, called "extended permutationally invariant 2-orthogonality" (EPI2O) conditions:

$$\forall i, j \in \{1, \dots, m\} \text{ distincts, } \forall l \in \{1, \dots, m'\}, \begin{cases} B_i^l = B_j^l \implies \lambda_i^l \lambda_j^l = 0 \\ B_{\theta_l, i}^l = B_{\theta_l, j}^l \implies \lambda_{\theta_l, i}^l \lambda_{\theta_l, j}^l = 0 \end{cases} , \quad (31)$$

In other words, two distinct geminals are not allowed to have the same matrix type at a given block position, corresponding to superscript, say l . So, in a 4-type 2D-block, there can be at most 4 nonzero λ_i^l coefficient, each in factor of one of the 4 different block types, and similarly, in a 3-type 2D-block, there can be at most 3 nonzero $\lambda_{\theta_l, i}^l$ coefficient, each in factor of one of the 3 possible block types.

Remark 1: A pair of successive 1D-blocks associated with say $\varphi_i \wedge \overline{\varphi}_i, \varphi_{i+1} \wedge \overline{\varphi}_{i+1}$ and occupied in a given geminal matrix, say C_k , can always be replaced by a 3-type 2D-block with a nonzero coefficient λ for this block, only in factor a G_θ in C_k . So instead of optimizing two coefficients λ_i and λ_{i+1} in factor of the 1D-blocks, one optimizes λ and θ , and identifies $\lambda_i = \lambda \times \sqrt{2} \sin \theta, \lambda_{i+1} = \lambda \times \sqrt{2} \cos \theta$.

B. Expression of the overlap for the EPI2O 2D-block geminal ansatz

Let Φ_1, \dots, Φ_n and Φ'_1, \dots, Φ'_n be geminals verifying the EPI2O conditions with for all l , each C'_k in the ket having the same matrix form as their C_k counterpart in the bra (same block dimensions, types and zero coefficients at the same places). One can anticipate a simplification of the overlap formula Eq.(14) since in every block there will be at most traces of length 4 (product of 4 pairs of block-submatrix) for the cases of

spin-unrestricted 4-type- $2D$ -blocks, of length 3 for the cases of spin-restricted 4-type- $2D$ -blocks or spin-unrestricted 3-type- $2D$ -blocks, of length 2 for the cases of spin-restricted 3-type- $2D$ -blocks, or of length 1 for the cases of $1D$ -blocks.

We shall prove that Eq.(14) becomes:

$$\langle \Phi_1 \wedge \cdots \wedge \Phi_n | \Phi'_1 \wedge \cdots \wedge \Phi'_n \rangle = \sum_{\substack{0 \leq j_1, \dots, j_n \leq m' \\ \text{distinct if nonzero}}} \zeta_{\Phi_1, \Phi'_1}^{j_1} \cdots \zeta_{\Phi_n, \Phi'_n}^{j_n} \quad (32)$$

$$\text{with } \zeta_{\Phi_u, \Phi'_u}^j = \begin{cases} \sum_{t=h_{u-1}+1}^{h_u} \overline{\lambda}_u^{-t} \lambda'_u{}^{-t} & \text{if } j = 0 \\ 2 \overline{\lambda}_{\theta_j, u}^j \lambda'_{\theta_j, u}{}^j & \text{if } j \in \{1, \dots, m''\} \\ 2 \overline{\lambda}_u^j \lambda'_u{}^j & \text{if } j \in \{m'' + 1, \dots, m'\} \end{cases} \quad (33)$$

Direct demonstration:

To prove the formula, we do not need to distinguish between the 3-type- $2D$ -blocks, and the 4-type- $2D$ -blocks, so we will omit the θ label of the coefficients and block matrices in the 3-type cases. We decompose each geminal Φ_i into its 1-orthogonal part and its EPI2O part:

$$\Phi_i = \Phi_i^0 + \sum_{x=1}^{m'} \Phi_i^x \quad (34)$$

where

$$\Phi_i^0 := \sum_{t=h_{i-1}+1}^{h_i} \lambda_i^{-t} \varphi_t \wedge \overline{\varphi}_t \quad , \quad (35)$$

and

$$\Phi_i^x = \sum_{p, q \in \{1, 2\}} \lambda_i^x (B_i^x)_{p, q} \varphi_{\mathfrak{J}(x, p)} \wedge \overline{\varphi}_{\mathfrak{J}(x, q)} \quad . \quad (36)$$

\mathfrak{J} denotes the function that maps component 1 or 2 of block $x \in \{1, \dots, m'\}$ to the corresponding orbital index:

$$\begin{cases} \mathfrak{J}(x, 1) = h_n + 2x - 1 \\ \mathfrak{J}(x, 2) = h_n + 2x \end{cases} \quad (37)$$

Then,

$$\bigwedge_{1 \leq i \leq n} \Phi_i = \sum_{k=1}^n \sum_{1 \leq i_1 < \dots < i_{n-k} \leq n} \sum_{1 \leq x_1, \dots, x_k \leq m'} \Phi_{i_1}^0 \wedge \dots \wedge \Phi_{i_{n-k}}^0 \wedge \Phi_{i_1}^{x_1} \wedge \dots \wedge \Phi_{i_k}^{x_k} \quad . \quad (38)$$

where $\{i_1, \dots, i_{n-k}\} \cup \{\bar{i}_1, \dots, \bar{i}_k\} = \{1, \dots, n\}$, and $\bar{i}_1 < \dots < \bar{i}_k$. If the same $2D$ -block is chosen, say $x_a = x_b$, for two different geminals a and b , then the EPI2O conditions imply $B_{i_a}^{x_a} \neq B_{i_b}^{x_b}$, so that $\Phi_{i_a}^{x_a} \wedge \Phi_{i_b}^{x_b} = 0$. Therefore, the internal sum can be restricted to x_1, \dots, x_k all distinct.

The 1-orthogonality between distinct blocks, whether $1D$ or $2D$ -blocks, implies that each term in the sum must be paired with its exact counterpart in $\bigwedge_{1 \leq i \leq n} \Phi'_i$ to give a nonzero contribution to the overlap. For the $1D$ -part, since there is an exact correspondence between geminal indices and nonzero $1D$ -blocks, 1-orthogonality is enough to associate a single partner for each block. In contrast, for a given $2D$ -block, say x , there could be up to 4 ket-geminal indices compatible with say $\Phi_{i_l}^x$. However, only the index corresponding to the same index \bar{i}_l will give a nonzero contribution, since the EPI2O conditions imply that different indices correspond to different matrix types and by 2-orthogonality of the submatrices: $\bar{i}_l \neq \bar{i}_{l'} \implies \langle \Phi_{i_l}^x | \Phi_{i_{l'}}^x \rangle = 0$. So we are left with:

$$\langle \bigwedge_{1 \leq i \leq n} \Phi_i | \bigwedge_{1 \leq i \leq n} \Phi'_i \rangle = \sum_{k=1}^n \sum_{1 \leq i_1 < \dots < i_{n-k} \leq n} \sum_{\substack{1 \leq x_1, \dots, x_k \leq m' \\ \text{distincts}}} \prod_{j=1}^{n-k} \langle \Phi_{i_j}^0 | \Phi_{i_j}^0 \rangle \prod_{l=1}^k \langle \Phi_{i_l}^{x_l} | \Phi_{i_l}^{x_l} \rangle \quad .$$

Now, if we do not distinguish the different terms according to their number k of $2D$ -blocks, we obtain formulas (32) and (33) where the indices j refers to a $1D$ -part if $j = 0$ or to $2D$ -block of index j , if $j > 0$.

In practice, it is often interesting to have $1D$ -parts, to build a Hartree-Fock component in the geminal product wave function, for example, or to have less terms in the overlap formula. However, let us take advantage of Remark 1 to simplify the discussion and assume that we have only $2D$ -blocks. Note that, if the number of orbital basis functions

is odd, one can always add an extra one at infinity. Then in Eq.(32), there will be only nonzero j -indices. Since the latter are necessarily all distinct, Eq.(32) can be recast in the form:

$$\begin{aligned}
\langle \Phi_1 \wedge \cdots \wedge \Phi_n | \Phi'_1 \wedge \cdots \wedge \Phi'_n \rangle &= \sum_{1 \leq j_1 < \cdots < j_n \leq m'} \sum_{\sigma \in \mathfrak{S}_n} \zeta_{\Phi_1, \Phi'_1}^{\sigma(j_1)} \cdots \zeta_{\Phi_n, \Phi'_n}^{\sigma(j_n)} \\
&= \sum_{1 \leq j_1 < \cdots < j_n \leq m'} \sum_{\sigma \in \mathfrak{S}_n} \zeta_{\Phi_{\sigma(1)}, \Phi'_{\sigma(1)}}^{j_1} \cdots \zeta_{\Phi_{\sigma(n)}, \Phi'_{\sigma(n)}}^{j_n} \\
&= \sum_{\sigma \in \mathfrak{S}_n} \sum_{1 \leq j_1 < \cdots < j_n \leq m'} \zeta_{\Phi_{\sigma(1)}, \Phi'_{\sigma(1)}}^{j_1} \cdots \zeta_{\Phi_{\sigma(n)}, \Phi'_{\sigma(n)}}^{j_n}, \quad (39)
\end{aligned}$$

where σ runs over all permutations of n -indices, \mathfrak{S}_n . For a block y , let $\mathcal{U}(y) = \{u \in \{1, \dots, n\}, \lambda_u^x \neq 0\}$ be the set of indices of the geminals which contain a submatrix with a nonzero coefficient for this block. In fact, in Eq.(39) there are always a large number of terms that are zero, since the permutation must fulfill $\forall i \in \{1, \dots, n\}, \sigma(i) \in \mathcal{U}(j_i)$ otherwise $\zeta_{\Phi_{\sigma(i)}, \Phi'_{\sigma(i)}}^{j_i} = 0$ and we recall that $Card(\mathcal{U}(j_i)) \leq 4$.

Let us set $\mathcal{I}_n^{m'}(\sigma) = \sum_{1 \leq x_1 < \cdots < x_n \leq m'} \zeta_{\Phi_{\sigma(1)}, \Phi'_{\sigma(1)}}^{x_1} \cdots \zeta_{\Phi_{\sigma(n)}, \Phi'_{\sigma(n)}}^{x_n}$ and more generally for p indices $u_1 < \cdots < u_p$, and q block indices $y_1 < \cdots < y_q$, let us denote $\mathcal{I}_{n \setminus \{u_1, \dots, u_p\}}^{m' \setminus \{y_1, \dots, y_q\}}(\sigma)$ the quantity:

$$\sum_{\substack{1 \leq x_1 < \cdots < x_{n-p} \leq m' \\ x_1, \dots, x_{n-p} \notin \{y_1, \dots, y_q\}}} \zeta_{\Phi_{\sigma(1)}, \Phi'_{\sigma(1)}}^{x_1} \cdots \zeta_{\Phi_{\sigma(u_1-1)}, \Phi'_{\sigma(u_1-1)}}^{x_{u_1-1}} \zeta_{\Phi_{\sigma(u_1+1)}, \Phi'_{\sigma(u_1+1)}}^{x_{u_1}} \cdots \zeta_{\Phi_{\sigma(u_p-1)}, \Phi'_{\sigma(u_p-1)}}^{x_{u_p-p}} \zeta_{\Phi_{\sigma(u_p+1)}, \Phi'_{\sigma(u_p+1)}}^{x_{u_p+1-p}} \cdots \zeta_{\Phi_{\sigma(n)}, \Phi'_{\sigma(n)}}^{x_{n-p}}$$

where q geminals and p blocks are excluded with respect to $\mathcal{I}_n^{m'}(\sigma)$. This allows us to write the following recursion formula:

$$\mathcal{I}_n^{m'}(\sigma) = \mathcal{I}_n^{m' \setminus \{y\}}(\sigma) + \sum_{\substack{k=1 \\ \sigma(k) \in \mathcal{U}(y)}}^n \zeta_{\Phi_{\sigma(k)}, \Phi'_{\sigma(k)}}^y \times \mathcal{I}_{n \setminus \{k\}}^{m' \setminus \{y\}}(\sigma), \quad (40)$$

which can be useful to compute the $\mathcal{I}_n^{m'}(\sigma)$'s efficiently.

C. Expression of the 2RDM for the EPI2O 2D-block geminal ansatz

To optimize variationally the expectation value of the electronic Coulomb Hamiltonian, we will make use of its contracted form

$${}^2\hat{h} = \frac{1}{n-1} \left(T_1 + T_2 + \sum_i Z_i \times \left(\frac{1}{\|R_i - r_1\|} + \frac{1}{\|R_i - r_2\|} \right) \right) + \frac{1}{\|r_1 - r_2\|}, \quad (41)$$

where T_i is the kinetic operator of electron i , r_i the position operator of electron i , Z_i the charge of nucleus i , and R_i the position operator of nucleus i . The expectation value over wave function Ψ_e is then computed according to:

$$E = \text{tr}({}^2\Gamma(\Psi_e) {}^2\hat{h}). \quad (42)$$

So, only the 2-electron reduced density matrix, ${}^2\Gamma(\Psi_e)$ is needed *in fine*.

In order to provide general expressions for the more general, 2-electron, reduced, transition matrix (2RTM) between wave functions Ψ_e and Ψ'_e , that we will denote ${}^2\Gamma(\Psi_e, \Psi'_e)$ (the 2RDM expression will be obtained by suppressing the primes), let us generalize our notation to 1D-blocks (using negative integers to number them):

$$\forall i \in \{1, \dots, n\}, \quad \forall t \in \{h_{i-1} + 1, \dots, h_i\}, \quad \text{we set } \Phi_i^{-t} = \lambda_i^{-t} \varphi_t \wedge \bar{\varphi}_t \text{ and } \mathcal{U}(-t) = \{i\}.$$

For any block j , we note $\bar{\mathcal{U}}(j)$ the complement of $\mathcal{U}(j)$ in $\{1, \dots, n\}$. We also set for p geminal indices, $1 \leq u_1 < \dots < u_p \leq n$, and q 2D-block indices

$$0 < y_1 < \dots < y_q < m',$$

$$\mathcal{S}_{n \setminus \{u_1, \dots, u_p\}}^{m' \setminus \{y_1, \dots, y_q\}} = \sum_{\substack{0 \leq x_1, \dots, x_{n-p} \leq m' \\ \text{all distinct if nonzero} \\ x_1, \dots, x_{n-p} \notin \{y_1, \dots, y_q\}}} \zeta_{\Phi_1, \Phi'_1}^{x_1} \cdots \zeta_{\Phi_{u_1-1}, \Phi'_{u_1-1}}^{x_{u_1-1}} \zeta_{\Phi_{u_1+1}, \Phi'_{u_1+1}}^{x_{u_1}} \cdots \zeta_{\Phi_{u_p-1}, \Phi'_{u_p-1}}^{x_{u_p-p}} \zeta_{\Phi_{u_p+1}, \Phi'_{u_p+1}}^{x_{u_p+1-p}} \cdots \zeta_{\Phi_n, \Phi'_n}^{x_{n-p}} \quad (43)$$

Since Ψ_e and Ψ'_e are product of $(S_z = 0)$ -geminals, the 2RTM can be decomposed into 4 \hat{S}_z -adapted blocks that we denote simply (omitting the (Ψ_e, Ψ'_e) dependency) ${}^2\Gamma^{\gamma, \delta}$ with

$\gamma, \delta \in \{\alpha, \beta\}$ referring to spin operator \hat{S}_z eigenstates.

Let us first consider ${}^2\Gamma^{\alpha,\beta}$, (the ${}^2\Gamma^{\beta,\alpha}$ block being obtained by symmetry). Denoting c_i (respectively \bar{c}_i) the annihilation operator of spin-orbital φ_i (respectively $\bar{\varphi}_i$)

$$\begin{aligned} & \forall i_1, j_1, i_2, j_2 \in \{1, \dots, m\}, \\ {}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} &= \langle c_{\bar{j}_1} c_{i_1} \Psi_e | c_{\bar{j}_2} c_{i_2} \Psi'_e \rangle \Leftrightarrow \\ {}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} &= \langle \varphi_{i_1} \wedge \bar{\varphi}_{j_1} \lrcorner \Phi_1 \wedge \dots \wedge \Phi_n | \varphi_{i_2} \wedge \bar{\varphi}_{j_2} \lrcorner \Phi'_1 \wedge \dots \wedge \Phi'_n \rangle \\ &= \sum_{\substack{-h_n \leq x_1, \dots, x_n \leq m' \\ \text{all distinct} \\ -h_n \leq y_1, \dots, y_n \leq m' \\ \text{all distinct}}} \langle \varphi_{i_1} \wedge \varphi_{\bar{j}_1} \lrcorner \Phi_1^{x_1} \wedge \dots \wedge \Phi_n^{x_n} | \varphi_{i_2} \wedge \varphi_{\bar{j}_2} \lrcorner \Phi_1^{y_1} \wedge \dots \wedge \Phi_n^{y_n} \rangle. \end{aligned}$$

Note that the summation can be restricted to block x_i and y_i such that $i \in \mathcal{U}(x_i) \cap \mathcal{U}(y_i)$ for all $i \in \{1, \dots, n\}$. We define the "block function", \mathfrak{B} , that associates to an orbital its block number:

$$\begin{aligned} \mathfrak{B} : \quad \{1, \dots, m\} &\longrightarrow \mathbb{Z}^* \\ i \longmapsto \mathfrak{B}(i) &= \begin{cases} -i & \text{if } i \leq h_n \\ \lfloor \frac{i-h_n+1}{2} \rfloor & \text{if } i > h_n \end{cases}. \end{aligned}$$

where $\lfloor z \rfloor$ is the floor of z , and the function, \mathfrak{C} , that associates to an orbital of a $2D$ -block its position in the block:

$$\begin{aligned} \mathfrak{C} : \quad \{h_n + 1, \dots, m\} &\longrightarrow \{1, 2\} \\ i \longmapsto \mathfrak{C}(i) &= i - h_n - 2(\mathfrak{B}(i) - 1). \end{aligned}$$

Finally, we introduce $\forall j \in \{1, 2\}$, the notation \bar{j} for the complementary of index j , that is to say, $\bar{j} = 2$ if $j = 1$, and $\bar{j} = 1$ if $j = 2$. Then for orbital i and geminal k , we associate the index $\mathfrak{D}(i, k)$ defined as follow:

$$\begin{cases} \mathfrak{D}(i, k) = \mathfrak{C}(i) & \text{if } B_k^{\mathfrak{B}(i)} \in \mathcal{D}(\mathfrak{B}(i)) \\ \mathfrak{D}(i, k) = \overline{\mathfrak{C}(i)} & \text{if } B_k^{\mathfrak{B}(i)} \in \overline{\mathcal{D}}(\mathfrak{B}(i)) \end{cases}.$$

Case 1: $\mathfrak{B}(i_1) \neq \mathfrak{B}(j_1)$

Two distinct blocks are broken in the bra, then, to have at least one ket with nonzero overlap, it is necessary that $\mathfrak{B}(i_2) = \mathfrak{B}(i_1)$ and $\mathfrak{B}(j_2) = \mathfrak{B}(j_1)$. Moreover, these two blocks have to appear within both x_1, \dots, x_n and y_1, \dots, y_n . The remaining $n - 2$ blocks in each set must be two by two identical because of 1-orthogonality between the blocks, and each pair must be associated with the same geminal because of the EPI2O conditions. This means also that the 2 broken blocks must be associated with the same pair of geminals in the bra and in the ket, of indices say (k_1, k_2) .

Subcase 1a: $\mathfrak{B}(i_1) < 0$ and $\mathfrak{B}(j_1) < 0$

Then, necessarily $i_2 = i_1$, $j_2 = j_1$ and $\mathcal{U}(\mathfrak{B}(i_1)) \neq \mathcal{U}(\mathfrak{B}(j_1))$ otherwise ${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} = 0$. k_1 is the unique element of $\mathcal{U}(\mathfrak{B}(i_1))$ and k_2 is the unique element of $\mathcal{U}(\mathfrak{B}(j_1))$

$${}^2\Gamma_{i_1 j_1, i_1 j_1}^{\alpha, \beta} = \bar{\lambda}_{k_1}^{-i_1} \lambda'_{k_1}{}^{-i_1} \bar{\lambda}_{k_2}^{-j_1} \lambda'_{k_2}{}^{-j_1} \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m'}. \quad (44)$$

Subcase 1b: $\mathfrak{B}(i_1) < 0$ and $\mathfrak{B}(j_1) > 0$

Then ${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} \neq 0$, implies $i_2 = i_1$ and k_1 is the unique element of $\mathcal{U}(\mathfrak{B}(i_1))$. The block $\mathfrak{B}(j_1)$ is associated with the same geminal $k_2 \in \mathcal{U}(\mathfrak{B}(j_1))$ in the bra and in the ket. The block matrix $B_{k_2}^{\mathfrak{B}(j_1)}$ being either diagonal or anti-diagonal, after annihilation of the β -spin-orbitals, one can only match the lone ket and bra α -spin-orbitals if $j_2 = j_1$. Then:

$${}^2\Gamma_{i_1 j_1, i_1 j_1}^{\alpha, \beta} = \sum_{k_2 \in \mathcal{U}(\mathfrak{B}(j_1))} \bar{\lambda}_{k_1}^{-i_1} \lambda'_{k_1}{}^{-i_1} \bar{\lambda}_{k_2}^{\mathfrak{B}(j_1)} \lambda'_{k_2}{}^{\mathfrak{B}(j_1)} (B_{k_2}^{\mathfrak{B}(j_1)})_{\mathfrak{D}(j_1, k_2), \mathfrak{C}(j_1)}^2 \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m' \setminus \{\mathfrak{B}(j_1)\}}$$

which can be decomposed as:

$$\begin{aligned} {}^2\Gamma_{i_1 j_1, i_1 j_1}^{\alpha, \beta} &= \sum_{k_2 \in \mathcal{D}(\mathfrak{B}(j_1))} \bar{\lambda}_{k_1}^{-i_1} \lambda'_{k_1}{}^{-i_1} \bar{\lambda}_{k_2}^{\mathfrak{B}(j_1)} \lambda'_{k_2}{}^{\mathfrak{B}(j_1)} (B_{k_2}^{\mathfrak{B}(j_1)})_{\mathfrak{C}(j_1), \mathfrak{C}(j_1)}^2 \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m' \setminus \{\mathfrak{B}(j_1)\}} \\ &+ \sum_{k_2 \in \bar{\mathcal{D}}(\mathfrak{B}(j_1))} \bar{\lambda}_{k_1}^{-i_1} \lambda'_{k_1}{}^{-i_1} \bar{\lambda}_{k_2}^{\mathfrak{B}(j_1)} \lambda'_{k_2}{}^{\mathfrak{B}(j_1)} \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m' \setminus \{\mathfrak{B}(j_1)\}}. \end{aligned} \quad (45)$$

where $\mathcal{D}(j)$ (resp. $\bar{\mathcal{D}}(j)$) is the subset of $\mathcal{U}(j)$ collecting only the indices corresponding to geminals having a diagonal (resp. anti-diagonal) submatrix for block j . $\forall j \in \{1, \dots, m'\}$, $\mathcal{U}(j) = \mathcal{D}(j) \cup \bar{\mathcal{D}}(j)$. Note that the factor $(B_{k_2}^{\mathfrak{B}(j_1)})_{\mathfrak{C}(j_1), \mathfrak{C}(j_1)}^2$ on the first line of Eq.(45) will be 1 in the 4-type case, and either $2\cos^2\theta$ or $2\sin^2\theta$ according to $\mathfrak{C}(j_1)$ being 1 or 2 in the 3-type case. On the second line, we used the fact that $\forall k_2 \in \bar{\mathcal{D}}(\mathfrak{B}(j_1)) \quad (B_{k_2}^{\mathfrak{B}(j_1)})_{1,2}^2 = (B_{k_2}^{\mathfrak{B}(j_1)})_{2,1}^2 = 1$.

Subcase 1c: $\mathfrak{B}(i_1) > 0$ and $\mathfrak{B}(j_1) < 0$

by symmetry with the previous subcase:

$$\begin{aligned} {}^2\Gamma_{i_1 j_1, i_1 j_1}^{\alpha, \beta} &= \sum_{k_1 \in \mathcal{D}(\mathfrak{B}(i_1))} \bar{\lambda}_{k_1}^{\mathfrak{B}(i_1)} \lambda'_{k_1}{}^{\mathfrak{B}(i_1)} (B_{k_1}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(i_1)}^2 \bar{\lambda}_{k_2}^{-j_1} \lambda'_{k_2}{}^{-j_1} \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m' \setminus \{\mathfrak{B}(i_1)\}} \\ &+ \sum_{k_1 \in \bar{\mathcal{D}}(\mathfrak{B}(i_1))} \bar{\lambda}_{k_1}^{\mathfrak{B}(i_1)} \lambda'_{k_1}{}^{\mathfrak{B}(i_1)} \bar{\lambda}_{k_2}^{-j_1} \lambda'_{k_2}{}^{-j_1} \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m' \setminus \{\mathfrak{B}(i_1)\}}. \end{aligned} \quad (46)$$

Subcase 1d: $\mathfrak{B}(i_1) > 0$ and $\mathfrak{B}(j_1) > 0$

This last case is more complicated than the previous ones, since there are 3 sub-subcases to consider:

- (i) $(k_1, k_2) \in \mathcal{U}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(j_1)) \times \mathcal{U}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(j_1))$,
- (ii) $(k_1, k_2) \in \mathcal{U}(\mathfrak{B}(i_1)) \times \bar{\mathcal{U}}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(j_1))$,
- (iii) $(k_1, k_2) \in \mathcal{U}(\mathfrak{B}(j_1)) \times \mathcal{U}(\mathfrak{B}(i_1)) \cap \bar{\mathcal{U}}(\mathfrak{B}(j_1))$.

Using the notation already defined, the following general formula can be obtained for case (i):

$$\begin{aligned}
{}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} &= \sum_{\substack{k_1, k_2 \in \mathcal{U}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(j_1)) \\ k_1 \neq k_2}} \mathcal{S}_{n \setminus \{k_1, k_2\}}^{m' \setminus \{\mathfrak{B}(i_1), \mathfrak{B}(j_1)\}} \left(\delta_{i_1, i_2} \delta_{j_1, j_2} \left(\bar{\lambda}_{k_1}^{\mathfrak{B}(i_1)} \lambda'_{k_1}{}^{\mathfrak{B}(i_1)} (B_{k_1}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{D}(i_1, k_1)}^2 \bar{\lambda}_{k_2}^{\mathfrak{B}(j_1)} \lambda'_{k_2}{}^{\mathfrak{B}(j_1)} \right. \right. \\
&\times \left. \left. (B_{k_2}^{\mathfrak{B}(j_1)})_{\mathfrak{D}(j_1, k_2), \mathfrak{C}(j_1)}^2 + \bar{\lambda}_{k_2}^{\mathfrak{B}(i_1)} \lambda'_{k_2}{}^{\mathfrak{B}(i_1)} (B_{k_2}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{D}(i_1, k_2)}^2 \bar{\lambda}_{k_1}^{\mathfrak{B}(j_1)} \lambda'_{k_1}{}^{\mathfrak{B}(j_1)} (B_{k_1}^{\mathfrak{B}(j_1)})_{\mathfrak{D}(j_1, k_1), \mathfrak{C}(j_1)}^2 \right) \right. \\
&+ \delta_{\mathfrak{D}(i_1, k_1), \mathfrak{D}(i_2, k_2)} \delta_{\mathfrak{D}(j_1, k_2), \mathfrak{D}(j_2, k_1)} \left(\bar{\lambda}_{k_1}^{\mathfrak{B}(i_1)} \lambda'_{k_2}{}^{\mathfrak{B}(i_1)} (B_{k_1}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{D}(i_1, k_1)} (B_{k_2}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_2), \mathfrak{D}(i_2, k_2)} \right. \\
&\times \left. \bar{\lambda}_{k_2}^{\mathfrak{B}(j_1)} \lambda'_{k_1}{}^{\mathfrak{B}(j_1)} (B_{k_2}^{\mathfrak{B}(j_1)})_{\mathfrak{D}(j_1, k_2), \mathfrak{C}(j_1)} B_{k_1}^{\mathfrak{B}(j_1)} \right)_{\mathfrak{D}(j_2, k_1), \mathfrak{C}(j_2)} \\
&+ \delta_{\mathfrak{D}(i_1, k_2), \mathfrak{D}(i_2, k_1)} \delta_{\mathfrak{D}(j_1, k_1), \mathfrak{D}(j_2, k_2)} \left(\bar{\lambda}_{k_2}^{\mathfrak{B}(i_1)} \lambda'_{k_1}{}^{\mathfrak{B}(i_1)} (B_{k_2}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{D}(i_1, k_2)} (B_{k_1}^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_2), \mathfrak{D}(i_2, k_1)} \right. \\
&\times \left. \bar{\lambda}_{k_1}^{\mathfrak{B}(j_1)} \lambda'_{k_2}{}^{\mathfrak{B}(j_1)} (B_{k_1}^{\mathfrak{B}(j_1)})_{\mathfrak{D}(j_1, k_1), \mathfrak{C}(j_1)} B_{k_2}^{\mathfrak{B}(j_1)} \right)_{\mathfrak{D}(j_2, k_2), \mathfrak{C}(j_2)} \Big). \tag{47}
\end{aligned}$$

The factor $\delta_{i_1, i_2} \delta_{j_1, j_2}$ comes from the argument used before, when the same $2D$ -block matrices are matched in the bra and in the ket. The other Kröneckers symbols can be worked out explicitly, according to i_1 and i_2 (resp. j_1 and j_2) being equal or not, and by detailing whether the $2D$ -block matrices are of \mathcal{D} or $\bar{\mathcal{D}}$ -type, as was done in Eqs.(45) and (46). This is not done for Eq.(47) to alleviate the presentation.

The formula for case (ii) can be deduced from Eq.(47) by eliminating the terms where either $\bar{\lambda}_{k_1}^{\mathfrak{B}(j_1)}$ or $\lambda'_{k_1}{}^{\mathfrak{B}(j_1)}$ appears, which leaves only the first term. Similarly, for case (iii), one must eliminate the terms where either $\bar{\lambda}_{k_1}^{\mathfrak{B}(i_1)}$ or $\lambda'_{k_1}{}^{\mathfrak{B}(i_1)}$ appears, which leaves only the second term on the second line. Care must be taken to count only once, the pairs $\{k_1, k_2\}$ such that one element is in $\bar{\mathcal{U}}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(j_1))$ and the other in $\mathcal{U}(\mathfrak{B}(i_1)) \cap \bar{\mathcal{U}}(\mathfrak{B}(j_1))$, which appear both in case (ii) and in case (iii).

Case 2: $\mathfrak{B}(i_1) = \mathfrak{B}(j_1)$

No broken block in the bra, then, there cannot be broken block in the ket too, so necessarily $\mathfrak{B}(i_2) = \mathfrak{B}(j_2)$.

Subcase 2a: $\mathfrak{B}(i_1) \neq \mathfrak{B}(i_2)$

$\mathfrak{B}(i_1)$ has to appear within x_1, \dots, x_n but not $\mathfrak{B}(i_2)$ since it will disappear from the ket

after annihilation of $\varphi_{i_2} \wedge \bar{\varphi}_{j_2}$ in it. Symmetrically, $\mathfrak{B}(i_2)$ has to appear within y_1, \dots, y_n but not $\mathfrak{B}(i_1)$. The remaining $n - 1$ blocks in each set must be two by two identical because of 1-orthogonality between the blocks, and each pair must be associated with the same geminal because of the EPI2O conditions. This means also that $\mathfrak{B}(i_1)$ in the bra and $\mathfrak{B}(i_2)$ in the ket must be associated with the same geminal of index, say k . This is only possible if $\mathcal{U}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(i_2)) \neq \emptyset$. When this is the case, we obtain for $2D$ -blocks:

$$\begin{aligned}
{}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} &= \delta_{i_1, j_1} \delta_{i_2, j_2} \sum_{k \in \mathcal{D}(\mathfrak{B}(i_1)) \cap \mathcal{D}(\mathfrak{B}(i_2))} \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(i_1)} (B_k^{\mathfrak{B}(i_2)})_{\mathfrak{C}(i_2), \mathfrak{C}(i_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1), \mathfrak{B}(i_2)\}} \\
&+ (1 - \delta_{i_1, j_1}) \delta_{i_2, j_2} \sum_{k \in \bar{\mathcal{D}}(\mathfrak{B}(i_1)) \cap \mathcal{D}(\mathfrak{B}(i_2))} \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(j_1)} (B_k^{\mathfrak{B}(i_2)})_{\mathfrak{C}(i_2), \mathfrak{C}(i_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1), \mathfrak{B}(i_2)\}} \\
&+ \delta_{i_1, j_1} (1 - \delta_{i_2, j_2}) \sum_{k \in \mathcal{D}(\mathfrak{B}(i_1)) \cap \bar{\mathcal{D}}(\mathfrak{B}(i_2))} \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(i_1)} (B_k^{\mathfrak{B}(i_2)})_{\mathfrak{C}(i_2), \mathfrak{C}(j_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1), \mathfrak{B}(i_2)\}} \\
&+ (1 - \delta_{i_1, j_1}) (1 - \delta_{i_2, j_2}) \sum_{k \in \bar{\mathcal{D}}(\mathfrak{B}(i_1)) \cap \bar{\mathcal{D}}(\mathfrak{B}(i_2))} \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(j_1)} (B_k^{\mathfrak{B}(i_2)})_{\mathfrak{C}(i_2), \mathfrak{C}(j_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1), \mathfrak{B}(i_2)\}}.
\end{aligned} \tag{48}$$

or more compactly (but at the price of summing terms that are zero),

$${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} = \sum_{k \in \mathcal{U}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(i_2))} \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(j_1)} (B_k^{\mathfrak{B}(i_2)})_{\mathfrak{C}(i_2), \mathfrak{C}(j_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1), \mathfrak{B}(i_2)\}}. \tag{49}$$

It is possible for some blocks to be $1D$. Then, $\mathcal{U}(\mathfrak{B}(i_1)) \cap \mathcal{U}(\mathfrak{B}(i_2))$ can contain at most one geminal index, say k . If $\mathfrak{B}(i_1) < 0$ and $\mathfrak{B}(i_2) > 0$ then necessarily $i_1 = j_1$ and :

$${}^2\Gamma_{i_1 i_1, i_2 j_2}^{\alpha, \beta} = \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_2)})_{\mathfrak{C}(i_2), \mathfrak{C}(j_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_2)\}}, \tag{50}$$

else if $\mathfrak{B}(i_2) < 0$ and $\mathfrak{B}(i_1) > 0$ then necessarily $i_2 = j_2$ and:

$${}^2\Gamma_{i_1 j_1, i_2 i_2}^{\alpha, \beta} = \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{C}(i_1), \mathfrak{C}(j_1)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1)\}}. \tag{51}$$

For both $\mathfrak{B}(i_1) < 0$ and $\mathfrak{B}(i_2) < 0$ to occur, it is necessary that $h_k - h_{k-1} > 1$ as $\mathfrak{B}(i_1), \mathfrak{B}(i_2) \in \{h_{k-1} + 1, \dots, h_k\}$, then

$${}^2\Gamma_{i_1 i_1, i_2 i_2}^{\alpha, \beta} = \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_2)} \mathcal{S}_{n \setminus \{k\}}^{m'}. \tag{52}$$

Subcase 2b: $\mathfrak{B}(i_1) = \mathfrak{B}(i_2)$

$\mathfrak{B}(i_1)$ has to appear within x_1, \dots, x_n and within y_1, \dots, y_n . The remaining $n - 1$ blocks in each set must be two by two identical because of 1-orthogonality between the blocks, and each pair must be associated with the same geminal because of the EPI2O conditions. This means also that $\mathfrak{B}(i_1)$ in the bra and in the ket must be associated with the same geminal of index, say $k \in \mathcal{U}(\mathfrak{B}(i_1))$. When this is the case, we obtain for $2D$ -blocks:

$${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \beta} = \sum_{k \in \mathcal{U}(\mathfrak{B}(i_1))} \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_1)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{c}(i_1), \mathfrak{c}(j_1)} (B_k^{\mathfrak{B}(i_1)})_{\mathfrak{c}(i_2), \mathfrak{c}(j_2)} \mathcal{S}_{n \setminus \{k\}}^{m' \setminus \{\mathfrak{B}(i_1)\}}. \quad (53)$$

If $\mathfrak{B}(i_1) < 0$ then necessarily $i_1 = j_1 = i_2 = j_2$ and let k be the unique element of $\mathcal{U}(\mathfrak{B}(i_1))$:

$${}^2\Gamma_{i_1 i_1, i_1 i_1}^{\alpha, \beta} = \bar{\lambda}_k^{\mathfrak{B}(i_1)} \lambda_k^{\mathfrak{B}(i_1)} \mathcal{S}_{n \setminus \{k\}}^{m'}. \quad (54)$$

Let us now turn to ${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \alpha}$, the case of ${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\beta, \beta}$ being analogous. Here, the Pauli principle makes it sufficient to consider indices $i_1, j_1, i_2, j_2 \in \{1, \dots, m\}$ such that $i_1 < j_1$ and $i_2 < j_2$:

$${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\alpha, \alpha} = \sum_{\substack{-h_n \leq x_1, \dots, x_n \leq m' \\ \text{all distinct} \\ -h_n \leq y_1, \dots, y_n \leq m' \\ \text{all distinct}}} \langle \varphi_{i_1} \wedge \varphi_{j_1} \leftarrow \Phi_1^{x_1} \wedge \dots \wedge \Phi_n^{x_n} | \varphi_{i_2} \wedge \varphi_{j_2} \leftarrow \Phi_1^{y_1} \wedge \dots \wedge \Phi_n^{y_n} \rangle.$$

After annihilation of the α -spin-orbitals, the lone β -spin-orbitals in the bra and in the ket must be paired, this imposes $\mathfrak{B}(i_1) = \mathfrak{B}(i_2)$ and $\mathfrak{B}(j_1) = \mathfrak{B}(j_2)$.

Case 1: $\mathfrak{B}(i_1) \neq \mathfrak{B}(j_1)$

Two distinct blocks are broken and must be associated with the same pair of geminals in the bra and in the ket, of indices say (k_1, k_2) . The two blocks being 1-orthogonal, the algebra is the same whether the spins of the orbitals in each block are the same or not. The formulas for ${}^2\Gamma^{\alpha, \alpha}$ are the same as those for ${}^2\Gamma^{\alpha, \beta}$. However, only the cases where $i_1 < j_1$ and $i_2 < j_2$ are relevant.

Case 2: $\mathfrak{B}(i_1) = \mathfrak{B}(j_1)$

It is not possible to annihilate two spin-orbitals of the same spin in a given block, since all the blocks only contains ($S_z = 0$)-geminals. So this case does not give any nonzero contribution.

The expressions obtained for ${}^2\Gamma$ are extremely simple to compute, provided that we have an efficient way of calculating the $\mathcal{S}_{n \setminus \{u_1, \dots, u_p\}}^{m' \setminus \{y_1, \dots, y_q\}}$'s with $p, q \leq 2$.

D. Expression of gradients for the EPI2O 2D-block geminal ansatz

In the EPI2O 2D-block geminal ansatz there are three types of continuous parameters:

- (i) The coefficients defining the orbitals ϕ_i 's in a given basis set. We will not attempt to optimize them in the present study.
- (ii) The block coefficients λ_k^x 's.
- (iii) In the case of 3-type-2D-blocks, the angles θ_x 's.

In order to optimize the last two kinds, we provide formulas to compute the gradients with respect to their variations. We will assume that the λ_k^x 's are real, so that $\bar{\lambda}_k^x = \lambda_k^x$.

1. Gradient with respect to block coefficients

We first note that all the formulas of the previous section for the 2RTM, hence for the 2RDM, are of the form:

$${}^2\Gamma_{i_1 j_1, i_2 j_2}^{\gamma_1, \gamma_2} = \sum_w \mathbf{B}_w \Lambda_w \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}, \quad (55)$$

where \mathbf{B}_w is a product of block matrix elements (reduced to 1 in the case of 1D-block), which can depends upon θ_x 's but not upon λ_k^x 's; Λ_w is a product of block matrix coefficients, λ_k^x 's, which do not depends upon θ_x 's; K_w is a set of at most two 2D-block indices and X_w is a set of at most two geminal indices. Like Λ_w , the quantity $\mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}$ only depends upon the λ_k^x 's.

Taking the derivatives with respect to λ_k^x in Eq.(55) gives:

$$\frac{\partial^2 \Gamma_{i_1 j_1, i_2 j_2}^{\gamma_1, \gamma_2}}{\partial \lambda_k^x} = \sum_w \mathbf{B}_w \left(\frac{\partial \Lambda_w}{\partial \lambda_k^x} \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w} + \Lambda_w \frac{\partial \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}}{\partial \lambda_k^x} \right). \quad (56)$$

However, for a given w , at most one of the two terms can be nonzero: $\frac{\partial \Lambda_w}{\partial \lambda_k^x}$ will be nonzero only if $k \in K_w$, whereas $\frac{\partial \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}}{\partial \lambda_k^x}$ will be nonzero only if $k \notin K_w$.

Let us consider $\frac{\partial \Lambda_w}{\partial \lambda_k^x}$:

In "case 1", it will be nonzero only if $k \in K_w$ and $x \in \mathfrak{B}(i_1) \cup \mathfrak{B}(j_1)$, then:

$$\frac{\partial \Lambda_w}{\partial \lambda_k^x} = \begin{cases} \frac{\Lambda_w}{\lambda_k^x} & \text{in the last two terms of case 1d(i)} \\ \frac{2\Lambda_w}{\lambda_k^x} & \text{in all other cases} \end{cases}. \quad (57)$$

In "case 2", it will be nonzero only if $k \in K_w$ and $x \in \mathfrak{B}(i_1) \cup \mathfrak{B}(i_2)$, then:

$$\frac{\partial \Lambda_w}{\partial \lambda_k^x} = \begin{cases} \frac{\Lambda_w}{\lambda_k^x} & \text{if } \mathfrak{B}(i_1) \neq \mathfrak{B}(i_2) \\ \frac{2\Lambda_w}{\lambda_k^x} & \text{if } \mathfrak{B}(i_1) = \mathfrak{B}(i_2) \end{cases}. \quad (58)$$

Regarding $\frac{\partial \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}}{\partial \lambda_k^x}$, it will be nonzero only if $k \notin K_w$, then:

$$\frac{\partial \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}}{\partial \lambda_k^x} = \begin{cases} 2\lambda_k^x \mathcal{S}_{n \setminus K_w \cup \{k\}}^{m' \setminus X_w} & \text{if } x < 0 \\ 4\lambda_k^x \mathcal{S}_{n \setminus K_w \cup \{k\}}^{m' \setminus X_w \cup \{x\}} & \text{if } x > 0 \end{cases}. \quad (59)$$

Note that these gradients make sense only for coefficients λ_k^x such that $k \in \mathcal{U}(x)$.

2. Gradient with respect to angle parameters

In the case of a 3-type-2D-block of index x , $1 \leq x \leq m''$, taking the derivatives with respect to θ_x in Eq.(55) gives:

$$\frac{\partial^2 \Gamma_{i_1 j_1, i_2 j_2}^{\gamma_1, \gamma_2}}{\partial \theta_x} = \sum_w \frac{\partial \mathbf{B}_w}{\partial \theta_x} \Lambda_w \mathcal{S}_{n \setminus K_w}^{m' \setminus X_w}. \quad (60)$$

The quantity $\frac{\partial \mathbf{B}_w}{\partial \theta_x}$ will be nonzero only if $k \in K_w$, and:

For case 1, if $B_k^x \in \mathcal{D}(\mathfrak{B}(i_1)) \cup \mathcal{D}(\mathfrak{B}(j_1))$, then

$$\frac{\partial \mathbf{B}_w}{\partial \theta_x} = \begin{cases} \text{in the last two terms of case 1d(i)} & \begin{cases} -\tan\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{1,1} \text{ appears in } \mathbf{B}_w \\ \cot\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{2,2} \text{ appears in } \mathbf{B}_w \end{cases} \\ \text{in all other cases} & \begin{cases} -2 \tan\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{1,1}^2 \text{ appears in } \mathbf{B}_w \\ 2 \cot\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{2,2}^2 \text{ appears in } \mathbf{B}_w \end{cases} \end{cases} \quad (61)$$

For case 2, if $B_k^x \in \mathcal{D}(\mathfrak{B}(i_1)) \cup \mathcal{D}(\mathfrak{B}(i_2))$, then,

$$\frac{\partial \mathbf{B}_w}{\partial \theta_x} = \begin{cases} \text{if } \mathfrak{B}(i_1) \neq \mathfrak{B}(i_2) & \begin{cases} -\tan\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{1,1} \text{ appears in } \mathbf{B}_w \\ \cot\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{2,2} \text{ appears in } \mathbf{B}_w \end{cases} \\ \text{if } \mathfrak{B}(i_1) = \mathfrak{B}(i_2) & \begin{cases} -2 \tan\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{1,1}^2 \text{ appears in } \mathbf{B}_w \\ 2 \cot\theta_x \mathbf{B}_w & \text{if } (B_k^x)_{2,2}^2 \text{ appears in } \mathbf{B}_w \\ (\cot\theta_x - \tan\theta_x) \mathbf{B}_w & \text{if } (B_k^x)_{1,1}(B_k^x)_{2,2} \text{ appears in } \mathbf{B}_w \end{cases} \end{cases} \quad (62)$$

V. PROOF OF CONCEPT: APPLICATION TO LINEAR HYDROGEN CHAINS

We have implemented our new geminal ansatz in the Nice-branch of the Tonto⁴⁹ quantum chemistry code.

We have found the following EPI2O geminal product wave function for the linear H₆ molecule in the 6-31G basis set for *H* nuclei equally spaced by one angström:

$$\Phi = \Phi_1 \wedge \Phi_2 \wedge \Phi_3 \quad , \quad (63)$$

with C_i matrices corresponding to the Φ_i geminals of the form:

$$C_i = \begin{pmatrix} \lambda_i^1 B_i^1 & & & & & \\ & \lambda_i^2 B_i^2 & & & & \\ & & \lambda_i^3 B_i^3 & & & \\ & & & \lambda_i^4 B_i^4 & & \\ & & & & \lambda_i^5 B_i^5 & \\ & & & & & \lambda_i^6 B_i^6 \end{pmatrix} \quad (64)$$

where the coefficients and submatrices are given below:

$$\left\{ \begin{array}{l} \lambda_1^1 B_1^1 = 0, 699585 G_{\theta_1} \\ \lambda_1^2 B_1^2 = -0, 021264 G_{\theta_2} \\ \lambda_1^3 B_1^3 = 0, 100629 \sigma_x \\ \lambda_1^4 B_1^4 = 0 \\ \lambda_1^5 B_1^5 = -0, 001567 \sigma_x \\ \lambda_1^6 B_1^6 = 0 \end{array} \right\}, \left\{ \begin{array}{l} \lambda_2^1 B_2^1 = 0, 113021 \sigma_x \\ \lambda_2^2 B_2^2 = 0 \\ \lambda_2^3 B_2^3 = 0, 697602 G_{\theta_3} \\ \lambda_2^4 B_2^4 = -0, 024046 G_{\theta_4} \\ \lambda_2^5 B_2^5 = 0 \\ \lambda_2^6 B_2^6 = -0, 000019 \sigma_x \end{array} \right\}, \left\{ \begin{array}{l} \lambda_3^1 B_3^1 = 0 \\ \lambda_3^2 B_3^2 = -0, 00076 \sigma_x \\ \lambda_3^3 B_3^3 = 0 \\ \lambda_3^4 B_3^4 = 0, 000016 \sigma_x \\ \lambda_3^5 B_3^5 = 0, 706656 G_{\theta_5} \\ \lambda_3^6 B_3^6 = -0, 025247 G_{\theta_6} \end{array} \right\} \quad (65)$$

the angles of the diagonal submatrices being:

$$\left\{ \begin{array}{l} \theta_1 = 1, 662992 \text{ rad} \\ \theta_2 = 1, 391623 \text{ rad} \\ \theta_3 = 1, 701670 \text{ rad} \\ \theta_4 = 1, 267600 \text{ rad} \\ \theta_5 = 1, 702433 \text{ rad} \\ \theta_6 = 1, 268724 \text{ rad} \end{array} \right\} \quad (66)$$

The energy of this wave function has been calculated to be $-3, 295805$ Ha, that is about 1 millihartree below that of the APSG ansatz in the same basis, which is $-3, 294840$ Ha

(see Table 1 of Ref.⁵). Si, this establishes a proof of concept that the EPI2O model can be more accurate than the APSG one. We note that there are significant coefficients in front of seniority 2 submatrices, such as λ_1^3 and λ_2^1 , which are responsible for the departure from strong orthogonality and at the same time, from a seniority 0 APG wave function.

For such a small system, the computer time is too small to be significant. We will devote a forthcoming article to an in-depth study of the EPI2O geminal ansatz, computational performances.

VI. CONCLUSION

After presenting an overview of existing APG-based methods, we have derived a general formula for the overlap of APG wave functions, which exhibits the full combinatorics complexity of such model and serves us as a starting point to elaborate new constraints able to make it more amenable to practical computations. This led us to propose the so-called “permutationally invariant 2-orthogonality constraints” and a further extension of the latter within a 2D-block geminal model, which makes their implementation simpler and more efficient. Then we have obtained explicit formulas for the quantities necessary to implement the EPI2O geminal ansatz and finally, we have provided a proof of principle that the latter is able to give strictly lower energies than strongly orthogonal geminal products while removing the seniority 0 restriction.

Further work is in progress to evaluate the performances of this new geminal wave function ansatz both on classical and hybrid quantum-classical computers.

ACKNOWLEDGEMENTS

This work was supported by the grant CARMA ANR-12-BS01-0017. This work has been supported by the French government, through the UCAJEDI Investments in the Future project managed by the National Research Agency (ANR) with the reference number ANR-15-IDEX-01. Prof. V. Rassolov is acknowledged for fruitful discussions and his help in calculating 1-orthogonal starting guesses.

VII. APPENDIX

A. Generalized seniority

The concept relies on a given decomposition of the one-electron Hilbert space, \mathcal{H} , into a direct sum of Hilbert subspaces called “shells”, $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \dots \oplus \mathcal{H}_n$, (where the dimension of \mathcal{H}_i $\dim \mathcal{H}_i := d_i$) or more conveniently, in practice, on a partition of the spin-orbital basis set into subsets, $(\chi_{i,j})_{j \in \{1, \dots, d_i\}}$, spanning the shells of the decomposition, $i \in \{1, \dots, n\}$ being the shell index⁵⁰.

The generalized seniority number of a Slater determinant built over these spin-orbitals is simply the number of incomplete shells present in the determinant i.e. shells that are neither empty nor fully-occupied. More precisely, let $\chi_{i_1, j_1} \wedge \dots \wedge \chi_{i_1, j_{k_1}} \wedge \dots \wedge \chi_{i_p, j_p} \wedge \dots \wedge \chi_{i_p, j_{k_p}}$ be such a Slater determinant (where \wedge the intrinsiquely antisymmetrical, “Grassmann” or “exterior” product⁵¹), one counts 1 for every k_m such that $0 < k_m < d_m$ (assuming all spinorbitals are distinct otherwise the determinant is zero). This definition extends over linear combinations of Slater determinants having same generalized seniority number. Then, note that the definition is invariant under arbitrary rotations (or more generally, unitary transformations) of the spinorbitals within the same shell. Hence, it is legitimate to talk of seniority with respect to the shell decomposition and not just with respect to the spinorbital partition.

When there are m orbitals, one can retrieve the usual notion of seniority^{52–55} by decomposing the Hilbert space into m two-dimensional Hilbert subspaces, $\mathcal{H} = \mathcal{H}_1 \oplus \cdots \oplus \mathcal{H}_m$, where each \mathcal{H}_i is spanned by two spin-orbitals of spin- α and spin- β , respectively. However, other partitions not only based on spin-degeneracy, can prove physically relevant, such as partitions into atomic shell orbitals (hence the name we have coined), or into spatially degenerate molecular orbitals.

B. p -orthogonality

Let $p > 0$ be an integer, Ψ_1 an n_1 -electron wave function, and Ψ_2 an n_2 -electron one, with $n_1 \geq p$ and $n_2 \geq p$. Let us consider, their p -particle reduced density matrices, ${}^p\Gamma_1$ and ${}^p\Gamma_2$ respectively. We call “ p -external space of Ψ_i ” the p -particle Hilbert subspace spanned by the unoccupied p -particle natural wave functions, that is to say, by the eigenfunctions of ${}^p\Gamma_i$ associated with the 0 eigenvalue, for $i \in \{1, 2\}$, and “ p -internal space of Ψ_i ”, noted $\mathcal{I}[\Psi_i]$ the orthogonal subspace. In other words, $\mathcal{I}[\Psi_i]$ is the Hilbert space spanned by the partially occupied p -particle natural wave functions. We will say that Ψ_1 and Ψ_2 are p -orthogonal if and only if their p -internal spaces are orthogonal: $\mathcal{I}[\Psi_1] \perp \mathcal{I}[\Psi_2]$. Note that 1-orthogonality is nothing but the so-called “strong orthogonality”^{56,57}, and when $n_1 = n_2 = n$, n -orthogonality is just the usual concept of orthogonality. For any integers $0 < p \leq q \leq n$, we have the important property that p -orthogonality implies q -orthogonality, which means that the lesser the integer p , the stronger the p -orthogonality.

C. Proof of the general APG overlap formula

We will prove our formula by mathematical induction on $k \geq 1$, assuming that the equality is true for a particular k (the case $k = 1$ being trivial). For the step $(k + 1)$, we want to calculate the following scalar product:

$$\langle \Phi_1 \wedge \cdots \wedge \Phi_{k+1} | \Phi'_1 \wedge \cdots \wedge \Phi'_{k+1} \rangle \quad .$$

In order to use the induction hypothesis, we transform the last exterior product in the bra into a right co-product, by Hopf duality. The left-hand side becomes:

$$\Phi_1 \wedge \cdots \wedge \Phi_k \otimes \Phi_{k+1} \quad ,$$

while the action of the co-product on the right-hand side gives the following expression:

$$\begin{aligned} & \Phi'_1 \wedge \cdots \wedge \Phi'_k \otimes \Phi'_{k+1} + \sum_{u=1}^k \Phi'_1 \wedge \cdots \wedge \Phi'_{u-1} \wedge \Phi'_{k+1} \wedge \Phi'_{u+1} \wedge \cdots \wedge \Phi'_k \otimes \Phi'_u \\ & - \sum_{1 \leq \alpha, \beta, \gamma, \delta \leq m} \sum_{1 \leq u < v \leq k+1} [(C'_u)_{\gamma, \beta} (C'_v)_{\alpha, \delta} + (C'_v)_{\gamma, \beta} (C'_u)_{\alpha, \delta}] \times \\ & \quad \times \Phi'_1 \wedge \cdots \wedge \Phi'_{u-1} \wedge \Phi'_{u+1} \wedge \cdots \wedge \Phi'_{v-1} \wedge \Phi'_{v+1} \wedge \cdots \wedge \Phi'_{k+1} \wedge \varphi_\alpha \wedge \overline{\varphi_\beta} \otimes (\varphi_\gamma \wedge \overline{\varphi_\delta}) \quad . \end{aligned}$$

With this new formulation, we can develop each expression and apply the induction hypothesis. The scalar product can now be written as a sum of two distinct terms: K_1 , composed by the terms in which the geminals of the right-hand side term have not been cut by the co-product, and K_2 , formed by the other terms.

For K_1 , we obtain:

$$\begin{aligned} K_1 &= \langle \Phi_1 \wedge \cdots \wedge \Phi_k | \Phi'_1 \wedge \cdots \wedge \Phi'_k \rangle \langle \Phi_{k+1} | \Phi'_{k+1} \rangle + \sum_{u=1}^k \langle \Phi_1 \wedge \cdots \wedge \Phi_k | \bigwedge_{\substack{1 \leq l \leq k+1 \\ l \neq u}} \Phi'_l \rangle \langle \Phi_{k+1} | \Phi'_u \rangle \\ &= \langle \Phi_1 \wedge \cdots \wedge \Phi_k | \Phi'_1 \wedge \cdots \wedge \Phi'_k \rangle \text{tr}(C_{k+1}^\dagger C'_{k+1}) + \sum_{u=1}^k \langle \Phi_1 \wedge \cdots \wedge \Phi_k | \bigwedge_{\substack{1 \leq l \leq k+1 \\ l \neq u}} \Phi'_l \rangle \text{tr}(C_{k+1}^\dagger C'_u) \quad . \end{aligned}$$

Applying the induction hypothesis on the two terms of K_1 (for the second one, by replacing Φ'_u with Φ'_{k+1} so C'_u with C'_{k+1}), we can write:

$$K_1 = \sum_{\substack{0 \leq N_{k,0}, \dots, N_{k,k} \leq k \\ \sum_{i=0}^k N_{k,i} = \sum_{i=0}^k i N_{k,i} = k}} Q_{N_{k,0}, \dots, N_{k,k}} \sum_{u=1}^k \text{tr}(C_{k+1}^\dagger C'_u) \sum_{\sigma, \sigma' \in \mathfrak{S}_k} \prod_{i=1}^k P_{N_{k,0}, \dots, N_{k,i-1}}^{N_{k,i}}(\sigma, \tau_{u, k+1} \sigma') \quad , \quad (67)$$

with $\tau_{u, k+1}$ the transposition $(u, k+1) \in \mathfrak{S}_{k+1}$.

For K_2 we have:

$$\begin{aligned}
K_2 &= - \sum_{1 \leq \alpha, \beta, \gamma, \delta \leq m} \sum_{1 \leq u < v \leq k+1} [(C'_u)_{\gamma, \beta} (C'_v)_{\alpha, \delta} + (C'_v)_{\gamma, \beta} (C'_u)_{\alpha, \delta}] \langle \Phi_1 \wedge \cdots \wedge \Phi_k | \left(\bigwedge_{\substack{1 \leq l \leq k+1 \\ l \neq u, v}} \Phi'_l \right) \wedge \varphi_\alpha \wedge \overline{\varphi_\beta} \rangle \langle \Phi_{k+1} | \varphi_\gamma \wedge \overline{\varphi_\delta} \rangle \\
&= - \sum_{1 \leq u < v \leq k+1} \sum_{1 \leq \alpha, \beta \leq m} \left[\langle \Phi_1 \wedge \cdots \wedge \Phi_k | \left(\bigwedge_{\substack{1 \leq l \leq k+1 \\ l \neq u, v}} \Phi'_l \right) \wedge \varphi_\alpha \wedge \overline{\varphi_\beta} \rangle \times \right. \\
&\quad \left. \times \sum_{1 \leq \gamma, \delta \leq m} \left((C'_v)_{\alpha, \delta} \langle \Phi_{k+1} | \varphi_\gamma \wedge \overline{\varphi_\delta} \rangle (C'_u)_{\gamma, \beta} + (C'_u)_{\alpha, \delta} \langle \Phi_{k+1} | \varphi_\gamma \wedge \overline{\varphi_\delta} \rangle (C'_v)_{\gamma, \beta} \right) \right] \\
&= - \sum_{1 \leq u < v \leq k+1} \sum_{1 \leq \alpha, \beta \leq m} \langle \Phi_1 \wedge \cdots \wedge \Phi_k | \left(\bigwedge_{\substack{1 \leq l \leq k+1 \\ l \neq u, v}} \Phi'_l \right) \wedge \varphi_\alpha \wedge \overline{\varphi_\beta} \rangle [(C'_v C'_{k+1}^\dagger C'_u)_{\alpha\beta} + (C'_u C'_{k+1}^\dagger C'_v)_{\alpha\beta}] \quad .
\end{aligned}$$

Let $E_{\alpha\beta}$ be the matrix where only the coefficient of line α and column β is nonzero and equal to 1. We can use the induction hypothesis by replacing Φ'_u with $\varphi_\alpha \wedge \overline{\varphi_\beta}$ and, for $v \neq k+1$, Φ'_v with Φ'_{k+1} (i.e. by substituting C'_u with $E_{\alpha\beta}$ and, for $v \neq k+1$, C'_v with C'_{k+1}). To take advantage of the simple form of $E_{\alpha\beta}$, we have to rewrite our recursion formula as:

$$\begin{aligned}
&\sum_{\substack{0 \leq N_{k,0}, \dots, N_{k,k} \leq k \\ \sum_{i=0}^k N_{k,i} = k \\ N_{k,x_u} \geq 1}} Q_{N_{k,0}, \dots, N_{k,k}} \sum_{x_u=1}^k N_{k,x_u} x_u \sum_{\substack{\sigma, \sigma' \in \mathfrak{S}_k \\ \sigma'(L_{x_u})=u}} \text{tr} \left[\left(\prod_{l=L_{x_u-1}+(N_{k,x_u}-1)x_u+1}^{L_{x_u}-1} C'_{\sigma(l)} C'_{\sigma'(l)} \right) C'_{\sigma(L_{x_u})} E_{\alpha\beta} \right] \times \\
&\quad \times P_{N_{k,0}, \dots, N_{k,x_u-1}}^{N_{k,x_u}-1}(\sigma, \sigma') \prod_{\substack{1 \leq i \leq k \\ i \neq x_u}} P_{N_{k,0}, \dots, N_{k,i-1}}^{N_{k,i}}(\sigma, \sigma') \quad ,
\end{aligned}$$

with $L_{x_u} = \sum_{p=0}^{x_u} p N_{k,p}$.

Note that a multiplying factor has appeared. Indeed, since we have distinguished the trace containing $E_{\alpha\beta}$, our formula is no longer completely symmetric under permutations. We had to multiply by N_{k,x_u} , the number of possible positions of that specific trace of length $2x_u$ in its product with the other traces of same length (consequence of the commutativity of multiplication in \mathbb{K}), and then by x_u , which is the number of possible locations for $E_{\alpha\beta}$ in our trace (consequence of the invariance of the trace under circular permutations of matrices).

We also notice that N_{k,x_u} cannot be zero since it is related to the trace containing $E_{\alpha\beta}$, which necessarily exists. This trace is actually reduced to a single matrix element:

$$\text{tr} \left[\left(\prod_{l=L_{x_u-1}+(N_{k,x_u}-1)x_u+1}^{L_{x_u}-1} C_{\sigma(l)}^\dagger C'_{\sigma'(l)} \right) C_{\sigma(L_{x_u})}^\dagger E_{\alpha\beta} \right] = \left[\left(\prod_{l=L_{x_u-1}+(N_{k,x_u}-1)x_u+1}^{L_{x_u}-1} C_{\sigma(l)}^\dagger C'_{\sigma'(l)} \right) C_{\sigma(L_{x_u})}^\dagger \right]_{\beta\alpha} . \quad (68)$$

Then, a *bona fide* trace re-appears in K_2 since sum:

$$\sum_{1 \leq \alpha, \beta \leq m} [(C'_v C_{k+1}^\dagger C'_u)_{\alpha\beta} + (C'_u C_{k+1}^\dagger C'_v)_{\alpha\beta}] \left[\left(\prod_{l=L_{x_u-1}+(N_{k,x_u}-1)x_u+1}^{L_{x_u}-1} C_{\sigma(l)}^\dagger C'_{\sigma'(l)} \right) C_{\sigma(L_{x_u})}^\dagger \right]_{\beta\alpha}$$

is nothing but:

$$\text{tr} \left[C_{\sigma(L_{x_u})}^\dagger (C'_v C_{k+1}^\dagger C'_u + C'_u C_{k+1}^\dagger C'_v) \prod_{l=L_{x_u-1}+(N_{k,x_u}-1)x_u+1}^{L_{x_u}-1} C_{\sigma(l)}^\dagger C'_{\sigma'(l)} \right] .$$

By using the induction hypothesis, K_2 can be written:

$$K_2 = - \sum_{1 \leq u < v \leq k+1} \sum_{x_u=1}^k \sum_{\substack{0 \leq N_{k,0}, \dots, N_{k,k} \leq k \\ \sum_{i=0}^k N_{k,i} = \sum_{i=0}^k i N_{k,i} = k \\ N_{k,x_u} \geq 1}} Q_{N_{k,0}, \dots, N_{k,k}} N_{k,x_u} x_u \sum_{\substack{\sigma, \sigma' \in \mathfrak{S}_k \\ \sigma'(L_{x_u})=u}} P_{N_{k,0}, \dots, N_{k,x_u-1}}^{N_{k,x_u}-1} (\sigma, \tau_{v,k+1} \sigma') \times \\ \times \text{tr} \left[C_{\sigma(L_{x_u})}^\dagger (C'_v C_{k+1}^\dagger C'_u + C'_u C_{k+1}^\dagger C'_v) \prod_{l=L_{x_u-1}+(N_{k,x_u}-1)x_u+1}^{L_{x_u}-1} C_{\sigma(l)}^\dagger C'_{\tau_{v,k+1} \sigma'(l)} \right] \prod_{\substack{1 \leq i \leq k \\ i \neq x_u}} P_{N_{k,0}, \dots, N_{k,i-1}}^{N_{k,i}} (\sigma, \tau_{v,k+1} \sigma') , \quad (69)$$

with $\tau_{v,k+1}$ the transposition $(v, k+1) \in \mathfrak{S}_{k+1}$.

Note that the trace which is factored in the general term K_2 and whose length has increased from $2x_u$ to $2(x_u+1)$ actually represents two traces of C -matrix products since we have kept the symmetrized form $(C'_v C_{k+1}^\dagger C'_u + C'_u C_{k+1}^\dagger C'_v)$ inside in order to reduce the size of the K_2 -expression.

K_1 and K_2 have now a very similar form. However, we still have to replace the sum on partitions of k with one on partitions of $k + 1$. To perform this, we define several mappings.

First, let $\mathcal{F}_{k,0}$ denote the following mapping:

$$\begin{aligned} \mathcal{F}_{k,0} : \quad \mathbb{N}^{k+1} &\longrightarrow \mathbb{N} \times \mathbb{N}^* \times \mathbb{N}^{k-1} \times \{0\} \\ (N_{k,0}, \dots, N_{k,k}) &\longmapsto (N_{k,0}, N_{k,1} + 1, N_{k,2}, \dots, N_{k,k}, 0) \quad . \end{aligned}$$

For $1 \leq i \leq k - 1$, we define $\mathcal{F}_{k,i}$ as:

$$\begin{aligned} \mathcal{F}_{k,i} : \mathbb{N}^i \times \mathbb{N}^* \times \mathbb{N}^{k-i} &\longrightarrow \mathbb{N}^* \times \mathbb{N}^i \times \mathbb{N}^* \times \mathbb{N}^{k-i-1} \times \{0\} \\ (N_{k,0}, \dots, N_{k,k}) &\longmapsto (N_{k,0} + 1, N_{k,1}, \dots, N_{k,i-1}, N_{k,i} - 1, N_{k,i+1} + 1, N_{k,i+2}, \dots, N_{k,k}, 0) \quad . \end{aligned}$$

Finally, we introduce $\mathcal{F}_{k,k}$ as follows:

$$\begin{aligned} \mathcal{F}_{k,k} : \quad \mathbb{N}^k \times \mathbb{N}^* &\longrightarrow \mathbb{N}^* \times \mathbb{N}^k \times \{1\} \\ (N_{k,0}, \dots, N_{k,k}) &\longmapsto (N_{k,0} + 1, N_{k,1}, \dots, N_{k,k-1}, N_{k,k} - 1, 1) \quad . \end{aligned}$$

These mappings transform a partition of k into a partition of $k + 1$. Moreover, they are invertible on their image and their inverses allow to map a partition of $k + 1$ onto a partition of k :

$$\begin{aligned} \mathcal{F}_{k,0}^{-1} : \quad \mathbb{N} \times \mathbb{N}^* \times \mathbb{N}^{k-1} \times \{0\} &\longrightarrow \mathbb{N}^{k+1} \\ (N_{k+1,0}, \dots, N_{k+1,k}, 0) &\longmapsto (N_{k+1,0}, N_{k+1,1} - 1, N_{k+1,2}, \dots, N_{k+1,k}) \quad ; \end{aligned}$$

$$\begin{aligned} \mathcal{F}_{k,i}^{-1} : \mathbb{N}^* \times \mathbb{N}^i \times \mathbb{N}^* \times \mathbb{N}^{k-i-1} \times \{0\} &\longrightarrow \mathbb{N}^i \times \mathbb{N}^* \times \mathbb{N}^{k-i} \quad (1 \leq i \leq k - 1) \\ (N_{k+1,0}, \dots, N_{k+1,k}, 0) &\longmapsto (N_{k+1,0} - 1, N_{k+1,1}, \dots, N_{k+1,i-1}, N_{k+1,i} + 1, N_{k+1,i+1} - 1, N_{k+1,i+2}, \dots, N_{k+1,k}) \quad ; \end{aligned}$$

$$\begin{aligned} \mathcal{F}_{k,k}^{-1} : \quad \mathbb{N}^* \times \mathbb{N}^k \times \{1\} &\longrightarrow \mathbb{N}^k \times \mathbb{N}^* \\ (N_{k+1,0}, \dots, N_{k+1,k}, 1) &\longmapsto (N_{k+1,0} - 1, N_{k+1,1}, \dots, N_{k+1,k-1}, N_{k+1,k} + 1) \quad . \end{aligned}$$

If we consider a partition of k which has q nonzero $N_{k,i}$'s ($i \geq 1$), then we can apply exactly $q + 1$ distinct mappings $\mathcal{F}_{k,i}$ (one for each nonzero $N_{k,i}$, and $\mathcal{F}_{k,0}$ is always applicable), each giving a different partition of $k + 1$.

Conversely, let $(N_{k+1,0}, \dots, N_{k+1,k+1})$ be an arbitrary partition of $k + 1$. Consider $q \geq 1$ and $i_1 < \dots < i_q$ such as the N_{k+1,i_j} 's are the nonzero integers determining our partition of $k + 1$ (we deduce the value $N_{k+1,0}$). This partition can then stem from exactly q different partitions of k , denoted by $(N_{k,0}^{i_j}, \dots, N_{k,k}^{i_j})$:

$$\left\{ \begin{array}{l} (N_{k,0}^{i_1}, \dots, N_{k,k}^{i_1}) = \mathcal{F}_{k,i_1}^{-1}(N_{k+1,0}, \dots, N_{k+1,k+1}) \\ (N_{k,0}^{i_2}, \dots, N_{k,k}^{i_2}) = \mathcal{F}_{k,i_2}^{-1}(N_{k+1,0}, \dots, N_{k+1,k+1}) \\ \quad \cdot \\ \quad \cdot \\ \quad \cdot \\ (N_{k,0}^{i_q}, \dots, N_{k,k}^{i_q}) = \mathcal{F}_{k,i_q}^{-1}(N_{k+1,0}, \dots, N_{k+1,k+1}) \end{array} \right. .$$

So, we have three possible forms for the preimages of $(N_{k+1,0}, \dots, N_{k+1,k+1})$.

(i) If $i_1 = 0$, then:

$$(N_{k,0}^0, \dots, N_{k,k}^0) = (N_{k+1,0}, N_{k+1,1} - 1, N_{k+1,2}, \dots, N_{k+1,k}) \quad . \quad (70)$$

We see from Eq.(67), that such a pre-image will arise from K_1 . (ii) If $i_q = k$, then $q = 1$ and:

$$(N_{k,0}^k, \dots, N_{k,k}^k) = (N_{k+1,0} - 1, N_{k+1,1}, \dots, N_{k+1,k-1}, N_{k+1,k} + 1) \quad . \quad (71)$$

(iii) Finally, for the general case $1 \leq i_j \leq k - 1$:

$$(N_{k,0}^{i_j}, \dots, N_{k,k}^{i_j}) = (N_{k+1,0} - 1, N_{k+1,1}, \dots, N_{k+1,i_j-1}, N_{k+1,i_j} + 1, N_{k+1,i_j+1} - 1, N_{k+1,i_j+2}, \dots, N_{k+1,k}) \quad . \quad (72)$$

We see from Eq.(69), that these two last cases will arise from the terms in K_2 .

So, all the products of traces required for our partition of $k + 1$, $(N_{k+1,0}, \dots, N_{k+1,k+1})$, will appear either in K_1 or in K_2 , C_{k+1}^\dagger being inserted in the trace that has been factored

out from the product of traces, and the transpositions insuring that C'_{k+1} will be inserted in all possible factors of the remaining product of traces. Now, we just have to examine the coefficients coming from the different pre-images. Let us denote by Q_{i_j} the factor coming from the $(N_{k,0}^{i_j}, \dots, N_{k,k}^{i_j})$ pre-image. The products of traces related to the latter are those, where C'_{k+1} appears in a trace of length $2(i_j + 1)$. We have to show that by transforming the summations in K_1 or K_2 into sums over \mathfrak{S}_{k+1} , these factors are all equal to the expected normalization factor: $Q_{N_{k+1,0}, \dots, N_{k+1,k+1}}$.

Let us first consider Q_0 (only present if $i_1 = 0$). The terms for which Q_0 is factored come from K_1 . To find the value of Q_0 from $Q_{N_{k,0}^0, \dots, N_{k,k}^0}$, we have to examine how the summations over σ , σ' and u can be compacted into a double summation over \mathfrak{S}_{k+1} . To each $\sigma' \in \mathfrak{S}_k$, corresponds $k + 1$ distinct permutations $\sigma'_{k+1,u} \in \mathfrak{S}_{k+1}$:

$$\begin{cases} \sigma'_{k+1,u}(\sigma'^{-1}(u)) = k + 1 \\ \sigma'_{k+1,u}(k + 1) = u \\ \sigma'_{k+1,u}(l) = \sigma'(l) \text{ for all } l \in \{1, \dots, k\} \setminus \{\sigma'^{-1}(u)\} \end{cases} \quad . \quad (73)$$

In this way, the summation over u from 1 to $k + 1$ and the one over $\sigma' \in \mathfrak{S}_k$ can be condensed into a sum over \mathfrak{S}_{k+1} without changing the normalization factor already present. To extend the summation over σ from \mathfrak{S}_k to \mathfrak{S}_{k+1} , we have to divide by the number of possibilities to interleave the new trace of length 2 (the one containing C'_{k+1}) within the $(N_{k+1,1} - 1)$ other traces, (that is to say by $(N_{k+1,1} - 1) + 1$). So, we obtain:

$$\begin{aligned} Q_0 &= \underbrace{Q_{N_{k+1,0}, N_{k+1,1}-1, N_{k+1,2}, \dots, N_{k+1,k}}}_{\text{factor } Q_{N_{k,0}^0, \dots, N_{k,k}^0} \text{ appearing in } K_1} \times \underbrace{\frac{1}{N_{k+1,1}}}_{\text{additional factor}} \\ &= \frac{(-1)^{N_{k+1,0}}}{\prod_{i=1}^{k+1} i^{N_{k+1,i}} N_{k+1,i}!} \\ &= Q_{N_{k+1,0}, \dots, N_{k+1,k+1}} \quad . \end{aligned}$$

We proceed likewise with the other Q_{i_j} 's, which will be factored in the terms coming

from K_2 when $x_u = i_j$ and also when $k = i_j$. The i_j 's are in fact the only possible values for x_u since, at a fixed partition of k , the values of x_u giving nonzero terms in the sum are those corresponding to nonzero N_{k,x_u} . This time, it is the double summation over u and v in K_2 which will allow us to transform the sum over $\sigma' \in \mathfrak{S}_k$ into a sum over \mathfrak{S}_{k+1} without additional normalization factor. To each permutation $\sigma' \in \mathfrak{S}_k$, we can associate the following permutations $\sigma'_{k+1,u,v} \in \mathfrak{S}_{k+1}$:

$$\left\{ \begin{array}{l} \sigma'_{k+1,u,v}(\sigma'^{-1}(v)) = k + 1 \text{ if } v \neq k + 1 \\ \sigma'_{k+1,u,v}(L_{i_j}) = u \\ \sigma'_{k+1,u,v}(k + 1) = v \\ \sigma'_{k+1,u,v}(l) = \sigma'(l) \text{ for all } l \in \{1, \dots, k\} \setminus \{\sigma'^{-1}(v), L_{i_j}\} \end{array} \right. . \quad (74)$$

These permutations are pairwise distinct when we consider all the possible values for u and v and all the permutations $\sigma' \in \mathfrak{S}_k$. In K_2 , the double summation on (u, v) contains $\frac{k(k+1)}{2}$ values and the sum over $\sigma' \in \mathfrak{S}_k$ contains $(k-1)!$ different permutations as $\sigma'(L_{i_j}) = u$ is fixed. Therefore, we arrive at $\frac{(k-1)!k(k+1)}{2} = \frac{(k+1)!}{2}$ distinct permutations in \mathfrak{S}_{k+1} . This number has to be multiplied by two, since, as already noticed, the trace of length $2(i_j + 1)$ containing C_{k+1}^\dagger actually counts as two (the previous $\frac{(k+1)!}{2}$ permutations can be composed with transposition $\tau_{u,v}$ to account for the symmetrized C -matrix product). So, we have a total of $(k+1)!$ distinct permutations, which means that we have all the permutations belonging to \mathfrak{S}_{k+1} . Regarding the extension of the summation over σ to \mathfrak{S}_{k+1} , we have to divide by the number of ways to interleave the new trace of length $2(i_j + 1)$ (containing C_{k+1}^\dagger) with the $(N_{k+1,i_j+1} - 1)$ other traces, but also by a factor $(i_j + 1)$, which arises from the invariance of the trace containing C_{k+1}^\dagger of length $2(i_j + 1)$

under circular permutation. We obtain as required:

$$\begin{aligned}
Q_{i_j} &= \underbrace{-Q_{N_{k+1,0}-1, N_{k+1,1}, \dots, N_{k+1, i_j-1}, N_{k+1, i_j}+1, N_{k+1, i_j+1}-1, N_{k+1, i_j+2}, \dots, N_{k+1, k}}}_{\text{factor } -Q_{N_{k,0}^{i_j}, \dots, N_{k,k}^{i_j}} N_{k, i_j}^{i_j} \text{ appearing in } K_2} (N_{k+1, i_j} + 1) i_j \times \underbrace{\frac{1}{N_{k+1, i_j+1} (i_j + 1)}}_{\text{additional factor}} \\
&= \frac{(-1)^{N_{k+1,0}}}{\prod_{i=1}^{k+1} i^{N_{k+1,i}} N_{k+1,i}!} \\
&= Q_{N_{k+1,0}, \dots, N_{k+1, k+1}} \quad .
\end{aligned}$$

To resume our achievements: the sum over partitions of k have been transformed into a sum over partitions of $k + 1$, and all the other sums have been condensed into a double sum over \mathfrak{S}_{k+1} while the normalization factors turned into the expected $Q_{N_{k+1,0}, \dots, N_{k+1, k+1}}$ factors. All the terms in K_1 and K_2 have been consumed along the way. This implies that $\langle \Phi_1 \wedge \dots \wedge \Phi_{k+1} | \Phi'_1 \wedge \dots \wedge \Phi'_{k+1} \rangle = K_1 + K_2$ verifies our induction property for Step $(k + 1)$. This concludes our proof by induction of the general APG overlap formula.

Remark: The number of redundant terms is the same for all the products of traces associated with a given partition of n and is equal to $\prod_{k=1}^n k^{N_{n,k}} N_{n,k}!$.

The number of products of traces we sum in $T_{N_{n,0}, \dots, N_{n,n}}$ is $\frac{(n!)^2}{\prod_{k=1}^n k^{N_{n,k}} N_{n,k}!}$.

So, by counting all distinct products of traces for all the partitions of n , we find:

$$(n!)^2 = \sum_{\substack{0 \leq N_{n,1}, \dots, N_{n,n} \leq n \\ \sum_{k=1}^n i N_{n,k} = n}} \frac{(n!)^2}{\prod_{k=1}^n k^{N_{n,k}} N_{n,k}!} \quad , \quad (75)$$

which gives the remarkable identity:

$$\forall n \in \mathbb{N}^*, \quad 1 = \sum_{\substack{0 \leq N_{n,1}, \dots, N_{n,n} \leq n \\ \sum_{k=1}^n k N_{n,k} = n}} \frac{1}{\prod_{k=1}^n k^{N_{n,k}} N_{n,k}!} \quad . \quad (76)$$

REFERENCES

- ¹Charles-Émile Fecteau, Samuel Cloutier, Jean-David Moisset, Jérémy Boulay, Patrick Bultinck, Alexandre Faribault, Paul A. Johnson, *The Journal of Chemical Physics* **156**, 194103, 2022.
- ²Rishab Dutta, Guo P. Chen, Thomas M. Henderson, and Gustavo E. Scuseria, *The Journal of Chemical Physics*, **154**, 114112, 2021.
- ³Haochuan Wei, Eric Neuscammann, *J. Chem. Phys.* **149**, 184106, 2018.
- ⁴Patrick Cassam-Chenaï, *Physical Review* **A77**, 032103, 2008.
- ⁵Patrick Cassam-Chenaï, Vitaly Rassolov, *Chemical Physics Letters* **487**, 147-152, 2010.
- ⁶R. McWeeny, B. Sutcliffe, *Proc. Roy. Soc. (London)* **A273**, 103-116, 1963.
- ⁷Patrick Cassam-Chenaï, *The Journal of Chemical Physics* **124**, 194109, 2006.
- ⁸Paul A. Johnson, Peter A.Limacher, Taewon D. Kim, Michael Richer, Ramón Alain Miranda-Quintana, Farnaz Heidar Zadeh, Paul W. Ayers, Patrick Bultinck, Stijn De Baerdemacker, Dimitri Van Neck, *Computational and Theoretical Chemistry* **1116**, 207-219, 2017.
- ⁹D. M. Silver, *J. Chem. Phys.* **50**, p5108, 1969.
- ¹⁰P. A Limacher, *J. Chem. Phys.* **145**, 194102, 2016.
- ¹¹Paul A. Johnson, Paul W. Ayers, Peter A.Limacher, Stijn De Baerdemacker, Dimitri Van Neck, Patrick Bultinck, *Computational and Theoretical Chemistry* **1003**, 101-113, 2013.
- ¹²Pawel Tecmer, Katharina Boguslawski, Paul A. Johnson, Peter A.Limacher, Matthew CHAN, Toon Verstraelen, Paul W. Ayers, *The Journal of Physical Chemistry A* **118**, 9058-9068, 2014.
- ¹³John Bardeen, Leon N. Cooper, and J. Robert Schrieffer, *Physical Review* **108**, 1175, 1957.
- ¹⁴Paul A. Johnson, Charles-Émile Fecteau, Frédéric Berthiaume, Samuel Cloutier, Laurie Carrier, Marianne Gratton, Patrick Bultinck, Stijn De Baerdemacker, Dimitri Van Neck, PeterLimacher, Paul W. Ayers, *The Journal of Chemical Physics* **153**, 104110,

2020.

- ¹⁵P. A. Limacher, P. W. Ayers, P. A. Johnson, S. De Baerdemacker, D. Van Neck and P. Bultinck, *J. Chem. Theory and Computation* **9**, 1394-1401, 2013.
- ¹⁶Peter A. Limacher, Taewon D. Kim, Paul W. Ayers, Paul A. Johnson, Stijn De Baerdemacker, Dimitri Van Neck, Patrick Bultinck, *Mol. Phys.* **112**, 853-862, 2014.
- ¹⁷Katharina Boguslawski, Pawel Tecmer, Paul W. Ayers, Patrick Bultinck, Stijn De Baerdemacker, Dimitri Van Neck, *Physical Review B* **89**, 201106, 2014.
- ¹⁸Katharina Boguslawski, Pawel Tecmer, Patrick Bultinck, Stijn De Baerdemacker, Dimitri Van Neck, Paul W. Ayers, *Journal of Chemical Theory and Computation* **10**, 4873-4882, 2014.
- ¹⁹Katharina Boguslawski, Pawel Tecmer, Peter A. Limacher, Paul A. Johnson, Paul W. Ayers, Patrick Bultinck, Stijn De Baerdemacker, Dimitri Van Neck, *The Journal of Chemical Physics* **140**, 214114, 2014.
- ²⁰A.C. Hurley, J.E. Lennard-Jones and J.A. Pople, *Proc. Roy. Soc. (London)* **A220**, 446-455, 1953.
- ²¹E. L. Mehler, K. Ruedenberg, D. M. Silver, *J. Chem. Phys.* **52**, 1181, 1970.
- ²²D. M. Silver, *J. Chem. Phys.* **52**, 299, 1970.
- ²³D. M. Silver, E. L. Mehler, K. Ruedenberg, *J. Chem. Phys.* **52**, 1174, 1970.
- ²⁴David M. Silver, Klaus Ruedenberg, Ernest L. MEHLER, *J. Chem. Phys.* **52**, 1206, 1970.
- ²⁵V. A. Rassolov, *J. Chem. Phys.* **117**, 5978, 2002.
- ²⁶Vitaly A. Rassolov, Feng Xu, Sophya Garashchuk, *J. Chem. Phys.* **120**, 10385, 2004.
- ²⁷Vitaly A. Rassolov, Feng Xu, *J. Chem. Phys.* **126**, 234112, 2007.
- ²⁸V. A. Rassolov, F. Xu *J. Chem. Phys.* **127**, 44104, 2007.
- ²⁹A. John Coleman, *The AGP model for fermion systems, International Journal of Quantum Chemistry* **63**, 23-30, 1997.
- ³⁰E. Rosta and P.R. Surján, *Int. J. Quantum Chem.* **80**, 96-104, 2000.
- ³¹E. Rosta and P.R. Surján, *J. Chem. Phys.* **116**, 878, 2002.

- ³²P.R. Surján, Ágnes Szabados, Péter Jeszenszki, Tamás Zoboki, *J. Math. Chem.* **50**, 534, 2012.
- ³³Péter Jeszenszki, Péter R. Nagy, Tamás Zoboki, Ágnes Szabados, Péter R. Surján, *Int. J. Quantum Chem.* **114**, 1048-1052, 2014.
- ³⁴Peter A. Limacher, Paul W. Ayers, Paul A. Johnson, Stijn De Baerdemacker, Dimitri Van Neck, Patrick Bultinck, *Physical Chemistry Chemical Physics* **16**, 5061-5065, 2014.
- ³⁵Dominic Földvári, Zsuzsanna Tóth, Péter R. Surján, Ágnes Szabados, *The Journal of Chemical Physics* **150**, 034103, 2019.
- ³⁶Katharina Boguslawski, Paul W. Ayers, *Journal of Chemical Theory and Computation* **11**, 5252-5261, 2015.
- ³⁷Tamás Zoboki, Ágnes Szabados, P.R. Surján, *Journal of Chemical Theory and Computation* **9**, 2602-2608, 2013.
- ³⁸Eric Neuscamman, *J. Chem. Phys.* **139**, 181101, 2013.
- ³⁹Eric Neuscamman, *J. Chem. Theory and Computation* **12**, 3149, 2016.
- ⁴⁰Eric Neuscamman, *Physical Review Letters* **109**, 203001, 2012.
- ⁴¹Eric Neuscamman, *J. Chem. Phys.* **139**, 194105, 2013.
- ⁴²L. Zhao, E. Neuscamman, *J. Chem. Theory and Computation* **12**, 3719, 2016.
- ⁴³Thomas M. Henderson, Gustavo E. Scuseria, *The Journal of Chemical Physics* **151**, 051101, 2019.
- ⁴⁴T. M. Henderson and G. E. Scuseria, *J. Chem. Phys.* **153**, 084111, 2020.
- ⁴⁵A. Khamoshi, F. A. Evangelista, G. E Scuseria, Correlating AGP on a quantum computer, *Quantum Science and Technology* **6**, 014004, 2020.
- ⁴⁶V. E. Elfving, M. Millaruelo, J. A. Gámez, C. Gogolin, *Phys. Rev.* **A103**, 032605, 2021.
- ⁴⁷A. Khamoshi, Thomas M. Henderson, G. E Scuseria, *The Journal of Chemical Physics* **151**, 184103, 2019.
- ⁴⁸Thomas Perez, *Problèmes d'algèbre extérieure liés au calcul de fonctions d'ondes électroniques produits de géminales*, Université Côte d'Azur, 2020.

<http://www.theses.fr/2020COAZ4060>

- ⁴⁹D. Jayatilaka, D. J. Grimwood, A. Lee, A. Lemay, A. J. Russel, C. Taylor, S. K. Wolff, P. Cassam-Chenaï, A. Whitton, TONTO - A system for computational chemistry (2005).
- ⁵⁰Thomas Perez, Patrick Cassam-Chenaï, *Journal of Mathematical Chemistry* **56**, 1428-1436, 2018.
- ⁵¹Patrick Cassam-Chenaï, Frédéric Patras, *Journal of Mathematical Physics* **44**, 4884-4906, 2003.
- ⁵²G. Racah, *Phys. Rev.* **63**, 367, 1943.
- ⁵³D. J. Dean, M. Hjorth-Jensen, *Rev. Mod. Phys.* **75**, 607, 2003.
- ⁵⁴V. Zelevinsky and A. Volya, *Physics of atomic nuclei* **66**, 1781, 2003.
- ⁵⁵Laimutis Bytautas, Thomas M. Henderson, Carlos A. Jiménez-Hoyos, Jason K. Ellis, Gustavo E. Scuseria, *The Journal of Chemical Physics*, **135**, 044119, 2011.
- ⁵⁶R. G. Parr, F. O. Ellison, P. G. Lykos *J. Chem. Phys.* **24**, 1106, 1956.
- ⁵⁷R. McWeeny, *Proc. Roy. Soc. (London)* **A253**, 242, 1959.
- ⁵⁸J. Stephen Binkley, John A. Pople, Warren J. Hehre, *Journal of the American Chemical Society* **102**, 939-947, 1980.
- ⁵⁹Nicolas Bourbaki, *Éléments de mathématique, Livre II : Algèbre, Chap. 3 : Algèbre multilinéaire*, Hermann, Paris, (1948).
- ⁶⁰Patrick Cassam-Chenaï, *Journal of Mathematical Chemistry* **15**, 303-321, 1994.
- ⁶¹Patrick Cassam-Chenaï, Giovanni Granucci, *Chemical Physics Letters* **450**, 151-155, 2007.
- ⁶²P. D. Dacre, R. McWeeny, *Proc. Roy. Soc. A* **317**, 435-454, 1970.
- ⁶³Thom H. Dunning Jr., *The Journal of Chemical Physics* **90**, 1007, 1989.
- ⁶⁴William A. Goddard III, *Physical Review* **157**, 81, 1967.
- ⁶⁵William A. Goddard III, Thom H. Dunning Jr., William J. Hunt, P. Jeffrey Hay *Accounts of Chemical Research* **6**, 11, 368-376, 1973.
- ⁶⁶Martin Head-Gordon et al., *Molecular Physics: An International Journal at the Inter-*

- face Between Chemistry and Physics* **113**, 184-215, 2015.
- ⁶⁷W. J. Hehre, R. F. Stewart, and J. A. Pople, *J. Chem. Phys.* **51**, 2657, 1969.
- ⁶⁸Warren J. Hehre, Robert Ditchfield, John A. Pople, *The Journal of Chemical Physics* **56**, 2257, 1972.
- ⁶⁹S. Huzinaga, A. A. Cantu, *J. Chem. Phys.* **55**, 5543-5549, 1971.
- ⁷⁰Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **9**, 237-239, 1958.
- ⁷¹Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **10**, 125-127, 1959.
- ⁷²Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **11**, 97, 1960.
- ⁷³Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **11**, 409-415, 1960.
- ⁷⁴Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **12**, 185-187, 1960.
- ⁷⁵Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **12**, 351, 1960.
- ⁷⁶Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **13**, 345, 1961.
- ⁷⁷Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **13**, 461, 1961.
- ⁷⁸Ede Kapuy, *Acta Physica Academiae Scientiarum Hungaricae* **15**, 177-183, 1962.
- ⁷⁹E. Kapuy, *J. Chem. Phys.* **44**, 956, 1965.
- ⁸⁰Peter B. Karadakov, David L. Cooper, Brian J. Duke, and Jiabo Li, *J. Phys. Chem.* **A116**, 7238, 2012.
- ⁸¹Airi Kawasaki, Osamu Sugino, *The Journal of Chemical Physics* **145**, 244110, 2016.
- ⁸²A. Khamoshi, Guo P. Chen, Thomas M. Henderson, G. E Scuseria, *The Journal of Chemical Physics* **154**, 074113, 2021.
- ⁸³W. Kutzelnigg, D. Mukherjee, *J. Chem. Phys.* **110**, 2800, 1999.
- ⁸⁴P. A Limacher, *J. Chem. Theory Comput.* **11**, 3629, 2015.
- ⁸⁵P. G. Lykos, R. G. Parr, *J. Chem. Phys.* **24**, 1166, 1956.
- ⁸⁶Peter G. Lykos, Robert G. Parr, *J. Chem. Phys.* **25**, 1301, 1956.
- ⁸⁷Roy McWeeny, Yukiko Mizuno, *Proc. Roy. Soc. (London)* **A259**, 554-577, 1961.
- ⁸⁸Eric Neuscammann, *Mol. Phys.* **114**, 577, 2016.
- ⁸⁹P.R. Surjàn, *Croatica Chemica Acta* **71**, 489-499, 1998.
- ⁹⁰A. M. Tokmachev, R. Dronkowski, *J. Comp. Chem.* **27**, 296-308, 2005.

- ⁹¹A. M. Tokmachev, A. L. Tchougréeff, *Int. J. Quantum Chem.* **106**, 571-587, 2006.
- ⁹²A. M. Tokmachev, R. Dronskowski, *Chem. Phys.* **322**, 423-432, 2006.
- ⁹³Andrey M. Tokmachev, *Int. J. Quantum Chem.* **116**, 265-269, 2016.
- ⁹⁴Matthias Troyer, Uwe-Jens Wiese, *Physical Review Letters* **94**, 170201, 2005.
- ⁹⁵M. Vivier, *Annales scientifiques de l'Ecole Normale Supérieure* **73**, 26, 1956.
- ⁹⁶C. Valdemoro, D. R. Alcoba, L. M. Tel, *Int. J. Quantum Chem.* **93**, 212, 2003.
- ⁹⁷S. Wilson, *J. Chem. Phys.* **64**, 1692, 1976.
- ⁹⁸S. Wilson, *Int. J. Quantum Chem.* **74**, 135, 1999.
- ⁹⁹S. Wilson, *Int. J. Quantum Chem.* **99**, 925-935, 2004.