



N and S cycles in crucifer-legume cover crop mixtures

Antoine Couëdel, Lionel Alletto, Eric Justes

► To cite this version:

Antoine Couëdel, Lionel Alletto, Eric Justes. N and S cycles in crucifer-legume cover crop mixtures. 20th Nitrogen Workshop, Jun 2018, Rennes, France. hal-03763632

HAL Id: hal-03763632

<https://hal.science/hal-03763632>

Submitted on 31 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

N AND S CYCLES IN CRUCIFER-LEGUME COVER CROP MIXTURES

Antoine COUËDEL^{1,*} - Lionel ALLETTO^{1,2} - Éric JUSTES^{1,3}




¹ AGIR, Université de Toulouse, INRA, INPT, INP-EI PURPAN, Castanet-Tolosan, France; ² Chambre Régionale d'Agriculture Occitanie, 31321 Castanet-Tolosan, France;
³ CIRAD, UMR SYSTEM, Univ. Montpellier, CIHEAM-IAMM, INRA, Montpellier SupAgro, 34980 Montpellier, France. *corresponding author: antoine.couedel@inra.fr

Background



Question : Is there a mutualisation of services related to N and S cycles in crucifer and legume cover crop mixtures ?




Material and methods



Catch crop services during the growing period

Results


- crucifers as sole crops
- legumes as sole crops
- mixtures
- bare soil



Nutrient recycling services after cover crop termination



N mineralised from cover crop residues 6 months after termination (kg ha⁻¹)



Conclusion

Crucifers	100%*	100%	18%	100%
Legumes	66%	30%	100%	23%
Mixtures	98%**	99%**	63%**	85%**

→ Our study confirms for a wide range of cover crop species and on four sites x years that crucifer-legume mixtures can provide multi-ecosystem services by **mutualising services of N and S cycles**.

References for more informations

- Couëdel, A., Alletto, L., Justes, É., 2018a. Plant Soil. doi:10.1007/s11104-018-3615-87
 Couëdel, A., Alletto, L., Tribouillois, H., Justes, E., 2018b. Agric. Ecosyst. Environ. 254, 50–59. doi:10.1016/j.agee.2017.11.017

*The service is provided in proportion of the best sole crop family (average of all species tested), which then is considered at 100%.

**Higher than the average of sole cover crops