



HAL
open science

Discrete spectrum of the magnetic Laplacian on perturbed half-planes

Virginie Bonnaillie-Noël, Søren Fournais, Ayman Kachmar, Nicolas Raymond

► **To cite this version:**

Virginie Bonnaillie-Noël, Søren Fournais, Ayman Kachmar, Nicolas Raymond. Discrete spectrum of the magnetic Laplacian on perturbed half-planes. 2022. hal-03763777

HAL Id: hal-03763777

<https://hal.science/hal-03763777>

Preprint submitted on 29 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DISCRETE SPECTRUM OF THE MAGNETIC LAPLACIAN ON PERTURBED HALF-PLANES

VIRGINIE BONNAILLIE-NOËL, SØREN FOURNAIS, AYMAN KACHMAR,
AND NICOLAS RAYMOND

ABSTRACT. The existence of bound states for the magnetic Laplacian in unbounded domains can be quite challenging in the case of a homogeneous magnetic field. We provide an affirmative answer for almost flat corners and slightly curved half-planes when the total curvature of the boundary is positive.

1. INTRODUCTION

We consider the Neumann magnetic Laplacian with constant magnetic field $B = 1$ on an open set Ω_δ , which is defined through the following (closed) quadratic form

$$\forall \psi \in H_{\mathbf{A}}^1(\Omega_\delta), \quad Q_\delta(\psi) = \int_{\Omega_\delta} |(-i\nabla + \mathbf{A})\psi|^2 d\mathbf{x}.$$

Here $\mathbf{A} = (-x_2, 0)$ is a vector potential associated with $B = 1$ in the sense that

$$\partial_1 A_2 - \partial_2 A_1 = 1.$$

Then, we consider the magnetic Laplacian as the self-adjoint operator \mathcal{L}_δ associated with Q_δ . We denote by $\lambda(\delta)$ the bottom of its spectrum

$$\lambda(\delta) = \inf_{\substack{\psi \in H_{\mathbf{A}}^1(\Omega_\delta) \\ \psi \neq 0}} \frac{\|(-i\nabla + \mathbf{A})\psi\|_{L^2(\Omega_\delta)}^2}{\|\psi\|_{L^2(\Omega_\delta)}^2}.$$

The study of the spectrum of such operators in different geometries has been the focus of much interest in recent decades. Below, we will give a short overview of results relevant for the present article. For more in depth coverage we refer to [7, 9].

In general, when Ω_δ is unbounded, we do not know if $\lambda(\delta)$ is an eigenvalue. For instance, in the case when $\Omega_\delta = \Omega_0 =: \mathbf{R} \times \mathbf{R}_+$, it is well-known that the spectrum is absolutely continuous and given by the half-line $[\Theta_0, +\infty)$, where $\Theta_0 \approx 0.59$ is a positive universal constant (see (2.2) below). In the present article Ω_δ will be a perturbation of the half-plane Ω_0 . Let us describe the two types of perturbations that we consider. The first type is a singular perturbation when Ω_δ is of the (corner) form

$$(1.1) \quad \Omega_\delta = \mathcal{C}_\delta := \{\mathbf{x} = (x_1, x_2) \in \mathbf{R} \times \mathbf{R}_+ : \frac{x_2}{\tan \delta} > -x_1\}, \quad \delta \in (0, \pi).$$

The second type of perturbation under consideration is regular. Consider a bounded continuous compactly supported function $\kappa : \mathbf{R} \rightarrow \mathbf{R}$ and, for all $\delta \geq 0$, a simple \mathcal{C}^2 curve $\gamma_\delta : \mathbf{R} \rightarrow \mathbf{R}^2$, parametrized by arc-length and such that its algebraic curvature κ_δ satisfies

$$\kappa_\delta(s) = \delta \kappa(s).$$

Note that $\gamma_\delta''(s) = \delta \kappa(s) \mathbf{n}_\delta(s)$ where $\mathbf{n}_\delta(s)$ is the unit normal such that $\det(\gamma_\delta', \mathbf{n}_\delta) = 1$. We also assume that $\gamma_0(s) = (s, 0)$, $\gamma_\delta(0) = (0, 0)$ and that $\delta \mapsto \gamma_\delta'$ is continuous

for the uniform topology at $\delta = 0$. Let us now define our perturbed half-space. We write $\mathbf{R}^2 \setminus \gamma_\delta = \Gamma_\delta^+ \sqcup \Gamma_\delta^-$ in such a way that \mathbf{n}_δ is the inward pointing normal to $\partial\Gamma_\delta^+$. We let $\Omega_\delta = \Gamma_\delta^+$. A typical example of such a configuration is given by $\gamma_\delta(s) = (s, f_\delta(s))$ where $f_\delta(s) = \delta \int_0^s g(\sigma)(1 - \delta^2 g(\sigma)^2)^{-1/2} d\sigma$ and $g(\sigma) = \int_0^\sigma \kappa(x) dx$, with δ chosen in $(0, \delta_0)$ for a sufficiently small δ_0 to ensure that $\delta g(\sigma) \in (-1, 1)$; in which case $\Gamma_\delta^+ = \{(x, y) \in \mathbf{R}^2 \mid y > f_\delta(x)\}$.

By the Persson's theorem (see for instance [1]), it is well-known in both cases that the essential spectrum of this operator \mathcal{L}_δ is the same as the spectrum of \mathcal{L}_0 , *i.e.*, $[\Theta_0, +\infty)$. Let us now state our main two results, both establishing the existence of a bound state.

Theorem 1.1 (Almost flat corner). *We assume that $\Omega_\delta = \mathcal{C}_\delta$. There exists $\delta_0 \in (0, \pi)$ such that, for all $\delta \in (0, \delta_0)$,*

$$\lambda(\delta) \leq \Theta_0 - \frac{C_1^2}{4} \delta^2 + o(\delta^2) < \Theta_0,$$

where $C_1 > 0$ is a universal constant (defined below in (2.3)). In particular, the bottom of the spectrum of \mathcal{L}_δ belongs to the discrete spectrum.

In the corner setting, $\lambda(\delta)$ is only known to be an eigenvalue for the non-flat situation when $\delta \in (\frac{\pi}{2} - \varepsilon, \pi)$, for a sufficiently small $\varepsilon > 0$, see [1, 5]. The regime $\delta \rightarrow 0$ considered in the present article tackles the subtle situation when we know that the discrete spectrum tends to disappear. Let us also underline that it is still an open question to know if $\delta \mapsto \lambda(\delta)$ is monotone non-increasing (as suggested by the numerical simulations in [1]). If one were able to establish this monotonicity, Theorem 1.1 would imply that $\lambda(\delta) < \Theta_0$ for all $\delta \in (0, \pi)$, and thus the existence of a bound state for all (non flat) convex corners.

We next state the result for the slightly curved half-plane.

Theorem 1.2 (Slightly curved half-plane). *We assume $\Omega_\delta = \Gamma_\delta^+$ and $\int_{\mathbf{R}} \kappa(s) ds > 0$. There exists $\delta_0 > 0$ such that, for all $\delta \in (0, \delta_0)$, the discrete spectrum of \mathcal{L}_δ is non-empty. Moreover,*

$$\lambda(\delta) \leq \Theta_0 - \left(\frac{C_1}{2} \int_{\mathbf{R}} \kappa(s) ds \right)^2 \delta^2 + o(\delta^2),$$

where C_1 is the same universal constant as in Theorem 1.1.

Our analysis suggests that there is only one simple eigenvalue below $\Theta_0 - C\delta^{\frac{1}{2}}$, for some constant $C > 0$. This would follow from a dimensional reduction as in [6, 8] or in the Grushin spirit (see [2]). The question of estimating the number of bound states (below Θ_0) remains open. This requires the derivation of a very precise operator near the threshold of the essential spectrum, what is a quite interesting problem sharing similar features as in [3].

Remark 1.3. Actually, Theorem 1.1 can be seen as a *formal* consequence of Theorem 1.2. Indeed, it is possible to exhibit a normal parametrization of $\partial\mathcal{C}_\delta$ by considering

$$\gamma_\delta(s) = (s\mathbb{1}_{\mathbf{R}_+}(s) + s \cos \delta \mathbb{1}_{\mathbf{R}_-}(s), -s \sin \delta \mathbb{1}_{\mathbf{R}_-}(s)).$$

In the sense of distributions, we have

$$\gamma_\delta'' = (1 - \cos \delta, \sin \delta) \mathfrak{d}_0 = 2 \sin \left(\frac{\delta}{2} \right) \mathbf{n}_\delta \mathfrak{d}_0, \quad \mathbf{n}_\delta = (\sin(\delta/2), \cos(\delta/2)),$$

where \mathfrak{d}_0 is the Dirac distribution at 0 and \mathbf{n}_δ is the direction of the bisector of \mathcal{C}_δ . Formally, the curvature is $\kappa_\delta = 2 \sin\left(\frac{\delta}{2}\right) \mathfrak{d}_0$.

Organization of the article. In Section 2, we introduce the constants Θ_0 and C_1 , related to the de Gennes operator (2.1). Section 3 is devoted to the proof of Theorem 1.1, whereas Section 4 deals with that of Theorem 1.2. In both situations, the proof follows by construction of an appropriate trial state. In the corner case, the phase of the trial state (see (3.5)) is reminiscent of the construction appearing in the non-linear setting of [4], while the amplitude (see (3.11)) is obtained by minimizing a new energy functional in (3.16). In the regular case, we use a tensorized trial state in curvilinear coordinates (see (4.1)) involving the bound state of a 1D model operator studied in [10] and revisited in Appendix A.

2. THE DE GENNES OPERATOR AND THE CONSTANT C_1

The material of this section is standard and only included for convenience and to fix notation. We refer to [7] for more material and reference to earlier works. The constants Θ_0 and C_1 in Theorems 1.1 and 1.2 are defined starting from a family of 1D harmonic oscillators on the half axis. For all $\xi \in \mathbf{R}$, let us denote by $\mu(\xi)$ the first eigenvalue of the operator

$$(2.1) \quad H_\xi = -\frac{d^2}{dt^2} + (t - \xi)^2 \quad \text{in } L^2(\mathbf{R}_+),$$

with Neumann boundary condition at 0, $u'(0) = 0$.

We introduce the de Gennes constant

$$(2.2) \quad \Theta_0 = \inf_{\xi \in \mathbf{R}} \mu(\xi).$$

We know that $\frac{1}{2} < \Theta_0 < 1$ and there exists a unique ξ_0 such that

$$\Theta_0 = \mu(\xi_0).$$

Furthermore, $\xi_0 = \sqrt{\Theta_0}$ and $\mu''(\xi_0) > 0$. Let us denote by f_\star the positive normalized ground state of Θ_0 , i.e.

$$H_{\xi_0} f_\star = \Theta_0 f_\star, \quad f_\star > 0, \quad f'_\star(0) = 0, \quad \int_{\mathbf{R}_+} |f_\star(t)|^2 dt = 1.$$

We introduce the constant C_1 as follows

$$(2.3) \quad C_1 = \frac{|f_\star(0)|^2}{3}.$$

The function f_\star belongs to $\mathcal{S}(\mathbf{R}_+)$ and decays exponentially at infinity. It satisfies the additional property (Feynman-Hellmann)

$$(2.4) \quad \int_{\mathbf{R}_+} (t - \xi_0) |f_\star(t)|^2 dt = 0.$$

Noticing that

$$\begin{aligned} \int_0^{+\infty} t |f'_\star(t)|^2 dt &= - \int_0^{+\infty} (t f'_\star)' f_\star dt = \int_0^{+\infty} (t(-f''_\star) f_\star - f'_\star f'_\star) dt \\ &= \int_0^{+\infty} (\Theta_0 t - t(\xi_0 - t)^2) |f_\star(t)|^2 dt + \frac{f_\star(0)^2}{2}, \end{aligned}$$

we get, using (2.3), the interesting identity

$$(2.5) \quad \int_{\mathbf{R}_+} (|f'_\star(t)|^2 + (\xi_0 - t)^2 |f_\star(t)|^2 - \Theta_0 |f_\star(t)|^2) t dt = \frac{3C_1}{2}.$$

Another interesting identity is

$$(2.6) \quad \int_{\mathbf{R}_+} (t - \xi_0) t (t - 2\xi_0) |f_\star(t)|^2 dt = \frac{C_1}{2},$$

which follows by writing

$$(t - \xi_0)(t - 2\xi_0)t = (t - \xi_0)^3 - \xi_0^2(t - \xi_0),$$

using (2.4) and the formula (see [7, Lemma 3.2.7]):

$$\int_{\mathbf{R}_+} (t - \xi_0)^3 |f_\star(t)|^2 dt = \frac{C_1}{2}.$$

For a positive number ℓ , let us introduce the function

$$(2.7) \quad f_\ell(t) = \zeta\left(\frac{t}{\ell}\right) f_\star(t)$$

where $\zeta \in \mathcal{C}_0^\infty(\mathbf{R})$ satisfies

$$(2.8) \quad 0 \leq \zeta \leq 1, \quad \text{supp } \zeta \subset [-1, 1], \quad \zeta = 1 \text{ on } \left[-\frac{1}{2}, \frac{1}{2}\right].$$

Consequently, as $\ell \rightarrow +\infty$, we have

$$(2.9) \quad \begin{aligned} \int_{\mathbf{R}_+} |f_\ell(t)|^2 dt &= 1 + \mathcal{O}(\ell^{-\infty}), \\ q(f_\ell) &:= \int_{\mathbf{R}_+} (|f'_\ell(t)|^2 + (t - \xi_0)^2 |f_\ell(t)|^2) dt = \Theta_0 + \mathcal{O}(\ell^{-\infty}), \\ \int_{\mathbf{R}_+} (t - \xi_0)^k |f_\ell(t)|^2 dt &= \int_{\mathbf{R}_+} (t - \xi_0)^k |f_\star(t)|^2 dt + \mathcal{O}(\ell^{-\infty}), \end{aligned}$$

where $\mathcal{O}(\ell^{-\infty})$ denotes a quantity equal to $\mathcal{O}(\ell^{-N})$ for all $N > 0$.

Since $f_\star \in \mathcal{S}(\mathbf{R}_+)$, we deduce the following two identities from (2.5), which will be useful below in our proof of Theorem 1.1,

$$(2.10) \quad \begin{aligned} \int_{\mathbf{R}_+} (|f'_\ell(t)|^2 + (\xi_0 - t)^2 |f_\ell(t)|^2 - \Theta_0 |f_\ell(t)|^2) t dt &= \frac{3C_1}{2} + \mathcal{O}(\ell^{-\infty}), \\ \int_{\mathbf{R}_+} (t - \xi_0)(t - 2\xi_0) |f_\ell(t)|^2 t dt &= \frac{C_1}{2} + \mathcal{O}(\ell^{-\infty}). \end{aligned}$$

3. ALMOST FLAT SECTORS

This section is devoted to the proof of Theorem 1.1, so $\Omega_\delta = \mathcal{C}_\delta$ hereafter. The proof is by construction of a quasi-mode having approximately the form (after truncation)

$$\Psi^{\text{tr}} \approx f_\star(t) e^{i\Phi(s,t)}$$

where s denotes the tangential variable along $\partial\mathcal{C}_\delta$ and t denotes the transversal variable. For instance, $(s, t) = (x_1, x_2)$ when $\delta = 0$ (in which case \mathcal{C}_δ is the half-plane $\mathbf{R} \times \mathbf{R}_+$).

The phase term Φ is, up to symmetry considerations, a perturbation of $i\xi_0 s$. As already mentioned, the idea of perturbing the phase term in an almost flat sector was first introduced in the non-linear framework of the Ginzburg-Landau functional [4]. We use the same construction and add to this by proving that the phase term proposed in [4] is rather the optimal choice. Interestingly, we determine the best truncation profile by minimizing a non-linear functional, which allows us to capture the δ^2 -term of Theorem 1.1.

3.1. Geometric framework. We denote by T^+ the following trapezoid:

$$T^+ := \{\mathbf{x} \in (0, +\infty) \times (0, \ell) : x_2 \tan(\delta/2) < x_1\}.$$

Consider the angle bisector

$$D_\delta = \{\mathbf{x} \in \mathbf{R}^2 : x_1 = x_2 \tan(\delta/2)\}.$$

We denote by S_δ the reflection in the line D_δ , whose matrix is

$$S_\delta = \begin{pmatrix} -\cos \delta & \sin \delta \\ \sin \delta & \cos \delta \end{pmatrix}.$$

We denote by T^- the reflection of T^+ , i.e.,

$$T^- := S_\delta T^+.$$




FIGURE 1. Geometric setting: The figure illustrates the sector cut-off at height ℓ . Here $T^+ = T_{\delta,\gamma}^+ \cup V_{\delta,\gamma}^+$ and similarly for T^- . The symmetry axis D_δ is drawn in blue.

3.2. Towards a test function. Let us try to define a test function compatible with the symmetry and the magnetic field. Let us consider a function u_+ such that

$$(3.1) \quad E_+ := \int_{T^+} |(-i\nabla + \mathbf{A})u_+|^2 d\mathbf{x} < +\infty.$$

Now, we want to extend u_+ by using the symmetry. We wish to do this in such a way that the magnetic energy on T^- coincides with the one on T^+ .

Lemma 3.1. *Considering*

$$(3.2) \quad \phi(\mathbf{y}) = \frac{\sin(2\delta)}{4}(y_1^2 - y_2^2) - y_1 y_2 \sin^2 \delta$$

and

$$(3.3) \quad u_-(\mathbf{x}) = e^{-i\phi(S_\delta \mathbf{x})} \bar{u}_+(S_\delta \mathbf{x}),$$

we have

$$(3.4) \quad \int_{T^+} |(-i\nabla + \mathbf{A})u_+|^2 d\mathbf{x} = \int_{T^-} |(-i\nabla + \mathbf{A})u_-|^2 d\mathbf{x}.$$

Proof. For a given function u_- on T^- , we use the change of variable given by $\mathbf{x} = S_\delta \mathbf{y}$ (remember that $S_\delta = S_\delta^{-1} = S_\delta^*$) and notice that

$$\int_{T^-} |(-i\nabla + \mathbf{A})u_-|^2 d\mathbf{x} = \int_{T^+} |(-i\nabla_{\mathbf{y}} + \tilde{\mathbf{A}}(\mathbf{y}))\overset{\circ}{u}_-|^2 d\mathbf{y},$$

with

$$\tilde{\mathbf{A}} = S_\delta(\mathbf{A} \circ S_\delta), \quad \overset{\circ}{u}_- = u_- \circ S_\delta(\mathbf{y}).$$

A straightforward computation gives

$$\tilde{\mathbf{A}}(\mathbf{y}) = S_\delta \begin{pmatrix} -y_1 \sin \delta - y_2 \cos \delta \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{y_1}{2} \sin(2\delta) + y_2 \cos^2 \delta \\ -y_1 \sin^2 \delta - \frac{y_2}{2} \sin(2\delta) \end{pmatrix}.$$

The magnetic field associated with $\tilde{\mathbf{A}}$ is

$$\partial_{y_1} \tilde{A}_2 - \partial_{y_2} \tilde{A}_1 = -1.$$

Let us consider the function ϕ defined in (3.2). It satisfies

$$\tilde{\mathbf{A}} - \begin{pmatrix} y_2 \\ 0 \end{pmatrix} = \nabla \phi.$$

Therefore,

$$\begin{aligned} \int_{T^-} |(-i\nabla + \mathbf{A})u_-|^2 d\mathbf{x} &= \int_{T^+} \left| \left(-i\nabla_{\mathbf{y}} + \begin{pmatrix} y_2 \\ 0 \end{pmatrix} \right) e^{i\phi(\mathbf{y})} \overset{\circ}{u}_-(\mathbf{y}) \right|^2 d\mathbf{y} \\ &= \int_{T^+} |(-i\nabla_{\mathbf{y}} + \mathbf{A}(\mathbf{y})) e^{-i\phi(\mathbf{y})} v_-(\mathbf{y})|^2 d\mathbf{y}, \quad \text{with } v_- = \overline{\overset{\circ}{u}_-}, \end{aligned}$$

so that (3.4) holds if

$$e^{-i\phi(\mathbf{y})} \overline{\overset{\circ}{u}_-}(\mathbf{y}) = u_+(\mathbf{y}),$$

or, equivalently,

$$u_-(\mathbf{x}) = e^{-i\phi(S_\delta \mathbf{x})} \bar{u}_+(S_\delta \mathbf{x}).$$

□

Remark 3.2. If we choose

$$u_+(x_1, x_2) = f(x_2) e^{i\xi_0 x_1},$$

with a real-valued f in the Schwartz class, (3.3) gives

$$u_-(\mathbf{x}) = f(x_1 \sin \delta + x_2 \cos \delta) e^{-i\xi_0(-x_1 \cos \delta + x_2 \sin \delta)} e^{-i\phi(\mathbf{x})},$$

since $\phi(S_\delta \mathbf{x}) = \phi(\mathbf{x})$. Of course, this choice of u_+ is not appropriate since (3.1) is not satisfied due to the lack of integrability with respect to x_1 . However, up to using

a cutoff function with respect to x_1 , this gives a rather good idea of the shape of our test function in T^+ . Now, if we consider

$$u(\mathbf{x}) = \begin{cases} f(x_2)e^{i\xi_0 x_1}, & \text{if } \mathbf{x} \in T^+ \\ f(x_1 \sin \delta + x_2 \cos \delta)e^{i\xi_0(x_1 \cos \delta - x_2 \sin \delta)}e^{-i\phi(\mathbf{x})}, & \text{if } \mathbf{x} \in T^- \end{cases},$$

we see that u does not belong to H^1 near the symmetry axis D_δ due to the phase shift, which does not vanish on D_δ . In the next section, we slightly modify this function u to solve this inconvenience.

3.3. Smoothing the transition near D_δ . For the given δ , we let $0 < \gamma < \pi - \delta$ and define the following trial state (in polar coordinates)

$$(3.5) \quad \Psi(r \cos \theta, r \sin \theta) = \begin{cases} f(r \sin \theta)e^{i\xi_0 r \cos \theta}, & \text{if } (r \cos \theta, r \sin \theta) \in T_{\delta, \gamma}^+, \\ f(r \sin \theta)e^{i\alpha(r, \theta)}, & \text{if } (r \cos \theta, r \sin \theta) \in V_{\delta, \gamma}^+, \\ f(r \sin(\theta + \delta))e^{i\alpha(r, \theta)}, & \text{if } (r \cos \theta, r \sin \theta) \in V_{\delta, \gamma}^-, \\ f(r \sin(\theta + \delta))e^{i\xi_0 r \cos(\theta + \delta) - ir^2 \phi(\cos \theta, \sin \theta)}, & \text{if } (r \cos \theta, r \sin \theta) \in T_{\delta, \gamma}^-, \\ 0, & \text{else,} \end{cases}$$

where (cf. Fig.1):

- $V_{\delta, \gamma}^+$ is the sector defined in polar coordinates by $0 < r < r_* := \frac{\ell}{\cos(\frac{\delta + \gamma}{2})}$ and $\theta \in (\theta_\delta - \frac{\gamma}{2}, \theta_\delta)$ and $V_{\delta, \gamma}^- = S_\delta V_{\delta, \gamma}^+$, with $\theta_\delta = \frac{\pi - \delta}{2}$.
- the trapezoids $T_{\delta, \gamma}^+$ and $T_{\delta, \gamma}^-$ are given by

$$T_{\delta, \gamma}^+ = \left\{ \mathbf{x} \in (0, +\infty) \times (0, \ell) : x_2 \tan\left(\frac{\delta - \gamma}{2}\right) < x_1 \right\}$$

$$\text{and } T_{\delta, \gamma}^- = S_\delta T_{\delta, \gamma}^+.$$

The function f is given by $f = f_\ell$, see (2.7), and the phase α is chosen so that the function Ψ belongs to $H_{\text{loc}}^1(\mathcal{C}_\delta)$. Let us give an explicit choice of α . Note that

$$\phi(\cos \theta, \sin \theta) = \frac{\sin \delta}{2} \cos(2\theta + \delta),$$

and consider the two phases

$$\alpha_+(r, \theta) = r\xi_0 \cos(\theta), \quad \alpha_-(r, \theta) = r\xi_0 \cos(\theta + \delta) - \frac{r^2}{2} \sin \delta \cos(2\theta + \delta).$$

Notice that the transition zone is simply given by $\theta \in (\theta_\delta - \frac{\gamma}{2}, \theta_\delta + \frac{\gamma}{2})$ and that we have

$$\alpha_+\left(r, \theta_\delta - \frac{\gamma}{2}\right) = r\xi_0 \sin\left(\frac{\delta + \gamma}{2}\right)$$

and

$$\alpha_-\left(r, \theta_\delta + \frac{\gamma}{2}\right) = -r\xi_0 \sin\left(\frac{\delta + \gamma}{2}\right) + \frac{r^2}{2} \sin \delta \cos \gamma.$$

This leads to the choice

$$(3.6) \quad \alpha(r, \theta) = br^2 - \chi_{\delta, \gamma}(\theta) (ar - br^2),$$

with

$$a = \xi_0 \sin\left(\frac{\delta + \gamma}{2}\right), \quad b = \frac{1}{4} \sin \delta \cos \gamma,$$

and

$$\chi_{\delta,\gamma}(\theta) = \chi\left(\frac{2(\theta - \theta_\delta)}{\gamma}\right),$$

where χ is a smooth odd function such that $\chi(t) = 1$ for $t \geq 1$.

Remark 3.3. The function Ψ has no decay in the x_1 -direction, so it will be complemented by a well-chosen cut-off in that variable.

3.4. Estimate of the energy. We consider a function $\eta_+ \in H^1(\mathbf{R}_+)$ equal to 1 in a fixed neighborhood of 0 (so that $(x_1, x_2) \mapsto \eta_+(x_1)$ equals 1 on $V_{\delta,\gamma}^+$ as soon as δ and γ are small enough). This function will be chosen later on in Section 3.5.

The aim of this section is to establish the following proposition valid in the regime where $\ell \rightarrow +\infty$ and $(\delta, \gamma) \rightarrow 0$ provided that $\delta = o(\gamma)$.

Proposition 3.4. *For small enough values of $\delta, \gamma, \ell^{-1}$ and under the assumption that $\delta \leq \gamma$, we have*

$$\begin{aligned} \int_{T^+} |\eta_+(x_1)|^2 |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} &\leq (\Theta_0 + \mathcal{O}(\ell^{-\infty})) \|\eta_+\|_{L^2(\mathbf{R}_+)}^2 - J \frac{\delta}{2} \\ &\quad + \mathcal{O}(\gamma^{-1}\delta^2) + \mathcal{O}(\gamma^3) + \mathcal{O}(\ell^{-\infty}), \end{aligned}$$

with

$$(3.7) \quad J = \int_0^{+\infty} (|f'|^2 + (t - \xi_0)^2 |f(t)|^2) t dt - \int_0^{+\infty} (t - \xi_0)(t - 2\xi_0)t |f(t)|^2 dt.$$

The remainder terms $\mathcal{O}(\gamma^{-1}\delta^2)$, $\mathcal{O}(\ell^{-\infty})$ and $\mathcal{O}(\gamma^3)$ are controlled uniformly w.r.t. the function η_+ .

Proposition 3.4 is a consequence of Lemmas 3.5, Remark 3.6, and 3.7.

Lemma 3.5. *We have the estimate*

$$\int_{V_{\delta,\gamma}^+} |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} = \frac{\gamma}{2} J_{\delta,\gamma} + \mathcal{O}(\gamma^3 + \gamma\ell^{-\infty})$$

where

$$J_{\delta,\gamma} := \int_0^{+\infty} \left(|f'(t)|^2 + \left(t - \left(\xi_0 \left(1 + \frac{\delta}{\gamma} \right) - \frac{\delta}{2\gamma} t \right) \right)^2 |f(t)|^2 \right) t dt.$$

Remark 3.6. Expanding $J_{\delta,\gamma}$, we observe that

$$\frac{\gamma}{2} J_{\delta,\gamma} = \frac{\gamma}{2} \int_0^{+\infty} (|f'|^2 + (t - \xi_0)^2 |f(t)|^2) t dt + \frac{\delta}{2} \int_0^{+\infty} (t - \xi_0)(t - 2\xi_0)t |f(t)|^2 dt + \mathcal{O}(\gamma^{-1}\delta^2),$$

where we used the exponential decay of f_* to control the remainder.

Proof of Lemma 3.5. We have

$$\int_{V_{\delta,\gamma}^+} |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} = \int_{V_{\delta,\gamma}^+} |(-i\nabla + \mathbf{A}_0)\tilde{\Psi}|^2 d\mathbf{x}, \quad \mathbf{A}_0 = \frac{1}{2}(-x_2, x_1),$$

where $\tilde{\Psi} = e^{-ix_1 x_2/2} \Psi$. We get

$$\int_{V_{\delta,\gamma}^+} |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} = \int_{\theta_\delta - \frac{\gamma}{2}}^{\theta_\delta} \int_0^{r^*} \left(|\partial_r \tilde{\Psi}|^2 + r^{-2} \left| \left(-i\partial_\theta + \frac{r^2}{2} \right) \tilde{\Psi} \right|^2 \right) r dr d\theta.$$

Then, we write

$$\tilde{\Psi} = e^{i(-r^2 \sin(2\theta)/4 + \alpha(r, \theta))} f(r \sin \theta),$$

and, using that f is real-valued, we find

$$|\partial_r \tilde{\Psi}|^2 = \sin^2 \theta |f'(r \sin \theta)|^2 + (-r \sin(2\theta)/2 + \partial_r \alpha)^2 |f(r \sin \theta)|^2,$$

and

$$\begin{aligned} \left| \left(-i\partial_\theta + \frac{r^2}{2} \right) \tilde{\Psi} \right|^2 &= \left(-\frac{r^2}{2} \cos(2\theta) + \partial_\theta \alpha + \frac{r^2}{2} \right)^2 |f(r \sin \theta)|^2 + r^2 \cos^2 \theta |f'(r \sin \theta)|^2 \\ &= (r^2 \sin^2(\theta) + \partial_\theta \alpha)^2 |f(r \sin \theta)|^2 + r^2 \cos^2 \theta |f'(r \sin \theta)|^2. \end{aligned}$$

It follows that

$$\begin{aligned} &\int_{V_{\delta, \gamma}^+} |(-i\nabla + \mathbf{A})\Psi|^2 dx \\ &= \int_{\theta_\delta - \frac{\gamma}{2}}^{\theta_\delta} \int_0^{r^*} \left(|f'(r \sin \theta)|^2 \right. \\ &\quad \left. + \left((-r \sin(2\theta)/2 + \partial_r \alpha)^2 + (r \sin^2(\theta) + r^{-1} \partial_\theta \alpha)^2 \right) |f(r \sin \theta)|^2 \right) r dr d\theta \\ (3.8) \quad &= \int_{\theta_\delta - \frac{\gamma}{2}}^{\theta_\delta} \int_0^{r^*} \left(|f'(r \sin \theta)|^2 + F(r, \theta, \chi_{\delta, \gamma}) |f(r \sin \theta)|^2 \right) r dr d\theta, \end{aligned}$$

where

$$F(r, \theta, \chi_{\delta, \gamma}) = (-r \sin(2\theta)/2 - \chi_{\delta, \gamma}(\theta)(a - 2br) - 2br)^2 + (r \sin^2(\theta) - \chi'_{\delta, \gamma}(\theta)(a - br))^2.$$

Notice that the interval of integration in θ has length $\gamma/2 \ll 1$ and $\theta \in [\theta_\delta - \frac{\gamma}{2}, \theta_\delta]$ implies

$$1 \geq \sin \theta \geq 1 - C(\delta - \gamma)^2, \text{ and } 0 \leq \sin(2\theta) \leq \delta + \gamma.$$

Furthermore,

$$a = \xi_0 \frac{\gamma + \delta}{2} + \mathcal{O}(\gamma^3), \quad b = \frac{\delta}{4} + \mathcal{O}(\gamma^3)$$

and

$$\chi'_{\delta, \gamma}(\theta) = \frac{2}{\gamma} \chi' \left(\frac{2(\theta - \theta_\delta)}{\gamma} \right).$$

Therefore, the first part of $F(r, \theta, \chi_{\delta, \gamma})$ is small, and we get, using the decay of f and f' ,

$$\begin{aligned}
& \int_{V_{\delta, \gamma}^+} |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} \\
&= \int_{\theta_{\delta - \frac{\gamma}{2}}}^{\theta_{\delta}} \int_0^{r^*} \left(|f'(r \sin \theta)|^2 + (r \sin^2(\theta) - \chi'_{\delta, \gamma}(\theta)(a - br))^2 |f(r \sin \theta)|^2 \right) r dr d\theta + \mathcal{O}(\gamma^3) \\
&= \int_{\theta_{\delta - \frac{\gamma}{2}}}^{\theta_{\delta}} \int_0^{+\infty} \left(|f'(t)|^2 + \left(t - \chi' \left(\frac{2(\theta - \theta_{\delta})}{\gamma} \right) \left(\xi_0 \left(1 + \frac{\delta}{\gamma} \right) - \frac{\delta}{2\gamma} t \right) \right)^2 |f(t)|^2 \right) t dt d\theta \\
&\quad + \mathcal{O}(\gamma^3 + \gamma \ell^{-\infty}) \\
&= \frac{\gamma}{2} \int_0^{+\infty} \left(|f'(t)|^2 + t^2 |f(t)|^2 \right) t dt - \gamma \int_0^{+\infty} \left(\xi_0 \left(1 + \frac{\delta}{\gamma} \right) - \frac{\delta}{2\gamma} t \right) |f(t)|^2 t^2 dt \\
(3.9) \quad &+ \frac{\gamma}{2} \left(\int_{-1}^0 \chi^2 d\theta \right) \int_0^{+\infty} \left(\xi_0 \left(1 + \frac{\delta}{\gamma} \right) - \frac{\delta}{2\gamma} t \right)^2 |f(t)|^2 t dt + \mathcal{O}(\gamma^3 + \gamma \ell^{-\infty}),
\end{aligned}$$

where we used that χ is odd with $\chi(-1) = -1$, to get the last equality. Therefore, the optimal choice is that $\chi(x) = x$ on $[-1, 1]$ and the result follows. \square

Lemma 3.7. *We have*

$$\begin{aligned}
\int_{T_{\delta, \gamma}^+} \eta_+(x_1)^2 |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} &\leq (\Theta_0 + \mathcal{O}(\ell^{-\infty})) \|\eta_+\|_{L^2(\mathbf{R}_+)}^2 \\
&\quad - \frac{\gamma + \delta}{2} \int_0^{+\infty} (f'(t))^2 + (t - \xi_0)^2 |f|^2 t dt \\
&\quad + \mathcal{O}(\gamma^3) + \mathcal{O}(\ell^{-\infty}).
\end{aligned}$$

Proof. We recall that $\Psi(\mathbf{x}) = u(\mathbf{x}) = f(x_2) e^{i\xi_0 x_1}$, on $T_{\delta, \gamma}^+$ and that $\eta_+ = 1$ on $[0, \varepsilon]$. For small enough γ, δ we can therefore write

$$\int_{T_{\delta, \gamma}^+} \eta_+(x_1)^2 |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} = \int_R \eta_+(x_1)^2 |(-i\nabla + \mathbf{A})u|^2 d\mathbf{x} - E,$$

with $R = (0, +\infty) \times (0, \ell)$ and

$$E = \int_{R \setminus T_{\delta, \gamma}^+} |(-i\nabla + \mathbf{A})u|^2 d\mathbf{x}.$$

The calculation of E is similar to the beginning of the proof of Lemma 3.5. We have

$$E = \int_{R \setminus T_{\delta, \gamma}^+} |(-i\nabla + \mathbf{A}_0)\tilde{u}|^2 d\mathbf{x}, \quad \tilde{u}(\mathbf{x}) = e^{-ix_1 x_2 / 2} u(\mathbf{x}),$$

with $\mathbf{A}_0(\mathbf{x}) = \frac{1}{2}(-x_2, x_1)$. We let $w(r, \theta) = \tilde{u}(r \cos \theta, r \sin \theta) = f(r \sin \theta) e^{i(\xi_0 r \cos \theta - \frac{r^2}{4} \sin(2\theta))}$ to get

$$E = \int_{\theta_{\delta - \frac{\gamma}{2}}}^{\frac{\pi}{2}} \int_0^{r^*(\theta)} \left(|\partial_r w|^2 + r^{-2} \left| \left(-i\partial_\theta + \frac{r^2}{2} \right) w \right|^2 \right) r dr d\theta, \quad r^*(\theta) = \frac{\ell}{\sin \theta} \geq \ell.$$

Since f is real-valued,

$$|\partial_r w|^2 = \sin^2 \theta |f'(r \sin \theta)|^2 + (\xi_0 \cos \theta - r \sin \theta \cos \theta)^2 |f(r \sin \theta)|^2$$

$$r^{-2} \left| (-i\partial_\theta + \frac{r^2}{2})w \right|^2 = \cos^2 \theta |f'(r \sin \theta)|^2 + (-\xi_0 \sin \theta + r \sin^2(\theta))^2 |f(r \sin \theta)|^2.$$

We get

$$(3.10) \quad E = \int_{\theta_\delta - \gamma/2}^{\pi/2} \int_0^{\ell/\sin \theta} (|f'(r \sin \theta)|^2 + (\xi_0 - r \sin \theta)^2 |f(r \sin \theta)|^2) r dr d\theta,$$

Thus, with the change of variable $r = t/\sin \theta$, we get for all $N > 0$,

$$E \geq \frac{\gamma + \delta}{2} \int_0^{+\infty} (t f'(t)^2 + t(\xi_0 - t)^2 |f|^2) dt - C\gamma^3 - C\gamma \ell^{-N}.$$

Moreover,

$$\begin{aligned} \int_R \eta_+(x_1)^2 |(-i\nabla + \mathbf{A})u|^2 d\mathbf{x} &= \int_R \eta_+(x_1)^2 (|(-i\partial_1 - x_2)u|^2 + |\partial_{x_2} u|^2) d\mathbf{x} \\ &= \int_R \eta_+(x_1)^2 (|(\xi_0 - x_2)f(x_2)|^2 + |f'(x_2)|^2) d\mathbf{x} \\ &= \|\eta_+\|_{L^2(\mathbf{R}_+)}^2 \int_0^\ell (|(\xi_0 - x_2)f(x_2)|^2 + |f'(x_2)|^2) dx_2 \\ &= (\Theta_0 + \mathcal{O}(\ell^{-\infty})) \|\eta\|_{L^2(\mathbf{R}_+)}^2. \end{aligned}$$

□

3.5. Proof of Theorem 1.1. We have now almost all the elements at hand to establish Theorem 1.1. Let us first truncate the function Ψ to produce a test function in $H_{\mathbf{A}}^1(\mathcal{C}_\delta)$. We introduce the function

$$\Psi^{\text{tr}} = \eta \Psi$$

where

$$(3.11) \quad \eta(\mathbf{x}) = \begin{cases} \eta_+(x_1) & \text{if } \mathbf{x} \in T^+ \\ \eta_+(-x_1 \cos \delta + x_2 \sin \delta) & \text{if } \mathbf{x} \in T^-. \end{cases}$$

and $\eta_+ \in H^1(\mathbf{R}_+)$ is equal to 1 on the interval $(0, \varepsilon)$, for a fixed (and arbitrary) $\varepsilon > 0$. The construction of η and Ψ^{tr} respects the symmetry considerations in Section 3.2 (since $\eta \circ S_\delta = \eta$). Notice that on T^+ ,

$$\Psi^{\text{tr}}(\mathbf{x}) = \eta_+(x_1) \Psi(\mathbf{x}).$$

Recall that our main (small) parameter is $\delta \in (0, \pi)$. We have also introduced another small parameter γ with $0 < \delta < \gamma$ and the large parameter ℓ in the definition of Ψ . Below we will need the condition that

$$(3.12) \quad \ell \delta \rightarrow 0.$$

We will make the choices

$$(3.13) \quad \gamma = \delta^{\frac{1}{2}}, \quad \ell = \delta^{-\frac{1}{2}},$$

the first one ensuring that $\gamma^{-1} \delta^2 = \gamma^3$ (see the remainders in Proposition 3.4) and the second one being rather arbitrary. The function η_+ will be chosen to depend on δ at the very end of the proof.

3.5.1. *Estimates of the L^2 -norm and of the energy of Ψ^{tr} .* Let us estimate the L^2 -norm of our test function Ψ^{tr} . For small enough δ , we have by (2.9), and using (3.12) in the second term,

$$\begin{aligned} \int_{T^+} |\Psi^{\text{tr}}|^2 d\mathbf{x} &= \int_{(0,+\infty) \times (0,\ell)} \eta_+(x_1)^2 |f_\ell(x_2)|^2 d\mathbf{x} - \int_{\frac{\pi-\delta}{2}}^{\frac{\pi}{2}} \int_0^{\frac{\ell}{\sin\theta}} |f(r \sin\theta)|^2 r dr d\theta \\ &= (1 + \mathcal{O}(\ell^{-\infty})) \|\eta_+\|_{L^2(\mathbf{R}_+)}^2 - \int_{\frac{\pi-\delta}{2}}^{\frac{\pi}{2}} \sin^{-2}\theta \int_0^\ell |f(t)|^2 t dt d\theta. \end{aligned}$$

We deduce that

$$(3.14) \quad \int_{T^+} |\Psi^{\text{tr}}|^2 d\mathbf{x} = (1 + \mathcal{O}(\ell^{-\infty})) \|\eta_+\|_{L^2(\mathbf{R}_+)}^2 - \frac{\delta}{2} \int_0^{+\infty} t |f(t)|^2 dt + \mathcal{O}(\delta^3).$$

Let us now estimate the energy of Ψ^{tr} . An integration by parts and a symmetry consideration yield the following identity ($\|\cdot\|$ denotes the norm in $L^2(\mathcal{C}_\delta)$):

$$\begin{aligned} Q_\delta(\Psi^{\text{tr}}) &= \int_{\mathcal{C}_\delta} |\eta|^2 |(-i\nabla + \mathbf{A})\Psi|^2 d\mathbf{x} - \int_{\mathcal{C}_\delta} \eta \Delta \eta |\Psi|^2 d\mathbf{x} \\ &= \|\eta(-i\nabla + \mathbf{A})\Psi\|^2 - 2 \int_{T^+} \eta \Delta \eta |\Psi|^2 d\mathbf{x} \\ &= \|\eta(-i\nabla + \mathbf{A})\Psi\|^2 - 2 \int_{T^+} \eta(x_1) \eta''(x_1) |\Psi|^2 d\mathbf{x} \\ &= \|\eta(-i\nabla + \mathbf{A})\Psi\|^2 + 2 \|f\|_{L^2(\mathbf{R}_+)}^2 \int_{\mathbf{R}_+} |\eta'_+(x_1)|^2 dx_1. \end{aligned}$$

Notice, that for the integration by parts we needed $\eta_+ \in H^2(\mathbf{R}_+)$, but a density argument gives the identity for all $\eta_+ \in H^1(\mathbf{R}_+)$, with $\eta_+ = 1$ on the interval $(0, \varepsilon)$.

3.5.2. *Upper bound and proof of Theorem 1.1.* Using the symmetry of our construction with respect to D_δ , we get (as in Lemma 3.1):

$$\|\eta(-i\nabla + \mathbf{A})\Psi\|^2 - \Theta_0 \|\Psi^{\text{tr}}\|^2 = 2(\|\eta_+(-i\nabla + \mathbf{A})\Psi\|_{L^2(T^+)}^2 - \Theta_0 \|\Psi^{\text{tr}}\|_{L^2(T^+)}^2).$$

With (3.13), (3.14) and Proposition 3.4, we get

$$\begin{aligned} Q_\delta(\Psi^{\text{tr}}) - \Theta_0 \|\Psi^{\text{tr}}\|^2 &\leq -\delta \left(J - \Theta_0 \int_0^{+\infty} t |f(t)|^2 dt \right) + 2 \|\eta'_+\|_{L^2(\mathbf{R}_+)}^2 \\ &\quad + \mathcal{O}(\delta^{\frac{3}{2}}) + \mathcal{O}(\delta^\infty) \|\eta_+\|_{H^1(\mathbf{R}_+)}^2. \end{aligned}$$

Recalling (3.7) and (2.10), we get

$$J - \Theta_0 \int_0^{+\infty} t |f(t)|^2 dt = C_1 + \mathcal{O}(\delta^\infty),$$

so that

$$Q_\delta(\Psi^{\text{tr}}) - \Theta_0 \|\Psi^{\text{tr}}\|^2 \leq -C_1 \delta + 2 \|\eta'_+\|_{L^2(\mathbf{R}_+)}^2 + \mathcal{O}(\delta^{\frac{3}{2}}) + \mathcal{O}(\delta^\infty) \|\eta_+\|_{H^1(\mathbf{R}_+)}^2.$$

Therefore,

$$\frac{Q_\delta(\Psi^{\text{tr}})}{\|\Psi^{\text{tr}}\|^2} \leq \Theta_0 + \frac{1}{\|\Psi^{\text{tr}}\|^2} \left(-C_1 \delta + 2 \|\eta'_+\|_{L^2(\mathbf{R}_+)}^2 + \mathcal{O}(\delta^{\frac{3}{2}}) + \mathcal{O}(\delta^\infty) \|\eta_+\|_{H^1(\mathbf{R}_+)}^2 \right),$$

and by (3.14) and (2.4), we have

$$\frac{1}{\|\Psi^{\text{tr}}\|^2} = \frac{1}{2\|\eta_+\|^2} \left(1 + \frac{\xi_0\delta}{2\|\eta_+\|^2} + \mathcal{O}\left(\frac{\delta^2}{\|\eta_+\|^4}\right) + \mathcal{O}(\delta^3) \right),$$

where we used that $\|\eta_+\|^2 \geq \varepsilon$. Then,

$$(3.15) \quad \frac{Q_\delta(\Psi^{\text{tr}})}{\|\Psi^{\text{tr}}\|^2} \leq \Theta_0 + \mathcal{I}_\delta(\eta_+) + \mathcal{O}(\mathcal{R}_\delta(\eta_+)) + \mathcal{O}(\delta^\infty \|\eta_+\|_{H^1(\mathbf{R}_+)}^2),$$

where

$$(3.16) \quad \mathcal{I}_\delta(\eta_+) = \frac{1}{\|\eta_+\|^2} \left(-\frac{C_1\delta}{2} + \|\eta'_+\|_{L^2(\mathbf{R}_+)}^2 \right),$$

and

$$\mathcal{R}_\delta(\eta_+) = \frac{\delta^{\frac{3}{2}}}{\|\eta_+\|^2} + \delta \frac{\|\eta'_+\|^2}{\|\eta_+\|^4} + \delta^3 \frac{\|\eta'_+\|^2}{\|\eta_+\|^2}.$$

The estimate (3.15) leads us to minimize the functional \mathcal{I}_δ over the $\eta_+ \in H^1(\mathbf{R}_+)$ such that $\eta_+ = 1$ on $(0, \varepsilon)$, $\varepsilon > 0$ being fixed. For our purpose, namely to finish the proof of Theorem 1.1, it suffices to come up with a sufficiently good trial η_+ , so we will be brief. The Euler-Lagrange equation

$$-\eta_+'' = \mathcal{I}_\delta(\eta_+)\eta_+, \quad \text{on } [\varepsilon, +\infty),$$

leads us to consider test functions $\eta_+ = \eta_\alpha$ of the form

$$\eta_\alpha(x) = \begin{cases} 1 & \text{on } (0, \varepsilon) \\ e^{\alpha\varepsilon} e^{-\alpha x} & x \geq \varepsilon, \end{cases}$$

where $\alpha > 0$. We notice that

$$\mathcal{I}_\delta(\eta_\alpha) = \frac{-\frac{C_1\delta}{2} + \|\eta'_\alpha\|_{L^2(\varepsilon, +\infty)}^2}{\varepsilon + \|\eta_\alpha\|_{L^2(\varepsilon, +\infty)}^2} = \frac{\alpha^2 - d\alpha}{1 + 2\varepsilon\alpha}, \quad d = C_1\delta.$$

This last quantity is minimal for $\alpha_\delta = \frac{1}{2\varepsilon} (-1 + \sqrt{1 + 2\varepsilon d})$, and we have

$$\alpha_\delta \underset{\delta \rightarrow 0}{=} \frac{C_1\delta}{2} + \mathcal{O}(\delta^2).$$

Therefore, we choose $\alpha = \frac{C_1\delta}{2}$ and we have

$$\mathcal{I}_\delta(\eta_\alpha) = -\frac{C_1^2}{4}\delta^2 + \mathcal{O}(\delta^3).$$

We notice that

$$\|\eta_\alpha\|_{L^2(\mathbf{R}_+)}^2 = \varepsilon + \frac{1}{C_1\delta}, \quad \|\eta'_\alpha\|_{L^2(\mathbf{R}_+)}^2 = \frac{C_1\delta}{4},$$

so that

$$\mathcal{R}_\delta(\eta_\alpha) = \mathcal{O}(\delta^{\frac{5}{2}}).$$

With the upper bound (3.15), this concludes the proof of Theorem 1.1.

4. REGULAR PERTURBATION

The purpose of this section is to prove Theorem 1.2, so $\Omega_\delta = \Gamma_\delta^+$ hereafter. We construct a test function supported in a tubular neighborhood of $\partial\Omega_\delta$ where we can use the Frenet coordinates.

We choose $\ell = \delta^{-\rho}$ with $\rho \in (0, 1)$ and we let $B_\ell = \mathbf{R} \times (0, \ell)$. For δ small enough the classical tubular coordinates,

$$\Phi_\delta : B_\ell \ni (s, t) \mapsto \gamma_\delta(s) + t\mathbf{n}_\delta(s),$$

are well-defined in the sense that Φ_δ is injective and induces a local (and then global) \mathcal{C}^1 -diffeomorphism (the Jacobian of which being $1 - t\delta\kappa(s)$). Indeed, by using $\gamma'_0 = (1, 0)$, the continuity of $\delta \mapsto \gamma'_\delta$, and the Taylor formula, we can check that there exist $c, \delta_0 > 0$ such that, for all $\delta \in (0, \delta_0)$ and all $(s_1, s_2) \in \mathbb{R}^2$,

$$|\Phi_\delta(s_2, t_2) - \Phi_\delta(s_1, t_1)| \geq c(|s_2 - s_1| + |t_2 - t_1|).$$

We let $\Omega_{\delta, \ell} = \Phi_\delta(B_\ell)$. We consider a function of the form

$$(4.1) \quad \psi(s, t) = \zeta(\ell^{-1}t)f_\star(t)g(s),$$

where ζ is a cutoff function (see (2.8)) and where g has to be determined and will be chosen realvalued and normalised in $L^2(\mathbb{R})$. There exists a suitable phase φ such that (see [7, Lemma F.1.1]) if we let

$$\Psi = (e^{i\varphi}\psi) \circ \Phi_\delta^{-1},$$

which is supported in $\Omega_{\delta, \ell}$, we have

$$Q_\delta(\Psi) = \tilde{Q}_\delta(\psi),$$

where (with $\kappa = \kappa(s)$)

$$\begin{aligned} \tilde{Q}_\delta(\psi) &= \int_{B_\ell} (1 - t\delta\kappa)|\partial_t\psi|^2 ds dt \\ &\quad + \int_{B_\ell} (1 - t\delta\kappa)^{-1} \left| \left(-i\partial_s + \xi_0 - t + \frac{\delta\kappa t^2}{2} \right) \psi \right|^2 ds dt. \end{aligned}$$

By the exponential decay of f_\star , we have

$$(4.2) \quad \begin{aligned} \int_{B_\ell} (1 - t\delta\kappa)|\partial_t\psi|^2 ds dt &= \int_{B_\ell} (1 - t\delta\kappa)|g|^2|\ell^{-1}\zeta'(\ell^{-1}t)f_\star + \zeta(\ell^{-1}t)f'_\star|^2 ds dt \\ &\leq \|f'_\star\|^2\|g\|^2 - \delta \left(\int_0^\infty t|f'_\star|^2 dt \right) \int_{\mathbb{R}} \kappa|g|^2 ds + \mathcal{O}(\ell^{-\infty})\|g\|^2. \end{aligned}$$

Also,

$$(4.3) \quad \int_{B_\ell} (1 - t\delta\kappa)^{-1} \left| \left(-i\partial_s + \xi_0 - t + \frac{\delta\kappa t^2}{2} \right) \psi \right|^2 ds dt \leq I + \mathcal{O}(\ell^{-\infty})(\|g\|^2 + \|g'\|^2)$$

where, using first that g is real-valued and then that $(1 - t\delta\kappa)^{-1} = 1 + \delta\kappa t + \mathcal{O}(\delta^2 t^2 \kappa^2)$,

$$\begin{aligned}
I &= \int_{\mathbb{R} \times \mathbb{R}_+} (1 - t\delta\kappa)^{-1} \left| -ig'f_* + \left(\xi_0 - t + \frac{\delta\kappa t^2}{2}\right) f_* g \right|^2 ds dt \\
&= \int_{B_\ell} (1 - t\delta\kappa)^{-1} \left(|g'|^2 f_*^2 + |g|^2 \left(\xi_0 - t + \frac{\delta\kappa t^2}{2}\right)^2 f_*^2 \right) ds dt. \\
&\leq (1 + C\delta) \|g'\|^2 + \|(t - \xi_0) f_*\|^2 \|g\|^2 \\
(4.4) \quad &+ \left(\int_{\mathbb{R}} \delta\kappa |g|^2 ds \right) \int_0^{+\infty} ((\xi_0 - t)t^2 + t(t - \xi_0)^2) f_*^2 dt + C \int_{\mathbb{R}} \delta^2 \kappa^2 |g|^2 ds.
\end{aligned}$$

Combining this with (4.2) and (4.3), we deduce that

$$\begin{aligned}
\tilde{Q}_\delta(\psi) &\leq \Theta_0 \|g\|^2 + (1 + C\delta) \|g'\|^2 \\
&+ \left(\int_0^{+\infty} [((\xi_0 - t)t^2 + t(t - \xi_0)^2) f_*^2 - t f_*'^2] dt \right) \left(\int_{\mathbb{R}} \delta\kappa |g|^2 ds \right) \\
&+ C \int_{\mathbb{R}} \delta^2 \kappa^2 |g|^2 ds + \mathcal{O}(\delta^\infty) \|g\|_{H^1(\mathbf{R})}^2.
\end{aligned}$$

Using the decay of f_* and (2.4), the norm of ψ is given by

$$\begin{aligned}
\|\psi\|^2 &= \int_{B_\ell} (1 - \delta t\kappa) |g(s)|^2 |f_*(t)|^2 \zeta(\ell^{-1}t)^2 dt ds \\
(4.5) \quad &= \|g\|^2 - \delta \xi_0 \int_{\mathbf{R}_+} \kappa |g(s)|^2 ds + \mathcal{O}(\delta^\infty) \|g\|^2.
\end{aligned}$$

It follows that

$$\begin{aligned}
(4.6) \quad \tilde{Q}_\delta(\psi) - \Theta_0 \|\psi\|^2 &\leq (1 + \tilde{C}\delta) \|g'\|^2 \\
&+ \delta \int_0^{+\infty} \left(((\xi_0 - t)t^2 + t(t - \xi_0)^2) f_*^2 - t f_*'^2 + t\Theta_0 f_*^2 \right) dt \left(\int_{\mathbb{R}} \kappa |g|^2 ds \right) \\
&+ \mathcal{O}(\delta^2) \int_{\mathbf{R}} \kappa^2 |g(s)|^2 ds + \mathcal{O}(\delta^\infty) \|g\|^2.
\end{aligned}$$

We have to investigate the sign of

$$(1 + \tilde{C}\delta) \|g'\|^2 + A \int_{\mathbb{R}} \delta\kappa |g|^2 ds + \mathcal{O}(\delta^2) \int_{\mathbf{R}} \kappa^2 |g(s)|^2 ds + \mathcal{O}(\delta^\infty) \|g\|^2,$$

with

$$A = \int_0^{+\infty} \left((\xi_0 - t)t^2 + t(t - \xi_0)^2 \right) f_*^2 - t f_*'^2 + t\Theta_0 f_*^2 dt.$$

By using (2.5) and (2.6), we have

$$A = -\frac{3C_1}{2} + \int_0^{+\infty} t(t - \xi_0)(t - 2\xi_0) f_*^2(t) dt = -\frac{3C_1}{2} + \frac{C_1}{2} = -C_1.$$

We choose g such that $\|g\|^2 = 1$ and observe that (4.5) yields for δ small enough,

$$1 + \xi_0 \delta \int_{\mathbf{R}_+} \kappa |g(s)|^2 ds - \mathcal{O}(\delta^\infty) \leq \|\psi\|^{-2} \leq 1 + \mathcal{O}(\delta),$$

where the foregoing lower bound results from the fact that $\xi_0 > 0$, $\delta > 0$ and, for every natural number $N \geq 3$,

$$\|\psi\|^{-2} = 1 + \xi_0 \delta G + \underbrace{\xi_0^2 \delta^2 G^2 \left[1 + \sum_{j=1}^{N-2} (2\xi_0 \delta)^j G^j \right]}_{=1+\mathcal{O}(\delta) \geq 0} + \mathcal{O}(\delta^N), \quad G = \int_{\mathbf{R}_+} \kappa |g(s)|^2 ds.$$

By the min-max principle, we deduce from (4.6)

$$\lambda(\delta) \leq \Theta_0 + F_\delta(g) + \mathcal{O}(\delta^\infty),$$

where

$$(4.7) \quad F_\delta(g) = (1 + \hat{C}\delta) \|g'\|^2 + \delta \int_{\mathbf{R}} (-C_1 \kappa + \hat{C}\delta \kappa^2) |g(s)|^2 ds.$$

Minimizing over g and using the analysis in Appendix A, we get if $\int_{\mathbf{R}} \kappa(s) ds > 0$,

$$\lambda(\delta) \leq \Theta_0 - \frac{C_1^2}{4} \left(\int_{\mathbf{R}} \kappa(s) ds \right)^2 \delta^2 + \mathcal{O}(\delta^3).$$

More precisely, it is enough to consider the trial function

$$(4.8) \quad g(s) = \sqrt{\delta |\langle V \rangle|} \exp\left(\frac{\delta \langle V \rangle}{2} |s|\right), \quad V(s) = -C_1 \kappa(s), \quad \langle V \rangle = \int_{\mathbf{R}} V(s) ds < 0.$$

Notice that, by dominated convergence, $\int_{\mathbf{R}} \kappa^2 |g(s)|^2 ds = \mathcal{O}(\delta) \int_{\mathbf{R}} \kappa^2 ds$.

Acknowledgments. Part of this work was carried out while AK visited the University of Angers in February 2022. AK and NR are grateful to the support of the Fédération de Recherche Mathématiques des Pays de Loire and of the University of Angers. AK would like to thank B. Helffer for enlightening discussions.

APPENDIX A. WEAK PERTURBATION LIMIT

In this appendix, we explain where the choice (4.8) is coming from.

Let $V \in \mathcal{C}_0^0(\mathbf{R})$ such that $\langle V \rangle = \int_{\mathbf{R}} V(x) dx < 0$ and $\delta \in \mathbf{R}$. Let us consider the self-adjoint operator L_δ associated with the following quadratic form

$$q_\delta(\psi) = \|\psi'\|^2 + \delta \int_{\mathbf{R}} V(x) |\psi(x)|^2 dx, \quad \forall \psi \in H^1(\mathbf{R}),$$

which is the main term in (4.7), with $V = -C_1 \kappa$.

By the rescaling $x = \delta^{-1}y$, we see that L_δ is unitarily equivalent to $\delta^2 M_\delta$ where

$$M_\delta = -\partial_y^2 + \delta^{-1} V(\delta^{-1}y).$$

At a formal level, we see that M_δ converges to $M^{\text{eff}} := -\partial_y^2 + \langle V \rangle \delta_0$, whose spectrum is

$$\left\{ -\frac{\langle V \rangle^2}{4} \right\} \cup [0, +\infty),$$

and with groundstate $\exp(\frac{\langle V \rangle}{2} y)$.

Let us make this heuristics rigorous by comparing the quadratic forms. The quadratic form associated with M_δ is

$$p_\delta(\psi) = \|\psi'\|^2 + \int_{\mathbf{R}} \delta^{-1} V(\delta^{-1}y) |\psi|^2 dy, \quad \forall \psi \in H^1(\mathbf{R}).$$

We denote by p^{eff} the quadratic form associated with M^{eff} :

$$p^{\text{eff}}(\psi) = \|\psi'\|^2 + \langle V \rangle |\psi(0)|^2, \quad \forall \psi \in H^1(\mathbf{R}).$$

Let us estimate the difference $p_\delta - p^{\text{eff}}$. For all $\psi \in H^1(\mathbf{R})$, we have

$$\begin{aligned} |p_\delta(\psi) - p^{\text{eff}}(\psi)| &\leq \left| \int_{\mathbf{R}} \delta^{-1} V(\delta^{-1}y) |\psi(y)|^2 dy - \langle V \rangle |\psi(0)|^2 \right| \\ &\leq \left| \int_{\mathbf{R}} V(y) |\psi(\delta y)|^2 dy - \langle V \rangle |\psi(0)|^2 \right| \\ &\leq \int_{\mathbf{R}} |V(y)| \left| |\psi(\delta y)|^2 - |\psi(0)|^2 \right| dy. \end{aligned}$$

It is well-known that

$$\left| |\psi(\delta y)|^2 - |\psi(0)|^2 \right| \leq \delta^{\frac{1}{2}} \left((|\psi|^2)' \right) \sqrt{|y|} \leq 2\delta^{\frac{1}{2}} \|\psi\psi'\| \sqrt{|y|},$$

and also that, by the usual Sobolev embedding,

$$\|\psi\|_{L^\infty(\mathbf{R})} \leq C \|\psi\|_{H^1(\mathbf{R})}.$$

Thus,

$$\left| |\psi(\delta y)|^2 - |\psi(0)|^2 \right| \leq C\delta^{\frac{1}{2}} \|\psi\|_{H^1(\mathbf{R})}^2 \sqrt{|y|},$$

so that

$$|p_\delta(\psi) - p^{\text{eff}}(\psi)| \leq C\delta^{\frac{1}{2}} \|\psi\|_{H^1(\mathbf{R})}^2.$$

Therefore,

$$p_\delta^-(\psi) \leq p_\delta(\psi) \leq p_\delta^+(\psi),$$

where

$$\begin{aligned} p_\delta^-(\psi) &= (1 - C\delta^{\frac{1}{2}}) \|\psi'\|^2 + \langle V \rangle |\psi(0)|^2 - C\delta^{\frac{1}{2}} \|\psi\|^2, \\ p_\delta^+(\psi) &= (1 + C\delta^{\frac{1}{2}}) \|\psi'\|^2 + \langle V \rangle |\psi(0)|^2 + C\delta^{\frac{1}{2}} \|\psi\|^2. \end{aligned}$$

If $\nu(\delta)$ is the bottom of the spectrum of M_δ , we have

$$\nu(\delta) = -\frac{\langle V \rangle^2}{4} + \mathcal{O}(\delta^{\frac{1}{2}}) < 0.$$

REFERENCES

- [1] V. Bonnaillie. On the fundamental state energy for a Schrödinger operator with magnetic field in domains with corners. *Asymptot. Anal.*, 41(3-4):215–258, 2005.
- [2] V. Bonnaillie-Noël, F. Hérau, and N. Raymond. Purely magnetic tunneling effect in two dimensions. *Invent. Math.*, 227(2):745–793, 2022.
- [3] V. Bruneau, P. Miranda, and G. Raikov. Dirichlet and Neumann eigenvalues for half-plane magnetic Hamiltonians. *Rev. Math. Phys.*, 26(2):1450003, 23, 2014.
- [4] M. Correggi and E. L. Giacomelli. Almost flat angles in surface superconductivity. *Nonlinearity*, 34(11):7633–7661, 2021.
- [5] P. Exner, V. Lotoreichik, and A. Pérez-Obiol. On the bound states of magnetic Laplacians on wedges. *Rep. Math. Phys.*, 82(2):161–185, 2018.
- [6] S. Fournais and B. Helffer. Accurate eigenvalue asymptotics for the magnetic Neumann Laplacian. *Ann. Inst. Fourier (Grenoble)*, 56(1):1–67, 2006.
- [7] S. Fournais and B. Helffer. *Spectral methods in surface superconductivity*, volume 77 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 2010.
- [8] B. Helffer and A. Kachmar. Eigenvalues for the Robin Laplacian in domains with variable curvature. *Trans. Amer. Math. Soc.*, 369(5):3253–3287, 2017.

- [9] N. Raymond. *Bound states of the magnetic Schrödinger operator*, volume 27 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2017.
- [10] B. Simon. The bound state of weakly coupled Schrödinger operators in one and two dimensions. *Ann. Physics*, 97(2):279–288, 1976.

(V. Bonnaillie-Noël) DÉPARTEMENT DE MATHÉMATIQUES ET APPLICATIONS, ÉCOLE NORMALE SUPÉRIEURE, CNRS, UNIVERSITÉ PSL, 75005, PARIS, FRANCE
Email address: `Virginie.Bonnaillie@ens.fr`

(S. Fournais) DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY, NY MUNKEGADE 118, 8000 AARHUS C, DENMARK
Email address: `fournais@math.au.dk`

(A. Kachmar) LEBANESE UNIVERSITY, DEPARTMENT OF MATHEMATICS, NABATIYE, LEBANON
CENTER FOR ADVANCED MATHEMATICAL SCIENCES (CAMS, AMERICAN UNIVERSITY OF BEIRUT)
Email address: `akachmar@ul.edu.lb`

(N. Raymond) UNIV ANGERS, CNRS, LAREMA, SFR MATHSTIC, F-49000 ANGERS, FRANCE
Email address: `nicolas.raymond@univ-angers.fr`