



HAL
open science

W boson mass in minimal Dirac gaugino scenarios

Karim Benakli, Mark Goodsell, Wenqi Ke, Pietro Slavich

► **To cite this version:**

Karim Benakli, Mark Goodsell, Wenqi Ke, Pietro Slavich. W boson mass in minimal Dirac gaugino scenarios. *European Physical Journal C: Particles and Fields*, 2023, 83 (1), pp.43. 10.1140/epjc/s10052-022-11132-z . hal-03764322

HAL Id: hal-03764322

<https://hal.science/hal-03764322>

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



W boson mass in minimal Dirac gaugino scenarios

Karim Benakli^{1,a}, Mark Goodsell^{1,b}, Wenqi Ke^{1,c}, Pietro Slavich^{1,d}

¹ Sorbonne Université, CNRS, Laboratoire de Physique Théorique et Hautes Energies (LPTHE), 75005 Paris, France

Received: 23 September 2022 / Accepted: 13 December 2022 / Published online: 19 January 2023
© The Author(s) 2023

Abstract We investigate the conditions for alignment in Dirac Gaugino models with minimal matter content. This leads to several scenarios, including an aligned Dirac Gaugino NMSSM that allows a light singlet scalar. We then investigate the compatibility of minimal Dirac Gaugino models with an enhanced W boson mass, using a new precise computation of the quantum corrections included in the code SARAH 4.15.0.

Contents

1 Introduction	1
2 Dirac gaugino models with automatic tree-level alignment	2
2.1 Field content and interactions	2
2.2 The electroweak scalar sector and alignment	4
2.3 The electroweak fermion sector	7
2.4 Scenarios	8
3 W mass in Dirac gaugino models	8
3.1 Numerical setup	9
3.2 MSSM without μ term	10
3.3 W mass in the MDGSSM	11
3.4 W mass in the aligned MDGSSM	13
3.5 W mass in the aligned DGNMSSM	13
3.6 W mass in the general DGNMSSM	15
4 Conclusions	15
References	17

1 Introduction

While constraints on heavy Higgs bosons in supersymmetric models are rather stringent at large $\tan\beta$, excluding masses above a TeV, at small to moderate $\tan\beta$ direct searches do not place significant limits; only indirect constraints from $B \rightarrow s\gamma$ limit a heavy charged Higgs boson to be above 568 GeV, roughly independent of $\tan\beta$ [1]. On the other hand, in the Minimal Supersymmetric Standard Model (MSSM), this region is likely excluded for an additional neutral Higgs boson below a few hundred GeV due to modifications to the SM-like Higgs boson couplings. This has led to a lot of interest in extensions of the MSSM (or variants of the Two Higgs Doublet Model) where *alignment without decoupling* is possible [2–9], that is where the mixing between the SM-like Higgs boson and other scalars is minimised so that it aligns with the expectation values and has SM-like couplings.

Dirac Gaugino models [10–19], (see also, for example, [20–66]) accommodate scenarios where alignment without decoupling is *automatic* at tree-level [16, 67–71]. Under the assumption of an $N = 2$ supersymmetry in the gauge sector at some scale, these models contain an approximate $SU(2)_R$ R-symmetry which guarantees the tree-level alignment [68]. An investigation of the effects of quantum corrections showed that it is even radiatively stable [67], with competing effects partially cancelling.

These models have many interesting phenomenological properties, and have been extensively studied in the literature. They involve, at a minimum, an extension of the MSSM by three adjoint chiral superfields, one for each gauge group; the fermions from these pair with the gauginos to give them a Dirac mass. This means the presence of new scalar fields, in singlet, triplet and octet representations.

Actually, it has not adequately been investigated to what extent the *singlet* could be light in such models. One condition for this to be the case is that it should not disturb the couplings of the light Higgs – in other words, we should have

^a e-mail: kbenakli@lpthe.jussieu.fr

^b e-mail: goodsell@lpthe.jussieu.fr (corresponding author)

^c e-mail: wke@lpthe.jussieu.fr

^d e-mail: slavich@lpthe.jussieu.fr

some amount of alignments without decoupling. ATLAS and CMS both give constraints on the overall signal strength of the Higgs boson μ to be [72, 73]:

$$\mu = 1.06 \pm 0.07 \text{ (ATLAS)}, \quad \mu = 1.02_{-0.06}^{+0.07} \text{ (CMS)}. \quad (1.1)$$

If the Higgs boson h mixes with an inert singlet s , then we can write the mass eigenstates \tilde{h} , S as

$$\begin{pmatrix} h \\ s \end{pmatrix} = \begin{pmatrix} S_{11} & S_{12} \\ -S_{12} & S_{11} \end{pmatrix} \begin{pmatrix} \tilde{h} \\ S \end{pmatrix} \quad (1.2)$$

then we will find that

$$\mu = |S_{11}|^2 \leq 1. \quad (1.3)$$

Hence if we allow a 3σ deviation from the ATLAS result, we require

$$1 - |S_{11}|^2 = |S_{12}|^2 \leq 0.15 \longrightarrow |S_{12}| < 0.39. \quad (1.4)$$

While this still allows a moderate amount of mixing, the larger the mixing between the flavour eigenstates, the stronger the direct search bounds on the singlet will be. Therefore in this work we will consider the conditions for an approximate alignment in which the light Higgs mixes neither with the Heavy Higgs nor with the singlet (the triplet being decoupled). This will lead to a scenario that we refer to as the *aligned DGNMSSM*.

Recently, the CDF experiment reported a new measurement of the mass of the W boson [74]. Compared to the SM prediction [74–76], this gives as averages (combined Tevatron + LEP [74, 77–84]):

$$\begin{aligned} M_W^{\text{Tevatron+LEP}} &= 80,424 \pm 9 \text{ MeV}, \\ M_W^{\text{SM}} &= 80,356 \pm 6 \text{ MeV}. \end{aligned} \quad (1.5)$$

If we take the central value of the top quark mass to be 172.89 GeV then the central SM prediction becomes 80,352 MeV [85]. These differ by 7 standard deviations, although measurements at the LHC [86, 87] also differ from the combination of Tevatron + LEP by 4 standard deviations, so at this stage confirmation is required by other experiments. Nevertheless, a modification to the W boson mass is one of the most generic effects of new light particles coupling to the electroweak sector, so such a hint is tantalising.

It has generally been assumed in Dirac Gaugino models that the adjoint scalars should be heavy; indeed, the requirement that the triplet scalar vacuum expectation value (vev) must be very small compared to the Standard Model Higgs one – otherwise it would generate a large ρ parameter – is usually ensured by giving the triplet a heavy mass. Amusingly, following the new measurement, the simplest explanation for

the enhanced W boson mass is exactly an expectation value for the neutral component of such a triplet. In this work we shall investigate that possibility in minimal Dirac Gaugino models.

Such a triplet scalar also comes along with electroweak fermions, which can modify the quantum corrections too. Therefore a *precise* computation is required. While a preliminary such computation was performed for the MRSSM [88] using an update to FLEXIBLESUSY [89, 90], and a related computation was performed for the same model in [91], that model lacks a natural enhancement to the W boson quantum corrections. In this work we introduce a similarly precise computation in the package SARAH-4.15.0 and use it to examine the compatibility of our aligned DGNMSSM, along with four other scenarios – the “MSSM without μ term,” the MDGSSM, the aligned MDGSSM and the general DGNMSSM – with the new measurement of the W boson mass, or a naive world average value of $M_W^{\text{world average}} = 80,411 \pm 15$ MeV.

This work is organised as follows. In Sect. 2 we summarise the essential details of the class of Dirac Gaugino models, including the vacuum minimisation conditions and mass matrices. The conditions for alignment are reviewed and a comparison is made with the cases of the MSSM and NMSSM. We also introduce the different variants we shall consider: the MSSM without μ -term; the MDGSSM, the DGNMSSM and the aligned DGNMSSM. In Sect. 3 we will study the predictions for all of these classes of models for the W boson mass, examining in particular the effects of a precise computation of the quantum effects. We present our conclusions in Sect. 4.

2 Dirac gaugino models with automatic tree-level alignment

2.1 Field content and interactions

We shall consider in this work the extension of the MSSM by a minimal matter content to allow Dirac Gaugino masses, as in [19, 36]. The additional superfields consist of three chiral multiplets, in adjoint representations of the SM gauge group factors (DG-adjoints): a singlet \mathbf{S} , an $SU(2)_W$ triplet \mathbf{T}^a , and an $SU(3)_C$ octet \mathbf{O}^a . If we require gauge-coupling unification, even more states should be added to the model. For instance, for an $(SU(3))^3$ Grand Unification, the minimal set of chiral multiplets includes also extra Higgs-like doublets $\mathbf{R}_{u,d}$ as well as two pairs of vector-like right-handed electron $\mathbf{E}'_{1,2}$ in $(\mathbf{1}, \mathbf{1})_1$ and $\tilde{\mathbf{E}}'_{1,2}$ in $(\mathbf{1}, \mathbf{1})_{-1}$. We will not consider these states here.

In order to develop an intuition for the different interactions involved, it is helpful to consider a simple picture where the model descends from a supersymmetric theory in

D dimensions. The different states can appear in different sectors: some live in the whole D -dimensional bulk, others are localised on four-dimensional hyper-surfaces (branes) at points of the extra dimensions of coordinates $x_i = \{x_i^a\}$, $a = 5, \dots, D$. The corresponding Lagrangian can be written as

$$\int d^D x \mathcal{L} = \int d^D x \left\{ \mathcal{L}_{bulk} + \sum_i \delta^{(D-4)}(x - x_i) \mathcal{L}_{boundaries}^{(i)} \right\}, \tag{2.1}$$

where we have not explicitly written the metric factors. The four-dimensional theory arises after a truncation keeping only the compactification zero modes:

$$\mathcal{L}^{Ad} = \mathcal{L}_{bulk}^{Ad} + \mathcal{L}_{boundaries}, \tag{2.2}$$

$$\mathcal{L}_{bulk}^{Ad} = \int d^{D-4} x \mathcal{L}_{bulk}, \tag{2.3}$$

$$\mathcal{L}_{boundaries} = \int d^{D-4} x \sum_i \delta^{(D-4)}(x - x_i) \mathcal{L}_{boundaries}^{(i)}. \tag{2.4}$$

A tree-level alignment in the Higgs sector appears in a class of models where the bulk theory leads to a four-dimensional Lagrangian with interactions governed by an $N = 2$ extended SUSY. In particular, the SM gauge fields and the DG-adjoint fields arise as $N = 2$ vector supermultiplets, and the two Higgs chiral superfields \mathbf{H}_d and \mathbf{H}_u form an $N = 2$ hypermultiplet, interacting through the superpotential [92]

$$\mathcal{L}_{bulk}^{Ad} \supset \int d^2 \theta \{ \mu \mathbf{H}_u \cdot \mathbf{H}_d + \lambda_S \mathbf{S} \mathbf{H}_u \cdot \mathbf{H}_d + 2\lambda_T \mathbf{H}_d \cdot \mathbf{T} \mathbf{H}_u \}, \tag{2.5}$$

where $\mathbf{T} \equiv \frac{1}{2} \sigma^a \mathbf{T}^a$, and the dot product is defined as

$$\mathbf{H}_u \cdot \mathbf{H}_d \equiv \epsilon_{ij} \mathbf{H}_u^i \mathbf{H}_d^j = \mathbf{H}_u^+ \mathbf{H}_d^- - \mathbf{H}_u^0 \mathbf{H}_d^0. \tag{2.6}$$

The $N = 2$ SUSY has a global $SU(2)_R$ R-symmetry that rotates between the generators of the two $N = 1$ supercharges. The scalar components S, T^a of \mathbf{S} and \mathbf{T}^a , respectively, are singlets of $SU(2)_R$. This R-symmetry rotates then between the auxiliary fields F_Σ^a of the adjoint superfields $\Sigma^a \in \{\mathbf{S}, \mathbf{T}^a\}$ and the auxiliary component D^a of the corresponding chiral gauge superfields $\mathcal{W}_{i\alpha}^a$ for $U(1)_Y$ and $SU(2)_W$. This implies that

$$(\text{Re}(F_\Sigma^a) , D^a , \text{Im}(F_\Sigma^a)) \tag{2.7}$$

form a triplet of $SU(2)_R$. As a consequence, in order that the interactions (2.5) of \mathbf{S} and \mathbf{T}^a with the two Higgs doublets preserve $SU(2)_R$, the couplings λ_S and λ_T must be related

through¹

$$\lambda_S = \frac{g_Y}{\sqrt{2}}, \quad \text{and} \quad \lambda_T = \frac{g_2}{\sqrt{2}} \tag{2.8}$$

to the couplings g_Y and g_2 of the $U(1)_Y$ and $SU(2)_W$ gauge groups, respectively. Below the scale where the $N = 2$ SUSY is broken to $N = 1$, these relationships are spoiled by a small amount through renormalisation group running, so in numerical evaluations we must treat the couplings λ_S and λ_T as independent parameters.

In addition to the $SU(2)_R$ R-symmetry which is broken in $N = 1$ (chiral) sectors, there is a global $U(1)_R$ R-symmetry under which the superspace coordinates θ^α carry a -1 charge. The $U(1)_R$ charges of the \mathbf{H}_u and \mathbf{H}_d superfields are R_{H_u} and R_{H_d} , respectively. They are arbitrary but subject to the constraint $R_{H_u} + R_{H_d} = 2$. The DG-adjoint superfields \mathbf{S}, \mathbf{T}^a , and \mathbf{O}^a are R-neutral. Below, we shall classify the different $N = 1$ interactions following whether they preserve or break the $U(1)_R$ symmetry.

The boundary Lagrangian can be split into different contributions:

$$\mathcal{L}_{boundaries}^{Ad} = \mathcal{L}_{localised}^{bulk} + \int d^2 \theta \{ W_{Yukawa} + W_{DG} + W_{NR} \} + \Delta \mathcal{L}^{soft}. \tag{2.9}$$

Here, we denote by $\mathcal{L}_{localised}^{bulk}$ kinetic and interaction terms already present in the bulk theory \mathcal{L}_{bulk}^{Ad} but appearing with relative coefficients that violate $N = 2$ supersymmetry. Such terms can a priori be present because the boundary theory preserves only $N = 1$ SUSY, thus the coefficients of these terms are less constrained. Here, for simplicity, we assume such terms to vanish at tree level, to be only generated by quantum loops after supersymmetry breaking, and will therefore be accounted for in our analysis, at least in part, through the radiative corrections.

Also in (2.9), we have the usual MSSM Yukawa superpotential W_{Yukawa} with the couplings responsible for the quark and lepton masses:

$$W_{Yukawa} = Y_u^{ij} \mathbf{U}^c_i \mathbf{Q}_j \cdot \mathbf{H}_u - Y_d^{ij} \mathbf{D}^c_i \mathbf{Q}_j \cdot \mathbf{H}_d - Y_e^{ij} \mathbf{E}^c_i \mathbf{L}_j \cdot \mathbf{H}_d, \tag{2.10}$$

which arises on the brane where the matter field supermultiplets are localised.

In this work, we consider a typical scale for the soft terms, for example squarks or gaugino masses, to be in the phenomenologically interesting range $m_{soft} \sim \text{TeV}$. If we denote by Λ a higher scale, for instance related to supersymmetry breaking messenger mass scale or to the Planck

¹ In the discussion of the W boson mass, we shall relax this condition and study also generic Minimal Dirac Gaugino models with arbitrary values for λ_S and λ_T . All of the description of the models presented in this section holds for these models except for the $N = 2$ SUSY and $SU(2)_R$ global R-symmetry that are broken.

scale, then we can consider the relative strength of the diverse SUSY-breaking terms as an expansion in powers of $\frac{m_{soft}}{\Lambda}$. We will assume that SUSY-breaking terms in the gauge sector preserve the $U(1)_R$ R-symmetry, giving rise to Dirac gaugino masses, while Majorana masses might be generated only by higher-order interaction terms, therefore suppressed by additional powers of hidden-sector couplings and/or $\frac{m_{soft}}{\Lambda}$ where Λ could be the Planck scale (for gravity-induced effects). The effective superpotential for the Dirac gaugino masses reads

$$W_{DG} = \sqrt{2}\theta^\alpha \left[m_{DY} \mathbf{S} \mathcal{W}_{1\alpha} + 2m_{D2} \text{tr}(\mathbf{T} \mathcal{W}_{2\alpha}) \right. \tag{2.11}$$

$$\left. + 2m_{D3} \text{tr}(\mathbf{O} \mathcal{W}_{3\alpha}) \right], \tag{2.12}$$

where $\mathbf{O} \equiv \frac{1}{2} \lambda^a \mathbf{O}^a$, and $\mathcal{W}_{i\alpha}$ are the chiral gauge-strength superfields. Finally, the superpotential W_{NR} contains terms that break explicitly the $U(1)_R$ R-symmetry:

$$W_{NR} = \xi_S \mathbf{S} + \frac{M_S}{2} \mathbf{S}^2 + \frac{\kappa}{3} \mathbf{S}^3 + \lambda_{ST} \mathbf{S} \text{tr}(\mathbf{T}\mathbf{T}) + \lambda_{SO} \mathbf{S} \text{tr}(\mathbf{O}\mathbf{O}) + M_T \text{tr}(\mathbf{T}\mathbf{T}) + M_O \text{tr}(\mathbf{O}\mathbf{O}) + \frac{\lambda_O}{3} \text{tr}(\mathbf{O}\mathbf{O}\mathbf{O}). \tag{2.13}$$

The soft SUSY-breaking Lagrangian $\Delta\mathcal{L}^{soft}$ can in turn be split in two parts. The first contains the scalar mass and interaction terms that preserve the $U(1)_R$ R-symmetry:

$$-\Delta\mathcal{L}_R^{soft} = m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 + 2m_T^2 \text{tr}(T^\dagger T) + 2m_O^2 \text{tr}(O^\dagger O) + (m_Q^2)^{ij} Q_i^\dagger Q_j + (m_U^2)^{ij} U_i^{c\dagger} U_j^c + (m_D^2)^{ij} D_i^{c\dagger} D_j^c + (m_L^2)^{ij} L_i^\dagger L_j + (m_E^2)^{ij} E_i^{c\dagger} E_j^c + \left(t_S S + \frac{1}{2} B_S S^2 + \frac{1}{3} T_\kappa S^3 + T_{ST} S \text{tr}(TT) + T_{SO} S \text{tr}(OO) + B_T \text{tr}(TT) + B_O \text{tr}(OO) + \frac{1}{3} T_O \text{tr}(OOO) + \text{h.c.} \right). \tag{2.14}$$

The second part of $\Delta\mathcal{L}^{soft}$ contains the scalar mass and interaction terms that break the $U(1)_R$ R-symmetry:

$$-\Delta\mathcal{L}_{NR}^{soft} \supset B_\mu H_u \cdot H_d + T_S S H_u \cdot H_d + 2T_T H_d \cdot T H_u + T_u^{ij} U^c_i Q_j \cdot H_u - T_d^{ij} D^c_i Q_j \cdot H_d - T_e^{ij} E^c_i L_j \cdot H_d + \text{h.c.}, \tag{2.15}$$

as well as the Majorana mass terms M_i (with $i = 1, 2, 3$) for the gauginos. In general, the mechanisms that break R-symmetry and SUSY could be independent of each other, hence in Eqs. (2.14) and (2.15) we refrained from defining the soft SUSY-breaking trilinear couplings as proportional to the corresponding superpotential couplings. In this work

we assume that the soft SUSY-breaking Higgs-sfermion-sfermion interactions in the second line of Eq. (2.15) are suppressed with respect to the R-conserving sfermion mass terms in Eq. (2.14). This can be realised in our D -dimensional picture if the quark and lepton superfields are localised on a brane that differs from the one where the breaking of the R-symmetry takes place.

Since our study focuses on the electroweak sector of Dirac Gaugino models, we assume for simplicity that the scalar octet O^a is heavy and can be integrated out of the theory. To insulate the singlet sector from threshold corrections involving the heavy octet, we also neglect the singlet-octet interaction term proportional to λ_{SO} in the R-violating part of the superpotential, Eq. (2.13), as well as the analogous term proportional to T_{SO} in the R-conserving part of the soft SUSY-breaking Lagrangian, Eq. (2.14). Similarly, since they cannot appear in some of our scenarios, for simplicity in the following we shall also neglect λ_{ST} , T_{ST} and the tadpole terms t_S , ξ_S .

2.2 The electroweak scalar sector and alignment

We can now discuss the neutral scalar sector of this class of models. The vacuum expectation values² (vevs) of the neutral components of the doublets H_d and H_u are related by $v_u^2 + v_d^2 = v^2$, where $v \simeq 246$ GeV is the electroweak scale, and we define $\tan\beta = v_u/v_d$. The neutral singlet and triplet scalars S and T^0 obtain vevs $\langle S \rangle = v_S/\sqrt{2}$ and $\langle T^0 \rangle = v_T/\sqrt{2}$, respectively. These lead to effective μ and B_μ parameters:

$$\mu_{\text{eff}} \equiv \mu + \frac{1}{\sqrt{2}}(\lambda_S v_S + \lambda_T v_T), \tag{2.16}$$

$$B_{\mu,\text{eff}} \equiv B_\mu + \frac{1}{\sqrt{2}}(\lambda_S M_S + T_S) v_S + \frac{1}{\sqrt{2}}(\lambda_T M_T + T_T) v_T + \frac{1}{2} \lambda_S \kappa v_S^2. \tag{2.17}$$

The vevs v_S and v_T are then determined as a solution for the coupled cubic equations:

$$\begin{aligned} &\kappa^2 v_S^3 + \frac{1}{\sqrt{2}}(T_\kappa + 3\kappa M_S) v_S^2 + \tilde{m}_{SR}^2 v_S \\ &+ \frac{v^2}{2} \left[\sqrt{2} \lambda_S \mu_{\text{eff}} - g_Y m_{DY} c_{2\beta} \right. \\ &\left. - \left(\frac{1}{\sqrt{2}} T_S + \frac{1}{\sqrt{2}} \lambda_S M_S + \lambda_S v_S \kappa \right) s_{2\beta} \right] = 0, \tag{2.18} \\ &\tilde{m}_{TR}^2 v_T + \frac{v^2}{2} \left[\sqrt{2} \lambda_T \mu_{\text{eff}} \right. \end{aligned}$$

² It should be emphasised that, throughout this work, we assume that CP symmetry is not spontaneously broken by the vacuum. Therefore, all vevs are real.

$$+g_2m_{D2}c_{2\beta} - \frac{1}{\sqrt{2}}(T_T + \lambda_T M_T)s_{2\beta}] = 0, \tag{2.19}$$

where

$$\begin{aligned} \tilde{m}_{SR}^2 &\equiv M_S^2 + m_S^2 + B_S + 4m_{DY}^2, \\ \tilde{m}_{TR}^2 &\equiv M_T^2 + m_T^2 + B_T + 4m_{D2}^2, \end{aligned} \tag{2.20}$$

are effective mass-squared parameters (at zero expectation value) for the real components S_R and T_R^0 of the neutral singlet and triplet scalars (the analogous masses for the imaginary components S_I and T_I^0 are $\tilde{m}_{SI}^2 = M_S^2 + m_S^2 - B_S$ and $\tilde{m}_{TI}^2 = M_T^2 + m_T^2 - B_T$, respectively). We know that v_T must be small – namely, less than a few GeV – to avoid an overlage tree-level $\Delta\rho$, so to a good approximation we can set $v_T = 0$ in the vacuum minimisation equation for v_S , Eq. (2.18), and decouple it from the one for v_T , Eq. (2.19); this would allow the cubic equation for v_S to be solved using standard techniques. However, the current state of technology for the computation of loop corrections assumes that we take expectation values as being valid for the true minimum of the full quantum-corrected potential, so in our numerical studies we must take them as *inputs*. Especially for v_T this can lead to complications; see [93] for a recent discussion of this issue.

To discuss the alignment in the Higgs sector, it is now convenient to introduce the so-called Higgs basis for the two doublets,

$$\Phi_1 \equiv \frac{v_d\Phi_d + v_u\Phi_u}{v}, \quad \Phi_2 \equiv \frac{-v_u\Phi_d + v_d\Phi_u}{v}, \tag{2.21}$$

where we defined for convenience two doublets with positive hypercharge, $\Phi_u^j \equiv H_u^j$ and $\Phi_d^j \equiv -\epsilon_{ij}H_d^{*j}$. In the Higgs basis the two doublets can be decomposed as

$$\Phi_1 = \begin{pmatrix} G^+ \\ (v+h+iG^0)/\sqrt{2} \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} H^+ \\ (H+iA)/\sqrt{2} \end{pmatrix}, \tag{2.22}$$

i.e., only the neutral component of Φ_1 has a non-zero vev, and the would-be-Goldstone bosons, G^\pm and G^0 , all lie in Φ_1 . In general, the neutral CP-even fields h and H mix with the neutral CP-even components of the singlet and the triplet. In the basis $\{h, H, S_R, T_R^0\}$, the tree-level mass matrix reads

$$(\mathcal{M}^2)_{\text{tree}} = \begin{pmatrix} M_Z^2 + \Delta_h s_{2\beta}^2 & \Delta_h s_{2\beta} c_{2\beta} & \Delta_{hs} & \Delta_{ht} \\ \Delta_h s_{2\beta} c_{2\beta} & M_A^2 - \Delta_h s_{2\beta}^2 & \Delta_{Hs} & \Delta_{Ht} \\ \Delta_{hs} & \Delta_{Hs} & \tilde{m}_S^2 & \lambda_S \lambda_T \frac{v^2}{2} \\ \Delta_{ht} & \Delta_{Ht} & \lambda_S \lambda_T \frac{v^2}{2} & \tilde{m}_T^2 \end{pmatrix}, \tag{2.23}$$

where

$$\begin{aligned} \Delta_h &= \frac{v^2}{2}(\lambda_S^2 + \lambda_T^2) - M_Z^2, \\ \Delta_{hs} &= v \left[\sqrt{2}\lambda_S\mu_{\text{eff}} - g_Y m_{DY} c_{2\beta} - \left(\frac{1}{\sqrt{2}}(T_S + \lambda_S M_S) \right. \right. \end{aligned} \tag{2.24}$$

$$\left. \left. + v_S \kappa \lambda_S \right) s_{2\beta} \right] \tag{2.25}$$

$$\begin{aligned} \Delta_{Hs} &= v \left[- \left(\frac{1}{\sqrt{2}}(T_S + \lambda_S M_S) + v_S \kappa \lambda_S \right) c_{2\beta} \right. \\ &\quad \left. + g_Y m_{DY} s_{2\beta} \right], \end{aligned} \tag{2.26}$$

$$\Delta_{ht} = v \left[\sqrt{2}\lambda_T\mu_{\text{eff}} + g_2m_{D2}c_{2\beta} - \frac{1}{\sqrt{2}}(T_T + \lambda_T M_T)s_{2\beta} \right], \tag{2.27}$$

$$\Delta_{Ht} = -v \left[\frac{1}{\sqrt{2}}(T_T + \lambda_T M_T)c_{2\beta} + g_2m_{D2}s_{2\beta} \right], \tag{2.28}$$

$$M_A^2 = \frac{2B_{\mu,\text{eff}}}{s_{2\beta}}, \tag{2.29}$$

$$\begin{aligned} \tilde{m}_S^2 &= \tilde{m}_{SR}^2 + \lambda_S^2 \frac{v^2}{2} - \kappa \lambda_S \frac{v^2}{2} s_{2\beta} \\ &\quad + 3\kappa^2 v_S^2 + \sqrt{2}v_S(T_\kappa + 3\kappa M_S), \end{aligned} \tag{2.30}$$

$$\tilde{m}_T^2 = \tilde{m}_{TR}^2 + \lambda_T^2 \frac{v^2}{2}. \tag{2.31}$$

Exact alignment in the Higgs sector is obtained when one of the eigenstates of the CP-even mass matrix – in this work, we take it to be the lightest one – is aligned in field space with the direction of the SM Higgs vev, and thus has SM-like couplings to gauge bosons and matter fermions. This is equivalent to requiring that h itself be an eigenstate of \mathcal{M}^2 , or in other words that $\mathcal{M}_{1j}^2 = 0$ with $j = 2, 3, 4$. If we make the reasonable assumption that the triplet is heavy, then this can be relaxed to just $j = 2, 3$. In addition, we will also refer in this work to cases where the singlet can be light without being potentially ruled out by direct searches. In this case we will require the supplementary condition that $\mathcal{M}_{23}^2 = 0$.

We start our discussion by focusing on the alignment between the two doublets. The use in Eq. (2.24) of the $N = 2$ condition for the singlet and triplet superpotential couplings, see Eq. (2.8), implies $\Delta_h = 0$ and $M_h = M_Z$, i.e. alignment is automatically realised at the tree level in this class of models, and the tree-level mass of the SM-like Higgs boson is independent of $\tan\beta$ but well below the value observed at the LHC. It is however well known that, in SUSY models, the radiative corrections to the Higgs mass matrix play a crucial role in lifting the prediction for the mass of the SM-like Higgs boson up to the observed value. Moreover, the radiative corrections to the condition in Eq. (2.8) for the superpotential couplings of the adjoint superfields can become relevant if the scale $M_{N=2}$ where the $N = 2$ SUSY is broken to $N = 1$ is much larger than the scale where the Higgs mass matrix is computed. All of these corrections inevitably affect also the condition for alignment in the Higgs sector. As was discussed in Ref. [67], in DG models the element that mixes the two doublets in the loop-corrected mass matrix can be recast as

$$\mathcal{M}_{12}^2 = \frac{1}{\tan\beta} \left[\mathcal{M}_{11}^2 - M_Z^2 \right] - \frac{v^2 \tan\beta}{1 + \tan^2\beta} \left[\left(\lambda_S^2 - \frac{g'^2}{2} \right) \right]$$

$$+ \left(\lambda_T^2 - \frac{g^2}{2} \right) \Big] + (\dots), \tag{2.32}$$

where \mathcal{M}_{11}^2 contains the dominant one-loop contribution from top and stops. The latter consists in a term enhanced by $y_t^4 \ln(M_{\text{SUSY}}^2/m_t^2)$, where y_t is the top Yukawa coupling and M_{SUSY} denotes for simplicity a common soft SUSY-breaking mass parameter for the stops. The second term in Eq. (2.32) accounts for the deviation of the superpotential couplings from the $SU(2)_R$ condition in Eq. (2.8). The ellipses denote one-loop top/stop contributions that are suppressed by small ratios of parameters – such as T_u^{33}/M_{SUSY} , see the comments after Eq. (2.15), or m_t^2/M_{SUSY}^2 – one-loop contributions that involve couplings other than y_t , and higher-loop contributions. Close to alignment, the loop-corrected mass-matrix element \mathcal{M}_{11}^2 can be empirically identified with the observed mass of the SM-like Higgs boson, $M_h^2 \approx 2 M_Z^2$. Therefore, Eq. (2.32) shows that the radiative corrections included in \mathcal{M}_{11}^2 tend to destroy the tree-level alignment in the Higgs sector of DG models. However, when $M_{N=2}$ is large the evolution of λ_S and λ_T down to the scale where the Higgs mass matrix is computed makes the second term in Eq. (2.32) negative, and partially compensates for the misalignment induced by the top/stop contributions.

It is instructive to compare the condition for doublet alignment in DG models with the analogous conditions in the MSSM and in the NMSSM. In the case of the MSSM, discussed e.g. in Refs. [6,94], one finds

$$\begin{aligned} \mathcal{M}_{12}^2 = & \frac{1}{\tan \beta} \left[\mathcal{M}_{11}^2 - M_Z^2 c_{2\beta} \right] \\ & + \frac{6 y_t^2 m_t^2 \mu A_t}{16\pi^2 M_{\text{SUSY}}^2} \left(1 - \frac{A_t^2}{6 M_{\text{SUSY}}^2} \right) + (\dots), \end{aligned} \tag{2.33}$$

where $A_t \equiv T_u^{33}/y_t$ is the soft SUSY-breaking Higgs-stop-stop interaction parameter, and again the ellipses denote subdominant terms. It appears that, in the MSSM, the alignment condition $\mathcal{M}_{12}^2 = 0$ can be realised radiatively when a large value of $\tan \beta$ suppresses the first term in Eq. (2.33), while the parameters M_{SUSY} , A_t and μ combine in such a way that the second term is large and negative. In contrast, in DG models the contributions to \mathcal{M}_{12}^2 analogous to the second term in Eq. (2.33) are suppressed by the assumption that $A_t \ll M_{\text{SUSY}}$, thus doublet alignment cannot be realised in this way. We remark however that, even with the $N = 2$ condition for the superpotential couplings, in DG models \mathcal{M}_{12}^2 is smaller by a factor of approximately $(2 - c_{2\beta})$ – i.e., between 2 and 3, depending on $\tan \beta$ which we assume to be greater than 1 – with respect to the case of the MSSM with small A_t .

In the case of the NMSSM, discussed e.g. in Refs. [7,9], the mixing between h and H is given by

$$\mathcal{M}_{12}^2 = \frac{1}{\tan \beta} \left[\mathcal{M}_{11}^2 - M_Z^2 c_{2\beta} - \lambda_S^2 v^2 s_\beta^2 \right]$$

$$+ \frac{6 y_t^2 m_t^2 \tilde{\mu} A_t}{16\pi^2 M_{\text{SUSY}}^2} \left(1 - \frac{A_t^2}{6 M_{\text{SUSY}}^2} \right) + (\dots), \tag{2.34}$$

where $\tilde{\mu} \equiv \mu + \lambda_S v_S/\sqrt{2}$. Comparing with the case of the MSSM, Eq. (2.33), we see that the condition $\mathcal{M}_{12}^2 = 0$ can be realised even in the absence of a large contribution from the terms proportional to $\mu_{\text{eff}} A_t/M_{\text{SUSY}}^2$, as long as the singlet-doublet superpotential coupling takes values in the range $\lambda_S^2 \approx (3-4) M_Z^2/v^2$, where the exact numerical coefficient depends on the value of $\tan \beta$. As first pointed out in Ref. [7], this condition singles out the region of the NMSSM parameter space where $\lambda_S \approx 0.7 \pm 0.05$, a much larger value than would be implied by the $SU(2)_R$ condition in DG models.

To summarise, the $SU(2)_R$ R-symmetry implies exact alignment at the tree level in the Higgs-doublet sector of the DG models, but the alignment is partially spoiled by the radiative corrections that are necessary to obtain a realistic value for the SM-like mass. Alignment in the MSSM can be realised only through radiative corrections, for large $\tan \beta$ and for specific choices of the parameters in the stop sector. Finally, doublet alignment in the NMSSM can be realised even without the help of radiative corrections for an appropriate choice of λ_S , which – differently from the DG case with $SU(2)_R$ R-symmetry – is treated as a free parameter.

The second condition for Higgs alignment in DG models is $\mathcal{M}_{13}^2 = 0$, i.e. vanishing mixing between h and S_R . Including the dominant contributions from stop loops, we find:

$$\mathcal{M}_{13}^2 = \Delta_{hs} - \frac{6 y_t \lambda_S c_\beta}{16\pi^2} m_t (A_t - \mu_{\text{eff}} \cot \beta) \ln \frac{M_{\text{SUSY}}^2}{Q^2} + (\dots), \tag{2.35}$$

where the tree-level mixing term Δ_{hs} is given in Eq. (2.25), μ_{eff} is given in Eq. (2.16), and Q is the renormalisation scale at which the parameters entering Δ_{hs} are expressed. We assumed again a common soft SUSY-breaking mass term M_{SUSY} for the stops, and we neglected terms suppressed by powers of m_t^2/M_{SUSY}^2 . The various terms that contribute to Δ_{hs} arise from different sectors of the D -dimensional picture discussed earlier in this section: namely, μ and λ_S enter the bulk superpotential in Eq. (2.5); m_{1D} enters the R-conserving boundary superpotential in Eq. (2.12); M_S and κ enter the R-violating boundary superpotential in Eq. (2.13); T_S enters the R-violating SUSY-breaking Lagrangian in Eq. (2.15). Therefore, even if we assume the $N = 2$ SUSY relation of Eq. (2.8) between λ_S and g_Y , a vanishing \mathcal{M}_{13}^2 can only result from an accidental cancellation between unrelated terms. We also note that, in contrast to the case of \mathcal{M}_{12}^2 , the radiative correction to \mathcal{M}_{13}^2 is not enhanced by $\tan \beta$ with respect to the tree-level part. Thus, its qualitative impact on our discussion of the alignment conditions is limited, as long as the scale Q is not too far from M_{SUSY} .

The minimum conditions of the scalar potential can be exploited to express the mass parameters for the doublets and the singlet in terms of the other Lagrangian parameters and of the vevs v_d, v_u and v_S . In particular, we obtain a relation between the mass parameter $\tilde{m}_{S_R}^2$ for the real component of the singlet, see Eq. (2.20), and the matrix element \mathcal{M}_{13}^2 given in Eq. (2.35):

$$\tilde{m}_{S_R}^2 = -\frac{v}{2v_S} \mathcal{M}_{13}^2 - \frac{v_S}{\sqrt{2}} \left(T_\kappa + \sqrt{2}\kappa^2 v_S + 3\kappa M_S \right). \tag{2.36}$$

Equation (2.36) above shows that the condition of vanishing mixing between h and S_R , however realised, carries implications for the mass of the singlet. The diagonal element for the singlet in the scalar mass matrix is

$$\begin{aligned} \mathcal{M}_{33}^2 = & \tilde{m}_{S_R}^2 + \frac{1}{2}\lambda_S^2 v^2 + \sqrt{2}T_\kappa v_S \\ & + \kappa \left(3\kappa v_S^2 + 3\sqrt{2}M_S v_S - \lambda_{SS\beta} c_\beta v^2 \right) \\ & + \frac{3y_t^2 \lambda_S^2 c_\beta^2}{32\pi^2} v^2 \ln \frac{M_{\text{SUSY}}^2}{Q^2} + (\dots), \end{aligned} \tag{2.37}$$

where we applied the same approximations as in Eq. (2.35) for the one-loop correction in the second line.

We now discuss the simplest case in which the global $U(1)_R$ R-symmetry is preserved in the superpotential but broken by soft SUSY-breaking terms, in which case we can set κ to zero. Since this implies a vanishing quartic self-coupling for the singlet, the stability of the scalar potential requires that we also assume $T_\kappa = 0$, even if the trilinear self-coupling of the singlet resides in the R-conserving part of the soft SUSY-breaking Lagrangian. In this scenario, which we shall refer to as the *aligned MDGSSM*, Eq. (2.36) shows that the alignment condition $\mathcal{M}_{13}^2 \approx 0$ requires $\tilde{m}_{S_R}^2 \approx 0$ or $v_S \ll v$. The first of these two options implies that the CP-even mass eigenstate that is mostly singlet is relatively light: setting $\tilde{m}_{S_R}^2 = 0$ and $\kappa = T_\kappa = 0$ in Eq. (2.37), and neglecting the small effect of the one-loop correction, we find that the value $\lambda_S \approx 0.7$ favored by the alignment condition for the Higgs doublets in the NMSSM, see Eq. (2.35), leads to $\mathcal{M}_{33}^2 \approx (122 \text{ GeV})^2$, whereas the value $\lambda_S \approx 0.25$ implied in our Dirac-gaugino model by the $N = 2$ SUSY relation of Eq. (2.8) leads to $\mathcal{M}_{33}^2 \approx (44 \text{ GeV})^2$. We remark that the mixing between S_R and the heavier, non-SM-like scalar H , which is controlled by \mathcal{M}_{23}^2 , is suppressed when $M_A^2 \gg \mathcal{M}_{33}^2$, and would in any case lower the mass of the singlet-like eigenstate.

The definition in Eq. (2.20) shows that even the vanishing of $\tilde{m}_{S_R}^2$ requires a cancellation between terms that arise from different sectors of our D -dimensional construction: m_{1D} from the R-conserving boundary superpotential in Eq. (2.12), M_S from the R-violating boundary superpotential in Eq. (2.13), m_S^2 and B_S from the R-conserving soft SUSY-breaking Lagrangian in Eq. (2.14). If such cancellation is not realised, the alternative requirement for alignment implied by Eq. (2.36) when $\kappa = 0$ is that $v_S \ll v$. This is not problematic as long as a suitable higgsino mass is provided by the μ term in the bulk superpotential, see Eq. (2.5).

Finally, the third condition for Higgs alignment in DG models is $\mathcal{M}_{14}^2 = 0$, i.e. vanishing mixing between h and T_R^0 . The formulas for the relevant mass-matrix elements and for the minimum condition, including the dominant one-loop corrections from top and stop loops, are similar to Eqs. (2.35)–(2.37), with the obvious singlet-to-triplet replacements but without terms analogous to those controlled by κ in the singlet case:

$$\mathcal{M}_{14}^2 = \Delta_{ht} - \frac{6y_t \lambda_T c_\beta}{16\pi^2} m_t (A_t - \mu_{\text{eff}} \cot \beta) \ln \frac{M_{\text{SUSY}}^2}{Q^2} + (\dots), \tag{2.38}$$

$$\mathcal{M}_{44}^2 = \tilde{m}_{T_R}^2 + \frac{1}{2}\lambda_T^2 v^2 + \frac{3y_t^2 \lambda_T^2 c_\beta^2}{32\pi^2} v^2 \ln \frac{M_{\text{SUSY}}^2}{Q^2} + (\dots), \tag{2.39}$$

$$\tilde{m}_{T_R}^2 = -\frac{v}{2v_T} \mathcal{M}_{14}^2. \tag{2.40}$$

The discussion of the constraints on the triplet mass induced by the condition of doublet-triplet alignment follows the lines of the discussion of singlet-triplet alignment for $\kappa = 0$, with the important difference that the condition $v_T \ll v$ must in any case be satisfied to avoid an excessive contribution to $\Delta\rho$. As a consequence, it is not necessary to require $\tilde{m}_{T_R}^2 \approx 0$ to obtain approximate alignment. Nevertheless, we remark that the condition of exact alignment $\mathcal{M}_{14}^2 = 0$ would imply $\mathcal{M}_{44}^2 \approx M_W^2$ through Eqs. (2.38)–(2.40).

2.3 The electroweak fermion sector

We now outline the mass spectrum of the electroweak fermions, which will be relevant for our discussion examining the W boson mass. The neutralino mass matrix, in the basis $\tilde{S}, \tilde{B}, \tilde{T}^0, \tilde{W}^0, \tilde{H}_d^0, \tilde{H}_u^0$ reads:

$$\begin{pmatrix}
 M_S + \sqrt{2}\kappa v_S & m_{DY} & 0 & 0 & -\frac{\sqrt{2}\lambda_S}{g_Y} M_{ZSWs\beta} & -\frac{\sqrt{2}\lambda_S}{g_Y} M_{ZSWc\beta} \\
 m_{DY} & M_1 & 0 & 0 & -M_{ZSWc\beta} & M_{ZSWs\beta} \\
 0 & 0 & M_T & m_{D2} & -\frac{\sqrt{2}\lambda_T}{g_2} M_{ZCWs\beta} & -\frac{\sqrt{2}\lambda_T}{g_2} M_{ZCWc\beta} \\
 0 & 0 & m_{D2} & M_2 & M_{ZCWc\beta} & -M_{ZCWs\beta} \\
 -\frac{\sqrt{2}\lambda_S}{g_Y} M_{ZSWs\beta} & -M_{ZSWc\beta} & -\frac{\sqrt{2}\lambda_T}{g_2} M_{ZCWs\beta} & M_{ZCWc\beta} & 0 & -\mu_{\text{eff}} \\
 -\frac{\sqrt{2}\lambda_S}{g_Y} M_{ZSWc\beta} & M_{ZSWs\beta} & -\frac{\sqrt{2}\lambda_T}{g_2} M_{ZCWc\beta} & -M_{ZCWs\beta} & -\mu_{\text{eff}} & 0
 \end{pmatrix} \tag{2.41}$$

The chargino masses, $-\frac{1}{2}((v^-)^T \mathcal{M}_{\chi^\pm} v^+ + h.c.)$ in the basis $v^+ = (\tilde{T}^+, \tilde{W}^+, \tilde{H}_u^+)$, $v^- = (\tilde{T}^-, \tilde{W}^-, \tilde{H}_d^-)$, are given by

$$\mathcal{M}_{\chi^\pm} = \begin{pmatrix}
 M_T & m_{D2} + g_2 v_T & \lambda_T v c_\beta \\
 m_{D2} - g_2 v_T & M_2 & \frac{g_2 v}{\sqrt{2}} s_\beta \\
 -\lambda_T v s_\beta & \frac{g_2 v}{\sqrt{2}} c_\beta & \mu_{\text{eff}} - \sqrt{2}\lambda_T v_T
 \end{pmatrix} \tag{2.42}$$

and do not depend on κ , but we have written for completeness the Majorana gaugino masses $M_{1,2}$ for the bino and wino respectively.

2.4 Scenarios

We have described the general features of minimal Dirac Gaugino models, and explained how some values of the couplings allow Higgs alignment to automatically occur at tree-level. Below, we shall consider the following different scenarios corresponding to specific choices of the model parameters:

- **General MDGSSM**

In the general MDGSSM, the only source of R -symmetry violation comes from a small B_μ term, which is a radiatively stable condition (the renormalisation group running will not generate other R -symmetry violating terms from a B_μ term). This excludes all of W_{NR} ; the only supersymmetric parameters beyond those of the MSSM retained are λ_S, λ_T , but those are allowed to have any value. Supersoft masses [14] are allowed, and soft supersymmetry-breaking masses are allowed, but squark/sfermion trilinears and T_S, T_T are not. On the other hand, trilinears involving the adjoint scalars may be allowed (so T_κ, T_{ST}, T_{SO}) but may be argued to be small in typical models. The couplings λ_S, λ_T enhance the Higgs mass and W mass at the same time; while large values of λ_T were previously deemed problematic for the ρ parameter, they are now a virtue.

- **MSSM without μ term (RIP)**

This model, described in [15], is identical to the general MDGSSM except that the μ -term is set to zero (although a small v_S generates a tiny effective μ). In the MSSM this

would yield massless higgsinos, but here the higgsinos obtain a mass through λ_T which causes them to mix with the triplet fermion. Thus the μ problem of the MSSM is solved, at the expense of an upper bound on the chargino masses, which, as we shall see, leads to the model being ruled out.

- **Aligned MDGSSM**

By taking λ_S, λ_T to their their $N = 2$ values given in Eq. (2.8) in the general MDGSSM we guarantee alignment with the heavy Higgs at tree level. If we further impose that $\Delta_{hs} = 0$, as described above, then the singlet also has negligible mixing with the SM-like Higgs. We shall refer to this scenario as the *aligned MDGSSM*.

- **DGNMSSM**

If we instead allow R -symmetry violation in the superpotential and the associated soft-breaking trilinears – in particular for κ, T_κ and T_S – we can generate μ and B_μ terms through a substantial expectation value for the singlet. The model thus resembles the NMSSM, especially if we set the μ, B_μ terms (and M_S, M_T) to zero; this was proposed in [19]. We shall therefore refer to this scenario as the DGNMSSM.

- **Aligned DGNMSSM**

We can achieve alignment in the DGNMSSM by setting λ_S, λ_T to their $N = 2$ values and taking $\Delta_{hs} = 0$. We shall refer to this scenario as the aligned DGNMSSM; in this work, when we consider the W boson mass, we shall also enforce $\Delta_{Hs} = 0$ which guarantees that the singlet couplings to SM fields are small, rendering light singlets safe from collider searches.

3 W mass in Dirac gaugino models

Dirac gaugino models offer two methods of explaining an enhancement of the W boson mass with respect to the SM: either quantum corrections or a tree-level expectation value for the triplet scalar. In the case of quantum corrections, there are new contributions to the W mass compared to the MSSM, again coming from interactions related to the adjoint triplet; in particular the coupling λ_T .

Recall the definition of the ρ parameter is $M_W^2 \equiv \rho c_W^2 M_Z^2$; through the presence of the adjoint triplet, DG mod-

els contain a tree-level modification to this relation compared to the SM:

$$\rho \equiv 1 + \Delta\rho_{\text{tree}} + \Delta\rho = 1 + 4\frac{v_T^2}{v^2} + \Delta\rho. \tag{3.1}$$

One could consider a modification to c_W instead of ρ as an explanation of an enhanced W mass, but this is discounted based on electroweak precision tests (see e.g. [95]); on the other hand, a triplet expectation value is one of the most generic and acceptable ways of enhancing the W mass at tree level. Naively we could then take the observed value of M_Z and the standard value of c_W and infer the value of $\Delta\rho$ to obtain a given value of M_W . However, in the SM and in any BSM theory it is necessary to take certain electroweak observables as input, and in our setup we will take the conventional choice of the Z mass, G_F and α . When we modify ρ this gives both a modification to M_W and a small modification to $\sin^2\theta_W$. At tree-level this is

$$\begin{aligned} \Delta_{\text{tree}} M_W^2 &= \frac{c_W^2}{c_W^2 - s_W^2} (M_W^2)_{\text{SM}} \Delta\rho_{\text{tree}}, \\ \Delta_{\text{tree}} s_W^2 &= -\frac{s_W^2 c_W^2}{c_W^2 - s_W^2} \Delta\rho_{\text{tree}}, \end{aligned} \tag{3.2}$$

so if we want to obtain $M_W = 80.424$ MeV we need $\Delta M_W^2 = 11$ (GeV)². Taking $s_W^2 = 0.23121$ this gives

$$\begin{aligned} \Delta_{\text{tree}}^{\text{CDF}} \rho_{\text{tree}} &= 0.0012 \\ \Delta_{\text{tree}}^{\text{CDF}} s_W^2 &= -4 \times 10^{-4}. \end{aligned} \tag{3.3}$$

Interpreted as a tree-level expectation value for the triplet, this yields

$$v_T \simeq 4 \text{ GeV}, \tag{3.4}$$

which was previously at the upper bound of what was acceptable.

In the minimal Dirac gaugino model, we have from equation (2.40)

$$\tilde{m}_{TR}^2 = -\frac{v}{2v_T} \Delta_{ht} = -\frac{\Delta_{ht}}{\sqrt{\Delta\rho_{\text{tree}}}}. \tag{3.5}$$

For a small v_T , if we do not tune $\Delta_{ht} \approx 0$, then triplets must be heavy. However, we also need heavy winos to evade collider bounds and this implies large m_{D2} : the connection between electroweakino masses and the expectation value of the triplet is a novel feature of this class of models. In [65] the conclusion was that for winos above 700 GeV there were essentially no constraints on the higgsinos beyond LEP. If

the contribution from m_{D2} dominates Δ_{ht} this implies

$$\begin{aligned} m_T^2 &\sim -\frac{g_2 v m_{D2} c_{2\beta}}{\sqrt{\Delta\rho_{\text{tree}}}} \sim (1.8 \text{ TeV})^2 \\ &\times \left(\frac{700 \text{ GeV}}{m_{D2}}\right) \times \left(\frac{c_{2\beta}}{-1}\right) \times \sqrt{\frac{0.0012}{\Delta\rho}}, \end{aligned} \tag{3.6}$$

which is a natural scale for supersymmetric scalars. Of course, we can have lighter winos provided that the neutralino is not too light, above around 200–300 GeV [65].

However, the model also contains ample room for quantum corrections to also enhance the W mass. In the following, we shall investigate this for the different scenarios described in Sect. 2.4.

3.1 Numerical setup

In order to accurately compute the quantum corrections to the W mass, we use a new EFT approach, closely related to that of [88], implemented in the spectrum-generator-generator SARAH. We use the expression:

$$\begin{aligned} M_W^2 &= (M_W^2)_{\text{SM}} \left(1 + \frac{s_W^2}{c_W^2 - s_W^2} \left[\frac{c_W^2}{s_W^2} (\Delta\rho_{\text{tree}} + \Delta\rho) \right. \right. \\ &\quad \left. \left. - \Delta r_W - \Delta\alpha \right] \right) \end{aligned} \tag{3.7}$$

where $(M_W^2)_{\text{SM}}$ is the full two-loop W mass in the SM, as computed in [76], depending on the pole masses of the Higgs boson, top quark, α_s and $\Delta\alpha_{\text{had}}^{(5)}$. We use the interpolating function from that paper. When we use the average values of the Higgs mass of 125.09 GeV and the top quark mass of 172.89 GeV this function gives us $M_W = 80.354$ GeV, just 2 MeV higher than the current world average for the W boson mass in the SM. The expressions computed in the square brackets are the differences between the high-energy theory (HET) and the SM:

$$\begin{aligned} \Delta\rho &\equiv \text{Re} \left[\frac{\Pi_{ZZ}^{\text{HET}}(M_Z^2)}{M_Z^2} - \frac{\Pi_{WW}^{\text{HET}}(M_W^2)}{M_W^2} \right] \\ &\quad - \text{Re} \left[\frac{\Pi_{ZZ}^{\text{SM}}(M_Z^2)}{M_Z^2} - \frac{\Pi_{WW}^{\text{SM}}(M_W^2)}{M_W^2} \right] \\ \Delta r_W &\equiv \left[\frac{\Pi_{WW}^{\text{HET}}(0)}{M_W^2} - \frac{\Pi_{WW}^{\text{HET}}(M_W^2)}{M_W^2} + \delta_{VB}^{\text{HET}} \right] \\ &\quad - \left[\frac{\Pi_{WW}^{\text{SM}}(0)}{M_W^2} - \frac{\Pi_{WW}^{\text{SM}}(M_W^2)}{M_W^2} + \delta_{VB}^{\text{SM}} \right]. \end{aligned} \tag{3.8}$$

$\Delta\alpha$ are now the gauge threshold corrections between the HET and the SM for the electromagnetic gauge coupling divided by α (so they do not now depend on $\Delta\alpha_{\text{had}}^{(5)}$).

In SARAH, we compute the expression in square brackets at the matching scale, which is the mass of the heavy particles. We are therefore ignoring the running from that scale down to the electroweak scale, which is of controllable size, but will nevertheless be included in a future development of the code. One could argue that we should instead perform the matching at the electroweak scale, but then all of the loop functions will contain large logarithms and there can be larger, spurious, running of the couplings of the high-energy theory which can spoil the results.

In order to be a strict one-loop matching between the HET and the SM, the weak mixing angle in the above must be the \overline{MS} or \overline{DR} value at the matching scale Q :

$$s_W^2 = \frac{g_Y^2(Q)}{g_2^2(Q) + g_Y^2(Q)}. \tag{3.9}$$

To extract this, we match the calculations of the Z-boson mass, $\alpha(Q)$ – and we also compute the decay of the muon at the matching scale. In practice, this means that we extract the couplings in the SM at the top mass scale without including the effects of new physics, then run them up to the matching scale. The threshold corrections to α between the two theories are simple to compute since it is unbroken and yield $\Delta\alpha$; the couplings $g_Y(Q)$, $g_2(Q)$ in the high-energy theory are chosen to solve the equation

$$c_W^2 s_W^2 = (c_W^2 s_W^2)_{SM} \times \frac{(1 + \Delta\alpha + \Delta\hat{r})}{1 + \Delta\rho_{tree}} \tag{3.10}$$

where

$$\Delta\hat{r} \equiv \left[(1 + \Delta\rho_{tree}) \frac{\Pi_{WW}^{HET}(0)}{M_W^2} - \frac{\Pi_{ZZ}^{HET}(M_Z^2)}{M_Z^2} + \delta_{VB}^{HET} \right] - \left[\frac{\Pi_{WW}^{SM}(0)}{M_W^2} - \frac{\Pi_{ZZ}^{SM}(M_Z^2)}{M_Z^2} + \delta_{VB}^{SM} \right], \tag{3.11}$$

which is done iteratively by progressively running up and down and updating at each step, along with all the other quantities in the high-energy theory. The value for $(c_W^2 s_W^2)_{SM}$ includes a compensatory term for corrections from \overline{MS} to \overline{DR} if needed.

In the SARAH model file, the couplings λ_S , λ_T , T_S , T_T are defined differently to the above: we have $1am = -\lambda_S$, $L T = \sqrt{2}\lambda_T$, $T[1am] = -T_S$, $T[L T] = \sqrt{2}T_T$. Hence in our plots and benchmark points we list the values in terms of $-\lambda_S$, $\sqrt{2}\lambda_T$, $-T_S$, which makes the correspondence with the numerical codes exact.

3.2 MSSM without μ term

The ‘‘MSSM without μ term’’ proposed in [15] was an intriguing solution to the μ problem. It was however chal-

lenged by the requirement of having a high enough Higgs mass, chargino mass and not too large ρ ; indeed the lack of intersection of points satisfying the latter two was demonstrated in [36]. It might therefore be tempting to revisit this model in light of the new data about the W mass. However, we shall demonstrate here that it is conclusively ruled out.

Putting aside the Higgs mass constraint, the see-saw effect on the charginos is a problem. LEP put a lower limit on the mass of the lightest chargino of 94 GeV [96–98]. In that model the chargino mass is

$$\mathcal{M}_{\chi^\pm} \xrightarrow{v_T \approx 0} \begin{pmatrix} 0 & m_{D2} & \frac{2\lambda_T}{g^2} M_{ZC} W c_\beta \\ m_{D2} & 0 & \sqrt{2} M_{ZC} W s_\beta \\ -\frac{2\lambda_T}{g} M_{ZC} W s_\beta & \sqrt{2} M_{ZC} W c_\beta & 0 \end{pmatrix} \tag{3.12}$$

It is known that it is possible to fulfil the LEP bound by a careful choice of λ_T and m_{D2} : a large value of λ_T as well as m_{D2} around 107 GeV is needed to maximize the mass of the lightest chargino. It would then be made of a higgsino-wino mixture with *two* charginos that are light and one (wino-like) somewhat heavier. It would be surprising if the LEP bound were still the most stringent, and indeed subsequent LHC searches are especially sensitive to winos up to about 800 GeV, see [65]. It might be possible (albeit difficult) to evade this constraint if the light wino and neutralino are close enough in mass so that decays such as $\tilde{\chi}^\pm \rightarrow \tilde{\chi}^0 + W^\pm$ are not possible, but without a detailed investigation we cannot exclude the possibility that some region of parameter space might evade direct LHC searches; we can only apply the LEP constraint as a hard lower bound on the chargino mass. Even so, just the LEP bound in combination with the measurement of the W boson mass proves fatal to the model, as we shall show below.

In [36] it was demonstrated that that the corrections to $\Delta\rho$ were correlated with the lightest chargino mass; in order to evade the LEP bound, λ_T has to be large and this drives large $\Delta\rho$. Here we can give a striking confirmation of this observation by plotting the W mass against the lightest chargino mass for a sample of $\mathcal{O}(70,000)$ spectra generated using SARAH. We fix the octet scalar, and all squark and slepton soft masses via $m_O^2 = m_q^2 = m_l^2 = 10 \text{ TeV}^2$, and fix the gluino mass to 3 TeV; this ensures that they are beyond all current and near-future bounds. Then we scan over the ranges:

$$\begin{aligned} m_{DY} &\in [100, 500] \text{ GeV}, & m_{D2} &\in [100, 250] \text{ GeV}, \\ v_S &\in [-50, 50] \text{ GeV}, & v_T &\in [-5, 5] \text{ GeV} \\ B_\mu &\in [10^4, 10^6] (\text{GeV})^2, & \lambda_T &\in [-1, 1], \\ \lambda_S &\in [-1.5, 1.5], & \tan\beta &\in [2, 50]. \end{aligned} \tag{3.13}$$

We use a Markov Chain Monte-Carlo (MCMC) algorithm to generate points (this helps to obtain points with non-tachyonic spectra with Higgs mass close to the observed




Fig. 1 W boson mass vs lightest chargino mass for points generated in the ‘MSSM without μ -term.’ The orange, purple and green horizontal bands show the SM, Tevatron and world average masses for the W boson mass respectively; the vertical green band shows the LEP constraint on the lightest chargino

value compared to a random scan). There is not intended to be a genuine statistical interpretation of the distribution of the points, but the overall envelope should show where valid sets of parameters exist. The results are shown in Fig. 1, where the LEP constraint is shown as a vertical green band. It can be clearly seen that it is not possible to both satisfy the LEP constraint and have an acceptable value for the W boson mass; from the W boson mass alone we would predict a chargino of mass below 65 GeV.

3.3 W mass in the MDGSSM

In the minimal Dirac Gaugino extension of the Standard Model (MDGSSM) the most typical scenario is to assume that R-symmetry is broken only via a B_μ term, i.e. the Higgs sector is special. It is identical to the previous model except that we allow a μ -term. This, however, makes all the difference: now the higgsino mass is not bounded from below, not requiring a large mixing with the winos; and further the enhancement to the Higgs mass is under control.

We perform a new MCMC scan with parameters allowed to vary within the ranges:

$$\begin{aligned}
 m_{D_Y} &\in [100, 700] \text{ GeV}, & m_{D_2} &\in [100, 1200] \text{ GeV}, \\
 v_S &\in [-50, 50] \text{ GeV}, & v_T &\in [-5, 5] \text{ GeV} \\
 \mu &\in [0, 1000] \text{ GeV}, & B_\mu &\in [10^4, 10^6] (\text{GeV})^2, \\
 \sqrt{2}\lambda_T &\in [-1.5, 1.5], & \lambda_S &\in [-1.5, 1.5], & \tan\beta &\in [2, 50].
 \end{aligned}
 \tag{3.14}$$

The range for m_{D_Y} is notably lower than that for m_{D_2} to favour a bino-like LSP and because, if we do not enforce an alignment condition, a larger m_{D_Y} drives down the Higgs mass through mixing between the singlet and doublet. We fix the octet scalar, and all squark and slepton soft masses via $m_O^2 = m_{\tilde{q}}^2 = m_{\tilde{l}}^2 = 10 \text{ TeV}^2$, and fix the gluino mass to 3 TeV. We choose a likelihood function to be a product of a gaussian in the Higgs mass with mean 125 GeV and standard deviation 3 GeV, a gaussian in the W mass with mean 80.413 GeV and standard deviation 20 MeV, and a sigmoid on the constraints (given as the maximum ratio of predicted cross-section to observed, across all channels) from HiggsBounds5 [99–102], which strongly suppresses the likelihood when the observed cross-section ratio is greater than one, but is otherwise close to unity. This choice of likelihood function is merely a device to select desirable points, and the distribution of the points is not meant to have a statistical interpretation in terms of their Bayesian likelihood; in particular, the theory uncertainty on the Higgs mass does not have a statistical interpretation, and a window of 3 GeV is a conservative estimate of the average error, since we use the latest two-loop corrections in the generalised effective potential and gaugeless limit [61, 103–105] with pole-mass matching onto the SM [106] (see [107] for a recent review). Such a conservative window of 3 GeV is employed because only a relatively small proportion of points actually generate a spectrum, and in principle a two-stage procedure pre-filtering points along those suggested in [65, 108] would probably be more efficient – or in addition the ability to invert the vacuum minimisation relations and compute v_S, v_T instead of treating them as inputs, but this is not yet automatically possible in the code in a way that would correctly incorporate the loop corrections to the Higgs masses [93].

We show plots in Fig. 2 for M_W against v_T (left plot), and then for M_W against λ_T (right plot), to show the points that benefit from large quantum corrections as the means of enhancing the W mass: the tree-level expectation just from modifying v_T is shown as a solid red curve on the left plot. We also show as a dashed red curve the value of M_W that would be obtained by insisting that the shift in $\Delta\rho$ only modifies M_W without changing $\sin\theta_W$ (if a different method of matching onto the SM parameters were used, for example).

All points shown satisfy all Higgs bounds; have a charged Higgs heavier than 600 GeV (so are safe from $B \rightarrow s\gamma$ constraints [1]); have charginos heavier than the LEP limit and winos heavier than 600 GeV. These baseline selections are marked as blue points in the plots; there are about 36,000, of which about 10,000 have $|v_T| < 1 \text{ GeV}$. Points shown in yellow further have charginos heavier than 200 GeV, $m_{D_2} > 700 \text{ GeV}$; while those marked in green have charginos heavier than 250 GeV and $m_{D_2} > 800 \text{ GeV}$ (about 1600 points in our sample survive these cuts). The green points are thus almost certainly guaranteed to be safe from current col-




Fig. 2 Left: W boson mass vs v_T in the MDGSSM, with the red curve showing the tree-level prediction. Right: W boson mass vs $\sqrt{2}\lambda_T$ in the MDGSSM. Colours of the points are described in the text, with

those obeying the strictest cuts shown in green. The colourful horizontal bands show the SM range in light orange; the Tevatron + LEP average in purple, and a conservative world average in green

Table 1 Benchmark points for the MDGSSM. Input parameters are given above the double line, and masses of the most important particles below

	MDG1	MDG2	MDG3
m_{D_Y} (GeV)	280	285	245
m_{D_2} (GeV)	983	941	940
μ (GeV)	276	255	353
$\tan \beta$	48	46	47
$-\lambda_S$	1.179	1.074	1.112
$\sqrt{2}\lambda_T$	-0.487	0.502	0.099
B_μ (GeV) ²	828,838	794,477	938,787
v_S (GeV)	5.0	4.3	1.4
v_T (GeV)	2.3	2.7	2.8
m_{h_1} (GeV)	125.4	124.7	124.7
m_{h_2} (GeV)	3120.9	2274.3	2405.8
m_{A_1} (GeV)	2394.2	1221.4	1456.5
$m_{H_1^\pm}$ (GeV)	2400.2	1213.0	1455.3
$m_{\tilde{\chi}_1^0}$ (GeV)	217.9	231.5	233.0
$m_{\tilde{\chi}_1^\pm}$ (GeV)	255.0	275.9	362.0
m_W (GeV)	80.425	80.420	80.421

lider bounds (although they may yet be probed in future). The requirement of heavy charged Higgs scalars sets the MSSM-like neutral and pseudoscalar masses to be heavy, and essentially guarantees the safety of all selected points from constraints on the couplings of the SM-like Higgs. Nevertheless, we also filtered the green points with constraints from HiggsSignals [109, 110]. The different categories

of points show the expected wider range of enhancements to the W mass as the charginos become lighter. We provide a selection of benchmark points, with the input parameters and crucial data, in Table 1. Note that m_{D_Y} and μ are close in value; the higgsinos drive the enhancement to the W mass and mixing between the bino and higgsino contribute to the mass splitting between neutral and charged components.

The first clear observation is that the quantum corrections to the W mass are at least as important as the tree-level contribution from the expectation value v_T , and the generic contribution to the W mass is positive with no points below the red curve. The asymmetry of the plot with v_T is due to the fact that we only take positive Dirac gaugino masses. It is important to note that in the red curves we take the SM value of the W boson mass to be 80.352 GeV, whereas the fitting function of [76] as employed in SARAH gives a value of 80.354 GeV for the central values; and gives 80.356 for a Higgs mass at the lower bound of our permitted range of 122 GeV. However, it is clear that the quantum corrections in our sample are generally more important than $\Delta\rho_{\text{tree}}$. Indeed, with the parameter ranges we have chosen, we are not within the range of masses required for decoupling of the quantum corrections (we have checked that the quantum corrections to M_W smoothly drop to near zero as the masses of all particles are raised to about 2 TeV or higher).

The second observation is that there is no clear correlation between the W mass and the parameter λ_T . Due to our requirement of a large wino mass, the selected points have light neutralinos/charginos of mixed bino/higgsino type. The mass splittings among the higgsinos – and thus the contribution to the W mass – can be driven large by λ_S and λ_T . These

couplings also enhance the Higgs mass at tree-level, so large values are favoured in the scans because we fixed the stop masses at $\sqrt{10}$ TeV. In this class of model there is therefore no particular preference for one or the other coupling.

Since the selected points generally have a mixed bino/higgsino LSP, they have good dark matter candidates, but it is expected that the relic density should be underdense. A detailed investigation of the dark matter-collider complementarity for scenarios satisfying the latest W mass data along the lines of [65] would be an interesting subject for future work, provided that the latest LHC analyses can be recast. As mentioned above, the possibility of a large λ_T (and the presence of the singlet scalar/fermion) distinguishes higgsinos in this scenario from those in the MSSM. It is clear that this class of models provides a very natural explanation for an enhancement to the W mass compared to the Standard Model.

3.4 W mass in the aligned MDGSSM

In the aligned MDGSSM (where the only source of R-symmetry breaking is the B_μ term, and we take $T_T = 0$), we choose the parameters to induce alignment at tree-level in the MDGSSM (so $\lambda_S = g_Y/\sqrt{2}, \lambda_T = g_2/\sqrt{2}$ and $m_{DY} = c_{2\beta}\mu_{\text{eff}}$) but taking $T_S = 0$ (which means the mixing of the singlet with the heavy Higgs cannot vanish unless it is heavy). To study this scenario we perform a scan with the same strategy as before except that now, since λ_S, λ_T are fixed, it is necessary to vary the masses of the stops/sbottoms to allow us to find the observed value of the Higgs mass; there is also therefore a preference for models with larger $\tan\beta$ since the tree-level contributions to the Higgs mass at low $\tan\beta$ are not sufficient. We therefore use a common mass for the third generation squarks M_{SUSY} ; we set $m_{Q,33}^2 = m_{U,33}^2 = m_{D,33}^2 = M_{\text{SUSY}}^2$ (the other squarks and sleptons we retain fixed at $\sqrt{10}$ TeV). The remaining parameters are scanned via the same MCMC algorithm in the ranges:

$$\begin{aligned} m_{D2} &\in [400, 1500] \text{ GeV}, & v_S &\in [-250, 250] \text{ GeV}, \\ v_T &\in [-5, 5] \text{ GeV} & \mu &\in [-1000, 1000] \text{ GeV}, \\ B_\mu &\in [10^4, 10^6] (\text{GeV})^2, \\ M_{\text{SUSY}} &\in [2, 10] \text{ TeV}, & \tan\beta &\in [2, 50]. \end{aligned} \tag{3.15}$$

We mostly find points with small μ/m_{DY} and very little enhancement to the W mass because the electroweakinos tend to be light. We give benchmark points in Table 2 and plots in Fig. 3 which demonstrate the lack of enhancement and scarcity of points (790 survived from a scan for one million).

3.5 W mass in the aligned DGNMSSM

We turn now to the case of the aligned DGNMSSM described in Sect. 2.4. We set the couplings λ_S, λ_T to their $N = 2$ values and then choose m_{DY} and T_S to make Δ_{h_s} and Δ_{H_s} vanish. This leads to

$$\begin{aligned} m_{DY} &= c_{2\beta}\mu_{\text{eff}} \\ T_S &= -g_Y(\kappa v_S - \sqrt{2}s_{2\beta}\mu_{\text{eff}}). \end{aligned} \tag{3.16}$$

This has an interesting consequence because in this model

$$\begin{aligned} B_{\mu, \text{eff}} &= \frac{1}{\sqrt{2}}T_S v_S + \frac{1}{2}\lambda_S \kappa v_S^2 \\ &= -\frac{1}{2\sqrt{2}}g_Y \kappa v_S^2 + g_Y s_{2\beta} v_S \mu_{\text{eff}}. \end{aligned} \tag{3.17}$$

Then

$$M_A^2 = g_Y v_S \left(2\mu_{\text{eff}} - \frac{\kappa v_S}{\sqrt{2}s_{2\beta}} \right) = g_Y v_S^2 \left(g_Y - \frac{\kappa}{\sqrt{2}s_{2\beta}} \right). \tag{3.18}$$

We perform a scan using the same strategy as the previous sections; we use a common mass for the third generation squarks M_{SUSY} ; we set $m_{Q,33}^2 = m_{U,33}^2 = m_{D,33}^2 = M_{\text{SUSY}}^2$ (the other squarks and sleptons we retain fixed at $\sqrt{10}$ TeV). Then we scan with the parameter ranges, using the same likelihood function as before:

$$\begin{aligned} M_{\text{SUSY}} &\in [2000, 10,000] \text{ GeV}, \\ m_{D2} &\in [400, 1500] \text{ GeV}, & v_S &\in [-1500, 1500] \text{ GeV}, \\ v_T &\in [-5, 5] \text{ GeV} & \kappa &\in [-1.5, 1.5], \\ T_k &\in [-2000, 2000] \text{ GeV}, & \tan\beta &\in [2, 50]. \end{aligned}$$

In our scans we impose that all Higgs searches are satisfied using `HiggsBounds` and `HiggsSignals`. We show the results for the W boson mass in Fig. 4 where the points have the same colour coding as in the previous sections. It is apparent that in this model it is complicated to enhance the W boson mass. This is because we have only a small quantum effect from λ_S, λ_T , but also because the lightest neutralinos are typically rather light: since $\mu_{\text{eff}} = g_Y v_S/2$ we need a large $v_S \gtrsim 500$ GeV (or 1 TeV for our more stringent points) to have heavy enough higgsinos. Then we need κ negative and not too small to avoid a too-small pseudoscalar/charged Higgs mass (if we neglect κ then M_A is bounded by $g_Y v_S$, so $M_A > 600$ GeV requires $v_S \gtrsim 1700$ GeV, at the limit of our search range). So this implies that the singlino is generally heavy compared to m_{DY} :

$$\frac{m_{DY}}{\sqrt{2}\kappa v_S} = \frac{g_Y c_{2\beta}}{2\sqrt{2}\kappa}. \tag{3.19}$$

Table 2 Benchmark points with a light singlet in the aligned MDGSSM

	AMDG1	AMDG2	AMDG3	AMDG4	AMDG5	AMDG6
M_{SUSY} (GeV)	8636.2	4550.9	5181.9	8436.1	7357.2	5454.9
m_{D2} (GeV)	1037	1249	1149	1107	1219	994
$\tan \beta$	2	2	3	10	3	2
μ (GeV)	158.2	160.4	165.8	174.1	159.5	201.6
v_S (GeV)	-19.0	-12.7	-8.4	-0.3	-5.0	-13.7
v_T (GeV)	1.5	1.5	1.1	0.3	1.9	1.1
m_{h_1} (GeV)	126.6	122.8	123.2	123.6	127.4	122.2
m_{h_2} (GeV)	457.7	312.3	308.5	421.8	465.8	487.5
m_{h_3} (GeV)	1282.2	799.7	935.2	1251.6	1176.5	762.3
m_{h_4} (GeV)	2667.0	2965.0	3350.8	7104.7	2847.1	2933.0
$m_{H_1^\pm}$ (GeV)	1284.0	793.5	933.4	1257.7	1174.3	765.4
$m_{\tilde{\chi}_1^0}$ (GeV)	106.3	110.4	120.5	172.6	128.0	134.5
$m_{\tilde{\chi}_1^\pm}$ (GeV)	168.4	169.6	177.3	193.1	173.1	211.5
m_W (GeV)	80.363	80.365	80.362	80.361	80.369	80.362




Fig. 3 Points in the aligned MDGSSM. Left: M_W against triplet expectation value. Right: M_W against singlet-like scalar mass. Colour coding as for previous plots

In Fig. 4 we show the scan results with the same colour coding as in previous sections. It is clear that in this scenario, models which can explain a large W mass are driven by a larger v_T with some modest quantum corrections enhancing the mass by $\mathcal{O}(10)$ MeV; there is very little spread due to the lack of variation in λ_T . However, it is difficult to find points with large enough v_T that satisfy other bounds.

Another feature of the selected points is that in almost all cases $v_T > 0$; considering $T_T = M_T = 0$ in equation (2.19) means that in order for $\tilde{m}_{T_R}^2 > 0$ (so that, at least, the pseudoscalar triplet should be non-tachyonic, since we take $B_T = 0$) and $v_T < 0$ we would need $\mu_{\text{eff}} > |m_{D2}c_{2\beta}|$. But we need large $\tan \beta$ to obtain the correct Higgs mass, and therefore $g_Y v_S/2 \gtrsim m_{D2}$; for our selected points we require

a minimum of $m_{D2} > 600$ GeV, and so v_S would again be beyond our search range.

Since we are interested here in alignment, we may have a light singlet scalar without falling foul of either light Higgs or heavy Higgs searches. In this limit we have

$$\tilde{m}_S^2 \rightarrow \frac{1}{4} \left[2v_S(\sqrt{2}T_k + 3\sqrt{2}\kappa M_S + 4v_S\kappa^2) + v^2 g_Y(g_Y - \sqrt{2}\kappa s_{2\beta}) \right]. \tag{3.20}$$

T_k of opposite sign to v_S allows the singlet to be made light while making M_A arbitrarily heavy. In the scan, we do not impose any likelihood bias to search for points with a light

Table 3 Benchmark points for the “aligned DGNMSSM”. Input parameters are given above the double line, and masses of the most important particles below

	A-DGN1	A-DGN2	A-DGN3	A-DGN4
M_{SUSY} (GeV)	6368.4	5186.5	8219.5	8702.9
m_{D2} (GeV)	802	991	932	923
$\tan \beta$	26	11	29	22
κ	-0.418	-0.332	-0.446	-0.418
v_S (GeV)	1417.4	1402.4	1337.0	1488.8
v_T (GeV)	0.2	0.1	0.1	0.8
T_κ (GeV)	-595.6	-267.5	-648.0	-639.4
m_{h_1} (GeV)	124.1	122.9	125.2	123.4
m_{h_2} (GeV)	353.5	417.6	347.2	353.9
m_{h_3} (GeV)	1838.5	1284.5	1843.0	1803.0
m_{h_4} (GeV)	6710.7	12226.0	10817.9	4056.2
$m_{H_1^\pm}$ (GeV)	1841.4	1288.5	1846.1	1806.2
$m_{\tilde{\chi}_1^0}$ (GeV)	72.2	84.2	64.5	75.2
$m_{\tilde{\chi}_1^\pm}$ (GeV)	280.1	274.6	265.1	291.9
m_W (GeV)	80.362	80.361	80.361	80.363

singlet, but we show benchmark points passing all constraints which have light singlet masses in Table 3. They show a W mass consistent with the SM prediction.

3.6 W mass in the general DGNMSSM

Finally we consider the general DGNMSSM, where we allow the values of λ_S, λ_T to vary and do not fix the values of m_{DY} or T_S to require alignment but scan over them. This means that we will not focus on light singlet (or doublet) scalars. Similar to the aligned case, there is still a see-saw effect on the lightest neutralino mass due to the non-zero singlino mass, which can drive down the quantum corrections to the W boson, but a large $|\lambda_T|$ can compensate for this and also help enhance the SM-like Higgs mass (Table 4).

We perform a scan using the strategy as in Sects. 3.2 and 3.3, with the parameter ranges:

$$\begin{aligned}
 m_{DY} &\in [100, 700] \text{ GeV}, & m_{D2} &\in [150, 1200] \text{ GeV}, \\
 v_S &\in [-700, 700] \text{ GeV}, & v_T &\in [-5, 5] \text{ GeV} \\
 \kappa &\in [-1.5, 1.5], & T_\kappa &\in [-2000, 2000] \text{ GeV}, \\
 T_S &\in [-4000, 4000] \text{ GeV}, & \lambda_S &\in [-1.5, 1.5], \\
 \sqrt{2}\lambda_T &\in [-1.5, 1.5], & \tan \beta &\in [2, 50].
 \end{aligned}
 \tag{3.21}$$

We give plots in Fig. 5 with the same colour coding as in the previous sections; the difference in the distribution to the previous examples is rather striking. It is clear that in this scenario a large negative λ_T and a positive v_T is favoured; this gives a tree-level enhancement to the Higgs mass and a loop-

Table 4 Benchmark points for the general DGNSSM. Input parameters are given above the double line, and masses of the most important particles below

	DGN1	DGN2	DGN3	DGN4	DGN5
m_{DY} (GeV)	392	298	410	380	292
m_{D2} (GeV)	927	971	841	1003	805
κ	1.391	-1.369	-1.266	-1.309	-1.361
$\tan \beta$	9	23	21	30	30
$-\lambda_S$	0.727	-0.893	-0.544	0.554	-0.677
$\sqrt{2}\lambda_T$	-1.426	1.496	1.463	-1.303	-1.296
$-T_S$ (GeV)	3077	3747	-2496	-3002	1183
T_κ (GeV)	1139	350	-1292	-571	728
v_S (GeV)	-658.1	-539.1	574.5	524.7	-482.8
v_T (GeV)	2.7	2.3	1.7	2.5	2.2
m_{h_1} (GeV)	125.1	125.3	125.8	124.6	124.7
m_{h_2} (GeV)	1017.6	937.1	665.0	831.8	740.1
m_{A_1} (GeV)	757.6	93.7	778.1	115.1	502.2
$m_{H_1^\pm}$ (GeV)	2793.1	3195.1	1281.7	778.5	806.5
$m_{\tilde{\chi}_1^0}$ (GeV)	115.0	87.2	123.6	110.9	90.7
$m_{\tilde{\chi}_1^\pm}$ (GeV)	278.8	273.6	265.5	254.8	268.8
m_W (GeV)	80.421	80.421	80.424	80.420	80.422

level enhancement to the W-boson mass. The singlino component will mix less with the higgsinos than in the MDGSSM because of the $\sqrt{2}\kappa v_S$ singlino mass, and thus the effect of λ_S on the W mass is diminished. The asymmetry in the signs of λ_T and v_T can be explained by the fact that we only take positive Dirac gaugino masses in the scans.

4 Conclusions

We have shown that an aligned Dirac Gaugino NMSSM is possible and compatible with current collider constraints; it can even lead to relatively light singlet scalars that may be of interest to future searches (although would be rather difficult to find directly as they are difficult to produce). Such a model favours a W boson mass compatible with or just above the SM prediction. We also showed how two different Dirac Gaugino scenarios can easily be compatible with an enhanced W boson mass, including a precise computation of the quantum corrections for the first time, which are now incorporated automatically in the package SARAH. We also used this computation to add more nails to the coffin of the “MSSM without μ term.”

We have been conservative in our application of collider constraints and concluded that the MDGSSM models would typically contain underdense dark matter densities. However, it would be interesting to examine the issue of dark matter and collider constraints again in all of these classes of




Fig. 4 W boson mass in the aligned DGMSSM. Left: W boson mass against triplet expectation value. Right: W boson mass against singlet-like Higgs mass. Colour coding of points is described in the text




Fig. 5 W boson mass in the general DGMSSM. Left: W boson mass against triplet expectation value. Right: W boson mass against λ_T . Colour coding of points as in previous figures

models when all the latest searches for electroweakinos have been recast; we have provided ample benchmark points for this purpose. In the DGMSSM or its aligned version, if we impose strict R-parity or have a heavy gravitino (by no means entirely obvious assumptions), it may be that we require a Higgs funnel to obtain the correct relic density, which would require a sophisticated search strategy to find allowed parameter ranges, along e.g. the lines of [108]. However, it is also likely that a light singlino in the aligned DGMSSM could fulfil the role of the Higgs funnel. We leave these questions to future work.

Acknowledgements The work of W. K. is supported by the Contrat Doctoral Spécifique Normalien (CDSN) of Ecole Normale Supérieure—PSL. K. B., P. S. and M. D. G. acknowledge support from the grant “HiggsAutomator” of the Agence Nationale de la Recherche (ANR) (ANR-15-CE31-0002). M. D. G. also acknowledges support from the ANR via grant “DMwithLLPatLHC”, (ANR-21-CE31-0013). M. D. G. thanks Werner Porod and Hannah Day for interesting discussions. W. K. thanks Pierre Fayet for illuminating discussions during the early stages of this work.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors’ comment: The manuscript has no associated data.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation,

distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- M. Misiak, M. Steinhauser, Weak radiative decays of the B meson and bounds on M_{H^\pm} in the two-Higgs-doublet model. *Eur. Phys. J. C* **77**(3), 201 (2017). <https://doi.org/10.1140/epjc/s10052-017-4776-y>. arXiv:1702.04571 [hep-ph]
- S. Davidson, H.E. Haber, Basis-independent methods for the two-Higgs-doublet model. *Phys. Rev. D* **72**, 035004 (2005). <https://doi.org/10.1103/PhysRevD.72.099902>. arXiv:hep-ph/0504050. [Erratum: *Phys. Rev. D* **72**, 099902 (2005)]
- P.S. Bhupal Dev, A. Pilaftsis, Maximally symmetric two Higgs doublet model with natural standard model alignment. *JHEP* **12** (2014) 024. [https://doi.org/10.1007/JHEP12\(2014\)024](https://doi.org/10.1007/JHEP12(2014)024). arXiv:1408.3405 [hep-ph]. [Erratum: *JHEP* **11**, 147 (2015)]
- J. Bernon, J.F. Gunion, H.E. Haber, Y. Jiang, S. Kraml, Scrutinizing the alignment limit in two-Higgs-doublet models: $m_h=125$ GeV. *Phys. Rev. D* **92**(7), 075004 (2015). <https://doi.org/10.1103/PhysRevD.92.075004>. arXiv:1507.00933 [hep-ph]
- J. Bernon, J.F. Gunion, H.E. Haber, Y. Jiang, S. Kraml, Scrutinizing the alignment limit in two-Higgs-doublet models. II. $m_H=125$ GeV. *Phys. Rev. D* **93**(3), 035027 (2016). <https://doi.org/10.1103/PhysRevD.93.035027>. arXiv:1511.03682 [hep-ph]
- M. Carena, I. Low, N.R. Shah, C.E.M. Wagner, Impersonating the standard model Higgs boson: alignment without decoupling. *JHEP* **04**, 015 (2014). [https://doi.org/10.1007/JHEP04\(2014\)015](https://doi.org/10.1007/JHEP04(2014)015). arXiv:1310.2248 [hep-ph]
- M. Carena, H.E. Haber, I. Low, N.R. Shah, C.E.M. Wagner, Alignment limit of the NMSSM Higgs sector. *Phys. Rev. D* **93**(3), 035013 (2016). <https://doi.org/10.1103/PhysRevD.93.035013>. arXiv:1510.09137 [hep-ph]
- H.E. Haber, S. Heinemeyer, T. Stefaniak, The impact of two-loop effects on the scenario of MSSM Higgs alignment without decoupling. *Eur. Phys. J. C* **77**(11), 742 (2017). <https://doi.org/10.1140/epjc/s10052-017-5243-5>. arXiv:1708.04416 [hep-ph]
- N.M. Coyle, C.E.M. Wagner, Dynamical Higgs field alignment in the NMSSM. *Phys. Rev. D* **101**(5), 055037 (2020). <https://doi.org/10.1103/PhysRevD.101.055037>. arXiv:1912.01036 [hep-ph]
- P. Fayet, Fermi–Bose hypersymmetry. *Phys. B. Nucl* **113**, 135 (1976). [https://doi.org/10.1016/0550-3213\(76\)90458-2](https://doi.org/10.1016/0550-3213(76)90458-2)
- P. Fayet, Massive gluinos. *Lett. B. Phys* **78**, 417–420 (1978). [https://doi.org/10.1016/0370-2693\(78\)90474-4](https://doi.org/10.1016/0370-2693(78)90474-4)
- J. Polchinski, L. Susskind, Breaking of supersymmetry at intermediate-energy. *Phys. Rev. D* **26**, 3661 (1982). <https://doi.org/10.1103/PhysRevD.26.3661>
- L.J. Hall, L. Randall, U(1)-R symmetric supersymmetry. *Nucl. Phys. B* **352**, 289–308 (1991). [https://doi.org/10.1016/0550-3213\(91\)90444-3](https://doi.org/10.1016/0550-3213(91)90444-3)
- P.J. Fox, A.E. Nelson, N. Weiner, Dirac gaugino masses and super-soft supersymmetry breaking. *JHEP* **08**, 035 (2002). <https://doi.org/10.1088/1126-6708/2002/08/035>. arXiv:hep-ph/0206096
- A.E. Nelson, N. Rius, V. Sanz, M. Unsal, The minimal supersymmetric model without a mu term. *JHEP* **08**, 039 (2002). <https://doi.org/10.1088/1126-6708/2002/08/039>. arXiv:hep-ph/0206102
- I. Antoniadis, K. Benakli, A. Delgado, M. Quiros, A new gauge mediation theory. *Adv. Stud. Theor. Phys.* **2**, 645–672 (2008). arXiv:hep-ph/0610265
- G. Belanger, K. Benakli, M. Goodsell, C. Moura, A. Pukhov, Dark matter with Dirac and majorana gaugino masses. *JCAP* **08**, 027 (2009). <https://doi.org/10.1088/1475-7516/2009/08/027>. arXiv:0905.1043 [hep-ph]
- K. Benakli, M.D. Goodsell, Dirac gauginos in general gauge mediation. *Nucl. Phys. B* **816**, 185–203 (2009). <https://doi.org/10.1016/j.nuclphysb.2009.03.002>. arXiv:0811.4409 [hep-ph]
- K. Benakli, M.D. Goodsell, A.-K. Maier, Generating mu and Bmu in models with Dirac gauginos. *Nucl. Phys. B* **851**, 445–461 (2011). <https://doi.org/10.1016/j.nuclphysb.2011.06.001>. arXiv:1104.2695 [hep-ph]
- G.D. Kribs, E. Poppitz, N. Weiner, Flavor in supersymmetry with an extended R-symmetry. *Phys. Rev. D* **78**, 055010 (2008). <https://doi.org/10.1103/PhysRevD.78.055010>. arXiv:0712.2039 [hep-ph]
- S.D.L. Amigo, A.E. Blechman, P.J. Fox, E. Poppitz, R-symmetric gauge mediation. *JHEP* **01**, 018 (2009). <https://doi.org/10.1088/1126-6708/2009/01/018>. arXiv:0809.1112 [hep-ph]
- K. Benakli, M.D. Goodsell, Dirac gauginos and kinetic mixing. *Nucl. Phys. B* **830**, 315–329 (2010). <https://doi.org/10.1016/j.nuclphysb.2010.01.003>. arXiv:0909.0017 [hep-ph]
- S.Y. Choi, J. Kalinowski, J.M. Kim, E. Popena, Scalar gluons and Dirac gluinos at the LHC. *Acta Phys. Polon. B* **40**, 2913–2922 (2009). arXiv:0911.1951 [hep-ph]
- K. Benakli, M.D. Goodsell, Dirac Gauginos, Gauge mediation and unification. *Nucl. Phys. B* **840**, 1–28 (2010). <https://doi.org/10.1016/j.nuclphysb.2010.06.018>. arXiv:1003.4957 [hep-ph]
- S.Y. Choi, D. Choudhury, A. Freitas, J. Kalinowski, J.M. Kim, P.M. Zerwas, Dirac neutralinos and electroweak scalar bosons of $N = 1/N = 2$ hybrid supersymmetry at colliders. *JHEP* **08**, 025 (2010). [https://doi.org/10.1007/JHEP08\(2010\)025](https://doi.org/10.1007/JHEP08(2010)025). arXiv:1005.0818 [hep-ph]
- L.M. Carpenter, Dirac Gauginos, negative supertraces and gauge mediation. *JHEP* **09**, 102 (2012). [https://doi.org/10.1007/JHEP09\(2012\)102](https://doi.org/10.1007/JHEP09(2012)102). arXiv:1007.0017 [hep-th]
- S. Abel, M. Goodsell, Easy Dirac gauginos. *JHEP* **06**, 064 (2011). [https://doi.org/10.1007/JHEP06\(2011\)064](https://doi.org/10.1007/JHEP06(2011)064). arXiv:1102.0014 [hep-th]
- R. Davies, J. March-Russell, M. McCullough, A supersymmetric one Higgs doublet model. *JHEP* **04**, 108 (2011). [https://doi.org/10.1007/JHEP04\(2011\)108](https://doi.org/10.1007/JHEP04(2011)108). arXiv:1103.1647 [hep-ph]
- K. Benakli, Dirac gauginos: a user manual. *Fortsch. Phys.* **59**, 1079–1082 (2011). <https://doi.org/10.1002/prop.201100071>. arXiv:1106.1649 [hep-ph]
- C. Frugiuele, T. Gregoire, Making the S neutrino a Higgs with a $U(1)_R$ lepton number. *Phys. Rev. D* **85**, 015016 (2012). <https://doi.org/10.1103/PhysRevD.85.015016>. arXiv:1107.4634 [hep-ph]
- H. Itoyama, N. Maru, D-term dynamical supersymmetry breaking generating split $N = 2$ gaugino masses of mixed majorana-Dirac type. *Int. J. Mod. Phys. A* **27**, 1250159 (2012). <https://doi.org/10.1142/S0217751X1250159X>. arXiv:1109.2276 [hep-ph]
- E. Bertuzzo, C. Frugiuele, Fitting neutrino physics with a $U(1)_R$ lepton number. *JHEP* **05**, 100 (2012). [https://doi.org/10.1007/JHEP05\(2012\)100](https://doi.org/10.1007/JHEP05(2012)100). arXiv:1203.5340 [hep-ph]
- R. Davies, Dirac gauginos and unification in F-theory. *JHEP* **10**, 010 (2012). [https://doi.org/10.1007/JHEP10\(2012\)010](https://doi.org/10.1007/JHEP10(2012)010). arXiv:1205.1942 [hep-th]

34. C. Frugiuele, T. Gregoire, P. Kumar, E. Ponton, $L=R$ - $U(1)_R$ as the origin of leptonic 'RPV. *JHEP* **03**, 156 (2013). [https://doi.org/10.1007/JHEP03\(2013\)156](https://doi.org/10.1007/JHEP03(2013)156). arXiv:1210.0541 [hep-ph]
35. C. Frugiuele, T. Gregoire, P. Kumar, E. Ponton, $L=R$ - $U(1)_R$ lepton number at the LHC. *JHEP* **05**, 012 (2013). [https://doi.org/10.1007/JHEP05\(2013\)012](https://doi.org/10.1007/JHEP05(2013)012). arXiv:1210.5257 [hep-ph]
36. K. Benakli, M.D. Goodsell, F. Staub, Dirac gauginos and the 125 GeV Higgs. *JHEP* **06**, 073 (2013). [https://doi.org/10.1007/JHEP06\(2013\)073](https://doi.org/10.1007/JHEP06(2013)073). arXiv:1211.0552 [hep-ph]
37. E. Dudas, M. Goodsell, L. Heurtier, P. Tziveloglou, Flavour models with Dirac and fake gluinos. *Nucl. Phys. B* **884**, 632–671 (2014). <https://doi.org/10.1016/j.nuclphysb.2014.05.005>. arXiv:1312.2011 [hep-ph]
38. K. Benakli, L. Darmé, M.D. Goodsell, P. Slavich, A fake split supersymmetry model for the 126 GeV Higgs. *JHEP* **05**, 113 (2014). [https://doi.org/10.1007/JHEP05\(2014\)113](https://doi.org/10.1007/JHEP05(2014)113). arXiv:1312.5220 [hep-ph]
39. K. Benakli, L. Darmé, M.D. Goodsell, (O)Mega split. *JHEP* **11**, 100 (2015). [https://doi.org/10.1007/JHEP11\(2015\)100](https://doi.org/10.1007/JHEP11(2015)100). arXiv:1508.02534 [hep-ph]
40. A. Arvanitaki, M. Baryakhtar, X. Huang, K. van Tilburg, G. Villadoro, The last vestiges of naturalness. *JHEP* **03**, 022 (2014). [https://doi.org/10.1007/JHEP03\(2014\)022](https://doi.org/10.1007/JHEP03(2014)022). arXiv:1309.3568 [hep-ph]
41. H. Itoyama, N. Maru, D-term triggered dynamical supersymmetry breaking. *Phys. Rev. D* **88**(2), 025012 (2013). <https://doi.org/10.1103/PhysRevD.88.025012>. arXiv:1301.7548 [hep-ph]
42. S. Chakraborty, S. Roy, Higgs boson mass, neutrino masses and mixing and keV dark matter in an $U(1)_R$ - lepton number model. *JHEP* **01**, 101 (2014). [https://doi.org/10.1007/JHEP01\(2014\)101](https://doi.org/10.1007/JHEP01(2014)101). arXiv:1309.6538 [hep-ph]
43. C. Csaki, J. Goodman, R. Pavesi, Y. Shirman, The $m_D - b_M$ problem of Dirac gauginos and its solutions. *Phys. Rev. D* **89**(5), 055005 (2014). <https://doi.org/10.1103/PhysRevD.89.055005>. arXiv:1310.4504 [hep-ph]
44. H. Itoyama, N. Maru, 126 GeV Higgs boson associated with D-term triggered dynamical supersymmetry breaking. *Symmetry* **7**(1), 193–205 (2015). <https://doi.org/10.3390/sym7010193>. arXiv:1312.4157 [hep-ph]
45. H. Beauchesne, T. Gregoire, Electroweak precision measurements in supersymmetric models with a $U(1)_R$ lepton number. *JHEP* **05**, 051 (2014). [https://doi.org/10.1007/JHEP05\(2014\)051](https://doi.org/10.1007/JHEP05(2014)051). arXiv:1402.5403 [hep-ph]
46. K. Benakli, (Pseudo)goldstinos, SUSY fluids, Dirac gravitino and gauginos. *EPJ Web Conf.* **71**, 00012 (2014). <https://doi.org/10.1051/epjconf/20147100012>. arXiv:1402.4286 [hep-ph]
47. E. Bertuzzo, C. Frugiuele, T. Gregoire, E. Ponton, Dirac gauginos, R-symmetry and the 125 GeV Higgs. *JHEP* **04**, 089 (2015). [https://doi.org/10.1007/JHEP04\(2015\)089](https://doi.org/10.1007/JHEP04(2015)089). arXiv:1402.5432 [hep-ph]
48. P. Dießner, J. Kalinowski, W. Kotlarski, D. Stöckinger, Higgs boson mass and electroweak observables in the MRSSM. *JHEP* **12**, 124 (2014). [https://doi.org/10.1007/JHEP12\(2014\)124](https://doi.org/10.1007/JHEP12(2014)124). arXiv:1410.4791 [hep-ph]
49. K. Benakli, M. Goodsell, F. Staub, W. Porod, Constrained minimal Dirac gaugino supersymmetric standard model. *Phys. Rev. D* **90**(4), 045017 (2014). <https://doi.org/10.1103/PhysRevD.90.045017>. arXiv:1403.5122 [hep-ph]
50. M.D. Goodsell, P. Tziveloglou, Dirac gauginos in low scale supersymmetry breaking. *Nucl. Phys. B* **889**, 650–675 (2014). <https://doi.org/10.1016/j.nuclphysb.2014.10.020>. arXiv:1407.5076 [hep-ph]
51. D. Busbridge, Constrained Dirac gluino mediation. arXiv:1408.4605 [hep-ph]
52. S. Chakraborty, A. Datta, S. Roy, $h \rightarrow \gamma\gamma$ in $U(1)_R$ -lepton number model with a right-handed neutrino. *JHEP* **02**, 124 (2015). [https://doi.org/10.1007/JHEP02\(2015\)124](https://doi.org/10.1007/JHEP02(2015)124). arXiv:1411.1525 [hep-ph]. [Erratum: *JHEP*09,077(2015)]
53. R. Ding, T. Li, F. Staub, C. Tian, B. Zhu, Supersymmetric standard models with a pseudo-Dirac gluino from hybrid F- and D-term supersymmetry breaking. *Phys. Rev. D* **92**(1), 015008 (2015). <https://doi.org/10.1103/PhysRevD.92.015008>. arXiv:1502.03614 [hep-ph]
54. D.S.M. Alves, J. Galloway, M. McCullough, N. Weiner, Goldstone gauginos. *Phys. Rev. Lett.* **115**(16), 161801 (2015). <https://doi.org/10.1103/PhysRevLett.115.161801>. arXiv:1502.03819 [hep-ph]
55. D.S.M. Alves, J. Galloway, M. McCullough, N. Weiner, Models of goldstone gauginos. *Phys. Rev. D* **93**(7), 075021 (2016). <https://doi.org/10.1103/PhysRevD.93.075021>. arXiv:1502.05055 [hep-ph]
56. L.M. Carpenter, J. Goodman, New calculations in Dirac Gaugino models: operators, expansions, and effects. *JHEP* **07**, 107 (2015). [https://doi.org/10.1007/JHEP07\(2015\)107](https://doi.org/10.1007/JHEP07(2015)107). arXiv:1501.05653 [hep-ph]
57. S.P. Martin, Nonstandard supersymmetry breaking and Dirac gaugino masses without supersoftness. *Phys. Rev. D* **92**(3), 035004 (2015). <https://doi.org/10.1103/PhysRevD.92.035004>. arXiv:1506.02105 [hep-ph]
58. M.D. Goodsell, M.E. Krauss, T. Müller, W. Porod, F. Staub, Dark matter scenarios in a constrained model with Dirac gauginos. *JHEP* **10**, 132 (2015). [https://doi.org/10.1007/JHEP10\(2015\)132](https://doi.org/10.1007/JHEP10(2015)132). arXiv:1507.01010 [hep-ph]
59. W. Kotlarski, Sgluons in the same-sign lepton searches. *JHEP* **02**, 027 (2017). [https://doi.org/10.1007/JHEP02\(2017\)027](https://doi.org/10.1007/JHEP02(2017)027). arXiv:1608.00915 [hep-ph]
60. K. Benakli, L. Darmé, M.D. Goodsell, J. Harz, The diphoton excess in a perturbative SUSY model. *Nucl. Phys. B* **911**, 127–162 (2016). <https://doi.org/10.1016/j.nuclphysb.2016.07.027>. arXiv:1605.05313 [hep-ph]
61. J. Braathen, M.D. Goodsell, P. Slavich, Leading two-loop corrections to the Higgs boson masses in SUSY models with Dirac gauginos. *JHEP* **09**, 045 (2016). [https://doi.org/10.1007/JHEP09\(2016\)045](https://doi.org/10.1007/JHEP09(2016)045). arXiv:1606.09213 [hep-ph]
62. S. Chakraborty, J. Chakraborty, Natural emergence of neutrino masses and dark matter from R -symmetry. *JHEP* **10**, 012 (2017). [https://doi.org/10.1007/JHEP10\(2017\)012](https://doi.org/10.1007/JHEP10(2017)012). arXiv:1701.04566 [hep-ph]
63. G. Chalons, M.D. Goodsell, S. Kraml, H. Reyes-González, S.L. Williamson, LHC limits on gluinos and squarks in the minimal Dirac gaugino model. *JHEP* **04**, 113 (2019). [https://doi.org/10.1007/JHEP04\(2019\)113](https://doi.org/10.1007/JHEP04(2019)113). arXiv:1812.09293 [hep-ph]
64. L.M. Carpenter, T. Murphy, M.J. Smylie, Exploring color-octet scalar parameter space in minimal R -symmetric models. *JHEP* **11**, 024 (2020). [https://doi.org/10.1007/JHEP11\(2020\)024](https://doi.org/10.1007/JHEP11(2020)024). arXiv:2006.15217 [hep-ph]
65. M.D. Goodsell, S. Kraml, H. Reyes-González, S.L. Williamson, Constraining electroweakinos in the minimal Dirac gaugino model. *SciPost Phys.* **9**(4), 047 (2020). <https://doi.org/10.21468/SciPostPhys.9.4.047>. arXiv:2007.08498 [hep-ph]
66. L.M. Carpenter, T. Murphy, Color-octet scalars in Dirac gaugino models with broken R symmetry. *JHEP* **05**, 079 (2021). [https://doi.org/10.1007/JHEP05\(2021\)079](https://doi.org/10.1007/JHEP05(2021)079). arXiv:2012.15771 [hep-ph]
67. K. Benakli, M.D. Goodsell, S.L. Williamson, Higgs alignment from extended supersymmetry. *Eur. Phys. J. C* **78**(8), 658 (2018). <https://doi.org/10.1140/epjc/s10052-018-6125-1>. arXiv:1801.08849 [hep-ph]
68. K. Benakli, Y. Chen, G. Lafforgue-Marmet, R -symmetry for Higgs alignment without decoupling. *Eur. Phys. J. C* **79**(2), 172 (2019). <https://doi.org/10.1140/epjc/s10052-019-6676-9>. arXiv:1811.08435 [hep-ph]

69. K. Benakli, Y. Chen, G. Lafforgue-Marmet, Predicting alignment in a two Higgs doublet model. *MDPI Proc.* **13**(1), 2 (2019). <https://doi.org/10.3390/proceedings2019013002>. arXiv:1812.02208 [hep-ph]
70. K. Benakli, Y. Chen, G. Lafforgue-Marmet, An emergent Higgs alignment. *PoS CORFU* **2018**, 014 (2019). <https://doi.org/10.22323/1.347.0014>. arXiv:1904.04248 [hep-ph]
71. J. Ellis, J. Quevillon, V. Sanz, Doubling up on supersymmetry in the Higgs sector. *JHEP* **10**, 086 (2016). [https://doi.org/10.1007/JHEP10\(2016\)086](https://doi.org/10.1007/JHEP10(2016)086). arXiv:1607.05541 [hep-ph]
72. CMS Collaboration, Combined Higgs boson production and decay measurements with up to 137 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13 \text{ TeV}$. <https://inspirehep.net/literature/1774836>
73. ATLAS Collaboration, A combination of measurements of Higgs boson production and decay using up to 139 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13 \text{ TeV}$ collected with the ATLAS experiment. <https://inspirehep.net/literature/1809954>
74. C.D.F. Collaboration, T. Aaltonen et al., High-precision measurement of the W boson mass with the CDF II detector. *376*(6589), 170–176 (2022). <https://doi.org/10.1126/science.abk1781Science>
75. M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Precise prediction for the W boson mass in the standard model. *Phys. Rev. D* **69**, 053006 (2004). <https://doi.org/10.1103/PhysRevD.69.053006>. arXiv:hep-ph/0311148
76. G. Degrandi, P. Gambino, P.P. Giardino, The m_{Higgs} – m_{W} interdependence in the Standard Model: a new scrutiny. *JHEP* **05**, 154 (2015). [https://doi.org/10.1007/JHEP05\(2015\)154](https://doi.org/10.1007/JHEP05(2015)154). arXiv:1411.7040 [hep-ph]
77. OPAL Collaboration, G. Abbiendi et al., Measurement of the mass and width of the W boson. *Eur. Phys. J. C* **45** (2006) 307–335. <https://doi.org/10.1140/epjc/s2005-02440-5>. arXiv:hep-ex/0508060
78. ALEPH Collaboration, S. Schael et al., Measurement of the W boson mass and width in e^+e^- collisions at LEP. *Eur. Phys. J. C* **47** (2006) 309–335. <https://doi.org/10.1140/epjc/s2006-02576-8>. arXiv:hep-ex/0605011
79. ALEPH Collaboration, S. Schael et al., Measurement of the W boson mass and width in e^+e^- collisions at LEP. *Eur. Phys. J. C* **47** (2006) 309–335. <https://doi.org/10.1140/epjc/s2006-02576-8>. arXiv:hep-ex/0605011
80. DELPHI Collaboration, J. Abdallah et al., Measurement of the Mass and Width of the W Boson in e^+e^- Collisions at $\sqrt{s} = 161\text{-}209\text{-GeV}$. *Eur. Phys. J. C* **55** (2008) 1–38. <https://doi.org/10.1140/epjc/s10052-008-0585-7>. arXiv:0803.2534 [hep-ex]
81. ALEPH, DELPHI, L3, OPAL, LEP Electroweak Collaboration, S. Schael et al., Electroweak measurements in electron-positron collisions at W-boson-pair energies at LEP. *Phys. Rep.* **532** (2013) 119–244. <https://doi.org/10.1016/j.physrep.2013.07.004>. arXiv:1302.3415 [hep-ex]
82. D0 Collaboration, V.M. Abazov et al., Measurement of the W boson mass with the D0 detector. *Phys. Rev. Lett.* **108** (2012) 151804. <https://doi.org/10.1103/PhysRevLett.108.151804>. arXiv:1203.0293 [hep-ex]
83. C.D.F. Collaboration, T. Aaltonen et al., Precise measurement of the W-boson mass with the CDF II detector. *Phys. Rev. Lett.* **108**, 151803 (2012). <https://doi.org/10.1103/PhysRevLett.108.151803>. arXiv:1203.0275 [hep-ex]
84. CDF, D0 Collaboration, T.A. Aaltonen et al., Combination of CDF and D0 W-boson mass measurements. *Phys. Rev. D* **88**(5), 052018 (2013). <https://doi.org/10.1103/PhysRevD.88.052018>. arXiv:1307.7627 [hep-ex]
85. S. Heinemeyer, W. Hollik, G. Weiglein, L. Zeune, Implications of LHC search results on the W boson mass prediction in the MSSM. *JHEP* **12**, 084 (2013). [https://doi.org/10.1007/JHEP12\(2013\)084](https://doi.org/10.1007/JHEP12(2013)084). arXiv:1311.1663 [hep-ph]
86. ATLAS Collaboration, M. Aaboud et al., Measurement of the W-boson mass in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector. *Eur. Phys. J. C* **78**(2), 110 (2018). <https://doi.org/10.1140/epjc/s10052-017-5475-4>. arXiv:1701.07240 [hep-ex]. [Erratum: *Eur.Phys.J.C* **78**, 898 (2018)]
87. LHCb Collaboration, R. Aaij et al., Measurement of the W boson mass. *JHEP* **01** (2022) 036. [https://doi.org/10.1007/JHEP01\(2022\)036](https://doi.org/10.1007/JHEP01(2022)036). arXiv:2109.01113 [hep-ex]
88. P. Athron, M. Bach, D.H.J. Jacob, W. Kotlarski, D. Stöckinger, A. Voigt, Precise calculation of the W boson pole mass beyond the Standard Model with FlexibleSUSY. arXiv:2204.05285 [hep-ph]
89. P. Athron, J.-H. Park, D. Stöckinger, A. Voigt, FlexibleSUSY—a meta spectrum generator for supersymmetric models. *Nucl. Part. Phys. Proc.* **273–275**, 2424–2426 (2016). <https://doi.org/10.1016/j.nuclphysbps.2015.09.413>. arXiv:1410.7385 [hep-ph]
90. P. Athron, M. Bach, D. Harries, T. Kwasnitza, J.-H. Park, D. Stöckinger, A. Voigt, J. Ziebell, FlexibleSUSY 2.0: extensions to investigate the phenomenology of SUSY and non-SUSY models. *Comput. Phys. Commun.* **230**, 145–217 (2018). <https://doi.org/10.1016/j.cpc.2018.04.016>. arXiv:1710.03760 [hep-ph]
91. P. Diessner, G. Weiglein, Precise prediction for the W boson mass in the MRSSM. *JHEP* **07**, 011 (2019). [https://doi.org/10.1007/JHEP07\(2019\)011](https://doi.org/10.1007/JHEP07(2019)011). arXiv:1904.03634 [hep-ph]
92. P. Fayet, Supergauge invariant extension of the Higgs mechanism and a model for the electron and its neutrino. *Nucl. Phys. B* **90**, 104–124 (1975). [https://doi.org/10.1016/0550-3213\(75\)90636-7](https://doi.org/10.1016/0550-3213(75)90636-7)
93. J. Braathen, M.D. Goodsell, S. Paßehr, E. Pinsard, Expectation management. *Eur. Phys. J. C* **81**(6), 498 (2021). <https://doi.org/10.1140/epjc/s10052-021-09285-4>. arXiv:2103.06773 [hep-ph]
94. M. Carena, H.E. Haber, I. Low, N.R. Shah, C.E.M. Wagner, Complementarity between nonstandard Higgs boson searches and precision Higgs boson measurements in the MSSM. *Phys. Rev. D* **91**(3), 035003 (2015). <https://doi.org/10.1103/PhysRevD.91.035003>. arXiv:1410.4969 [hep-ph]
95. M. Du, Z. Liu, P. Nath, CDF W mass anomaly from a dark sector with a Stueckelberg–Higgs portal. arXiv:2204.09024 [hep-ph]
96. ALEPH Collaboration, A. Heister et al., Search for charginos nearly mass degenerate with the lightest neutralino in e^+e^- collisions at center-of-mass energies up to 209-GeV. *Phys. Lett. B* **533**, 223–236 (2002). [https://doi.org/10.1016/S0370-2693\(02\)01584-8](https://doi.org/10.1016/S0370-2693(02)01584-8). arXiv:hep-ex/0203020
97. J. Abdallah et al., Searches for supersymmetric particles in e^+e^- collisions up to 208 GeV, and interpretation of the results within the MSSM. <https://inspirehep.net/literature/1660152>
98. LEP2 SUSY Working Group, ALEPH, DELPHI, L3 and OPAL experiments, note LEPSUSYWG/02-04.1. <http://lepsusy.web.cern.ch/lepsusy/Welcome.html>. Accessed 10 Aug 2022
99. P. Bechtel, O. Brein, S. Heinemeyer, G. Weiglein, K.E. Williams, HiggsBounds: confronting arbitrary Higgs sectors with exclusion bounds from LEP and the tevatron. *Comput. Phys. Commun.* **181**, 138–167 (2010). <https://doi.org/10.1016/j.cpc.2009.09.003>. arXiv:0811.4169 [hep-ph]
100. P. Bechtel, O. Brein, S. Heinemeyer, G. Weiglein, K.E. Williams, HiggsBounds 2.0.0: confronting neutral and charged Higgs sector predictions with exclusion bounds from LEP and the tevatron. *Comput. Phys. Commun.* **182**, 2605–2631 (2011). <https://doi.org/10.1016/j.cpc.2011.07.015>. arXiv:1102.1898 [hep-ph]
101. P. Bechtel, O. Brein, S. Heinemeyer, O. Stål, T. Stefaniak, G. Weiglein, K.E. Williams, *HiggsBounds* – 4: improved tests of extended Higgs sectors against exclusion bounds from LEP, the tevatron and the LHC. *Eur. Phys. J. C* **74**(3), 2693 (2014). <https://doi.org/10.1140/epjc/s10052-013-2693-2>. arXiv:1311.0055 [hep-ph]

102. P. Bechtle, D. Dercks, S. Heinemeyer, T. Klingl, T. Stefaniak, G. Weiglein, J. Wittbrodt, HiggsBounds-5: testing Higgs sectors in the LHC 13 TeV era. *Eur. Phys. J. C* **80**(12), 1211 (2020). <https://doi.org/10.1140/epjc/s10052-020-08557-9>. [arXiv:2006.06007](https://arxiv.org/abs/2006.06007) [hep-ph]
103. M. Goodsell, K. Nickel, F. Staub, Generic two-loop Higgs mass calculation from a diagrammatic approach. *Eur. Phys. J. C* **75**(6), 290 (2015). <https://doi.org/10.1140/epjc/s10052-015-3494-6>. [arXiv:1503.03098](https://arxiv.org/abs/1503.03098) [hep-ph]
104. J. Braathen, M.D. Goodsell, Avoiding the Goldstone boson catastrophe in general renormalisable field theories at two loops. *JHEP* **12**, 056 (2016). [https://doi.org/10.1007/JHEP12\(2016\)056](https://doi.org/10.1007/JHEP12(2016)056). [arXiv:1609.06977](https://arxiv.org/abs/1609.06977) [hep-ph]
105. J. Braathen, M.D. Goodsell, F. Staub, Supersymmetric and non-supersymmetric models without catastrophic Goldstone bosons. *Eur. Phys. J. C* **77**(11), 757 (2017). <https://doi.org/10.1140/epjc/s10052-017-5303-x>. [arXiv:1706.05372](https://arxiv.org/abs/1706.05372) [hep-ph]
106. F. Staub, W. Porod, Improved predictions for intermediate and heavy Supersymmetry in the MSSM and beyond. *Eur. Phys. J. C* **77**(5), 338 (2017). <https://doi.org/10.1140/epjc/s10052-017-4893-7>. [arXiv:1703.03267](https://arxiv.org/abs/1703.03267) [hep-ph]
107. P. Slavich et al., Higgs-mass predictions in the MSSM and beyond. *Eur. Phys. J. C* **81**(5), 450 (2021). <https://doi.org/10.1140/epjc/s10052-021-09198-2>. [arXiv:2012.15629](https://arxiv.org/abs/2012.15629) [hep-ph]
108. M.D. Goodsell, A. Joury, Active learning BSM parameter spaces. [arXiv:2204.13950](https://arxiv.org/abs/2204.13950) [hep-ph]
109. P. Bechtle, S. Heinemeyer, O. Stål, T. Stefaniak, G. Weiglein, *HiggsSignals*: confronting arbitrary Higgs sectors with measurements at the tevatron and the LHC. *Eur. Phys. J. C* **74**(2), 2711 (2014). <https://doi.org/10.1140/epjc/s10052-013-2711-4>. [arXiv:1305.1933](https://arxiv.org/abs/1305.1933) [hep-ph]
110. P. Bechtle, S. Heinemeyer, T. Klingl, T. Stefaniak, G. Weiglein, J. Wittbrodt, HiggsSignals-2: probing new physics with precision Higgs measurements in the LHC 13 TeV era. *Eur. Phys. J. C* **81**(2), 145 (2021). <https://doi.org/10.1140/epjc/s10052-021-08942-y>. [arXiv:2012.09197](https://arxiv.org/abs/2012.09197) [hep-ph]