



## Heterolytic O–O bond cleavage upon single electron transfer to a nonheme Fe(III)–OOH complex

Antoine Bohn, Katell Sénéchal-David, Jean-Noël Rebilly, Christian Herrero, Winfried Leibl, Elodie Anxolabéhère-Mallart, Frédéric Banse

### ► To cite this version:

Antoine Bohn, Katell Sénéchal-David, Jean-Noël Rebilly, Christian Herrero, Winfried Leibl, et al.. Heterolytic O–O bond cleavage upon single electron transfer to a nonheme Fe(III)–OOH complex. Chemistry - A European Journal, 2022, 2022, pp.e202201. 10.1002/chem.202201600 . hal-03766585

**HAL Id: hal-03766585**

**<https://hal.science/hal-03766585>**

Submitted on 1 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Heterolytic O-O Bond Cleavage Upon Single Electron Transfer to a Nonheme Fe(III)-OOH Complex.

Antoine Bohn,<sup>[a]</sup> Katell Sénéchal-David,<sup>[a]</sup> Jean-Noël Rebilly,<sup>[a]</sup> Christian Herrero,<sup>[a]</sup> Winfried Leibl,<sup>[b]</sup> Elodie Anxolabéhère-Mallart,<sup>\*[c]</sup> Frédéric Banse<sup>\*[a]</sup>

[a] Dr. A. Bohn, Dr. K. Sénéchal-David, Dr. J.-N. Rebilly, Dr. C. Herrero, Pr. F. Banse  
Université Paris-Saclay, CNRS, Institut de Chimie Moléculaire et des Matériaux d'Orsay, 91405, Orsay, France.  
E-mail: frederic.banse@universite-paris-saclay.fr

[b] Dr. W. Leibl  
Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France

[c] Dr. E. Anxolabéhère-Mallart  
Université Paris Cité, CNRS, Laboratoire d'Electrochimie Moléculaire, F-75013 Paris, France

Supporting information for this article is given via a link at the end of the document.

**Abstract:** The one-electron reduction of the nonheme iron(III)-hydroperoxo complex,  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  ( $\text{L}_5^2 = N$ -methyl- $N,N',N'$ -tris(2-pyridylmethyl)ethane-1,2-diamine), carried out at  $-70^\circ\text{C}$  results in the release of dioxygen and in the formation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  following a bimolecular process. This reaction can be performed either with cobaltocene as chemical reductant, or electrochemically. These experimental observations are consistent with the disproportionation of the hydroperoxo group in the putative  $\text{Fe}^{\text{II}}(\text{OOH})$  intermediate generated upon reduction of the  $\text{Fe}^{\text{III}}(\text{OOH})$  starting complex. One plausible mechanistic scenario is that this disproportionation reaction follows an O-O heterolytic cleavage pathway via a  $\text{Fe}^{\text{IV}}$ -oxo species.

## Introduction

Oxygen is used as direct oxidant by heme and nonheme oxygenases to carry out a variety of biochemical transformations of organic metabolites. Thanks to extensive studies, the mechanism by which cytochromes P450 activate dioxygen to transfer one oxygen atom to their substrate while releasing the other as water has been precisely determined and serves as a paradigm. Along their catalytic cycle, these enzymes first generate a  $\text{Fe}^{\text{III}}$ -superoxo ( $\text{Fe}^{\text{III}}(\text{OO}\cdot)$ ) intermediate resulting from coordination of  $\text{O}_2$  to the reduced  $\text{Fe}^{\text{II}}$  active site. Electron and proton transfer to this  $\text{Fe}^{\text{III}}(\text{OO}\cdot)$  species yields a  $\text{Fe}^{\text{III}}$ -hydroperoxo ( $\text{Fe}^{\text{III}}(\text{OOH})$ ) complex which evolves via heterolytic O-O cleavage into Compound I, a formally  $\text{Fe}^{\text{V}}$ -oxo ( $\text{Fe}^{\text{V}}(\text{O})$ ) as the catalytic active species for the target reaction, frequently being an oxygen atom transfer to the substrate.<sup>[1-3]</sup> A similar Compound I is generated following the heterolytic activation of hydrogen peroxide by heme catalase but it is directed towards the oxidation of a second equivalent of  $\text{H}_2\text{O}_2$ , leading to an overall disproportionation of hydrogen peroxide into dioxygen and water.<sup>[4]</sup> In contrast, the exact mechanism of nonheme Mn Catalases is precluded due to their fast kinetics, but it is admitted that their dinuclear active site cycles between the  $\text{Mn}^{\text{II}}_2$  and  $\text{Mn}^{\text{III}}_2$  states for  $\text{H}_2\text{O}_2$  removal, even though higher oxidation states are accessible.<sup>[5]</sup> Nonetheless, the implication of higher valent intermediates is supported by studies with synthetic functional models of Mn Catalases.<sup>[5-8]</sup> Therefore, it is not surprising that disproportionation of hydrogen peroxide is a common competitive reaction in  $\text{H}_2\text{O}_2$ -dependent hydrocarbon oxidations catalyzed by metal complexes.<sup>[9-11]</sup>


It is now well-documented that the reaction of nonheme iron complexes with hydrogen peroxide generates  $\text{Fe}^{\text{III}}(\text{OOH})$  intermediates.<sup>[12-14]</sup> Studies performed with  $(N_4)\text{Fe}^{\text{III}}(\text{OOH})$  complexes have demonstrated that the heterolytic O-O cleavage is favored when the tetradentate  $N_4$  ligand leaves two *cis* labile positions in the coordination sphere of the metal center,<sup>[15-18]</sup> thus leading to *cis*- $\text{Fe}^{\text{V}}(\text{O})(\text{OH})$  species,<sup>[19-21]</sup> which are efficient metal-based oxidants.<sup>[22]</sup> In contrast,  $(N_5)\text{Fe}^{\text{III}}(\text{OOH})$  intermediates evolve via homolytic O-O cleavage yielding  $\text{Fe}^{\text{IV}}(\text{O}) + \text{OH}\cdot$  species.<sup>[11,23-25]</sup> Interestingly, several studies report that the reduction of  $(N_4)\text{Fe}^{\text{III}}$ -hydroperoxo,  $(N_4)\text{Fe}^{\text{III}}$ -alkylperoxo<sup>[26]</sup> or  $(N_4)\text{Fe}^{\text{III}}$ -peroxo binding a non-redox cation<sup>[27]</sup> (where  $N_4$  is a macrocyclic ligand of the tetramethylcyclam family) generates a  $(N_4)\text{Fe}^{\text{IV}}(\text{O})$  species, suggesting an heterolytic cleavage occurs after reduction. Along the same line, indirect evidence for inter-related  $\text{Fe}^{\text{II}}(\text{OOH})$  /  $\text{Fe}^{\text{IV}}(\text{O})(\text{OH})$  species have also been reported.<sup>[28-30]</sup>

We have analyzed both the one-electron chemical and electrochemical reduction of the non-porphyrinic  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  intermediate stabilized at  $-70^\circ\text{C}$  ( $\text{L}_5^2 = N$ -methyl- $N,N',N'$ -tris(2-pyridylmethyl)ethane-1,2-diamine, Figure 1), and report in this paper that it leads to the release of  $\text{O}_2$  and the formation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  in a bimolecular process. This reaction formally corresponds to the disproportionation of the hydroperoxo group, which suggests that the transient  $\text{Fe}^{\text{II}}(\text{OOH})$  species generated upon reduction of the  $\text{Fe}^{\text{III}}(\text{OOH})$  complex evolves via heterolytic O-O cleavage. Alternative mechanistic scenarios following the initial electron transfer are discussed.

## Results and Discussion

In order to investigate the redox properties of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$ , the complex was prepared and isolated following the method we reported previously (Supporting Information, Experimental Section).<sup>[23,31]</sup>  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  was formed in MeOH following reaction of  $[\text{Fe}^{\text{II}}\text{Cl}(\text{L}_5^2)](\text{PF}_6)$  with an excess (100 equiv.) of  $\text{H}_2\text{O}_2$ , and precipitated at  $-80^\circ\text{C}$  upon addition of cold  $\text{Et}_2\text{O}$ . The purple solid,  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)](\text{PF}_6)_2$ , was isolated by removing the solvent and the excess of  $\text{H}_2\text{O}_2$  via a cannula equipped with a glass filter. The obtained solid was carefully washed with a large volume of cold  $\text{Et}_2\text{O}$  and redissolved in butyronitrile (PrCN) at  $-80^\circ\text{C}$  yielding stock solutions with concentrations ranging between 1 and 4 mM as quantified by both the hydroperoxo-to-


$\text{Fe}^{\text{III}}$  charge transfer band centered at 530 nm ( $1000 \text{ M}^{-1}\text{cm}^{-1}$ )<sup>[12,32]</sup> and double integration of the X band EPR signal of this low spin ( $S=1/2$ ) complex (Figure 1 and Supporting Information, Figure S1 for assignment). All experiments were performed at low temperature ( $< -50^\circ\text{C}$ ) to minimize the degradation of the  $\text{Fe}^{\text{III}}(\text{OOH})$  intermediate.



**Figure 1.** (A) Electronic absorption spectrum of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)](\text{PF}_6)_2$  with its schematic representation (1 mM, PrCN + 0.2 M TBAPF<sub>6</sub> at  $-70^\circ\text{C}$ ). (B) X-band EPR spectrum of the same sample recorded at 90 K. The inset shows the full spectral range (0–5000 Gauss). The complete assignment is given in the Supporting Information, Figure S1. Microwave frequency 9.65 GHz, microwave power 1.0 mW, modulation amplitude 8 G, gain 50 dB, modulation frequency 100 kHz.


The cyclic voltammogram (CV) of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  exhibits a cathodic peak at  $E_{\text{p,c}} = -0.20 \text{ V}$  vs SCE (Figure 2), a value consistent with the reduction potential measured for the similar  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{TPEN})]^{2+}$  complex,  $E_{\text{p,c}} = -0.08 \text{ V}$  vs SCE.<sup>[33–35]</sup> A second cathodic peak is observed at  $-0.94 \text{ V}$  vs SCE, which is assigned to the reduction of  $\text{O}_2$  into  $\text{O}_2^-$  (Supporting Information, Figure S2). Interestingly, comparison between the CVs of independently prepared samples reveals that the intensity of the cathodic peak of  $\text{O}_2$  is dependent on the intensity of the  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  peak (Supporting Information, Figure S2), thus indicating that the presence of  $\text{O}_2$  is a consequence of the reduction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$ . On the reverse scan, the anodic peak detected at  $E_{\text{p,a}} = 0.62 \text{ V}$  vs SCE is assigned to the oxidation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$ , in agreement with the CV of  $[\text{Fe}^{\text{II}}(\text{MeCN})(\text{L}_5^2)]^{2+}$  recorded under alkaline conditions (Supporting Information, Figure S3) and reported data.<sup>[33,36–38]</sup> Note that the  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  oxidation wave is observed only when the sample is first scanned

towards negative potentials (Figure 2 and Supporting Information, Figure S2A). Hence,  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  and  $\text{O}_2$  are formed as a result of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  reduction.




**Figure 2.** Cyclic Voltammogram of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)](\text{PF}_6)_2$  (1 mM) in PrCN + 0.2 M TBAPF<sub>6</sub> at  $-70^\circ\text{C}$ ,  $0.1 \text{ V}\cdot\text{s}^{-1}$ . The arrows indicate the initial direction of the scan.

To confirm this hypothesis, this reaction was carried out using cobaltocene ( $E^\circ = -0.95 \text{ V}$  vs SCE)<sup>[39]</sup> as a single electron reductant and followed by UV-visible spectroscopy (Figure 3A). Immediately after the introduction of cobaltocene (1 equiv vs Fe), the 530-nm band rapidly decreased in intensity and vanished while an absorption at ca. 390 nm progressively developed (Figure 3A). This 390-nm band is assigned to a  $t_{2g}(\text{Fe}^{\text{II}})$ -to- $\pi^*(\text{pyridine})$  charge transfer transition consistent with the formation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  (Supporting Information, Figure S3A). Aliquots from this reaction mixture were taken and analyzed by X-band EPR spectroscopy in parallel mode detection. The progressive formation of  $\text{O}_2$  following the reduction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  was confirmed by detecting its characteristic signature over the 0–5000 G spectral window (Figure 3B and Supporting Information, Figure S4).<sup>[40]</sup> UV/vis experiments indicate that the complete reduction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  requires 1 equivalent of reductant. As seen in Figure 3A, the extinction of the  $\text{Fe}^{\text{III}}(\text{OOH})$  LMCT at 530 nm is achieved before the MLCT of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  at 390 nm stops developing. This observation indicates that the reduction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  is faster than the formation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$ , and as a result, of  $\text{O}_2$ , and suggests that these species are formed in a secondary step (Figure 3A).  $\text{O}_2$  production was also confirmed by locating a fluorescent probe in the headspace of a UV-visible cuvette (Supporting Information, Figure S5). Introduction of cobaltocene in a solution of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  led to a decrease of the fluorescence lifetime of the probe chromophore. This observation indicates that the triplet excited state of the chromophore is quenched by reaction with the triplet state of released dioxygen. Performing the reaction by introducing only 0.5 equiv. reductant vs the initial amount of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  intermediate followed by another portion of 0.5 equiv. reductant after the initial release of  $\text{O}_2$  resulted in the production of a second identical amount of  $\text{O}_2$ . To summarize, the 1:1 reaction between  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  and the one electron reductant results in the formation of  $\text{O}_2$  and  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$ .





**Figure 3.** (A) UV-visible monitoring of the reaction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)](\text{PF}_6)_2$  ( $-50^\circ\text{C}$ ,  $0.15\text{ mM}$  in  $\text{PrCN}/\text{MeCN}$   $1:1$ , blue trace) with  $1\text{ equiv. Co}(\text{Cp})_2$  (black to grey to red trace). The inset shows the time traces of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  disappearance at  $530\text{ nm}$  (blue) and  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  formation at  $390\text{ nm}$  (red). (B) Parallel mode X-band EPR spectra at  $10\text{ K}$  of aliquots taken from the same reaction mixture (with corresponding color lines). Microwave frequency  $9.38\text{ GHz}$ , microwave power  $2.0\text{ mW}$ , modulation amplitude  $8\text{ G}$ , gain  $50\text{ dB}$ , modulation frequency  $100\text{ kHz}$ , temperature  $10\text{ K}$ . Experiments were done in the presence of  $2\text{ equivalents}$  of potassium tetrakis(pentafluorophenyl)borate to avoid cobaltocenium precipitation.

Formally, the generation of  $\text{O}_2$  and  $\text{HO}^-$  (*i. e.*  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$ ) corresponds to the disproportionation of the hydroperoxo ligand upon reduction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$ , which can be summarized with the reaction proposed in Equation 1.




By comparison, addition of cobaltocene to  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  (obtained by deprotonation of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$ , see Supporting Information, Experimental Section)<sup>[33,41,42]</sup> led to the disappearance of the iron-peroxo complex but did not lead to the evolution of  $\text{O}_2$  (Supporting Information, Figure S6).<sup>[43]</sup>

With  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  being susceptible to exogenous ligand exchange in solution (Supporting Information, Figure S3), its amount is difficult to determine. Hence, in order to validate the stoichiometry of the reaction written in Equation 1 we estimated the quantity of  $\text{O}_2$  produced by simulating the CV of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  (Supporting Information, Electrochemical Methodology and Figure S16).<sup>[44]</sup> To obtain this information the overall reaction described in Equation 1 was decomposed into electrochemical and chemical steps (Equations 2 and 3). The  $\text{O}_2 / \text{O}_2^{\cdot-}$  redox couple, which is expected to contribute to the global current, was considered as well (Equation 4).



A good agreement between the experimental CV and the simulation resulting from equations 2-4 (Supporting Information, Figure S16) supports this series of reactions and confirms the global reaction (Equation 1) in which the one electron reduction of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  generates one equivalent of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  and half an equivalent of  $\text{O}_2$ .

Further support for the reaction in Equation 1 was obtained by analyzing the absorbance vs time trace at  $390\text{ nm}$  (Figure 3A, inset). This trace representing the formation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$ , was fitted to second-order kinetics, as suggested by Equation 1, and to first order kinetics for comparison. As shown in Supporting Information, Figure S7, the agreement with the second order rate law is excellent, whereas first-order kinetics appear inconsistent. Monitoring this reaction at lower  $T$  ( $-60^\circ\text{C}$ ) and different initial concentrations of  $[\text{Fe}^{\text{III}}(\text{OOH})(\text{L}_5^2)]^{2+}$  confirmed that the formation of  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$  is a second order reaction (Supporting Information, Figure S8). Additionally, the observation that the complete reduction of the  $\text{Fe}^{\text{III}}(\text{OOH})$  intermediate occurs before the formation of the  $\text{Fe}^{\text{II}}(\text{OH})$  complex is achieved (compare blue and red traces in Figure 3A, inset) suggests that after the one-electron reduction process, the generated  $[\text{Fe}^{\text{II}}(\text{OOH})(\text{L}_5^2)]^+$  engages in a global bimolecular rate determining reaction to produce  $\text{O}_2$  and  $[\text{Fe}^{\text{II}}(\text{OH})(\text{L}_5^2)]^+$ . Overall, these observations also imply that the O-O bond of the peroxide must be cleaved heterolytically at one point to ultimately produce the  $\text{Fe}^{\text{II}}(\text{OH})$  complex.



**Scheme 1.** Possible mechanisms accounting for the disproportionation reaction of the peroxo group without (A) or with (B) accumulation of a high-valent iron-oxo intermediate.

Two alternative mechanisms can be evoked to account for the disproportionation of the peroxide. Either a dimerization step of the generated  $\text{Fe}^{\text{II}}(\text{OOH})$  occurs first, followed by O-O bond cleavage; or the O-O bond cleavage occurs in the early steps of the mechanism yielding a high-valent iron-oxo species which reacts in a bimolecular process in the later stages.

In proposition A (Scheme 1 A and Supporting Information, Figure S9), the  $\text{Fe}^{\text{II}}(\text{OOH})$  species would engage in a rate-determining bimolecular reaction to form a (speculative) diferrous species with

$\mu$ -1,2-peroxo and  $\mu$ -1,1-hydroperoxo bridging ligands which would release a hydroxide and generate a ( $\mu$ -1,2-peroxo)diferric intermediate. This latter intermediate would finally release O<sub>2</sub> and two Fe<sup>II</sup>(OH) complexes. Such a mechanistic scenario is reminiscent to that of Mn Catalases with two metal centers cycling between the +II and +III oxidation states.

In this proposition, the uncommon  $\mu$ -1,1 binding mode of one peroxo ligand has recently been observed in the Nonheme N<sup>□</sup> Oxygenase CmlI.<sup>[45]</sup> A key intermediate in the catalytic cycle of  $\Delta^9$  Desaturase has also been proposed to display such a motif.<sup>[46]</sup>

Alternatively, in proposition B (Scheme 1 A and Supporting Information, Figure S10), the heterolytic cleavage of the putative Fe<sup>II</sup>(OOH) intermediate would result in the formation of Fe<sup>IV</sup>(O) and an OH<sup>-</sup> which we propose to bind the high valent metal center upon substitution of one of the (L<sub>5</sub><sup>2</sup>) pyridine groups. The nascent [Fe<sup>IV</sup>(O)(OH)(L<sub>5</sub><sup>2</sup>)]<sup>+</sup> would then engage in a rate-determining reaction to form a ( $\mu$ -1,2-peroxo)diferric intermediate, which would produce O<sub>2</sub> and two equivalents of [Fe<sup>II</sup>(OH)(L<sub>5</sub><sup>2</sup>)]<sup>2+</sup> in the final step.

For this latter proposition, which relies on the formation of a Fe<sup>IV</sup>(O)(OH) intermediate upon reduction of [Fe<sup>III</sup>(OOH)(L<sub>5</sub><sup>2</sup>)]<sup>2+</sup>, consistent examples can be found in the literature. Bang et al.<sup>[47]</sup> and Bae et al.<sup>[48]</sup> demonstrated that the reduction of [Fe<sup>III</sup>(OO)(TMC)]<sup>+</sup>-M<sup>n+</sup> (where M<sup>n+</sup> is a hard Lewis acid) led to [Fe<sup>IV</sup>(O)(TMC)]<sup>2+</sup> upon heterolytic cleavage of the peroxo group. Heterolytic O-O cleavage of Fe<sup>II</sup>-H<sub>2</sub>O<sub>2</sub> adducts was shown to be promoted by the presence of a general base which shuttles the proximal proton to the distal oxygen to generate Fe<sup>IV</sup>(O) while releasing H<sub>2</sub>O.<sup>[49-51]</sup> Paine and coworkers also proposed the formation of a reactive Fe<sup>IV</sup>(O)(OH) species from a putative Fe<sup>II</sup>(OOH) intermediate formed by oxygenation of the [(TpPh<sub>2</sub>)Fe<sup>II</sup>(benzilate)] complex and decarboxylation of benzilate.<sup>[28]</sup> Another key feature of this mechanism is the formation of the O-O bond between the two Fe<sup>IV</sup>(O)(OH) entities to yield a peroxo-bridged diiron(III) intermediate. A reversible equilibrium between such type of intermediates has been reported by Kodera et al.<sup>[52,53]</sup> Furthermore, formation of O-O bonds between two Fe<sup>IV</sup>(O) units in a hexanuclear iron complex supported on a stannoxane core was proposed by Ray and coworkers.<sup>[54]</sup> These authors suggested that the O-O bond was readily formed *via* radical coupling yielding a peroxo bridged diiron(III) species. A radical coupling step has also been proposed in the case of O<sub>2</sub> production catalyzed by a ruthenium complex.<sup>[55]</sup> Using both of these mechanisms, we performed a complete kinetic analysis based on the decay of [Fe<sup>III</sup>(OOH)(L<sub>5</sub><sup>2</sup>)]<sup>2+</sup> at 530 nm and the formation of [Fe<sup>II</sup>(OH)(L<sub>5</sub><sup>2</sup>)]<sup>+</sup> at 390 nm (Supporting Information, Figures S9 & S10). The results obtained are consistent with both proposition. In each mechanism, only the species which is engaged in the bimolecular rate determining step accumulates to a significant amount *i.e.* Fe<sup>II</sup>(OOH) in the first case (Supporting Information, Figure S9) and the Fe<sup>IV</sup>(O)(OH) complex in the second case (Supporting Information, Figure S10).

A closer inspection of the UV-visible monitoring of the reaction in the NIR region reveals the growth and decay of a broad absorption around 900 nm with an apparent extinction coefficient of ca. 250 M<sup>-1</sup>cm<sup>-1</sup> (Supporting Information, Figure S11). To our knowledge, no Fe<sup>II</sup>(OOH) species has been spectroscopically characterized yet and only one Fe<sup>IV</sup>(O)(OH) complex, namely [Fe<sup>IV</sup>(O)(TMC)(OH)]<sup>+</sup>, has been reported.<sup>[56,57]</sup> This complex exhibited two weak absorptions at 830 (100 M<sup>-1</sup>cm<sup>-1</sup>) and 1060 nm (110 M<sup>-1</sup>cm<sup>-1</sup>), which is similar to what we observed. Based on

proposition B, implying the transient formation of a Fe<sup>IV</sup>(O)(OH) species, we could extract by a global kinetic fit species spectra corresponding to the initial Fe<sup>III</sup>(OOH) complex, the final Fe<sup>II</sup>(OH) and the proposed transient Fe<sup>IV</sup>(O)(OH) species, supporting the heme catalase-based mechanism as the most likely scenario. Note that [Fe<sup>IV</sup>(O)(L<sub>5</sub><sup>2</sup>)]<sup>2+</sup>, namely the Fe<sup>IV</sup>(O) species supported by the *pentadentate* L<sub>5</sub><sup>2</sup> ligand, displays an absorption band at 730 nm (300 M<sup>-1</sup>cm<sup>-1</sup>) and a much longer lifetime than the fast decaying species observed at 900 nm. Thus, this latter species cannot be formulated to be the same.

While the stoichiometry of the global disproportionation of the peroxide has been demonstrated, there is still a question regarding the exact mechanism. An important progress would be to determine whether a high valent iron-oxo species indeed forms upon heterolytic cleavage of the O-O bond, by intercepting it with a substrate. Additionally, confirming (with labelling studies for example) that the O-O bond formation results from coupling between two iron-oxo species would be significant. However, these studies are made difficult by the instability of the reagent and the stoichiometric nature of the reaction.

## Conclusion

The chemical and electrochemical single-electron reduction of [Fe<sup>III</sup>(OOH)(L<sub>5</sub><sup>2</sup>)]<sup>2+</sup> at low temperature (-50°C — -70°C) yields a putative Fe<sup>II</sup>(OOH) species which disproportionates to give one equivalent of Fe<sup>II</sup>(OH) complex and half an equivalent of O<sub>2</sub>, indicating that the Fe<sup>II</sup>(OOH) species evolves *via* heterolytic O-O cleavage. To account for the global reaction, two kinetically valuable alternative mechanisms can be envisioned. One involves dinuclear Fe<sup>II</sup><sub>2</sub> and Fe<sup>III</sup><sub>2</sub> species as key intermediates, in a mechanism which is reminiscent of Mn Catalases. The other one involves the generation of a Fe<sup>IV</sup>(O)(OH) intermediate and the coupling between two equivalents of this latter species to release O<sub>2</sub>. This latter proposition is supported by the transient formation of a chromophore which we tentatively assign to such high valent iron(IV)-oxo species. The determination of the exact mechanism requires to identify the precise nature of this intermediate. Studies in that direction are under investigation in our laboratories.

## Acknowledgements

This work was supported by the LABEX CHARMMMAT (ANR-11-LABX-0039). Dr Zakaria Halime is thanked for his kind help.


**Keywords:** Fe complex • O-O bond breaking • mechanism • N ligand • Rieske dioxygenases

- [1] B. Meunier, S. P. de Visser, S. Shaik, *Chem. Rev.* **2004**, *104*, 3947–3980.
- [2] I. G. Denisov, T. M. Makris, S. G. Sligar, I. Schlichting, *Chem. Rev.* **2005**, *105*, 2253–2278.
- [3] T. L. Poulos, *Chem. Rev.* **2014**, *114*, 3919–3962.
- [4] P. Nicholls, *Arch. Biochem. Biophys.* **2012**, *525*, 95–101.
- [5] S. Signorella, C. Hureau, *Coord. Chem. Rev.* **2012**, *256*, 1229–1245.
- [6] Y. G. Abashkin, S. K. Burt, *Journal of Physical Chemistry B* **2004**, *108*, 2708–2711.
- [7] Y. G. Abashkin, S. K. Burt, *Inorg. Chem.* **2005**, *44*, 1425–1432.
- [8] M. L. Merlini, G. J. P. Britovsek, M. Swart, P. Belanzoni, *ACS Catal.* **2018**, *8*, 2944–2958.
- [9] C. Zondervan, R. Hage, Ben L. Feringa, *Chem. Commun.* **1997**, 419–420.

- [10] J. J. Dong, D. Unjaroen, F. Mecozzi, E. C. Harvey, P. Saisaha, D. Pijper, J. W. de Boer, P. Alsters, B. L. Feringa, W. R. Browne, *ChemSusChem* **2013**, *6*, 1774–1778.
- [11] J. Chen, A. Draksharapu, D. Angelone, D. Unjaroen, S. K. Padamati, R. Hage, M. Swart, C. Duboc, W. R. Browne, *ACS Catal.* **2018**, *8*, 9665–9674.
- [12] J.-J. Girerd, F. Banse, A. J. Simaan, *Struct. Bond.* **2000**, *97*, 145–177.
- [13] M. Costas, M. P. Mehn, M. P. Jensen, L. Que, *Chem. Rev.* **2004**, *104*, 939–986.
- [14] K. P. Bryliakov, E. P. Talsi, *Coord. Chem. Rev.* **2014**, *276*, 73–96.
- [15] K. Chen, L. Que, *J. Am. Chem. Soc.* **2001**, *123*, 6327–6337.
- [16] K. Chen, M. Costas, J. Kim, A. K. Tipton, L. Que, *J. Am. Chem. Soc.* **2002**, *124*, 3026–3035.
- [17] W. N. Oloo, A. J. Fielding, L. Que Jr., *J. Am. Chem. Soc.* **2013**, *135*, 6438–6441.
- [18] I. Prat, A. Company, V. Postils, X. Ribas, L. Que Jr., J. M. Luis, M. Costas, *Chem. Eur. J.* **2013**, *19*, 6724–6738.
- [19] S. Xu, J. J. Veach, W. N. Oloo, K. C. Peters, J. Wang, R. H. Perry, L. Que, *Chem. Commun.* **2018**, *54*, 8701–8704.
- [20] I. Prat, J. S. Mathieson, M. Güell, X. Ribas, J. M. Luis, L. Cronin, M. Costas, *Nature Chem* **2011**, *3*, 788–793.
- [21] M. Borrell, E. Andris, R. Navrátil, J. Roithová, M. Costas, *Nature Communications* **2019**, *10*, 901–9.
- [22] G. Olivo, O. Cussó, M. Borrell, M. Costas, *J Biol Inorg Chem* **2017**, *22*, 425–452.
- [23] A. Thibon, V. Jollet, C. Ribal, K. Sénéchal-David, L. Billon, A. B. Sorokin, F. Banse, *Chem. Eur. J.* **2012**, *18*, 2715–2724.
- [24] A. S. Faponle, M. G. Quesne, C. V. Sastri, F. Banse, S. P. de Visser, *Chem. Eur. J.* **2015**, *21*, 1221–1236.
- [25] J.-N. Rebilly, W. Zhang, C. Herrero, H. Dridi, K. Sénéchal-David, R. Guillot, F. Banse, *Chem. Eur. J.* **2020**, *26*, 659–668.
- [26] S. Bang, S. Park, Y.-M. Lee, S. Hong, K.-B. Cho, W. Nam, *Angew. Chem. Int. Ed.* **2014**, *53*, 7843–7847.
- [27] Y.-M. Lee, S. Bang, Y. M. Kim, J. Cho, S. Hong, T. Nomura, T. Ogura, O. Troeppner, I. Ivanović-Burmazović, R. Sarangi, et al., *Chem. Sci.* **2013**, *4*, 3917–7.
- [28] S. Paria, L. Que Jr., T. K. Paine, *Angew. Chem. Int. Ed.* **2011**, *50*, 11129–11132.
- [29] S. Paria, S. Chatterjee, T. K. Paine, *Inorg. Chem.* **2014**, *53*, 2810–2821.
- [30] S. Chatterjee, T. K. Paine, *Angew. Chem. Intl. Ed.* **2015**, *54*, 9338–9342.
- [31] M. Martinho, P. Dorlet, E. Rivière, A. Thibon, C. Ribal, F. Banse, J.-J. Girerd, *Chem. Eur. J.* **2008**, *14*, 3182–3188.
- [32] A. J. Simaan, S. Döpner, F. Banse, S. Bourcier, G. Bouchoux, A. Boussac, P. Hildebrandt, J.-J. Girerd, *Eur. J. Inorg. Chem.* **2000**, *2000*, 1627–1633.
- [33] N. Ségaud, E. Anxolabéhère-Mallart, K. Sénéchal-David, L. Acosta-Rueda, M. Robert, F. Banse, *Chem. Sci.* **2015**, *6*, 639–647.
- [34] From a simple electrostatic model based on the Born equation, it can be shown that the redox potential value of similar species increases with their size, as observed when going from  $L_5^2$  (Me substituent) to TPEN (pyridyl substituent). see reference 35.
- [35] J. Simaan, S. Poussereau, G. Blondin, J. J. Girerd, D. Defaye, C. Philouze, J. Guilhem, L. Tchertanov, *Inorg. Chim. Acta* **2000**, *299*, 221–230.
- [36] N. Ségaud, J.-N. Rebilly, K. Sénéchal-David, R. Guillot, L. Billon, J.-P. Baltaze, J. Farjon, O. Reinaud, F. Banse, *Inorg. Chem.* **2013**, *52*, 691–700.
- [37] P. Comba, H. Wadepohl, A. Waleska, *Aust. J. Chem.* **2014**, *67*, 398–404.
- [38] C. Herrero, A. Quaranta, M. Sircoglou, K. Sénéchal-David, A. Baron, I. M. Marín, C. Buron, J.-P. Baltaze, W. Leibl, A. Aukauloo, et al., *Chem. Sci.* **2015**, *6*, 2323–2327.
- [39] N. G. Connelly, W. E. Geiger, *Chem. Rev.* **1996**, *96*, 877–910.
- [40] W. R. Hagen, *Biomolecular EPR Spectroscopy*, CRC Press, **2008**.
- [41] A. J. Simaan, F. Banse, P. Mialane, A. Boussac, S. Un, T. Kargar-Grisel, G. Bouchoux, J. J. Girerd, *Eur. J. Inorg. Chem.* **1999**, 993–996.
- [42] A. J. Simaan, F. Banse, J.-J. Girerd, K. Wiegardt, E. Bill, *Inorg. Chem.* **2001**, *40*, 6538–6540.
- [43] This reaction and its products were not analyzed further.
- [44] The amount of dissolved dioxygen could not be estimated by analysis of the fluorescence quenching due to our particular experimental condions. See Supporting Information for details.
- [45] A. J. Jasniewski, A. J. Komor, J. D. Lipscomb, L. Que Jr., *J. Am. Chem. Soc.* **2017**, *139*, 10472–10485.
- [46] J. Chalupský, T. A. Rokob, Y. Kurashige, T. Yanai, E. I. Solomon, L. Rulišek, M. Srnc, *J. Am. Chem. Soc.* **2014**, *136*, 15977–15991.
- [47] S. Bang, Y.-M. Lee, S. Hong, K.-B. Cho, Y. Nishida, M. S. Seo, R. Sarangi, S. Fukuzumi, W. Nam, *Nature Chem* **2014**, *6*, 934–940.
- [48] S. H. Bae, Y.-M. Lee, S. Fukuzumi, W. Nam, *Angew. Chem. Int. Ed.* **2017**, *56*, 801–805.
- [49] F. Li, J. England, L. Que Jr., *J. Am. Chem. Soc.* **2010**, *132*, 2134–2135.
- [50] K. Cheaib, M. Q. E. Mubarak, K. Sénéchal-David, C. Herrero, R. Guillot, M. Clémancey, J.-M. Latour, S. P. de Visser, J.-P. Mahy, F. Banse, et al., *Angew. Chem. Int. Ed.* **2019**, *58*, 854–858.
- [51] A. Bohn, C. Chinaux-Chaix, K. Cheaib, R. Guillot, C. Herrero, K. Sénéchal-David, J.-N. Rebilly, F. Banse, *Dalton Trans.* **2019**, *48*, 17045–17051.
- [52] M. Kodera, Y. Kawahara, Y. Hitomi, T. Nomura, T. Ogura, Y. Kobayashi, *J. Am. Chem. Soc.* **2012**, *134*, 13236–13239.
- [53] M. Kodera, T. Tsuji, T. Yasunaga, Y. Kawahara, T. Hirano, Y. Hitomi, T. Nomura, T. Ogura, Y. Kobayashi, P. K. Sajith, Y. Shiota, K. Yoshizawa, *Chem. Sci.* **2014**, *5*, 2282–2292.
- [54] S. Kundu, E. Matito, S. Walleck, F. F. Pfaff, F. Heims, B. Rábay, J. M. Luis, A. Company, B. Braun, T. Glaser, K. Ray, *Chem. Eur. J.* **2012**, *18*, 2787–2791.
- [55] S. Romain, F. Bozoglian, X. Sala, A. Llobet, *J. Am. Chem. Soc.* **2009**, *131*, 2768–2769.
- [56] T. A. Jackson, J.-U. Rohde, M. S. Seo, C. V. Sastri, R. DeHont, A. Stubna, T. Ohta, T. Kitagawa, E. Münck, W. Nam, L. Que, *J. Am. Chem. Soc.* **2008**, *130*, 12394–12407.
- [57] J. E. M. N. Klein, A. Draksharapu, A. Shokri, C. J. Cramer, L. Que, *Chem. Eur. J.* **2018**, *24*, 5373–5378.

---

## Entry for the Table of Contents



The one-electron reduction of a nonheme Fe<sup>III</sup>(OOH) intermediate results in the release of the Fe<sup>II</sup>(OH) complex and O<sub>2</sub>, following a bimolecular reaction consistent with the disproportionation of the hydroperoxo ligand in a putative Fe<sup>II</sup>(OOH) species. One plausible mechanistic scenario is that this disproportionation reaction follows an O-O heterolytic cleavage pathway *via* a Fe<sup>IV</sup>-oxo species.