

**Appendix 3.** On the relationship between the newly proposed parametric measures of community dissimilarity and a number of classical single-point measures of resemblance.

Given two plots  $U$  and  $V$ , let  $x_{Uj}$  and  $x_{Vj}$  be the abundances of species  $j$  ( $j = 1, 2, \dots, N$ ) in both plots. First, note that as  $x_{Uj}$  and  $x_{Vj}$  are both nonnegative, we have for  $\alpha > 0$   $\sum_j |x_{Uj} - x_{Vj}|^\alpha \leq \sum_j (\max_j(x_{Uj}, x_{Vj}))^\alpha$  and thus  $\sum_j |x_{Uj} - x_{Vj}|^\alpha \leq \sum_j x_{Uj}^\alpha + \sum_j x_{Vj}^\alpha \leq \sum_j (x_{Uj} + x_{Vj})^\alpha$ , and for  $\alpha \geq 1$ ,  $\sqrt[\alpha]{\sum_j (x_{Uj} + x_{Vj})^\alpha} \leq \sqrt[\alpha]{\sum_j x_{Uj}^\alpha} + \sqrt[\alpha]{\sum_j x_{Vj}^\alpha}$  (Minkowski's inequality).

Therefore, the following inequalities are satisfied:  $D^\alpha \leq \Delta^\alpha \leq M^\alpha$  for  $\alpha > 0$  and  $L^\alpha \leq D^\alpha$  for  $\alpha \geq 1$  (for the definition of the parametric dissimilarities, see the main text).

For  $\alpha = 2$ , the dissimilarity index  $\Delta^\alpha$  is related to several resemblance measures published in the literature:

- Morisita-Horn similarity index (Morisita 1959; Horn 1966):

$$S_{MH} = \frac{\sum_j 2x_{Uj}x_{Vj}}{\left[ \left( \sum_j x_{Uj}^2 / X_U^2 \right) + \left( \sum_j x_{Vj}^2 / X_V^2 \right) \right] X_U X_V} \quad (\text{A1})$$

where  $X_U = \sum_j x_{Uj}$  and  $X_V = \sum_j x_{Vj}$  are the total abundances in plots  $U$  and  $V$ , respectively. The Morisita-Horn similarity index thus corresponds to

$$S_{MH} = \frac{\sum_j 2p_{Uj}p_{Vj}}{\sum_j p_{Uj}^2 + \sum_j p_{Vj}^2} \quad (\text{A2})$$

where  $x_{Uj}$  and  $x_{Vj}$  are the *relative* abundances of species  $j$  in plots  $U$  and  $V$ , respectively.

Accordingly, a dissimilarity index can be defined as  $D_{MH} = \sqrt{1 - S_{MH}}$ :

$$\begin{aligned} D_{MH} &= \sqrt{1 - \frac{\sum_j 2p_{Uj}p_{Vj}}{\sum_j p_{Uj}^2 + \sum_j p_{Vj}^2}} = \sqrt{\frac{\sum_j p_{Uj}^2 + \sum_j p_{Vj}^2 - \sum_j 2p_{Uj}p_{Vj}}{\sum_j p_{Uj}^2 + \sum_j p_{Vj}^2}} \\ &= \sqrt{\frac{\sum_j (p_{Uj} - p_{Vj})^2}{\sum_j p_{Uj}^2 + \sum_j p_{Vj}^2}} \end{aligned} \quad (\text{A3})$$

The index  $D_{MH}$  thus corresponds to index  $\Delta^\alpha$  (eq. 6b of the main text) computed from species relative abundances with  $\alpha = 2$ .

- Similarity ratio (e.g. Wishart 1969; Janssen 1972):

$$S_{SR} = \frac{\sum_j x_{Uj}x_{Vj}}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 - \sum_j x_{Uj}x_{Vj}} \quad (\text{A4})$$

A dissimilarity index can be defined as  $D_{SR} = \sqrt{1 - S_{SR}}$  :

$$\begin{aligned} D_{SR} &= \sqrt{1 - \frac{\sum_j x_{Uj}x_{Vj}}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 - \sum_j x_{Uj}x_{Vj}}} = \sqrt{\frac{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 - \sum_j x_{Uj}x_{Vj} - \sum_j x_{Uj}x_{Vj}}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 - \sum_j x_{Uj}x_{Vj}}} \\ &= \sqrt{\frac{\sum_j (x_{Uj} - x_{Vj})^2}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 - \sum_j x_{Uj}x_{Vj}}} \end{aligned} \quad (\text{A5})$$

- A similarity index derived from Sokal and Sneath (1963) as shown in Pavoine and Ricotta (2014):

$$S_{SN} = \frac{\sum_j x_{Uj}x_{Vj}}{2\sum_j x_{Uj}^2 + 2\sum_j x_{Vj}^2 - 3\sum_j x_{Uj}x_{Vj}} \quad (\text{A6})$$

with the corresponding dissimilarity index which is defined as  $D_{SN} = \sqrt{1 - S_{SN}}$  :

$$\begin{aligned} D_{SN} &= \sqrt{1 - \frac{\sum_j x_{Uj}x_{Vj}}{2\sum_j x_{Uj}^2 + 2\sum_j x_{Vj}^2 - 3\sum_j x_{Uj}x_{Vj}}} = \sqrt{\frac{2\sum_j x_{Uj}^2 + 2\sum_j x_{Vj}^2 - 3\sum_j x_{Uj}x_{Vj} - \sum_j x_{Uj}x_{Vj}}{2\sum_j x_{Uj}^2 + 2\sum_j x_{Vj}^2 - 3\sum_j x_{Uj}x_{Vj}}} \\ &= \sqrt{\frac{2\sum_j (x_{Uj} - x_{Vj})^2}{2\sum_j x_{Uj}^2 + 2\sum_j x_{Vj}^2 - 3\sum_j x_{Uj}x_{Vj}}} \end{aligned} \quad (\text{A7})$$

All these dissimilarity measures satisfy the complementarity requirement *sensu* Clarke et al. (2006). That is, the measure takes its maximum value (1) when the two plots  $U$  and  $V$  have no species in common.

In addition, starting from  $D^\alpha$  (eq. 6a of the main text), for  $\alpha = 2$  we can define a new dissimilarity index  $D_{RP} = D^2$ . The corresponding similarity index can be thus defined as:

$$\begin{aligned} S_{RP} &= 1 - D_{RP} = 1 - \frac{\sum_j (x_{Uj} - x_{Vj})^2}{\sum_j (x_{Uj} + x_{Vj})^2} \\ &= \frac{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 + 2\sum_j x_{Uj}x_{Vj} - \sum_j x_{Uj}^2 - \sum_j x_{Vj}^2 + 2\sum_j x_{Uj}x_{Vj}}{\sum_j (x_{Uj} + x_{Vj})^2} \\ &= \frac{4\sum_j x_{Uj}x_{Vj}}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 + 2\sum_j x_{Uj}x_{Vj}} \end{aligned} \quad (\text{A8})$$

Generalizing the previous indices we obtain:

$$S(x_U, x_V; m) = \frac{\frac{m}{2} \left( 2 \sum_j x_{Uj} x_{Vj} \right)}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 + \left( \frac{m}{2} - 1 \right) \left( 2 \sum_j x_{Uj} x_{Vj} \right)} \quad (\text{A9})$$

with

$$\begin{aligned} S(x_U, x_V; 0.5) &= S_{SN}; \\ S(x_U, x_V; 1) &= S_{SR}; \\ S(x_U, x_V; 2) &= S_{MH} \text{ (if relative abundances are used);} \\ S(x_U, x_V; 4) &= S_{RP} \end{aligned}$$

and

$$D(x_U, x_V; m) = \sqrt{1 - S(x_U, x_V; m)} = \sqrt{\frac{\sum_j (x_{Uj} - x_{Vj})^2}{\sum_j x_{Uj}^2 + \sum_j x_{Vj}^2 + \left( \frac{m}{2} - 1 \right) \left( 2 \sum_j x_{Uj} x_{Vj} \right)}} \quad (\text{A10})$$

where  $D(x_U, x_V; 2)$  and  $D(x_U, x_V; 4)$  correspond to eq. 6b and eq. 6a of the main text for  $\alpha = 2$ , respectively.

The parametric dissimilarities  $D^\alpha$ ,  $\Delta^\alpha$  and the other dissimilarity measures presented above can be further put into an even wider context with the general formulas:

$$S(x_U, x_V; m, \alpha) = \frac{\frac{m}{2} \left( \sum_j (x_{Uj} + x_{Vj})^\alpha - \sum_j x_{Uj}^\alpha - \sum_j x_{Vj}^\alpha \right)}{\sum_j x_{Uj}^\alpha + \sum_j x_{Vj}^\alpha + \left( \frac{m}{2} - 1 \right) \left( \sum_j (x_{Uj} + x_{Vj})^\alpha - \sum_j x_{Uj}^\alpha - \sum_j x_{Vj}^\alpha \right)} \quad (\text{A11})$$

$$D(x_U, x_V; m, \alpha) = \sqrt[\alpha]{\frac{\sum_j |x_{Uj} - x_{Vj}|^\alpha}{\sum_j x_{Uj}^\alpha + \sum_j x_{Vj}^\alpha + \left( \frac{m}{2} - 1 \right) \left( \sum_j (x_{Uj} + x_{Vj})^\alpha - \sum_j x_{Uj}^\alpha - \sum_j x_{Vj}^\alpha \right)}} \quad (\text{A12})$$

$D(x_U, x_V; m, \alpha)$  satisfies the complementarity requirement as for two plots without shared species

$$\sum_j (x_{Uj} + x_{Vj})^\alpha - \sum_j x_{Uj}^\alpha - \sum_j x_{Vj}^\alpha = 0.$$

For  $\alpha = 2$ , we find again the generalizations given above ( $S(x_U, x_V; m)$  and  $D(x_U, x_V; m)$ ). Note also

$$\text{that } \sum_j (x_{Uj} + x_{Vj})^\alpha - \sum_j x_{Uj}^\alpha - \sum_j x_{Vj}^\alpha = \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k}.$$

In addition,  $D(x_U, x_V; 2, \alpha)$  and  $D(x_U, x_V; 4, \alpha)$  correspond to eq. 6b and eq. 6a of the main text, respectively.

For  $m = 0.5$ , we obtain a generalization of the Sokal and Sneath similarity index (presented above) and its associated generalized dissimilarity index:

$$S(x_U, x_V; 0.5, \alpha) = \frac{0.5 \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k}}{2 \sum_j x_{Uj}^\alpha + 2 \sum_j x_{Vj}^\alpha - 3 \left( 0.5 \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k} \right)} \quad (\text{A13})$$

$$D(x_U, x_V; 0.5, \alpha) = \sqrt[\alpha]{\frac{2 \sum_j |x_{Uj} - x_{Vj}|^\alpha}{2 \sum_j x_{Uj}^\alpha + 2 \sum_j x_{Vj}^\alpha - 3 \left( 0.5 \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k} \right)}} \quad (\text{A14})$$

Finally, for  $m = 1$ , we have a generalization of the similarity ratio (presented above) and its associated dissimilarity index:

$$S(x_U, x_V; 1, \alpha) = \frac{\frac{1}{2} \left( \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k} \right)}{\sum_j x_{Uj}^\alpha + \sum_j x_{Vj}^\alpha - \frac{1}{2} \left( \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k} \right)} \quad (\text{A15})$$

$$D(x_U, x_V; 1, \alpha) = \sqrt[\alpha]{\frac{\sum_j |x_{Uj} - x_{Vj}|^\alpha}{\sum_j x_{Uj}^\alpha + \sum_j x_{Vj}^\alpha - \frac{1}{2} \left( \sum_j \sum_{k=1}^{\alpha-1} C_\alpha^k x_{Uj}^k x_{Vj}^{\alpha-k} \right)}} \quad (\text{A16})$$

## References

- Clarke, K.R., Somerfield, P.J., Chapman, M.G. (2006) On resemblance measures for ecological studies, including taxonomic dissimilarities and a zero-adjusted Bray-Curtis coefficient for denuded assemblages. *Journal of Experimental Marine Biology and Ecology* 330: 55–80.
- Horn, H.S. (1966) Measurement of “overlap” in comparative ecological studies. *American Naturalist* 100: 419–424.
- Janssen, J.G.M. (1972) Detection of some micropatterns of winter annuals in pioneer communities of dry sandy soils. *Acta Botanica Neerlandica* 21: 609–616.
- Morisita, M. (1959) Measuring of interspecific association and similarity between communities. *Memoirs of the Faculty of Science, Kyushu University Series E (Biology)* 3: 65–80.
- Pavoine, S., Ricotta, C. (2014) Functional and phylogenetic similarity among communities. *Methods in Ecology and Evolution* 5: 666–675.
- Sokal, R.R., Sneath, P.H.A. (1963) *Principles of Numerical Taxonomy*. W.H. Freeman, San Francisco.
- Wishart, D. (1969) An algorithm for hierarchical classification. *Biometrics* 25: 165–170.