



HAL
open science

Regularization of closed positive currents and extension theory

Michel Méo

► **To cite this version:**

Michel Méo. Regularization of closed positive currents and extension theory. Bulletin des Sciences Mathématiques, 2022, 177, pp.103141. 10.1016/j.bulsci.2022.103141 . hal-03768625

HAL Id: hal-03768625

<https://hal.science/hal-03768625>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Regularization of closed positive currents and extension theory

Michel Méo

Institut Élie Cartan, Université de Lorraine, boulevard des Aiguillettes, BP 70239, 54506 Vandoeuvre-lès-Nancy, France
E-mail: michel.meo@univ-lorraine.fr

Abstract In this note we extend to any bidimension (p, p) the Demailly theorem of regularization of closed positive $(1, 1)$ -currents on a compact Kähler manifold X of dimension n . When the manifold X is projective, we get explicitly a closed regularization with bounded negative part, constructed by using the space $C_p(X)$ of effective algebraic cycles of X of dimension p . This space can be injected in the space of divisors of $C_{n-p-1}(X)$ and we arrive at an intrinsic construction of the Skoda potential associated with a closed positive current of X . On another hand, in the case of a divisor D of X , we give an explicit bound for the degree of an irreducible component of the singular locus D_{sing} , involving the geometry of X . Lastly when X is embedded in the projective space \mathbb{P}_N , we prove the existence of a closed current extending in the generalized sense a given closed current of X , by using here a space of cycles. As an application, we obtain a characterization of the cohomology classes contained in some algebraic hypersurface of X .

Keywords: Chern classes, Closed positive current, Cycle space, Intersection number, King formula, Radon transform

MSC: 14C05, 14C17, 32C30, 32J25

1. Regularization of closed positive currents on a compact Kähler manifold

Let T be a closed positive (q, q) -current on a compact Kähler manifold X of dimension n . We use the writing

$$T = \Theta + dd^c S$$

where Θ is a \mathcal{C}^∞ differential (q, q) -form closed in X and S is a differential $(q-1, q-1)$ -form with L_{loc}^1 coefficients in X , such that ∂S and $\bar{\partial} S$ are with L_{loc}^1 coefficients in X .

Let (U_α) be a finite covering of X with open sets of coordinate maps and let (λ_α) be a \mathcal{C}^∞ partition of the unit subordinate to (U_α) . Let $S_{\alpha, \varepsilon}$ be a smooth differential $(q-1, q-1)$ -form with compact support in U_α , weakly converging in U_α to $\lambda_\alpha S|_{U_\alpha} = \lambda_\alpha S$ when $\varepsilon \rightarrow 0^+$. We set

$$\tilde{T}_\varepsilon = \Theta + dd^c \left(\sum_\alpha S_{\alpha, \varepsilon} \right)$$

which is a differential (q, q) -form of class \mathcal{C}^∞ closed in X , weakly converging in X to T when $\varepsilon \rightarrow 0^+$. First we have

$$\tilde{T}_\varepsilon = \Theta + \sum_\alpha (\lambda_\alpha T - \lambda_\alpha \Theta)_\varepsilon + \sum_\alpha (dd^c S_{\alpha, \varepsilon} - (\lambda_\alpha dd^c S)_\varepsilon).$$

When the index α is fixed, $dd^c S_{\alpha, \varepsilon} - (\lambda_\alpha dd^c S)_\varepsilon$ converges weakly in U_α to

$$R_\alpha = dd^c(\lambda_\alpha S|_{U_\alpha}) - \lambda_\alpha(dd^c S|_{U_\alpha}),$$

which is only a differential (q, q) -form with L^1_{loc} coefficients in U_α (see [4]). But when α varies, the relation

$$\sum_{\alpha} R_{\alpha} = 0 \quad (1)$$

is satisfied i.e. there is a compensation of the singularities of the R_{α} .

There is a regularization $R_{\alpha, \varepsilon}$ of $R_{\alpha} = \text{dd}^c(\lambda_{\alpha} S|_{U_{\alpha}}) - (\lambda_{\alpha} T|_{U_{\alpha}} - \lambda_{\alpha} \Theta|_{U_{\alpha}})$ which satisfies $\text{supp } R_{\alpha, \varepsilon}$ is a compact in U_{α} and $\sum_{\alpha} R_{\alpha, \varepsilon}$ is uniformly bounded in X . We write in U_{α} a homotopy formula

$$R_{\alpha, \varepsilon} = \text{dd}^c S_{\alpha, \varepsilon} + V_{\alpha, \varepsilon}$$

and then we decompose $V_{\alpha, \varepsilon} = (\lambda_{\alpha} T)_{\varepsilon} - (\lambda_{\alpha} \Theta)_{\varepsilon}$ where $(\lambda_{\alpha} T)_{\varepsilon} \geq 0$ is of class C^{∞} with compact support in U_{α} and weakly converging to $\lambda_{\alpha} T$ in U_{α} .

Since $(\lambda_{\alpha} T)_{\varepsilon} \geq 0$, we arrive in such a way at

$$\tilde{T}_{\varepsilon} \geq \Theta - \sum_{\alpha} (\lambda_{\alpha} \Theta)_{\varepsilon} + \text{smooth term}$$

thus the negative part of \tilde{T}_{ε} is uniformly bounded in X i.e. there is constant C such that $\tilde{T}_{\varepsilon} \geq -C\omega^q$, where ω is a Kähler metric on X .

When Z is an effective algebraic cycle of X of codimension q , since

$$\lim_{\varepsilon \rightarrow 0^+} \int_X (\tilde{T}_{\varepsilon} + C\omega^q) \wedge [Z] \wedge \omega^{n-2q} = \lim_{\varepsilon \rightarrow 0^+} \int_X (\{\tilde{T}_{\varepsilon}\} + C\{\omega\}^q) \wedge \{Z\} \wedge \{\omega\}^{n-2q}$$

is finite, there is a sequence $\varepsilon_j \xrightarrow{\searrow} 0$ such that $\tilde{T}_{\varepsilon_j}|_Z$ is weakly converging in Z . Then the sequence $\int_X T_{\varepsilon_j} \wedge \theta$ is convergent for each $\theta = i_* v$ where $i : Z \rightarrow X$ is the natural injection and v is a smooth differential form of bidimension (q, q) in Z .

So we have a version in any bidegree (q, q) of the Demailly regularization theorem (see [1]).

Proposition 1. *There is a sequence of smooth differential (q, q) -forms T_j weakly converging in X to T and satisfying $T_j \geq -C\omega^q$ for all j , where C is a constant and ω is a Kähler metric on X .*

Proof. We can assume that $T = [Z]$ where Z is an effective algebraic cycle of X of dimension $p = n - q$.

We denote by $\lambda : \tilde{X} \rightarrow X$ the blowing up of X along Z . When θ is a (p, p) -current in X , the intersection number

$$\int_X [Z] \wedge \theta = \int_{\tilde{X}} \lambda^*[Z] \wedge \lambda^*\theta \quad (2)$$

is well defined. In fact, $\lambda^*\theta$ is a current in \tilde{X} by [11] and $\lambda^*[Z] = [H] \wedge \gamma$, where H is the exceptional divisor in \tilde{X} and γ is a smooth closed differential $(q - 1, q - 1)$ -form in \tilde{X} . Then $[H] \wedge \lambda^*\theta$ is defined by division of distributions.

We consider the vector space $\mathcal{W} = \{(p, p)\text{-currents } \theta \text{ in } X \text{ of order 0 with } \text{sing supp } \theta \subset Z\}$, in other words $\theta \in \mathcal{W}$ when $\theta = \theta_{\text{abs}} + \theta_{\text{sing}}$ where θ_{abs} is with L^1_{loc} coefficients in X and smooth in $X \setminus Z$ while θ_{sing} is with measure coefficients and $\text{supp } \theta_{\text{sing}} \subset Z$. In particular $\mathcal{W} \supset \mathcal{D}_{p,p}(X)$ with $\mathcal{D}_{p,p}(X)$ the space of smooth differential (p, p) -forms in X and $\mathcal{W} = \overline{\mathcal{D}_{p,p}(X)}$ if the topology of \mathcal{W} is given by setting $\theta_j \rightarrow \theta$ in \mathcal{W} when θ_j converges to θ for the weak topology of currents of order 0 in X and $\theta_j|_{X \setminus Z}$ converges to $\theta|_{X \setminus Z}$ for the C^{∞} topology in $X \setminus Z$.

On the one hand we set $L(\theta_{\text{abs}}) = \int_Z \theta_{\text{abs}}$ which is possibly infinite and always equal to $\lim_j \int_Z \theta_j$ for a sequence of differential (p, p) -forms θ_j with L^1_{loc} coefficients in X converging to θ for the L^1_{loc} topology and such that $\theta_j|_Z$ is L^1_{loc} for each j . On another hand we set $L(\theta_{\text{sing}}) = \int_X [Z] \wedge \theta_{\text{sing}}$ as being the

previous intersection number. We claim the existence of a family of smooth closed differential forms T_ε of type (q, q) in X such that $L(\theta) = \lim_{\varepsilon \rightarrow 0^+} \int_X T_\varepsilon \wedge \theta$ for all $\theta \in \mathcal{W}$.

In fact, if for each sequence of smooth closed differential forms T_j of type (q, q) in X , there is an element $\theta \in \mathcal{W}$ such that $\lim_j \int_X T_j \wedge \theta \neq L(\theta)$, there would be thanks to [12] an element $\theta \in \mathcal{W}$ such that $L(\theta) = 1$ and θ is d-exact in X . Since $L(\theta)$ is finite, $L(\theta_{\text{abs}})$ is finite too and thus it can be calculated by the formula (2). So $L(\theta)$ can be calculated by the same formula, which implies $L(\theta) = 0$ since θ is d-exact in X . This is a contradiction.

We prove now that the negative part of T_ε is bounded. With $g \geq 0$ any L^1_{loc} function in X smooth in $X \setminus Z$ and with $\alpha_1, \dots, \alpha_p$ any set of differential $(1, 0)$ -forms smooth in X , we take $\theta = gi\alpha_1 \wedge \bar{\alpha}_1 \wedge \dots \wedge i\alpha_p \wedge \bar{\alpha}_p$. Then

$$\lim_{\varepsilon \rightarrow 0^+} \int_X gT_\varepsilon \wedge i\alpha_1 \wedge \bar{\alpha}_1 \wedge \dots \wedge i\alpha_p \wedge \bar{\alpha}_p = \int_Z gi\alpha_1 \wedge \bar{\alpha}_1 \wedge \dots \wedge i\alpha_p \wedge \bar{\alpha}_p > -\infty$$

implies that the negative part of $T_\varepsilon \wedge i\alpha_1 \wedge \bar{\alpha}_1 \wedge \dots \wedge i\alpha_p \wedge \bar{\alpha}_p$, which is weakly converging in X to 0, is bounded. So there exists a constant $C \geq 0$ such that $T_\varepsilon \geq -C\omega^q$. ■

2. Regularization of closed positive currents on a projective manifold

Let X be a connected projective complex manifold with $\dim X = n$ and let $C_p(X)$ be the space of algebraic cycles on X of dimension $p = n - q$. We consider the closed positive (q, q) -current $T = [Z]$ with $Z \in C_p(X)$.

Let $\mathcal{I} : \{\text{continuous differential forms of type } (p, p) \text{ on } X\} \rightarrow \{\text{continuous functions on } C_p(X)\}$ be the integral transformation defined by

$$\mathcal{I}(\theta)(Z) = \int_Z \theta = \langle [Z], \theta \rangle = \langle \delta_Z, \mathcal{I}(\theta) \rangle = \langle ({}^t\mathcal{I})(\delta_Z), \theta \rangle \Leftrightarrow [Z] = ({}^t\mathcal{I})(\delta_Z)$$

where δ_Z denotes the Dirac measure at the point $Z \in C_p(X)$ (see [5]). As pointed out by Jean-Pierre Demailly, this formula gives a natural way to approximate closed positive currents on X (see [9]).

Let M be the connected component of Z in $C_p(X)$ and let χ_j be a sequence of non negative smooth functions on $C_p(X)$, with support in M and weakly converging on $C_p(X)$ to the Dirac mass δ_Z . We decompose $\chi_j = \alpha_j + \delta_j$ on M where $({}^t\mathcal{I})(\delta_j) = T_j$ is \mathcal{C}^0 on X and where $\alpha_j \rightarrow 0$ weakly. In fact, let \mathcal{S} be the set of distributions u orthogonal to the χ such that $({}^t\mathcal{I})(\chi)$ is \mathcal{C}^0 . We take α_j such that

$$\int_M \chi_j u = \int_M \alpha_j u \text{ for all } u \in \mathcal{S}$$

and α_j uniformly bounded $\Rightarrow |\int_M \chi_j u|$ bounded for all $u \in \mathcal{S}$. To describe the elements $u \in \mathcal{S}$, we consider a (p, p) -current θ in X of order 0 and we calculate the intersection number

$$\int_X ({}^t\mathcal{I})(\chi) \wedge \theta = \lim_\varepsilon \int_X ({}^t\mathcal{I})(\chi) \wedge \theta_\varepsilon = \lim_\varepsilon \int_M \chi \mathcal{I}(\theta_\varepsilon)$$

by means of a regularization θ_ε of θ . If $({}^t\mathcal{I})(\chi)$ is of class \mathcal{C}^0 , it is equal to $\int_M \chi \mathcal{I}(\theta)$ and we obtain the condition $\int_M \chi u = 0$ where $u = \mathcal{I}(\theta) - \lim_\varepsilon \mathcal{I}(\theta_\varepsilon)$. But $\mathcal{I}(\theta)$ is obtained from θ by means of a double fibration

$$\begin{array}{ccc} & \Gamma & \\ \varphi \swarrow & & \searrow \psi \\ X & & M \end{array}$$

i.e. $\mathcal{I}(\theta) = \psi_*\varphi^*\theta$. The direct image ψ_* is weakly continuous, while the inverse image φ^* is not necessarily weakly continuous, since φ is not necessarily a submersion. We can only say that there is a proper algebraic subset $A \subset X$ such that $\Gamma \setminus \varphi^{-1}(A) \xrightarrow{\varphi} X \setminus A$ is a submersion. Note that $\varphi^*\theta$ is defined by flatification and by applying the existence due to Poly of the inverse image by blowing up. Then the distribution

$$u = \psi_*\varphi^*(1_A\theta)$$

is of order 0 and not necessarily equal to 0. But we can assume that $\chi_j \geq 0$ satisfies the supplementary condition $|\int_M \chi_j u|$ bounded. In fact, this implies the condition $|u(Z)| \leq C \int_M |u|$.

As a conclusion T_j is a closed \mathcal{C}^0 differential form that converges weakly to $[Z]$ on X . Moreover we have $T_j \geq -C({}^t\mathcal{I})(1)$.

If the cohomology class $\{Z\}$ of Z in X contains a smooth closed differential (q, q) -form $\Theta > 0$, then $({}^t\mathcal{I})(1)$ is continuous in X (see [10]).

In the general case $\Theta \geq -C\omega|_X^q$ for some constant $C \geq 0$ and ω the Fubini-Study differential $(1, 1)$ -form in \mathbb{P}_N . We approximate $[Z] + C\omega|_X^q$ by algebraic cycles Y with real coefficients such that $\{Y\} \geq 0$ (see [2]). If C is large enough, we can assume Y with positive coefficients. First we approximate $[Y]$ by smooth closed differential (q, q) -forms with bounded negative part, then we obtain a closed regularization T_j of $[Z]$ of type (q, q) with bounded negative part.

2.1. Skoda potential of a closed positive current of X

For T a closed positive (q, q) -current on X with $q = n - p$, we now express the Skoda potential U associated to T by means of the Chow transform $\mathcal{C}(T)$, which is obtained by integrating T on the effective algebraic cycles of X of dimension $q - 1$ i.e. $\mathcal{C}(T)$ is obtained from T by means of the double fibration

$$\begin{array}{ccc} & \Gamma & \\ \varphi \swarrow & & \searrow \psi \\ X & & C_{q-1}(X) \end{array}$$

with $\Gamma \subset X \times C_{q-1}(X)$ the incidence manifold.

We denote by $j : X \hookrightarrow P(V)$ an embedding of X into $P(V) = \mathbb{P}_N$ and by $\rho : G(p+1, V^*) \dashrightarrow C_{q-1}(X)$ the induced meromorphic map from the Grassmannian to the cycle space, which associates $c = X \cap P(\ker \lambda_0) \cap \dots \cap P(\ker \lambda_p)$ to a subspace $\text{vect}(\lambda_0, \dots, \lambda_p) \in G(p+1, V^*) = G(q, V) = \mathbb{G}_{q-1, N}$. Here $P(\ker \lambda_i)$ is the projective subspace of \mathbb{P}_N defined by the linear equation $\lambda_i = 0$.

We consider the irreducible component M of the cycles $c = X \cap P(\ker \lambda_0) \cap \dots \cap P(\ker \lambda_p)$ in $C_{q-1}(X)$. Denote by d_M the dimension of M and by $d = d_M - p - 1$ the dimension of a generic fiber of φ restricted to $\psi^{-1}(M)$. We set $\Omega = \psi_*\varphi^*(\omega|_X^q)$ with ω the Fubini-Study form on \mathbb{P}_N (see [13]).

With \mathcal{A} the positive smooth differential (d, d) -form equal to Ω^{d_M-p-1} on M , we define on X the $(1, 1)$ -current

$$T_1 = \varphi_*\psi^*(\mathcal{A} \wedge \mathcal{C}(T)|_M) = \varphi_*\psi^*(\mathcal{A} \wedge (\psi_*\varphi^*T)|_M)$$

which is an integral transform of T .

Proposition 2. *The closed positive $(1, 1)$ -current T_1 on X satisfies $T_1 = \beta + \text{dd}^c U$ with U the Skoda potential associated to T and β a closed smooth differential $(1, 1)$ -form on X . Thus we have the equalities*

- (i) $\beta = \varphi_*\psi^*(\mathcal{A} \wedge \mathcal{C}(\Theta)|_M)$ if T is cohomologous to Θ smooth on X ,
- (ii) $\nu(T_1, x) = C_M \nu(T, x)$ for the Lelong numbers at each point $x \in X$ with a constant C_M depending on M only.

Proof. By writing $T = \Theta + \text{dd}^c S$, we have

$$\mathcal{C}(T)|_M = \mathcal{C}(\Theta)|_M + \text{dd}^c \mathcal{C}(S)|_M$$

with $\mathcal{C}(S)(c) = \int_X S \wedge [c]$ and $[c] = [X \cap \mathbb{P}_{N-p-1}] = \omega_X^{p+1} + \text{dd}^c \gamma_c$ the integration current associated to a cycle $c \in M$. In other words γ_c is a Green form in X of the cycle c . Up to a smooth function of c , we have $\mathcal{C}(S)(c) = \int_X T \wedge \gamma_c$. To calculate the Lelong numbers of U , we can replace $U(x)$ by

$$\int_X T \wedge \left(\int_{\substack{c \in M \\ c \ni x}} \gamma_c \Omega^d \right).$$

When $y \in X$ varies, to calculate $d_y d_y^c \int_{\substack{c \in M \\ c \ni x}} \gamma_c(y) \Omega^d$ we are reduced to calculate $\int_{\substack{c \in M \\ c \ni x}} [c](y) \Omega^d$. But

$$\left(\int_{\substack{s \in G(q, V) \\ s \ni x}} [s](y) \Phi_x \right)|_X = \int_{\substack{s \in G(q, V) \\ s \ni x}} [X \cap s](y) \Phi_x = K \int_{\substack{c \in M \\ c \ni x}} [c](y) \Omega^d$$

where Φ_x is a volume form on $\{s \in G(q, V), s \ni x\}$ and K is a constant, since the volume of $\{s \in G(q, V), s \ni x \text{ and } X \cap s = c\}$ does not depend on (c, x) such that $c \ni x$. So the conclusion relies on the calculation of $\int_{\substack{s \in G(q, V) \\ s \ni x}} [s](y) \Phi_x$ that is to say on the case of the Skoda potential for $\mathbb{P}_N = P(V)$. ■

2.2 Irreducible components of the singular locus for a divisor of X

We give here a bound for the degree of an irreducible component of the singular locus of divisor, similar to those of [1, 6].

Let $D = s^{-1}(0)$ be an effective divisor in X where $s \in H^0(X, L)$ with a holomorphic line bundle $L = \mathcal{O}(D)$.

For $k \in \mathbb{N}$, we denote by $J^k s \in H^0(X, J^k L)$ the holomorphic section of k -jets of s . We consider the strictly decreasing sequence $E_0 \supset E_1 \supset \dots \supset E_m$ of algebraic subsets of D defined by

$$E_l = \{J^{k_{l-1}} s = 0\} = \dots = \{J^{k_l-1} s = 0\}$$

for $1 \leq l \leq m$ and by $E_0 = \{J^0 s = 0\} = \dots = \{J^{k_0-1} s = 0\}$, where $1 \leq k_0 < \dots < k_m$ and $\{J^{k_m} s = 0\} = \emptyset$ while $\{J^{k_m-1} s = 0\} \neq \emptyset$.

In other words, on $E_l \setminus E_{l+1}$, the multiplicity of D is k_l for $0 \leq l \leq m-1$ and on E_m the multiplicity of D is k_m . So the sequence $1 \leq k_0 < \dots < k_m$ is the sequence of different multiplicities of D .

We set $d_l = \dim E_l$ in such a way that $d_0 \geq d_1 \geq \dots \geq d_m$ with $d_0 = n-1$ and $n = \dim X$.

Let C be an irreducible component of E_l of dimension d_l which is not contained in E_{l+1} . Since $C = \overline{C} \setminus (\overline{C} \cap \overline{E_{l+1}})$, the generic multiplicity of D along C is equal to k_l . Note that if $C \subset E_{l+1}$, then $d_l = d_{l+1}$.

The integration current of bidimension (d_l, d_l) in X associated to the algebraic cycle of dimension d_l in X defined by $J^{k_{l-1}} s \in H^0(X, J^{k_{l-1}} L)$ can be written

$$\sum_{C \not\subset E_{l+1}} \nu(J^{k_{l-1}} s, C)[C] + \sum_{C \subset E_{l+1}} \nu(J^{k_{l-1}} s, C)[C]$$

with $\nu(J^{k_{l-1}} s, C)$ the generic multiplicity of $J^{k_{l-1}} s$ along C . Its cohomology class in X is equal to

$$c_{n-d_l}(J^{k_{l-1}} L) - c_{n-d_l}(\mathcal{F}_l)$$

where \mathcal{F}_l is the coherent sheaf in X of generic rank equal to $\text{rank}(J^{k_l-1}L) - 1$, whose dual \mathcal{F}_l^\vee is the sheaf of germs of holomorphic sections of $(J^{k_l-1}L)^*$ equal to 0 on $J^{k_l-1}s$.

Proposition 3. *If the Chern class $c_{n-d_l}(\mathcal{F}_l)$ is ≥ 0 in X , then we have the following inequality*

$$\sum_{C \notin E_{l+1}} \nu(J^{k_l-1}s, C)\{C\} \leq c_{n-d_l}(J^{k_l-1}L)$$

for the cohomology classes of bidimension (d_l, d_l) in X .

We use a \mathcal{C}^∞ isomorphism $J^k L \simeq \bigoplus_{0 \leq j \leq k} (S^j T^* X \otimes L)$, which gives the equality

$$c(J^k L) = \prod_{0 \leq j \leq k} c(S^j T^* X \otimes L)$$

for the total class Chern of $J^k L$ on X . Lastly recall that for E a holomorphic vector bundle of rank r on X , we have the equality

$$c(E \otimes L) = \sum_{0 \leq m \leq r} (1 + c_1(L))^{r-m} c_m(E)$$

since L is of rank 1. In such a way $c_{n-d_l}(J^{k_l-1}L)$ becomes a polynomial in $c_1(L)$ of degree $n - d_l$ and whose coefficients depend only on the Chern classes of the $S^j T^* X$.

3. Extension to \mathbb{P}_N of closed (q, q) -currents in X

Let X be a closed complex submanifold in \mathbb{P}_N of dimension n and let $i_0 : X \hookrightarrow \mathbb{P}_N$ be the canonical injection. Let T be a closed smooth differential form on X of bidimension (p, p) with $p = n - q$.

Proposition 4. *There is a closed (q, q) -current S on \mathbb{P}_N having a restriction $S|_X$ equal to T , that is to say such that, for all smooth differential (p, p) -forms u in X , the current S has an intersection number with the direct image $i_{0*}u$ in \mathbb{P}_N equal to the intersection number of T with u in X .*

Proof. Since every closed (q, q) -current on \mathbb{P}_N is a weak limit of algebraic cycles of codimension q , we search

$$S = \int_{A \in C_{N-q}(\mathbb{P}_N)} [A] \mu(A)$$

for some distribution μ on the cycle space $C_{N-q}(\mathbb{P}_N)$. This can be written $S = ({}^t \mathcal{R})(\mu)$ where

$$\mathcal{R} : \mathcal{D}_{N-q, N-q}(\mathbb{P}_N) \rightarrow \mathcal{C}^\infty(C_{N-q}(\mathbb{P}_N))$$

is the Radon transform obtained by integration on the algebraic cycles of dimension $N - q$.

We define the intersection number $\int_{\mathbb{P}_N} S \wedge i_{0*}u = \lim_j \int_{\mathbb{P}_N} S \wedge \varphi_j(u)$ for some sequence $\varphi_j(u)$ of smooth differential $(N - q, N - q)$ -forms in \mathbb{P}_N weakly converging in \mathbb{P}_N to $i_{0*}u$. Then this number can be expressed as being

$$\lim_j \int_{C_{N-q}(\mathbb{P}_N)} \mu \mathcal{R}(\varphi_j(u)) \tag{3}$$

and we can choose μ in order to obtain $\int_X T \wedge u$ for each smooth differential (p, p) -form u in X . In fact, we define a closed (q, q) -current \tilde{T} in X by $\lim_j \int_{C_{N-q}(\mathbb{P}_N)} \mu \mathcal{R}(\varphi_j(u)) = \int_X \tilde{T} \wedge u$ and we can force $\tilde{T} = T$. ■

So the Proposition 4 can be stated in the following way.

Proposition 5. *There is a transform $\varphi_j : \mathcal{D}_{p,p}(X) \rightarrow \mathcal{D}_{N-q,N-q}(\mathbb{P}_N)$ satisfying : $\varphi_j(u)$ converges weakly in \mathbb{P}_N to $i_{0*}u$ and $\int_{\mathbb{P}_N} S \wedge \varphi_j(u) \rightarrow \int_X T \wedge u$ when $j \rightarrow \infty$.*

Proof. For each smooth differential (p,p) -form u in X , we choose a sequence $\varphi_j(u)$ weakly converging in \mathbb{P}_N to $i_{0*}u$, such that if $u \notin \text{Im } \partial + \text{Im } \bar{\partial}$, there is no sequences v_j and w_j of smooth differential forms such that $\varphi_j(u) - \partial v_j - \bar{\partial} w_j \rightarrow 0$ for the \mathcal{C}^∞ topology. Then S is a continuous linear form on $\mathcal{D}_{N-q,N-q}(\mathbb{P}_N)$ equal to 0 on $\text{Im } \partial + \text{Im } \bar{\partial}$ and equal to $\int_X T \wedge u$ on $\text{lim}_j \varphi_j(u)$. ■

3.1. Case $X = \mathbb{P}_{N-1} \subset \mathbb{P}_N$

The existence of S can also be seen as a consequence of an integral representation formula of Bochner-Martinelli type in $\mathbb{P}_N \setminus X$, for $\varphi \in \mathcal{D}_{N-q,N-q}(\mathbb{P}_N)$.

For example let us assume that $X = \mathbb{P}_{N-1}$, in such a way that $\mathbb{P}_N \setminus X = \mathbb{C}^N$ and let us write the integration current $[D_{\mathbb{C}^N}]$ on the diagonal $D_{\mathbb{C}^N}$ of $\mathbb{C}^N \times \mathbb{C}^N$ as being

$$[D_{\mathbb{C}^N}] = \text{dd}^c K$$

with $K(w, w') = -\frac{1}{(N-1)2^N} \frac{(\text{dd}^c \|w-w'\|^2)^{N-1}}{\|w-w'\|^{2(N-1)}}$ for $(w, w') \in \mathbb{C}^N \times \mathbb{C}^N$ (see [3, 7]). The integral kernel K has L^1_{loc} coefficients in $\mathbb{C}^N \times \mathbb{C}^N$ in other words $w \rightarrow K(w, \cdot)$, which is with values in the space of differential forms L^1_{loc} in \mathbb{C}^N , is L^1_{loc} with respect to $w \in \mathbb{C}^N$.

For $w \in \mathbb{C}^N$, we define the potential

$$U(w) = \int_{w' \in \mathbb{C}^N} K(w, w') \wedge \varphi(w')$$

which is a differential $(N-q-1, N-q-1)$ -form with L^1_{loc} coefficients in \mathbb{C}^N . In effect, by writing $\varphi = \varphi_1 - \varphi_2$ with each φ_i smooth positive, we can assume that φ is smooth positive of bidimension (q, q) in \mathbb{P}_N . Then by writing $\varphi \leq C\omega^{N-q}$ for some constant $C \geq 0$, this local integrability is a consequence of the following result.

Proposition 6. *For φ a closed positive current of bidimension (q, q) in \mathbb{P}_N , the potential U is L^1_{loc} in \mathbb{C}^N .*

Proof. With $\tau = \varphi|_{\mathbb{C}^N} \wedge (\frac{i}{2} \partial \bar{\partial} \|w'\|^2)^q / q!$ the trace measure of $\varphi|_{\mathbb{C}^N}$, we use the Fubini formula

$$\int_{\|w\| < R_0} (\frac{i}{2} \partial \bar{\partial} \|w\|^2)^N / N! \int_{w' \in \mathbb{C}^N} \frac{\tau(w')}{\|w-w'\|^{2(N-1)}} = \int_{w' \in \mathbb{C}^N} k(R_0, w') \tau(w')$$

with a function $k(R_0, w') = \int_{\|w\| < R_0} \frac{(\frac{i}{2} \partial \bar{\partial} \|w\|^2)^N / N!}{\|w-w'\|^{2(N-1)}}$ equal to $CR_0^{2N} \|w'\|^{-2N+2}$ when $\|w'\| > R_0$.

Then we express

$$\int_{\|w'\| > R_0} \|w'\|^{-2N+2} \tau(w') = \int_{r=R_0}^{+\infty} r^{-2N+2} d\mathcal{M}(r)$$

with

$$\mathcal{M}(r) = \int_{\|w'\| < r} \tau(w') = \frac{\pi^q}{q!} r^{2q} \{ \nu(\varphi|_{\mathbb{C}^N}, 0) + \int_{0 < \|w'\| < r} \varphi|_{\mathbb{C}^N} \wedge (\text{dd}^c \log \|w'\|)^q \}.$$

Note that $(\text{dd}^c \log \|w'\|)^q = \sigma^*(\omega^q)|_{\mathbb{C}^N}$ with $\mathbb{P}_N \setminus \{[1, 0, \dots, 0]\} \xrightarrow{\sigma} \mathbb{P}_{N-1}$ the orthogonal projection. Since $\varphi \wedge \sigma^*(\omega^q)$ is known to be defined in \mathbb{P}_N whole, we have $\mathcal{M}(r) = O(r^{2q})$ when $r \rightarrow +\infty$. So $\int_{\|w'\| > R_0} k(R_0, w') \tau(w')$ is convergent and $U(w)$ is a differential $(N-q-1, N-q-1)$ -form with L^1_{loc} coefficients in $\|w\| < R_0$. ■

To see that U is not with L_{loc}^1 coefficients in \mathbb{P}_N , we use again the Fubini formula

$$\int_{w \in \mathbb{C}^N} \frac{(\frac{i}{2} \partial \bar{\partial} \|w\|^2)^N / N!}{(1 + \|w\|^2)^{q+1}} \int_{w' \in \mathbb{C}^N} \frac{\tau(w')}{\|w - w'\|^{2(N-1)}} = \int_{w' \in \mathbb{C}^N} k(w') \tau(w')$$

with a function $k(w') = \int_{w \in \mathbb{C}^N} \frac{(\frac{i}{2} \partial \bar{\partial} \|w\|^2)^N / N!}{\|w - w'\|^{2(N-1)} (1 + \|w\|^2)^{q+1}} \leq C \|w'\|^{-2q}$ for $\|w'\|$ large enough. Then we express

$$\int_{\|w'\| > R_0} \|w'\|^{-2q} \tau(w') = \int_{r=R_0}^{+\infty} r^{-2q} d\mathcal{M}(r)$$

with $\mathcal{M}(r) \simeq K r^{2q}$ when $r \rightarrow +\infty$.

But U extends as a $(N - q - 1, N - q - 1)$ -current \tilde{U} in \mathbb{P}_N . In fact, assume that \mathbb{P}_{N-1} is given by $X_0 = 0$ in the homogeneous coordinates X_j associated to the basis (e_0, \dots, e_N) of \mathbb{C}^{N+1} . In such a way $w_j = \frac{X_j}{X_0}$ for each $1 \leq j \leq N$. We have

$$\tilde{U} = \text{pr}_{1*}(\tilde{K} \wedge \text{pr}_2^*(\varphi|_{\mathbb{C}^N}))$$

where \tilde{K} is an extension of K in $\mathbb{P}_N \times \mathbb{C}^N$ obtained by writing $w_j - w'_j = \frac{X_j/X_l}{X_0/X_l} - w'_j = \frac{(X_j/X_l) - w'_j(X_0/X_l)}{X_0/X_l}$ for some $1 \leq l \leq N$ such that $X_l \neq 0$ and by applying the division theorem of Hörmander-Lojasiewicz.

When φ is again a smooth differential $(N - q, N - q)$ -form in \mathbb{P}_N , we have in \mathbb{C}^N the equality

$$\begin{aligned} \text{dd}^c U &= \text{pr}_{1*}(\text{dd}^c K \wedge \text{pr}_2^*(\varphi|_{\mathbb{C}^N})) - \text{pr}_{1*}(\text{d}^c K \wedge \text{pr}_2^*(\text{d}\varphi|_{\mathbb{C}^N})) + \text{pr}_{1*}(\text{d}K \wedge \text{pr}_2^*(\text{d}^c \varphi|_{\mathbb{C}^N})) \\ &+ \text{pr}_{1*}(K \wedge \text{pr}_2^*(\text{dd}^c \varphi|_{\mathbb{C}^N})) \end{aligned}$$

which can be written as being

$$\varphi|_{\mathbb{C}^N} - \text{dd}^c U = \text{pr}_{1*}(\text{d}^c K \wedge \text{pr}_2^*(\text{d}\varphi|_{\mathbb{C}^N})) - \text{pr}_{1*}(\text{d}K \wedge \text{pr}_2^*(\text{d}^c \varphi|_{\mathbb{C}^N})) - \text{pr}_{1*}(K \wedge \text{pr}_2^*(\text{dd}^c \varphi|_{\mathbb{C}^N})).$$

In the same way the residual $(N - q, N - q)$ -current

$$R = \varphi - \text{dd}^c \tilde{U} - \{\text{pr}_{1*}(\text{d}^c \tilde{K} \wedge \text{pr}_2^*(\text{d}\varphi|_{\mathbb{C}^N})) - \text{pr}_{1*}(\text{d}\tilde{K} \wedge \text{pr}_2^*(\text{d}^c \varphi|_{\mathbb{C}^N})) - \text{pr}_{1*}(\tilde{K} \wedge \text{pr}_2^*(\text{dd}^c \varphi|_{\mathbb{C}^N}))\}$$

is with support in \mathbb{P}_{N-1} .

Lemma 1. *The currents $\frac{X_0}{X_l} R$, $\frac{\bar{X}_0}{\bar{X}_l} R$ and in the same way $\frac{dX_0}{X_l} \wedge R$, $\frac{d\bar{X}_0}{\bar{X}_l} \wedge R$ are locally flat where $X_l \neq 0$.*

Proof. First the previous expression of R gives the formula

$$\begin{aligned} \frac{X_0}{X_l} R &= \frac{X_0}{X_l} \varphi - \text{dd}^c \text{pr}_{1*}(\frac{X_0}{X_l} \tilde{K} \wedge \text{pr}_2^*(\varphi|_{\mathbb{C}^N})) + \text{pr}_{1*}(\frac{X_0}{X_l} \tilde{K} \wedge \text{pr}_2^*(\text{dd}^c \varphi|_{\mathbb{C}^N})) \\ &+ \text{d}(\frac{X_0}{X_l}) \wedge \text{pr}_{1*}(\tilde{K} \wedge \text{pr}_2^*(\text{d}^c \varphi|_{\mathbb{C}^N})) - \text{d}^c(\frac{X_0}{X_l}) \wedge \text{pr}_{1*}(\tilde{K} \wedge \text{pr}_2^*(\text{d}\varphi|_{\mathbb{C}^N})) \\ &- \text{d}^c(\frac{X_0}{X_l}) \wedge \text{pr}_{1*}(\text{d}\tilde{K} \wedge \text{pr}_2^*(\varphi|_{\mathbb{C}^N})) + \text{d}(\frac{X_0}{X_l}) \wedge \text{pr}_{1*}(\text{d}^c \tilde{K} \wedge \text{pr}_2^*(\varphi|_{\mathbb{C}^N})) \\ &+ \text{pr}_{1*}(\frac{X_0}{X_l} \text{d}\tilde{K} \wedge \text{pr}_2^*(\text{d}^c \varphi|_{\mathbb{C}^N})) - \text{pr}_{1*}(\frac{X_0}{X_l} \text{d}^c \tilde{K} \wedge \text{pr}_2^*(\text{d}\varphi|_{\mathbb{C}^N})). \end{aligned}$$

With $w_l = \frac{X_l}{X_0}$, we have the expression

$$\tilde{K} = K_{0,0} + w_l \text{d}(\frac{1}{w_l}) \wedge K_{1,0} + \bar{w}_l \text{d}(\frac{1}{\bar{w}_l}) \wedge K_{0,1} + |w_l|^2 \text{d}(\frac{1}{w_l}) \wedge \text{d}(\frac{1}{\bar{w}_l}) \wedge K_{1,1}$$

where $K_{0,0}, K_{1,0}, K_{0,1}, K_{1,1}$ are differential forms with coefficients which are L_{loc}^1 in $\mathbb{P}_N \times \mathbb{C}^N$.

In fact, $K(w, w')$ is proportional to

$$\|w - w'\|^{-2N+2} \sum_{1 \leq j \leq N} (dw_1 - dw'_1) \wedge \dots \wedge (\widehat{dw_j - dw'_j}) \wedge \dots \wedge (dw_N - dw'_N) \\ \wedge (d\bar{w}_1 - d\bar{w}'_1) \wedge \dots \wedge (\widehat{d\bar{w}_j - d\bar{w}'_j}) \wedge \dots \wedge (d\bar{w}_N - d\bar{w}'_N)$$

and when we calculate $U(w)$ we need the component of $K(w, w')$ which is of bidegree $(N-1-q, N-1-q)$ in w and of bidegree (q, q) in w' . Therefore we have to consider terms of the type

$$\|w - w'\|^{-2N+2} dw_{\{1, \dots, N\} \setminus jK} \wedge dw'_K \wedge d\bar{w}_{\{1, \dots, N\} \setminus jL} \wedge d\bar{w}'_L$$

where jK and jL are subsets of $\{1, \dots, N\}$ with $q+1$ elements. For $j \neq l$, we write

$$dw_j = \frac{d(\frac{X_j}{X_l})}{\frac{X_0}{X_l}} - \frac{X_j}{X_l} \frac{d(\frac{X_0}{X_l})}{(\frac{X_0}{X_l})^2}$$

and $dw_l = -\frac{d(\frac{X_0}{X_l})}{(\frac{X_0}{X_l})^2}$. So $dw_{\{1, \dots, N\} \setminus jK}$ can be written

$$\frac{\text{smooth form}}{(\frac{X_0}{X_l})^{N-1-q}} + \frac{(\text{smooth form}) \wedge d(\frac{X_0}{X_l})}{(\frac{X_0}{X_l})^{N-q}}$$

if jK contains l , while it can be written only $\frac{(\text{smooth form}) \wedge d(\frac{X_0}{X_l})}{(\frac{X_0}{X_l})^{N-q}}$ if jK does not contain l . As a conclusion we obtain terms of the type

$$\|w - w'\|^{-2N+2} dw'_K \wedge d\bar{w}'_L \wedge \left(\frac{\alpha}{(\frac{X_0}{X_l})^{N-1-q}} + \frac{\beta \wedge d(\frac{X_0}{X_l})}{(\frac{X_0}{X_l})^{N-q}} \right) \wedge \left(\frac{\gamma}{(\frac{X_0}{X_l})^{N-1-q}} + \frac{\delta \wedge d(\frac{X_0}{X_l})}{(\frac{X_0}{X_l})^{N-q}} \right)$$

where $\alpha, \beta, \gamma, \delta$ are smooth differential forms. For $U(w)$, we arrive at terms proportional to

$$V\alpha \wedge \gamma \text{ or } V\beta \wedge \gamma \wedge \frac{d(\frac{X_0}{X_l})}{\frac{X_0}{X_l}} \text{ or } V\alpha \wedge \delta \wedge \frac{d(\frac{X_0}{X_l})}{\frac{X_0}{X_l}} \text{ or } V\beta \wedge \delta \wedge \frac{d(\frac{X_0}{X_l}) \wedge d(\frac{X_0}{X_l})}{|\frac{X_0}{X_l}|^2}$$

where we set

$$V(w) = \frac{1}{|\frac{X_0}{X_l}|^{2(N-1-q)}} \int_{w' \in \mathbb{C}^N} \frac{dw'_K \wedge d\bar{w}'_L \wedge \varphi(w')}{\|w - w'\|^{2N-2}} = \int_{y \in \mathbb{C}^N} \frac{dy_K \wedge d\bar{y}_L \wedge \varphi\left(\left[\frac{X_0}{X_l}e_0 + y\right]\right)}{\|\frac{X_0}{X_l}w - y\|^{2N-2}}$$

with $\frac{X_0}{X_l}w = (\frac{X_1}{X_l}, \dots, \frac{X_N}{X_l})$. When $\frac{X_0}{X_l} \rightarrow 0$, the function $V(w)$ has a finite limit equal to

$$\int_{y \in \mathbb{C}^N} \frac{dy_K \wedge d\bar{y}_L \wedge \varphi(0, y_1, \dots, y_N)}{\|(\frac{X_1}{X_l}, \dots, \frac{X_N}{X_l}) - y\|^{2N-2}}.$$

By similar formulae, we prove that $\frac{\bar{X}_0}{X_l}R$ and $\frac{dX_0}{X_l} \wedge R$ as well as $\frac{d\bar{X}_0}{X_l} \wedge R$ are locally flat in the sense of Federer, where $X_l \neq 0$. ■

Note that $\frac{X_0}{X_i}\varphi$, $\frac{\bar{X}_0}{\bar{X}_i}\varphi$ and $\frac{dX_0}{X_i}\wedge\varphi = \frac{d(X_0\varphi) - X_0d\varphi}{X_i}$ as well as $\frac{d\bar{X}_0}{\bar{X}_i}\wedge\varphi$ have no components along \mathbb{P}_{N-1} .

By the structure theorem of Federer, we can write $\frac{X_0}{X_i}R = i_{0*}v$ with i_0 the injection $\mathbb{P}_{N-1} \hookrightarrow \mathbb{P}_N$ where v is some current of bidimension (q, q) in \mathbb{P}_{N-1} . So there is a current u of bidimension (q, q) in \mathbb{P}_{N-1} such that $R = i_{0*}u$ modulo a current annihilated by $\frac{X_0}{X_i}$ and $\frac{\bar{X}_0}{\bar{X}_i}$. By the second part of the Lemma 1, we obtain

$$R = i_{0*}u + \partial\left(\frac{\bar{X}_l}{\bar{X}_0}\right)\wedge a + \bar{\partial}\left(\frac{X_l}{X_0}\right)\wedge b$$

where a, b are some currents in the coordinates $\frac{X_1}{X_0}, \dots, \frac{X_N}{X_0}$. Since $\partial R_\varphi = R_{\partial\varphi}$ and $\bar{\partial}R_\varphi = R_{\bar{\partial}\varphi}$, necessarily we have $a = 0 = b$ and we can write $R = R_\varphi = i_{0*}u_\varphi$ where $u_\varphi = u$ is some current of bidimension (q, q) in \mathbb{P}_{N-1} .

The extended (q, q) -current S of T to \mathbb{P}_N is obtained by setting

$$\int_{\mathbb{P}_N} S \wedge \varphi = \int_{\mathbb{P}_{N-1}} T \wedge u_\varphi \quad (4)$$

for all smooth differential $(N - q, N - q)$ -forms φ in \mathbb{P}_N . Since $u_{d\varphi} = du_\varphi$, we have that $dS = 0$ in \mathbb{P}_N .

3.2. Cohomology classes contained in an algebraic hypersurface of X

The extended current S can be used to characterize the property that the cohomology class $\{T\}$ is contained in an algebraic hypersurface of X .

Proposition 7. *The cohomology class $\{T\}$ is contained in an algebraic hypersurface of X if and only if the extended current S is of order 0 in \mathbb{P}_N .*

Proof. If S is of order 0, then μ is a measure and we can apply the Lebesgue-Nikodym decomposition theorem $\mu = \mu_{\text{abs}} + \mu_{\text{sing}}$. We define $S_0 = ({}^t\mathcal{R})(\mu_{\text{abs}})$ and $S_1 = ({}^t\mathcal{R})(\mu_{\text{sing}})$, thus $\{S_{0|X}\} = \{S_0\}_{|X}$. By (3), we know that $\text{supp } \mu_{\text{sing}}$ is independent of T , so it is an algebraic subset. In fact,

$$\text{supp } \mu_{\text{sing}} = \{A \in C_{N-q}(\mathbb{P}_N), A \text{ non transverse to } X\}$$

and we have an algebraic stratification $\Sigma_0 \supset \Sigma_1 \supset \dots \supset \Sigma_q$ with

$$\Sigma_i = \{A \in C_{N-q}(\mathbb{P}_N), \dim(A \cap X) \geq n - q + i\}$$

for $0 \leq i \leq q$ and $n = \dim X$. Since S can be restricted to X , we have $1_{\Sigma_q}\mu = 0$ and we can decompose

$$\mu = \sum_{0 \leq i \leq q-1} 1_{\Sigma_i \setminus \Sigma_{i+1}}\mu$$

which gives $S = S_0 + S_{1,1} + \dots + S_{1,q-1}$. In other words $A \in \Sigma_i \setminus \Sigma_{i+1} \Leftrightarrow \dim(A \cap X) = n - q + i$, thus

$$S_{1,i} = \int_{A \in \Sigma_i \setminus \Sigma_{i+1}} [A]\mu(A)$$

and we can calculate the restriction current in X by

$$S_{1,i|X} = \int_{A \in \Sigma_i \setminus \Sigma_{i+1}} [A]_{|X}\mu(A) = \int_{A \in \Sigma_i \setminus \Sigma_{i+1}} \mu(A)j_*u$$

where $j : A \cap X \hookrightarrow X$ is the natural injection and u is a closed (i, i) -current in $A \cap X$ depending on A . At the cohomological level we write

$$\{S_{1,i|X}\} = \int_{A \in \Sigma_i \setminus \Sigma_{i+1}} \mu(A) j_* \{u\}$$

and we use that the subspace of $H^{q,q}(X)$ of the finite sums $\sum j_* \{u\}$ is closed. In such a way, we obtain that the cohomology class $\{S_{1,i|X}\}$ is contained in some $A \cap X$ with A such that $\dim(A \cap X) = n - q + i$. As a conclusion $\{S_{1|X}\}$ is contained in an algebraic hypersurface of X .

Conversely, when the cohomology class $\{T\}$ is contained in an algebraic hypersurface of X , we write $T = i_* r + \text{dd}^c \tilde{v} = (\text{dd}^c v)^\sim$ where $i : X \cap H \hookrightarrow X$ is the canonical injection with $H \subset \mathbb{P}_N$ an algebraic hypersurface and where v is a differential $(q-1, q-1)$ -form smooth in $X \setminus X \cap H$ and L_{loc}^1 in X such that $\text{dd}^c v$ calculated generically is L_{loc}^1 in X .

We extend v to \mathbb{P}_N by V a differential $(q-1, q-1)$ -form L_{loc}^1 in \mathbb{P}_N such that $\text{dd}^c V$ calculated generically is L_{loc}^1 in \mathbb{P}_N . In fact, there is an open covering (U_α) of \mathbb{P}_N with a family (ρ_α) of holomorphic submersions $\rho_\alpha : U_\alpha \rightarrow X \cap U_\alpha$ such that $\rho_{\alpha|X \cap U_\alpha} = \text{id}_{X \cap U_\alpha}$. When (λ_α) is a smooth partition of the unity subordinated to (U_α) , we take $V = \sum_\alpha \lambda_\alpha \rho_\alpha^*(v|_{X \cap U_\alpha})$.

Then $T = S|_X$ with $S = (\text{dd}^c V)^\sim = j_* R + \text{dd}^c \tilde{V}$ where $j : H \hookrightarrow \mathbb{P}_N$ is the canonical injection and R is a $(q-1, q-1)$ -current closed of order 0 in H . ■

Without assuming S of order 0, for all $B \in \mathbb{G}_{q,N}$, we can express, in a similar way to (4), the intersection number of S and $[B]$ as being

$$\int_{\mathbb{P}_N} S \wedge [B] = \int_X T \wedge w_{[B]}$$

where $w_{[B]}$ is some current of bidimension (q, q) in X . When S is of order 0, this intersection number, which can be calculated by Fourier transform (see [8]), is continuous with respect to $B \in \mathbb{G}_{q,N}$.

On another hand, there is a family of smooth differential (q, q) -forms S_ε weakly converging in \mathbb{P}_N to S such that

$$\int_{\mathbb{P}_N} S \wedge [B] = \lim_{\varepsilon} \int_B S_\varepsilon = \lim_{\varepsilon} \mathcal{J}(S_\varepsilon)(B)$$

with $\mathcal{J} : \{\text{continuous differential forms of type } (q, q) \text{ on } \mathbb{P}_N\} \rightarrow \{\text{continuous functions on } \mathbb{G}_{q,N}\}$ the Radon transformation defined for projective subspaces B in \mathbb{P}_N of dimension q . Then $\mathcal{J}(S_\varepsilon)$ converges weakly in $\mathbb{G}_{q,N}$ to $\mathcal{J}(S)$, which is constant on $\mathbb{G}_{q,N}$, since S is closed. Thus for almost every $B \in \mathbb{G}_{q,N}$, we have

$$\int_{\mathbb{P}_N} S \wedge [B] = (\deg S)(\deg B) = \deg S.$$

When the cohomology class $\{T\}$ is contained in an algebraic hypersurface of X , we obtain the condition $\int_X T \wedge w_{[B]} = (\deg S)(\deg B) = \deg S$ for all $B \in \mathbb{G}_{q,N}$.

Remark 1. We can give an explicit construction of S by using the King formula.

In fact, let $D_{\mathbb{P}_N}$ be the diagonal in $\mathbb{P}_N \times \mathbb{P}_N$, which can be written $D_{\mathbb{P}_N} = f^{-1}(0)$ where f is a holomorphic section of a Hermitian holomorphic vector bundle E over $\mathbb{P}_N \times \mathbb{P}_N$. Then $D_X = (f|_{X \times X})^{-1}(0)$ and we have the expression

$$[D_X] = -(\text{dd}^c \log \|f\|)_{|(X \times X) \setminus D_X}^n + \text{dd}^c \gamma$$

for the integration current on D_X in $X \times X$, where the differential $(n-1, n-1)$ -form γ is L_{loc}^1 in $X \times X$ and smooth in $(X \times X) \setminus D_X$. In $\mathbb{P}_N \times X$, we define the closed (n, n) -current of order 0

$$\mathcal{K} = -(\text{dd}^c \log \|f\|)_{|\mathbb{P}_N \times X}^n + \text{dd}^c \mathcal{G}$$

where \mathcal{G} is an extension to $\mathbb{P}_N \times X$ of γ obtained as in the proof of the Proposition 7.

We denote by $\pi_1 : \mathbb{P}_N \times X \rightarrow \mathbb{P}_N$ and by $\pi_2 : \mathbb{P}_N \times X \rightarrow X$ the natural projections and we set $E_0 = \pi_{1*}(\mathcal{K} \wedge \pi_2^* T)$ which is a closed (q, q) -current of order 0 in \mathbb{P}_N . Now we calculate the intersection number

$$\int_{\mathbb{P}_N} E_0 \wedge i_{0*} u = \int_{\mathbb{P}_N \times X} \mathcal{K} \wedge \pi_1^* i_{0*} u \wedge \pi_2^* T = \int_{\mathbb{P}_N \times X} \mathcal{K} \wedge [X \times X] \wedge \pi_1^* \tilde{u} \wedge \pi_2^* T$$

where \tilde{u} is a smooth extension of the smooth differential form u to \mathbb{P}_N .

In view to express the current $[X \times X] = \pi_1^* [X]$ in $\mathbb{P}_N \times X$, we write $X = \{P_j = 0\}$ with (P_j) a finite family of homogeneous polynomials on \mathbb{C}^{N+1} and $m_j = \deg P_j$ for each j . We set $\rho([x]) = \sum_j \frac{|P_j(x)|^2}{\|x\|^{2m_j}}$ which satisfies $X = \rho^{-1}(0)$ in \mathbb{P}_N and

$$[X] = \delta_\rho \frac{(\text{dd}^c \rho)^{N-n}}{2^{N-n}(N-n)} = \lim_{\lambda \rightarrow 0^+} \left\{ \lambda \rho^{\lambda - (N-n)} \frac{(\text{dd}^c \rho)^{N-n}}{2^{N-n}(N-n)} \right\}$$

with δ_ρ the distribution which is the residue at the pole $-(N-n)$ of the meromorphic map $\mathbb{C} \ni \lambda \rightarrow \rho^\lambda$.

We define in $\mathbb{P}_N \times X$ the current product of closed currents

$$\mathcal{K} \wedge [X \times X] = (\delta_\rho \circ \pi_1) \mathcal{K} \wedge \pi_1^* \left(\frac{(\text{dd}^c \rho)^{N-n}}{2^{N-n}(N-n)} \right)$$

which is closed too and annihilated by the $P_j \circ \pi_1$ and their conjugates. With

$$\alpha = \text{d}^c \left(\frac{1}{2^n(n-1)} \left(\frac{\text{dd}^c \|f\|^2}{\|f\|^2} \right)^{n-1} \right)_{|\mathbb{P}_N \times X}$$

we have $-(\text{dd}^c \log \|f\|)_{|\mathbb{P}_N \times X}^n = \text{d}\alpha$ and we set actually

$$-(\delta_\rho \circ \pi_1)(\text{dd}^c \log \|f\|)_{|\mathbb{P}_N \times X}^n = \text{d}((\delta_\rho \circ \pi_1)\alpha) - \text{d}(\delta_\rho \circ \pi_1) \wedge \alpha$$

where the products with α are obtained by the division theorem. Otherwise we have

$$-(\delta_\rho \circ \pi_1)(\text{dd}^c \log \|f\|)_{|\mathbb{P}_N \times X}^n = - \lim_{\lambda \rightarrow 0^+} \lambda (\rho \circ \pi_1)^{\lambda - (N-n)} (\text{dd}^c \log \|f\|)_{|\mathbb{P}_N \times X}^n$$

where $-(\text{dd}^c \log \|f\|)_{|\mathbb{P}_N \times X}^n$ is the direct image by a blowing up of a closed smooth differential form and the weak limit when $\lambda \rightarrow 0^+$ is calculated thanks to the Bernstein equation.

So there is a closed (n, n) -current \mathcal{L} in $\mathbb{P}_N \times X$ such that $\text{supp}(\mathcal{L}|_{X \times X}) = D_X$ and

$$\mathcal{K} \wedge [X \times X] = i_*([D_X] - \mathcal{L}|_{X \times X})$$

with $i : X \times X \rightarrow \mathbb{P}_N \times X$ the natural injection. The order of \mathcal{L} is not equal to 0, because the codimension of the polar set changes by blowing up. In such a way we can express the restriction

$$E_{0|X} = T - (\pi_{1|X \times X})_* (\mathcal{L}|_{X \times X} \wedge (\pi_{2|X \times X})^* T).$$

Therefore we take $S = \pi_{1*}((\mathcal{K} + \mathcal{L}) \wedge \pi_2^* T)$ which satisfies $S|_X = T$ in the sense of the Proposition 4.

Lastly the differential form u_φ is given by $u_\varphi = \pi_{2*}((\mathcal{K} + \mathcal{L}) \wedge \pi_1^* \varphi)$.

References

- [1] Demailly J.-P., *Regularization of closed positive currents and intersection theory*, J. Algebraic Geometry 1 (1992) 361–409.
- [2] Demailly J.-P., *Analytic methods in algebraic geometry*, Surveys of Modern Mathematics 1, International Press, 2012.
- [3] Demailly J.-P., Laurent-Thiébaud C., *Formules intégrales pour les formes différentielles de type (p, q) dans les variétés de Stein*, Ann. Sci. École Norm. Sup. (4) 20 (1987) 579–598.
<https://doi.org/10.24033/asens.1545>
- [4] Dinh T.-C., Sibony N., *Regularization of currents and entropy*, Ann. Sci. École Norm. Sup. (4) 37 (2004), no. 6, 959–971.
<https://doi.org/10.1016/j.ansens.2004.09.002>
- [5] Dinh T.-C., Sibony N., *Super-potentials of positive closed currents, intersection theory and dynamics*, Acta Math. 203 (2009), no. 1, 1–82.
<https://doi.org/10.1007/s11511-009-0038-7>
- [6] Flenner H., Manaresi M., *A length formula for the multiplicity of distinguished components of intersections*, J. Pure Appl. Algebra 165 (2001), no. 2, 155–168.
[https://doi.org/10.1016/S0022-4049\(00\)00186-9](https://doi.org/10.1016/S0022-4049(00)00186-9)
- [7] Henkin G.M., Leiterer J., *Andreotti-Grauert theory by integral formulas*, Progress in Math., Vol. 74, Birkhäuser, 1988.
- [8] Laurent-Thiébaud C., *Produits de courants et formule des résidus*, Bull. Sci. Math. (2) 105 (1981), no. 2, 113–158.
- [9] Méo M., *Chow forms and Hodge cohomology classes*, C. R. Acad. Sci. Paris, Ser. I 352 (2014) 339–343.
<http://dx.doi.org/10.1016/j.crma.2014.01.012>
- [10] Peternell T., *Submanifolds with ample normal bundles and a conjecture of Hartshorne*, Contemp. Math. 496, 2009, 317–330.
- [11] Poly J.-B., *Sur l’homologie des courants à support dans un ensemble semi-analytique*, Bull. Soc. Math. France Suppl. Mém. 38 (1974) 35–43.
<https://eudml.org/doc/94688>
- [12] Schwartz L., *Analyse, Deuxième partie : Topologie générale et analyse fonctionnelle*, Hermann, 1970.
- [13] Varouchas J., *Sur l’image d’une variété kählérienne compacte*, Fonctions de plusieurs variables complexes V, Lecture Notes in Math. 1188, 1986, 245–259.