



**HAL**  
open science

## Hydrogen phenylphosphonate phenylphosphonic acid water 1:1:1 co-crystal and hydrogen phenylarsonate cyclohexylammonium salts: synthesis and crystal elucidation

Mouhamadou Birame Diop, Modou Sarr, Mouhamadou Sembene Boye, Aminata Diasse-Sarr, Libasse Diop, Stanislav Pechev, Philippe Guionneau

### ► To cite this version:

Mouhamadou Birame Diop, Modou Sarr, Mouhamadou Sembene Boye, Aminata Diasse-Sarr, Libasse Diop, et al.. Hydrogen phenylphosphonate phenylphosphonic acid water 1:1:1 co-crystal and hydrogen phenylarsonate cyclohexylammonium salts: synthesis and crystal elucidation. International journal of advanced research, 2022, 10 (05), pp.690-700. 10.21474/IJAR01/14756 . hal-03770005

**HAL Id: hal-03770005**

**<https://hal.science/hal-03770005>**

Submitted on 6 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Journal Homepage: [-www.journalijar.com](http://www.journalijar.com)

## INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/14756

DOI URL: <http://dx.doi.org/10.21474/IJAR01/14756>



### RESEARCH ARTICLE

#### HYDROGEN PHENYLPHOSPHONATE PHENYLPHOSPHONIC ACID WATER 1:1:1 CO-CRYSTAL AND HYDROGEN PHENYLARSONATE CYCLOHEXYLAMMONIUM SALTS: SYNTHESIS AND CRYSTAL ELUCIDATION

Mouhamadou Birame Diop<sup>1</sup>, Modou Sarr<sup>1</sup>, Mouhamadou Sembene Boye<sup>1,2</sup>, Aminata Diassé-Sarr<sup>1</sup>, Libasse Diop<sup>1</sup>, Stanislav Pechev<sup>3</sup> and Philippe Guionneau<sup>3</sup>

1. Laboratoire de Chimie Minérale et Analytique (LA.CHI.MI.A.), Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar, Sénégal.
2. Département de Physique-Chimie, Faculté des Sciences et Technologies de l'Education et de la Formation (FASTEF), Université Cheikh Anta Diop, Dakar, Sénégal.
3. Institut de Chimie de la Matière Condensée de Bordeaux, CNRS-Université de Bordeaux, 87 Avenue du Docteur A. Schweitzer 33608 Pessac, France.

#### Manuscript Info

##### Manuscript History

Received: 19 March 2022

Final Accepted: 25 April 2022

Published: May 2022

##### Key words:-

Single Crystal X-ray,  
Phenylphosphonate, Phenylarsonate,  
Cyclohexylammonium, 2D Layer-Like  
Structure

#### Abstract

One pot reactions of phenylphosphonic acid or phenylarsonic acid with cyclohexylamine led to two new organic-inorganic hybrid salts  $[\text{C}_6\text{H}_{14}\text{N}][\text{C}_6\text{H}_5\text{PO}_3\text{H.C}_6\text{H}_5\text{PO}_3\text{H}_2.\text{H}_2\text{O}]$  (**1**) and  $[(\text{C}_6\text{H}_{14}\text{N})(\text{C}_6\text{H}_5\text{AsO}_3\text{H})]$  (**2**) which were investigated by single-crystal X-ray diffraction analysis. Salt **1** crystallizes in the monoclinic system, space group  $\text{P}2_1/n$  with  $a = 12.3963$  (4),  $b = 6.1123$  (2),  $c = 28.2081$  (8) Å,  $\beta = 90.426$  (1)°,  $V = 2137.27$  (12) Å<sup>3</sup> and  $Z = 4$ . Salt **2** crystallizes in the monoclinic system, space group  $\text{P}2_1/c$  with  $a = 12.3991$  (5),  $b = 6.5048$  (2),  $c = 18.5466$  (7) Å,  $\beta = 105.824$  (2)°,  $V = 1439.16$  (9) Å<sup>3</sup> and  $Z = 4$ . In the co-crystal **1**, the hydrogen phenylphosphonate anions are organized into dimers *via* O–H···O H-bonds, whereas the phenylphosphonic acid and the water molecules are connected, through O<sub>water</sub>–H···O and O–H···O hydrogen bonding interactions, to form dimers. Inter-species O–H···O hydrogen bonds connect the alternating dimers into anionic infinite chains. The 1D infinite ribbons are linked into an anionic 2D layer-like structure, by means of water molecules through O<sub>water</sub>–H···O and O–H···O<sub>water</sub>. The cyclohexylammonium interacts *via* N–H···O hydrogen bonds, to consolidate the compactness of the 2D layer-like structure. In the crystal of salt **2**, the hydrogen phenylarsonate anions are organized into dimers through O–H···O H-bonds. Hydrogen bonded dimers are connected by the cyclohexylammonium cations, through N–H···O hydrogen bonds, leading to the formation of an infinite ribbon. The 1D hydrogen bonded ribbons are interlinked through N–H···O hydrogen bonding interactions giving rise to a 2D layer-like structure.

Copy Right, IJAR, 2022.. All rights reserved.

**Corresponding Author:- Mouhamadou Birame Diop**

Address:- Laboratoire de Chimie Minérale et Analytique (LA.CHI.MI.A.), Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar, Sénégal.

## Introduction:-

In the past, several phosphonate based metal complexes investigations exhibited catalysis (Ranocchiaro and van Bokhoven, 2011; Hu and Zhao, 2017; White *et al.*, 2009), gas sorption and separation (Llewellyn *et al.*, 2015; Taddei *et al.*, 2016; Hermer *et al.*, 2017; Oschatz and Antonietti, 2018; Li and Chen, 2016), energy storage and conversion applications (Winter and Brodd, 2004). The solid state luminescent properties of a mixed phenylphosphonic acid/singly protonated phenylphosphonate copper complex (Latham *et al.*, 2009) as well as magnetic properties (Bao and Zheng, 2016; Hamchaoui *et al.*, 2013) and several metal phosphonates surface photovoltage, semi conductor and luminescent properties (Cai *et al.*, 2018; Zhu *et al.*, 2016; Shi *et al.*, 2014; Zhang *et al.*, 2020; Firmino *et al.*, 2020; Ayhan *et al.*, 2020; Jiao *et al.*, 2016; Chu *et al.*, 2013; Dai *et al.*, 2014; Yang *et al.*, 2016; Zhao *et al.*, 2016) have also been reported. New materials either usable as electrodes or electrolyte components has been reported too (Peng and Chen, 2009). Related phenylarsonates with diverse substituents at the ortho, meta or para position, have widely been investigated, though hydrogen phenylarsonate compounds are most scarcely encountered in the literature (Smith and Wermuth, 2010; Reck and Schmitt, 2012; Yi *et al.*, 2009; Padalwar and Vidyasagar, 2016; Padalwar *et al.*, 2013; Reddy *et al.*, 2006). Indeed, several singly protonated phenylarsonate crystal compounds are reported to date, according to the Cambridge Structural Database CSD (Smith and Wermuth, 2010; Reck and Schmitt, 2012; Yi *et al.*, 2009; Padalwar and Vidyasagar, 2016; Padalwar *et al.*, 2013; Reddy *et al.*, 2006). Tetrabutylammonium hydrogen phenylarsonate–phenylarsonic acid (1/1) exhibits an acidic anionic infinite chain with alternating species grown from the junction of the dimers  $\text{PhAsO}_3\text{H}/\text{PhAsO}_3\text{H}_2$  through  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds (Reck and Schmitt, 2012).  $\text{A}(\text{HO}_3\text{PPh})(\text{H}_2\text{O}_3\text{PPh})$  (A = Li, Na, K, Rb, Cs, Tl,  $\text{NH}_4$ ) phenylphosphonates as well as  $\text{Na}(\text{HO}_3\text{AsPh})(\text{H}_2\text{O}_3\text{AsPh})$  phenylarsonate crystal structures have been reported (Padalwar *et al.*, 2013); they display hydrogen bonding interaction patterns between the singly protonated forms, nevertheless the hydrogen bonding pattern investigations between species have not been deepened. In 2010, Smith and Wermuth reported the crystal structure of guanidinium phenylarsonate–guanidine–water (1/1/2) which describes extended hydrogen bonding patterns, but without usually encountered hydrogen bonding interactions between the singly protonated phenylarsonate ions. Several phosphonate based metal complexes with mixed phenylphosphonic acid/singly protonated phenylphosphonate such as  $[\text{CuX}(\text{phen})_2][(\text{C}_6\text{H}_5\text{PO}(\text{OH})_2)(\text{C}_6\text{H}_5\text{PO}_2(\text{OH}))]$  (where X = Cl, Br, I, NCS) and  $[\text{Cu}^{\text{II}}(\text{phen})_2\text{X}][((\text{OH})_2\text{OPC}_6\text{H}_5)((\text{OH})\text{O}_2\text{PC}_6\text{H}_5)]$  (where X = I, Br, Cl, NCS) have earlier been reported, describing hydrogen bonded acidic ribbons (Clarke *et al.*, 2005; Latham *et al.*, 2007; Rao and Vidyasagar, 2005). The Dakar group has been involved in single crystal X-ray diffraction investigations of phosphonates and cyclohexylammonium co-crystallization, by describing various phenylphosphonate crystal structures exclusive dealing (Diop *et al.*, 2019, 2013, 2012, 2011; Sarr *et al.*, 2012), but also some cyclohexylammonium salts (Sarr *et al.*, 2020, 2015a, 2015b, 2013). Continuing to widen our contribution in this field, one pot reactions of phenylphosphonic acid,  $\text{C}_6\text{H}_5\text{PO}_3\text{H}_2$  and nickel(II) chloride hexahydrate,  $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$  or phenylarsonic acid,  $\text{C}_6\text{H}_5\text{AsO}_3\text{H}_2$  with cyclohexylamine,  $\text{C}_6\text{H}_{11}\text{N}$  were undertaken in solution media. These reactions led to the isolation of namely,  $[\text{C}_6\text{H}_{14}\text{N}][\text{C}_6\text{H}_5\text{PO}_3\text{H}\cdot\text{C}_6\text{H}_5\text{PO}_3\text{H}_2\cdot\text{H}_2\text{O}]$  **1** and  $[(\text{C}_6\text{H}_{14}\text{N})(\text{C}_6\text{H}_5\text{AsO}_3\text{H})]$  **2** whose structure elucidations were carried out by single-crystal X-ray diffraction analysis.

## Experimental section:-

Reagents phenylphosphonic acid,  $\text{C}_6\text{H}_5\text{PO}(\text{OH})_2$  (98 % purity), cyclohexylamine,  $\text{C}_6\text{H}_{11}\text{NH}_2$  (99 % purity), nickel(II) chloride hexahydrate,  $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$  (99.9 % purity) and phenylarsonic acid,  $\text{C}_6\text{H}_5\text{AsO}(\text{OH})_2$  (97 % purity) were purchased from Sigma-Aldrich, Steinheim am Albuch, Germany and used without any further purification. The X-ray crystallographic data for compounds **1** and **2** were collected using a Bruker Kappa APEX-II diffractometer operating at 150 K and a Nonius Kappa CCD working at 293 K, respectively.

## Synthesis and isolation of $[\text{C}_6\text{H}_{14}\text{N}][\text{C}_6\text{H}_5\text{PO}_3\text{H}\cdot\text{C}_6\text{H}_5\text{PO}_3\text{H}_2\cdot\text{H}_2\text{O}]$ (**1**):-

The pooling of methanolic solutions containing two equivalents of phenylphosphonic acid,  $\text{C}_6\text{H}_5\text{PO}_3\text{H}_2$  (0.816 g; 5.16 mmol), one equivalent of cyclohexylamine,  $\text{C}_6\text{H}_{11}\text{NH}_2$  (0.8547 g/cm<sup>3</sup>; 0.3 mL; 2.58 mmol) and one equivalent of nickel(II) chloride hexahydrate,  $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$  (0.613 g; 2.58 mmol) has afforded a cloudy solution. The resultant solution was subsequently stirred at room temperature for more than 2h then filtered. After some days of slow evaporation of the filtrate, colorless plate-like crystals suitable for single-crystal X-ray structure determination were collected within the solvent and characterized as  $[\text{C}_6\text{H}_{14}\text{N}][\text{C}_6\text{H}_5\text{PO}_3\text{H}\cdot\text{C}_6\text{H}_5\text{PO}_3\text{H}_2\cdot\text{H}_2\text{O}]$  (**1**).

Compound **1** can also be synthesized by directly reacting two equivalents of phenylphosphonic acid,  $\text{C}_6\text{H}_5\text{PO}_3\text{H}_2$  with one equivalent of cyclohexylamine,  $\text{C}_6\text{H}_{11}\text{NH}_2$ .

**Synthesis and isolation of [(C<sub>6</sub>H<sub>14</sub>N)(C<sub>6</sub>H<sub>5</sub>AsO<sub>3</sub>H)] (2):-**

Salt **2** has been isolated by reacting one equivalent of phenylarsonic acid, C<sub>6</sub>H<sub>5</sub>AsO<sub>3</sub>H<sub>2</sub> (3.124 g; 15 mmol) with one equivalent of cyclohexylamine, C<sub>6</sub>H<sub>11</sub>NH<sub>2</sub> (1.7 mL; 15 mmol), in water. The resultant solution was subsequently stirred about 2h. Colorless prism-like crystals suitable for a single-crystal X-ray diffraction study were obtained, after some days of slow solvent evaporation at room temperature and characterized as [(C<sub>6</sub>H<sub>14</sub>N)(C<sub>6</sub>H<sub>5</sub>AsO<sub>3</sub>H)] (**2**).

**X-ray crystallography of [C<sub>6</sub>H<sub>14</sub>N][C<sub>6</sub>H<sub>5</sub>PO<sub>3</sub>H.C<sub>6</sub>H<sub>5</sub>PO<sub>3</sub>H<sub>2</sub>.H<sub>2</sub>O] (1):-**

A crystal of approximate dimensions 0.246×0.103×0.050 mm was used for data collection. The X-ray crystallographic data were collected using a Bruker Kappa Apex II diffractometer operating at T = 150 (2) K. Data were measured using  $\varphi$  and  $\omega$  scans using MoK $\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ) using a collection strategy to obtain a hemisphere of unique data determined by *Apex2* (Bruker AXS Inc., Madison, WI 53719-1173., 2012). Cell parameters were determined and refined using the *SAINT* program (*SAINT* (2012)). Data were corrected for absorption by multi-scan absorption using *SADABS* (*SADABS2012/1*). The structure of **1** was solved by dual-space analysis using *SHELXT* (Sheldrick, 2015) and refined using least-squares minimization with *SHELXL97* (Sheldrick, 2008).

Programs used for the representation of the molecular and crystal structures: *Olex2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2008). The Crystallographic data and experimental details for structural analyses are summarized in Table 1. Selected bond lengths and angles are listed in the caption of Figure 1.

CCDC 2172774 (**1**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by e-mailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44(0)1223-336033.

**X-ray crystallography of [(C<sub>6</sub>H<sub>14</sub>N)(C<sub>6</sub>H<sub>5</sub>AsO<sub>3</sub>H)] (2):-**

A crystal of approximate dimensions 0.22×0.12×0.02 mm was used for data collection. The X-ray crystallographic data were collected using a Nonius Kappa CCD diffractometer operating at T = 293 (2) K. Data were measured using  $\varphi$  and  $\omega$  scans using MoK $\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ) using a collection strategy to obtain a hemisphere of unique data determined by *COLLECT* (Hooft, Nonius Kappa CCD, Nonius BV: Delft (The Netherlands), 2003). Cell parameters were determined and refined using *DIRAX* (Duisenberg, 1992). Data were corrected for absorption correction using *EVAL CCD* (Duisenberg *et al.*, 2003). The structure of **2** was solved using *SHELXS* (Sheldrick, 2008) and refined using least-squares minimization with *SHELXL97* (Sheldrick, 2008).

Programs used for the representation of the molecular and crystal structures: *Olex2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2008). The Crystallographic data and experimental details for structural analyses are summarized in Table 1. Selected bond lengths and angles are listed in the caption of Figure 5.

CCDC 2172775 (**2**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by e-mailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44(0)1223-336033.

**Table 1:-** Crystal data and structure refinement for compounds (**1**) and (**2**).

Parameters	Compound	
	(1)	(2)
Empirical formula	C <sub>18</sub> H <sub>29</sub> NO <sub>7</sub> P <sub>2</sub>	C <sub>12</sub> H <sub>20</sub> AsNO <sub>3</sub>
Formula weight	433.36	301.21
Temperature/K	150.00	293(2)
Crystal system	monoclinic	monoclinic
Space group	P2 <sub>1</sub> /n	P2 <sub>1</sub> /c
a/Å	12.3963(4)	12.3991(5)
b/Å	6.1123(2)	6.5048(2)
c/Å	28.2081(8)	18.5466(7)
$\alpha$ /°	90	90



$\beta/^\circ$	90.4260(10)	105.824(2)
$\gamma/^\circ$	90	90
Volume/ $\text{\AA}^3$	2137.27(12)	1439.16(9)
Z	4	4
$\rho_{\text{calc}}/\text{g/cm}^3$	1.347	1.390
$\mu/\text{mm}^{-1}$	0.242	2.359
F(000)	920	624
Crystal size/ $\text{mm}^3$	$0.246 \times 0.103 \times 0.05$	$0.22 \times 0.12 \times 0.02$
Radiation	MoK $\alpha$ ( $\lambda = 0.71073$ )	MoK $\alpha$ ( $\lambda = 0.71073$ )
2 $\theta$ range for data collection/ $^\circ$	3.58–62.442	6.406–54.996
Index ranges	$-18 \leq h \leq 18$ , $-8 \leq k \leq 8$ , $-41 \leq l \leq 41$	$-14 \leq h \leq 16$ , $-8 \leq k \leq 8$ , $-24 \leq l \leq 24$
Reflections collected	77999	9364
Independent reflections	6887 [ $R_{\text{int}} = 0.0543$ , $R_{\text{sigma}} = 0.0270$ ]	3306 [ $R_{\text{int}} = 0.0333$ , $R_{\text{sigma}} = 0.0350$ ]
Data/restraints/parameters	6887/7/369	3306/453/252
Goodness-of-fit on $F^2$	1.069	1.060
Final R indexes [ $I > 2\sigma(I)$ ]	$R_1 = 0.0379$ , $wR_2 = 0.0871$	$R_1 = 0.0397$ , $wR_2 = 0.1003$
Final R indexes [all data]	$R_1 = 0.0543$ , $wR_2 = 0.0937$	$R_1 = 0.0540$ , $wR_2 = 0.1074$
Largest diff. peak/hole / $e \text{\AA}^{-3}$	0.43/-0.36	0.71/-0.55

<sup>a</sup>  $R_1 = \Sigma(|F_o| - |F_c|) / \Sigma |F_o|$ ; <sup>b</sup>  $wR_2 = [\Sigma w(F_o^2 - F_c^2)^2 / \Sigma w(F_o^2)^2]^{1/2}$  where  $w = 1/[\sigma^2(F_o^2) + (0.0435P)^2 + 0.7237P]$  for **1** and  $w = 1/[\sigma^2(F_o^2) + (0.0506P)^2 + 0.9955P]$  for **2** with  $P = (F_o^2 + 2F_c^2)/3$ ; <sup>c</sup> goodness of fit =  $[\Sigma w(F_o^2 - F_c^2)^2 / (N_o - N_v)]^{1/2}$ .


## Results and Discussion:-

Compound **1** was isolated from the reaction in methanol, from a 1:1 molar ratio between phenylphosphonic acid and cyclohexylamine in the presence of nickel(II) chloride hexahydrate. Colorless single crystals grown from the filtrate (Eq. 1) and were characterized as **1**,  $[\text{C}_6\text{H}_{14}\text{N}][\text{C}_6\text{H}_5\text{PO}_3\text{H.C}_6\text{H}_5\text{PO}_3\text{H}_2.\text{H}_2\text{O}]$ . Several X-ray investigations involving both phenylphosphonic acid and singly protonated phenylphosphonate were previously investigated (Smith and Wermuth, 2010; Wilke *et al.*, 2016a, 2016b).

The compound **2** was isolated from the reaction in water, from a 1:1 molar ratio between amounts of phenylarsonic acid and cyclohexylamine. This yielded the compound  $[(\text{C}_6\text{H}_{14}\text{N})(\text{C}_6\text{H}_5\text{AsO}_3\text{H})]$  (**2**) collected as colorless crystals (Eq. 2). To our knowledge, very few examples of non-substituted hydrogen arsonate containing compounds have been isolated and characterized by X-ray study (Yi *et al.*, 2009; Padalwar and Vidyasagar, 2016; Padalwar *et al.*, 2013; Adelani *et al.*, 2013). The yields of the reactions reported herein are 53 and 95%, respectively.




Compound **1**,  $[\text{C}_6\text{H}_{14}\text{N}][\text{C}_6\text{H}_5\text{PO}_3\text{H.C}_6\text{H}_5\text{PO}_3\text{H}_2.\text{H}_2\text{O}]$ , crystallizes as colorless plate-like crystals in the monoclinic space group  $P2_1/n$ . The asymmetric unit is comprised of a cyclohexylammonium, a hydrogen phenylphosphonate anion, a molecule of phenylphosphonic acid, and a water molecule both connected through N–H $\cdots$ O and O–H $\cdots$ O hydrogen bonds. The molecular structure of the compound **1** is represented in Figure 1.




**Fig. 1:-** Molecular view of **1** showing 50% probability ellipsoids for atoms and the crystallographic numbering scheme [atom color code: C, black; H, white; O, red; N, blue; P, pink]. Selected bond lengths and angles (Å, deg): P1–O1 1.5097 (9), P1–O2 1.5646 (10), P1–O3 1.5152 (9), P1–C1 1.7959 (13), P2–O4 1.5509 (10), P2–O5 1.5511 (10), P2–O6 1.4915 (10), P2–C7 1.7912 (13), C13–N1 1.4987 (16); O1–P1–O2 112.42 (5), O1–P1–O3 115.21 (5), O3–P1–O2 105.20 (5), O1–P1–C1 108.64 (6), O2–P1–C1 106.95 (6), O3–P1–C1 105.20 (5), O5–P2–O4 105.84 (5), O6–P2–O5 114.05 (5), O6–P2–O4 111.38 (6), O4–P2–C7 106.95 (6), O5–P2–C7 108.06 (6), O6–P2–C7 110.21 (6), N1–C13–C18 110.25 (10), N1–C13–C14 109.72 (10).

The P–O length values are in accordance with the published values (Diop *et al.*, 2013, 2012, 2011; Sarr *et al.*, 2012, 2020; Beckmann *et al.*, 2003) and evidence single (O1, O3) and double (O2, O4) bonds. The geometric parameters within the cyclohexylammonium well corroborate the yet reported values (Sarr *et al.*, 2020, 2015a, 2015b; Lemmerer, 2011; Grimball *et al.*, 2017; Tang *et al.*, 2015). In the crystal structure of compound **1**, the hydrogen phenylphosphonate anions are organized into dimers through strong O–H···O hydrogen bonds forming  $R_2^2(8)$  rings (see Figure 2). The phenylphosphonic acid and the water molecules likewise form dimers through three strong  $O_{\text{water}}\text{--H}\cdots\text{O}$  and one  $\text{O--H}\cdots O_{\text{water}}$  hydrogen bonds affording  $R_4^3(10)$  rings (see Figure 2). Inter species O–H···O hydrogen bonding interactions connect these dimers into anionic infinite chains in which they alternate. Each hydrogen bonded anionic chain is linked to its neighbours *via*  $O_{\text{water}}\text{--H}\cdots\text{O}$  and  $\text{O--H}\cdots O_{\text{water}}$  hydrogen bonds involving the phenylphosphonic acid and water molecules, leading to the formation of an anionic layer (see Figure 2 and Table 2).




**Fig. 2:-** Molecular view of **1** showing 50% probability ellipsoids for atoms, showing the two different dimers and chains connected into an anionic layer network. Phenyl groups are omitted for clarity.

The cyclohexylammonium cations slipped in between the chains that they link *via* N–H···O hydrogen bonds, to consolidate the compactness and the stability of the 2D network. These hydrogen bonding interactions enable H-bonded cyclic patterns: 8-membered  $R_2^2(8)$  and 10-membered  $R_4^3(10)$  rings alternating within the chains and, 8-membered  $R_3^3(8)$ , 10-membered  $R_4^3(10)$ , 16-membered  $R_6^4(16)$ , and 14-membered  $R_5^4(14)$  rings from the junction of chains and cyclohexylammonium ions, leading to a layer-like structure along [101] plane (Figure 3). However, this compound exhibits a curve pleated sheet whose molecular packing diagram is depicted in Figure 4.



**Fig. 3:-** Molecular view of **1** showing 50% probability ellipsoids for atoms, showing the interactions of cations slipped in between the chains growing the layer network. Phenyl and cyclohexyl groups are omitted for clarity.




**Fig. 4:-** Molecular partial packing diagram of **1** showing 50% probability ellipsoids for atoms, showing the pleated layer network.

**Table 2:-** Hydrogen bond and interaction geometries (Å, deg) in the crystal of **1** (symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x, y-1, z$ ; (iii)  $x, y+1, z$ ; (iv)  $-x+1/2, y+1/2, -z+3/2$ ).

D–H···A	d(D–H)	d(H···A)	d(D···A)	$\angle$ (D–H···A)
O4–H4···O3	0.88 (2)	1.60 (2)	2.4691 (14)	171 (3)
O2–H2···O1 <sup>i</sup>	0.85 (2)	1.75 (2)	2.6001 (13)	177 (2)
N1–H1A···O4	0.90 (1)	2.20 (2)	2.9570 (15)	141 (1)

N1–H1B...O3 <sup>ii</sup>	0.91 (1)	1.89 (1)	2.7982 (14)	172 (2)
N1–H1C...O1	0.91 (1)	1.86 (1)	2.7662 (14)	175 (2)
O7–H7A...O6 <sup>iii</sup>	0.87 (2)	1.81 (2)	2.6824 (14)	176 (2)
O7–H7B...O6 <sup>iv</sup>	0.88 (2)	1.84 (2)	2.7126 (13)	174 (2)
O5–H5...O7	0.83 (2)	1.68 (2)	2.4995 (14)	173 (3)


The compound **2**, [(C<sub>6</sub>H<sub>14</sub>N)(C<sub>6</sub>H<sub>5</sub>AsO<sub>3</sub>H)], crystallizes as colorless prism-like crystals in the monoclinic space group P2<sub>1</sub>/c. The asymmetric unit contains one cyclohexylammonium and one hydrogen phenylarsonate ions. The molecular structure of the compound **2** is shown in Figure 5.




**Fig. 5:-** Molecular view of **2** showing 50% probability ellipsoids for atoms and the crystallographic numbering scheme adopted [atom color code: C, black; H, white; O, red; N, blue; As, pink]. Only one fragment of the disordered species is shown. Selected bond lengths and angles (Å, deg): O1–As1 1.662 (2), O2–As1 1.649 (2), O3–As1 1.719 (2), As1–C1A 1.927 (5), As1–C1B 1.890 (5), N1A–C7A 1.477 (7), N1B–C7B 1.477 (7); O1–As1–O2 115.53 (11), O1–As1–O3 111.13 (11), O1–As1–C1A 107.3 (2), O1–As1–C1B 108.5 (2), O2–As1–O3 103.84 (11), O2–As1–O1 104.03 (11), O2–As1–C1A 110.6 (2), O2–As1–C1B 110.2 (2), O3–As1–C1A 108.3 (2), O3–As1–C1B 107.3 (2), N1A–C7A–C8A 109.3 (6), N1A–C7A–C12A 109.3 (6), N1B–C7B–C8B 117.1 (19), N1B–C7B–C12B 115.8 (19).

The As–O distances indicate single and double bonds and are in the expected range (Smith and Wermuth, 2010; Reck and Schmitt, 2012; Yi *et al.*, 2009; Padalwar and Vidyasagar, 2016; Padalwar *et al.*, 2013; Reddy *et al.*, 2006). The geometric parameters within the cyclohexylammonium are closely similar to those in compound **1**, and are in accordance with the earlier published values (Sarr *et al.*, 2020, 2015a, 2015b; Lemmerer, 2011; Grimball *et al.*, 2017; Tang *et al.*, 2015). In the crystal of compound **2**, the hydrogen phenylarsonate anions are organized into dimers through strong O–H...O hydrogen bonds exhibiting a R<sub>2</sub><sup>2</sup>(8) ring. Each hydrogen bonded dianionic dimer is linked to its neighbor *via* two cyclohexylammonium cations, through N–H1A...O and N–H1B...O hydrogen bonds, leading to the formation of an infinite chain which describes two alternating and fused rings R<sub>2</sub><sup>2</sup>(8) and R<sub>6</sub><sup>4</sup>(16) (see Figure 6 and Table 3). In the chains, the cyclohexylammonium cations and the dianionic dimers alternate. The chains are then linked *via* the remaining nitrogen H atoms of the cations through N–H1C...O2 hydrogen bonds, leading to a layer-like structure along [100] plane (Figure 6). The interconnections between chains exhibit only 10-membered hydrogen bonded rings R<sub>4</sub><sup>3</sup>(10) (see Figure 6). The molecular packing diagram of the compound **2** is depicted in Figure 7.





**Fig. 6:-** Molecular view of **2** showing 50% probability ellipsoids for atoms, showing the propagation of the infinite chain network. Phenyl and cyclohexyl groups are omitted for clarity.



**Fig. 7:-** Molecular packing of **2**, approximately along [010] direction, showing 50% probability ellipsoids for atoms, showing the propagation of the layer network.

**Table 3:-** Hydrogen bonds in crystals of **2** (symmetry codes: (i)  $-x+2, -y, -z$ ; (ii)  $x, y+1, z$ ; (iii)  $-x+2, y+1/2, -z+1/2$ ).

D-H...A	d(D-H)	d(H...A)	d(D...A)	$\angle$ (D-H...A)
O3-H3...O1 <sup>i</sup>	0.82	1.80	2.618 (3)	175
N1A-H1AA...O1 <sup>ii</sup>	0.89	1.83	2.719 (12)	177
N1A-H1AB...O2 <sup>iii</sup>	0.89	1.85	2.740 (13)	176
N1A-H1AC...O2	0.89	1.85	2.728 (12)	169
N1B-H1BA...O1 <sup>ii</sup>	0.89	1.95	2.83 (4)	170
N1B-H1BB...O2 <sup>iii</sup>	0.89	1.97	2.81 (5)	157
N1B-H1BC...O2	0.89	1.95	2.83 (4)	172

### Conclusion:-

The desired nickel phosphonate complex has not been isolated by carrying the reaction between cyclohexylamine, phenylphosphonic acid and nickel(II) dichloride hexahydrate. Two cyclohexylammonium salts of phenylphosphonic acid and phenylarsonic acid have been synthesized by one pot reaction and investigated by X-ray diffraction analysis. Both compounds **1** and **2** exhibit extended hydrogen bonds, even those in **1** are more extensive, evidencing five ring types *ie*  $R_2^2(8)$ ,  $R_3^3(8)$ ,  $R_4^3(10)$ ,  $R_5^4(14)$  and  $R_6^4(16)$  whereas **2** describes only three ring types *viz*  $R_2^2(8)$ ,  $R_3^3(10)$  and  $R_6^4(16)$ . In **1**, phenylphosphonate anions as well as phenylphosphonic acid and the water molecules lonely form H-bonded dimers connected into infinite ribbons which are linked into an anionic 2D layer-like structure. The cyclohexylammonium cations fit in the anionic 2D layer, strengthening its stability and compactness. Pooling to form dimers, the phenylarsonate ions interconnections with cyclohexylammonium in **2** describes infinite ribbons whose hydrogen bonding linkages give rise to a 2D layer-like structure. In the attempt to synthesize and characterize new complexes of various metallic reagents, further works involving salts **1** and **2** are in progress.

### Acknowledgment:-

The authors gratefully thank Prof. Dr. Thierry Maris, Université de Montréal, Québec (Canada), the Cheikh Anta Diop University – Dakar (Senegal) and the University of Bordeaux (France) for equipment facilities.

### References:-

1. Ranocchiari, M. and van Bokhoven, J. A. (2011): Catalysis by metal-organic frameworks: Fundamentals and opportunities. *Phys. Chem. Chem. Phys.*, 13: 6388–6396.
2. Hu, Z. and Zhao, D. (2017): Metal-organic frameworks with Lewis acidity: Synthesis, characterization, and catalytic applications. *CrystEngComm*, 19, 4066–4081.
3. White, R. J., Luque, R., Budarin, V. L., Clark, J. H. and MacQuarrie, D. J. (2009): Supported metal nanoparticles on porous materials. *Methods and applications. Chem. Soc. Rev.*, 38, 481–494.
4. Llewellyn, P. L., Garcia-Rates, M., Gaberová, L., Miller, S. R., Devic, T., Lavalley, J. C., Bourrelly, S., Bloch, E., Filinchuk, Y., Wright, P. A., Serre, C., Vimont, A. and Maurin, G. (2015): Structural origin of unusual CO<sub>2</sub> adsorption behavior of a small-pore aluminum bisphosphonate MOF. *J. Phys. Chem. C*, 119: 4208–4216.
5. Taddei, M., Costantino, F. and Vivani, R. (2016): Robust Metal-Organic Frameworks Based on Tritopic Phosphonoaromatic Ligands. *Eur. J. Inorg. Chem.*, 2016: 4300–4309.
6. Hermer, N., Wharmby, M. T. and Stock, N. (2017): Re-Determination of the Crystal Structure of MIL-91(Al). *Z. Anorg. Allg. Chem.*, 643: 137–140.
7. Oschatz, M. and Antonietti, M. (2018): A search for selectivity to enable CO<sub>2</sub> capture with porous adsorbents. *Energy Environ. Sci.*, 11:57–70.
8. Li, B. and Chen, B. (2016): Fine-Tuning Porous Metal-Organic Frameworks for Gas Separations at Will. *Chem*, 1: 669–671.
9. Winter, M. and Brodd, R. J. (2004): What Are Batteries, Fuel Cells, and Supercapacitors? *Chem. Rev.*, 104: 4245–4270.
10. Latham, K., White, K. F., Szpakolski, K. B., Rix, C. J. and White, J. M. (2009): Synthesis, crystal structure and luminescent behaviour of coordination complexes of copper with bi- and tridentate amines and phosphonic acids. *Inorg. Chim. Acta*, 362: 1872–1886.
11. Bao, S.-S. and Zheng, L.-M. (2016): Magnetic materials based on 3d metal phosphonates. *Coord. Chem. Rev.*, 319: 63–85.
12. Hamchaoui, F., Alonzo, V., Venegas-Yazigi, D., Rebbah, H. and LeFur, E. (2013): Six novel transition-metal hydrogenphosphonate compounds, with structure related to yavapaiite: Crystal structures and magnetic and thermal properties of  $A^I[M^{III}(HPO_3)_2]$  (A=K, NH<sub>4</sub>, Rb and M=V, Fe). *J. Solid State Chem.*, 198: 295–302.
13. Cai, X.-O., Sun, M., Shao, Y.-J., Liu, F., Liu, Q.-L., Zhu, Y.-Y., Sun, Z.-G., Dong, D.-P. and Li, J. (2018): Two Highly Stable Luminescent Lead Phosphonates Based on Mixed Ligands: Highly Selective and Sensitive Sensing for Thymine Molecule and VO<sub>3</sub><sup>-</sup> Anion. *ACS Omega*, 3: 16443–16452.
14. Zhu, Y.-Y., Wang, M.-L., Ma, M.-X., Sun, Z.-G., Jiao, C.-Q., Ma, C. and Li, H.-Y. (2016): Transition metal phosphonates with supramolecular structures: syntheses, structures, surface photovoltage and luminescence properties. *New J. Chem.*, 40: 578–588.
15. Shi, S.-P., Zhu, Y.-Y., Sun, Z.-G., Zhou, W., Dai, L.-L., Ma, M.-X., Li, W.-Z., Luo, H. and Sun, T. (2014): Synthesis, crystal structures, and surface photovoltage and molecular recognition properties of three novel metal carboxyphosphonates with a 3D pillared-layered structure. *Cryst. Growth Des.*, 14: 1580–1590.

16. Zhang, N., Guo, Y.-H., Yu, Y.-Z., Wang, Z., Niu, Y.-S. and Wu, X.-L. (2020): Solvothermal Synthesis, Crystal Structure and Luminescence Property of a 1D Silver(I) Coordination Polymer. *Chin. J. Struct. Chem.*, 39: 2009–2015.
17. Firmino, A. D. G., Mendes, R. F., Ananias, D., Figueira, F., Tome, J. P. C., Rocha, J. and Paz, F. A. A. (2020): pyrenetetrakisphosphonate-based metal-organic framework: structure and photoluminescence. *Eur. J. Inorg. Chem.*, 2020: 3565–3572.
18. Ayhan, M. M., Bayraktar, C., Yu, K. B., Hanna, G., Yazaydin, A. O., Zorlu, Y. and Yucesan, G. (2020): A Nanotubular Metal–Organic Framework with a Narrow Bandgap from Extended Conjugation. *Chem. Eur. J.*, 26: 14813–14816.
19. Jiao, C. Q., Zhao, Z., Ma, C., Sun, Z. G., Dong, D. P., Zhu, Y. Y. and Li, J. (2016): Chiral and achiral copper (II) carboxyphosphonates supramolecular structures: synthesis, structures, surface photovoltage, and magnetic properties. *Cryst. Growth Des.*, 16: 5624–5635.
20. Chu, W., Sun, Z. G., Jiao, C. Q., Zhu, Y. Y., Sun, S. H., Tian, H. and Zheng, M. J. (2013): Two novel lead (II) carboxyphosphonates with a layered and a 3D framework structure: syntheses, crystal structures, reversible dehydration/hydration, and luminescence properties. *Dalton Trans.*, 42: 8009–8017.
21. Dai, L. L., Zhu, Y. Y., Jiao, C. Q., Sun, Z. G., Shi, S. P., Zhou, W., Li, W. Z., Sun, T., Luo, H. and Ma, M. X. (2014): Syntheses, structures, luminescence and molecular recognition properties of four new cadmium carboxyphosphonates with 2D layered and 3D supramolecular structures. *CrystEngComm.*, 16: 5050–5061.
22. Yang, W. T., Tian, H. R., Li, J. P., Hui, Y. F., He, X., Li, J. Y., Dang, S., Xie, Z. G. and Sun, Z. M. (2016): Photochromic terbium phosphonates with photomodulated luminescence and metal ion sensitive detection. *Chem. - Eur. J.*, 22: 15451–15457.
23. Zhao, Z., Yang, D., Xing, B., Ma, C., Sun, Z. G., Zhu, Y. Y., Li, H. Y. and Li, J. (2016): Cadmium (II) carboxyphosphonates based on mixed ligands: syntheses, crystal structures and recognition properties toward amino acids. *RSC Adv.*, 6: 92175–92185.
24. Peng, B. and Chen, J. (2009): Functional materials with high-efficiency energy storage and conversion for batteries and fuel cells. *Coord. Chem. Rev.*, 253: 2805–2813.
25. Smith, G. and Wermuth, U. D. (2010): Guanidinium phenylarsonate–guanidine–water (1/1/2). *Acta Crystallogr.*, E66: o1893–o1894.
26. Reck, L. and Schmitt, W. (2012): Tetra-butyl-ammonium hydrogen phenyl-arsonate–phenyl-arsonic acid (1/1). *Acta Crystallogr.*, E68: m1212–m1213.
27. Yi, F.-Y., Zhao, N., Wu, W. and Mao, J.-G. (2009): Syntheses and Crystal Structures of Novel Manganese(II) or Cadmium(II) Arsonates with Dinuclear Clusters or 1D Arrays. *Inorg. Chem.*, 48: 628–637.
28. Padalwar, N. B. and Vidyasagar, K. (2016): A Cobalt Arylphosphonate MOF – Superior Stability, Sorption and Curie Magnetism Properties. *J. Solid State Chem.*, 243: 83–94.
29. Padalwar, N. B., Pandu, C. and Vidyasagar K. (2013): Monovalent metal phenylphosphonates and phenylarsonates: Single crystal X-ray structures of  $A(\text{HO}_3\text{PPh})(\text{H}_2\text{O}_3\text{PPh})$  ( $A = \text{K}, \text{Rb}, \text{Cs}, \text{Tl}$ ) and  $\text{Na}(\text{HO}_3\text{AsPh})(\text{H}_2\text{O}_3\text{AsPh})$  and methylamine intercalation of  $A(\text{HO}_3\text{PPh})(\text{H}_2\text{O}_3\text{PPh})$  ( $A = \text{Li}, \text{Na}, \text{K}, \text{Tl}$ ). *J. Solid State Chem.*, 203: 321–325.
30. Reddy, B. K., Rao, K. P. and Vidyasagar K. (2006): Syntheses, structure and intercalation properties of low-dimensional phenylarsonates,  $A(\text{HO}_3\text{AsC}_6\text{H}_5)(\text{H}_2\text{O}_3\text{AsC}_6\text{H}_5)$  ( $A = \text{Tl}, \text{Na}, \text{K}$  and  $\text{Rb}$ ). *J. Chem. Sci.*, 118: 117–126.
31. Clarke, R., Latham, K., Rix, C., Hobday, M. and White, J. (2005): Novel copper materials based on the self-assembly of organophosphonic acids and bidentate amines. *CrystEngComm*, 7: 28–36.
32. Latham, K., Coyle, A. M., Rix, C. J., Fowless, A. and White, J. M. (2007): Effect of ring substituents on crystal packing and H-bonding in a series of halobis(phen)copper(II) arylphosphonic acid complexes. *Polyhedron*, 26: 222–236.
33. Rao, K. P. and Vidyasagar, K. (2005): Syntheses, Structure and Intercalation Properties of Low-Dimensional Phenylphosphonates,  $A(\text{HO}_3\text{PC}_6\text{H}_5)(\text{H}_2\text{O}_3\text{PC}_6\text{H}_5)$  ( $A = \text{Alkali Metal}, \text{NH}_4$  and  $\text{Tl}$ ). *Eur. J. Inorg. Chem.*, 2005: 4936–4943.
34. Diop, M. B., Boye, M. S., Diassé-Sarr, A., Diop, L., Guionneau, P. and Maris, T. (2019): Triorganotinphosphonates polymeric chains – Synthesis, Infrared, Mössbauer and Single Crystal Characterization: the First organotin(IV)  $\text{PH}_2^-$  bridged. *Int. J. Adv. Res.*, 7: 266–278.
35. Diop, T., Diop, L., Kociock-Köhn, G., Molloy, K. C. and Ardisson, J. D. (2013): Synthesis, spectroscopic characterization and crystal and molecular structures of phenylphosphonato  $\text{SnR}_3$  ( $R = \text{Ph}, \text{Me}$ ) derivatives. *Main Group Met. Chem.*, 36: 29–34.

36. Diop, T., Diop, L., Da Silva, J. G. and Fall, D. (2012): Supramolecular architecture in crystalline  $\text{Bu}_2\text{NH}_2(\text{PhPO}_3\text{H})_2\text{SnMe}_3$ . *Main Group Met. Chem.*, 35: 63–65.
37. Diop, T., Diop, L., Molloy, K. C., Kociock-Köhn, G. and Stoeckli-Evans, H. (2011): catena - Poly[[triphenyltin(IV)]- $\mu$ -phenylphosphinato- $\kappa^2\text{O}:\text{O}'$ ]. *Acta Crystallogr.*, E67: m1674–m1675.
38. Sarr, M., Boye, M. S., Diassé-Sarr, A., Grosjean, A. and Guionneau, P. (2012): Diisopropylammonium hydrogen phenylphosphonate. *Acta Crystallogr.*, E68: o3078.
39. Sarr, M., Diop, M. B., Boye, M. S., Diassé-Sarr, A. and Guionneau, P. (2020): Crystal structures of two phosphonate salts: monocyclohexylammonium hydrogen phosphonate and monocyclohexylammonium phenylphosphonate. *J. Eng. Stud. Res.*, 26: 50–56.
40. Sarr, M., Diassé-Sarr, A., Diop, L., Plasseraud, L. and Cattey, H. (2015a): Crystal structure of bis(cyclohexylammonium) diphenyl-dioxalato-stannate(IV). *Acta Crystallogr.*, E71: 151–153.
41. Sarr, M., Diassé-Sarr, A., Diop, L., Plasseraud, L. and Cattey, H. (2015b): Crystal structure of bis(cyclohexylammonium) succinate succinic acid salt adduct. *Acta Crystallogr.*, E71: 899–901.
42. Sarr, M., Diallo, W., Diassé-Sarr, A., Diop, L., Plasseraud, L. and Cattey, H. (2013): Tris(cyclohexylammonium) cis-dichlorido-bis-(oxalato- $\kappa^2\text{O}^1\text{O}^2$ )stannate(IV) chloride monohydrate. *Acta Crystallogr.*, E69: m581–m582.
43. Apex2, Crystallographic Software, Suite, Bruker AXS Inc., Madison, Wisconsin (USA) 2012.
44. SAINT, Area Detector Integration Software, Bruker AXS Inc., Madison, Wisconsin, USA, 2012.
45. SADABS (Version 2012/1), Bruker AXS Inc., Madison, Wisconsin, USA, 2012.
46. Sheldrick, G. M. (2015): SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallogr.*, A71: 3–8.
47. Sheldrick, G. M. (2008): A short history of SHELX. *Acta Crystallogr.*, A64: 112–122.
48. Hoof, R. W. W. Collect, Nonius Kappa CCD Data Collection Software, Nonius BV: Delft (The Netherlands), 2003.
49. Duisenberg, A. J. M. (1992): Indexing in single-crystal diffractometry with an obstinate list of reflections. *J. Appl. Crystallogr.*, 25: 92–96.
50. Duisenberg, A. J. M., Kroon-Batenburg, L. M. J. and Schreurs, A. M. M. (2003): An intensity evaluation method: EVAL-14. *J. Appl. Crystallogr.*, 36: 220–229.
51. Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. and Puschmann, H. (2009): OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.*, 42: 339–341.
52. Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. and Wood, P. A. (2008): Mercury CSD 2.0 – new features for the visualization and investigation of crystal structures. *J. Appl. Crystallogr.*, 41: 466–470.
53. Wilke, M., Buzanich, A. G., Reinholz, U., Rademann, K. and Emmerling, F. (2016a): The structure and in situ synthesis investigation of isomorphous mononuclear molecular metal phenylphosphonates. *Dalton Trans.*, 45: 9460–9467.
54. Wilke, M., Bätzdorf, L., Fischer, F., Rademann, K. and Emmerling, F. (2016b): Cadmium phenylphosphonates: Preparation, characterisation and in situ investigation. *RSC Advance*, 6, 36011–36019.
55. Adelani, P. O., Sigmon, G. E. and Burns, P. C. (2013): Hybrid uranyl arsonate coordination nanocages. *Inorg. Chem.*, 52: 6245–6247.
56. Beckmann, J., Dakternieks, D. and Duthie, A. (2003): Dimethylammonium phenylphosphonate 2(phenylphosphonic acid). *Appl. Organomet. Chem.*, 17: 817–818.
57. Lemmerer, A. (2011): Two-Dimensional Hydrogen-Bonding Patterns in a Series of Salts of Terephthalic Acid and the Cyclic Amines  $\text{C}_n\text{H}_{2n-1}\text{NH}_2$ ,  $n = 3, 4, 5, 6, 7, 8$ , and 12. *Cryst. Growth Des.*, 11: 583–593.
58. Grimball, B., Veillo, L., Calhoun, T., Fronczek, F. R., Arceneaux, E. and Laine, R. A. (2017): Cyclohexylamine inexplicably induces antennae loss in Formosan subterranean termites (*Coptotermes formosanus* Shiraki): cyclohexylamine hydrogen phosphate salts are novel termiticides. *Pest Manag. Sci.*, 73: 2039–2047.
59. Tang, Y.-Z., Yu, Y.-M., Xiong, J.-B., Tan, Y.-H. and Wen, H.-R. (2015): Unusual High-Temperature Reversible Phase-Transition Behavior, Structures, and Dielectric–Ferroelectric Properties of Two New Crown Ether Clathrates. *J. Am. Chem. Soc.*, 137: 13345–13351.